

Research Article

The Interplay between Linear Representations of the Braid Group

Mohammad N. Abdulrahim and Nibal H. Kassem

Received 19 May 2007; Accepted 10 July 2007

Recommended by Howard E. Bell

We consider Wada's representation as a twisted version of the standard action of the braid group, B_n , on the free group with n generators. Constructing a free group, G_{nm} , of rank nm , we compose Cohen's map $B_n \rightarrow B_{nm}$ and the embedding $B_{nm} \rightarrow \text{Aut}(G_{nm})$ via Wada's map. We prove that the composition factors of the obtained representation are one copy of Burau representation and $m - 1$ copies of the standard representation after changing the parameter t to t^k in the definitions of the Burau and standard representations. This is a generalization of our previous result concerning the standard Artin representation of the braid group.

Copyright © 2007 M. N. Abdulrahim and N. H. Kassem. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

There are many kinds of representations of B_n , the braid group on n strings. The earliest was the Artin representation, which is an embedding $B_n \rightarrow \text{Aut}(F_n)$, the automorphism group of a free group on n generators [1, page 25]. A certain type of representation, introduced by F. R. Cohen and studied by him and others, is the map $B_n \rightarrow B_{nm}$ which is defined on geometric braids by replacing each string with m strings [2, page 208].

In Section 2 of this paper, we present an infinite series of representations generalizing the standard Artin representation, which were discovered by M. Wada [3]. More precisely, for an arbitrary nonzero integer k , the automorphism corresponding to the braid generator σ_i takes x_i to $x_i^k x_{i+1} x_i^{-k}$, x_{i+1} to x_i , and fixes all other free generators. Utilizing Fox derivatives, we have a twisted version of the Burau representation. Shpilrain has shown that these representations are indeed faithful [3, page 773]. In [4], it was shown that Wada's representations are unitary.

In Section 3, we compose Cohen’s map with Wada’s representation and we get a linear representation of degree nm which has a subrepresentation isomorphic to the Burau representation, and the quotient is isomorphic to the direct sum of $m - 1$ copies of the standard representation, which was studied by Sysoeva [5]. This is done after we change the indeterminate t to t^k in the definitions of the Burau and standard representations. As a corollary, by letting $k = 1$, we get our previous result concerning the standard Artin representation of the braid group. For more details, see [6].

2. Notation and preliminaries

The braid group on n strings, B_n , is an abstract group which has a presentation with generators

$$\sigma_1, \dots, \sigma_{n-1} \tag{2.1}$$

and defining relations

$$\begin{aligned} \sigma_i \sigma_{i+1} \sigma_i &= \sigma_{i+1} \sigma_i \sigma_{i+1} \quad \text{for } i = 1, 2, \dots, n - 2, \\ \sigma_i \sigma_j &= \sigma_j \sigma_i \quad \text{if } |i - j| \geq 2. \end{aligned} \tag{2.2}$$

The generators $\sigma_1, \dots, \sigma_{n-1}$ are called the standard generators of B_n . Let t be an indeterminate and let $\mathbb{C}[t^{\pm 1}]$ represent the Laurent polynomial ring over complex numbers.

Definition 2.1. The Burau representation $\beta_n(t) : B_n \rightarrow GL_n(\mathbb{C}[t^{\pm 1}])$ is defined by

$$\beta_n(t)(\sigma_i) = \left(\begin{array}{c|cc|c} I_{i-1} & 0 & 0 & \\ \hline 0 & 1-t & t & 0 \\ & 1 & 0 & \\ \hline 0 & 0 & 0 & I_{n-i-1} \end{array} \right) \quad \text{for } i = 1, \dots, n - 1. \tag{2.3}$$

The standard representation $\gamma_n(t) : B_n \rightarrow GL_n(\mathbb{C}[t^{\pm 1}])$ is defined by

$$\gamma_n(t)(\sigma_i) = \left(\begin{array}{c|cc|c} I_{i-1} & 0 & 0 & \\ \hline 0 & 0 & t & 0 \\ & 1 & 0 & \\ \hline 0 & 0 & 0 & I_{n-i-1} \end{array} \right) \quad \text{for } i = 1, \dots, n - 1. \tag{2.4}$$

For more details about the standard representation, see [5].

There is a well-known standard representation (due to Artin) of group B_n in group $\text{Aut}(F_n)$ of automorphisms of the free group F_n generated by x_1, \dots, x_n . The automorphism $\bar{\sigma}_i$ corresponding to the braid generator σ_i takes $x_i \rightarrow x_i x_{i+1} x_i^{-1}$; $x_{i+1} \rightarrow x_i$, and fixes all other free generators.

A twisted version of the standard action of the braid group on the free group is Wada’s representation; thus we have the following definition.

Definition 2.2. Wada’s representations are generalizations of the standard Artin representation, discovered by M. Wada, and assert that the automorphism corresponding to σ_i

takes

$$\begin{aligned}
 x_i &\longrightarrow x_i^k x_{i+1} x_i^{-k}, \\
 x_{i+1} &\longrightarrow x_i, \\
 x_j &\longrightarrow x_j \quad \text{for } j \neq i, i + 1.
 \end{aligned}
 \tag{2.5}$$

Definition 2.3 [7, page 104]. Let G be an arbitrary group and let $\mathbb{Z}G$ be the group ring of G with respect to the ring of integers \mathbb{Z} . A mapping $D : \mathbb{Z}G \rightarrow \mathbb{Z}G$ is said to be a *derivative* if and only if

- (1) $D(f + h) = Df + Dh$ and
- (2) $D(fh) = (Df)(\epsilon h) + f(Dh)$ (product rule) for all f and h in $\mathbb{Z}G$.

Here, ϵ is the augmentation homomorphism: $\mathbb{Z}G \rightarrow \mathbb{Z}$ defined by $\epsilon(\sum_{g \in G} n_g g) = \sum_{g \in G} n_g$.

Let F_n be a free group of rank n , with free basis x_1, \dots, x_n . We define for $j = 1, 2, \dots, n$ the *free derivatives* on the group $\mathbb{Z}F_n$ by

$$\begin{aligned}
 \frac{\partial}{\partial x_j} (x_{\mu_1}^{\epsilon_1} \cdots x_{\mu_r}^{\epsilon_r}) &= \sum_{i=1}^r \epsilon_i \delta_{\mu_i, j} x_{\mu_1}^{\epsilon_1} \cdots x_{\mu_i}^{(1/2)(\epsilon_i - 1)} \cdots x_{\mu_r}^{\epsilon_r}, \\
 \frac{\partial}{\partial x_j} \left(\sum a_g g \right) &= \sum a_g \frac{\partial g}{\partial x_j}, \quad g \in F_n, a_g \in \mathbb{Z},
 \end{aligned}
 \tag{2.6}$$

where $\epsilon_i = \pm 1$ and $\delta_{i,j}$ is the Kronecker symbol.

The following properties hold true.

- (i) $\partial x_i / \partial x_j = \delta_{i,j}$.
- (ii) $\partial x_i^{-1} / \partial x_j = -\delta_{i,j} x_i^{-1}$.
- (iii) $(\partial / \partial x_j)(uv) = (\partial u / \partial x_j)\epsilon(v) + u(\partial v / \partial x_j)$ $u, v \in \mathbb{Z}F_n$.

Note that if $v \in F_n$, then $\epsilon(v) = 1$. For simplicity, we denote $\partial / \partial x_j$ by d_j .

Using the Magnus representation, the automorphism σ_i under Wada's representation is mapped onto the $n \times n$ matrix $[\phi((\partial / \partial x_r)\sigma_i(x_j))]$ which differs from the identity only by a 2×2 block with the top left corner in the (i, i) th place. More precisely,

$$\sigma_i(t) = \left(\begin{array}{c|cc|c} I_{i-1} & 0 & 0 & 0 \\ \hline 0 & 1 - t^k & t^k & 0 \\ & 1 & 0 & \\ \hline 0 & 0 & & I_{n-i-1} \end{array} \right) \quad \text{for } i = 1, 2, \dots, n - 1. \tag{2.7}$$

Given a positive integer k , we introduce indeterminates y_1, \dots, y_n defined as $y_1 = x_1^k$, $y_2 = x_2^k, \dots, y_n = x_n^k$ and let G_n be the free group of rank n with free basis y_1, \dots, y_n .

If ϕ is an arbitrary homomorphism acting on F_n defined as $\phi(x_i) = t$, then $\phi(y_i) = t^k$ for $i = 1, \dots, n$. Let G_n^ϕ denote the image of G_n under ϕ .

Under Wada's representation, the action of the generators of B_n on the free group F_n induces an action on the free subgroup G_n . That is, we have a faithful representation of B_n as a subgroup of $\text{Aut}(G_n)$.

LEMMA 2.4. *Under Wada's representation, the action of σ_i on the basis of G_n , namely, $\{y_1, \dots, y_n\}$, is given by*

$$\begin{aligned} y_i &\longrightarrow y_i y_{i+1} y_i^{-1}, \\ y_{i+1} &\longrightarrow y_i, \\ y_r &\longrightarrow y_r, \quad r \neq i, i+1. \end{aligned} \tag{2.8}$$

Proof. $\sigma_i(y_i) = \sigma_i(x_i^k) = (\sigma_i(x_i))^k = x_i^k x_{i+1} x_i^{-k} x_i^k x_{i+1} x_i^{-k} \cdots x_i^k x_{i+1} x_i^{-k} = x_i^k x_{i+1}^k x_i^{-k} = y_i y_{i+1} y_i^{-1}$.

The action of σ_i on the other generators follows easily. □

Using Lemma 2.4 and the Magnus representation of B_n as a subgroup of $\text{Aut}(G_n)$, the automorphism σ_i is mapped onto the $n \times n$ matrix $[\phi((\partial/\partial y_r)\sigma_i(y_s))]$. Direct computations show that it is the same matrix as in (2.7). Therefore, we get the following corollary.

COROLLARY 2.5. *Under Wada's representation, the $n \times n$ matrices obtained by letting B_n act on F_n or on G_n are exactly the same.*

Proof. This follows easily from Lemma 2.4 and the fact that we have defined $\phi(y_i) = t^k$. □

3. Automorphisms of G_{nm}

Definition 3.1 [2, page 208]. The *Cohen representation* is the map $B_n \rightarrow B_{nm}$ defined as follows:

$$\sigma_i \longrightarrow 1 \times \sigma_i = (\sigma_{mi} \sigma_{mi+1} \cdots \sigma_{mi+m-1}) (\sigma_{mi-1} \sigma_{mi} \cdots \sigma_{mi+m-2}) \cdots (\sigma_{mi-m+1} \sigma_{mi-m+2} \cdots \sigma_{mi}). \tag{3.1}$$

Here, $1 \times \sigma_i$ is the braid obtained by replacing each string of the geometric braid, σ_i , with m parallel strings. Cohen called $1 \times \sigma_i$ a tensor product.

Putting $k = 1$ in the definition of Wada's map, we get the result in [6], which asserts that by composing Cohen's map with Artin's representation of the braid group, we get a linear representation: $B_n \rightarrow B_{nm} \rightarrow GL_{nm}(\mathbb{Z}[t^{\pm 1}])$ which has a subrepresentation isomorphic to the Burau representation, and the quotient is isomorphic to the direct sum of $m - 1$ copies of the standard representation, which was studied by Sysoeva [5].

In this paper, we generalize the result by taking any positive integer k and consider Wada's representation, which is a twisted version of the standard action of the braid group on the free group.

Given the free generators x_1, \dots, x_{nm} , we let $y_i = x_i^k$ for $i = 1, \dots, nm$. We take G_{nm} to be the free group generated by y_1, \dots, y_{nm} .

Let τ_i be the image of the braid generator σ_i of B_n under the Cohen map. Using Lemma 2.4, there is an induced action of τ_i on the free subgroup G_{nm} . As in Section 2, we show that the $(nm) \times (nm)$ matrix obtained by letting τ_i as act on F_{nm} with generators x_1, \dots, x_{nm} is exactly the same as that obtained by having τ_i act on G_{nm} with generators x_1^k, \dots, x_{nm}^k instead. Therefore, we get the following theorem.

THEOREM 3.2. *The action of the image of the generator of B_n under Cohen's map, namely, τ_i , on F_{nm} gives an $(nm) \times (nm)$ matrix which is the same as the one obtained under the action of τ_i on the free subgroup G_{nm} .*

Proof. Let

$$\tau_i = (\sigma_{mi}\sigma_{mi+1} \cdots \sigma_{mi+m-1})(\sigma_{mi-1}\sigma_{mi} \cdots \sigma_{mi+m-2}) \cdots (\sigma_{mi-m+1}\sigma_{mi-m+2} \cdots \sigma_{mi}). \quad (3.2)$$

Let us see the action of τ_i on F_{nm} with generators x_1, \dots, x_{nm} .

It is clear that we need to see the action of τ_i especially on the $2m$ elements, namely,

$$x_{mi-m+1}, x_{mi-m+2}, \dots, x_{mi}, x_{mi+1}, x_{mi+2}, \dots, x_{mi+m}. \quad (3.3)$$

As for the other elements, the action of τ_i is trivial. Direct computations show that

$$\tau_i(x_{mi-m+s}) = (x_{mi-m+1}^k \cdots x_{mi}^k)x_{mi+s}(x_{mi-m+1}^k \cdots x_{mi}^k)^{-1} \quad \text{for } s = 1, \dots, m. \quad (3.4)$$

Also, we have that

$$\tau_i(x_{mi+s}) = x_{mi+s-m} \quad \text{for } s = 1, \dots, m. \quad (3.5)$$

The action of τ_i on the free subgroup G_{nm} with generators y_1, \dots, y_{nm} , where $y_j = x_j^k$ for $j = 1, \dots, nm$, is given by

$$\tau_i(y_{mi-m+s}) = (y_{mi-m+1} \cdots y_{mi})y_{mi+s}(y_{mi-m+1} \cdots y_{mi})^{-1} \quad \text{for } s = 1, \dots, m. \quad (3.6)$$

Also, we have that

$$\tau_i(y_{mi+s}) = y_{mi+s-m} \quad \text{for } s = 1, \dots, m. \quad (3.7)$$

Next, we apply Magnus representation to get the matrices corresponding to τ_i , namely, $[\phi((\partial/\partial x_r)\tau_i(x_s))]$ and $[\phi((\partial/\partial y_r)\tau_i(y_s))]$. Using Fox derivatives and having defined $\phi(x_j) = t$ and $\phi(y_j) = t^k$ for $j = 1, \dots, nm$, we get that the matrices are the same. To see this, we make some computations.

For fixed values of i and m , we denote $\phi((\partial/\partial y_r)\tau_i(y_{mi-m+s}))$ or $\phi((\partial/\partial x_r)\tau_i(x_{mi-m+s}))$ by $d_r(\tau_i(y_{mi-m+s}))$ or $d_r(\tau_i(x_{mi-m+s}))$. Direct computations show that these derivatives are

equal. More precisely, we have that

$$\begin{aligned}
 d_{mi-m+1}(\tau_i(y_{mi-m+s})) &= 1 - t^k, & d_{mi-m+2}(\tau_i(y_{mi-m+s})) &= t^k - t^{2k}, \\
 d_{mi-m+3}(\tau_i(y_{mi-m+s})) &= t^{2k} - t^{3k}, \dots, & d_{mi}(\tau_i(y_{mi-m+s})) &= t^{(m-1)k} - t^{mk}.
 \end{aligned}
 \tag{3.8}$$

For $2 \leq s \leq m$, we have

$$d_{mi+1}(\tau_i(y_{mi-m+s})) = \dots = d_{mi+s-1}(\tau_i(y_{mi-m+s})) = 0. \tag{3.9}$$

Also, we have that for $1 \leq s \leq m$

$$d_{mi+s}(\tau_i(y_{mi-m+s})) = t^{mk}. \tag{3.10}$$

If $s \leq m - 1$, then

$$d_{mi+s+1}(\tau_i(y_{mi-m+s})) = \dots = d_{mi+m}(\tau_i(y_{mi-m+s})) = 0. \tag{3.11}$$

As for the elements y_{mi+s} , we have that

$$d_p(\tau_i(y_{mi+s})) = \delta_{p,mi+s-m} \tag{3.12}$$

($\delta_{i,j}$ is the Kronecker symbol). □

Notice that for $m = 1$, we get Corollary 2.5.

Throughout our work, we will then treat the generators of B_n as automorphisms of the free group G_{nm} with generators y_1, \dots, y_{nm} , where $y_i = x_i^k$ rather than automorphisms of F_{nm} .

Next, we proceed as in [6] by choosing elements $z_{i,j}$ of G_{nm} , each of which is a word in these y_i 's. More precisely, for $1 \leq i \leq m$ and $1 \leq j \leq n$ we define $z_{i,j}$ as follows:

$$z_{i,j} = y_{1+mj-m} y_{2+mj-m} \dots y_{mj-i+1}. \tag{3.13}$$

It is then clear that for fixed choices of a positive integer, m , and an integer $i : 1 \leq i \leq m$, the length of $z_{i,j}$ is $m - i + 1$. In other words, the generators $\{z_{i,j}\}$ are defined as follows:

$$\begin{aligned}
 z_{1,1} &= y_1 \dots y_m, & z_{2,1} &= y_1 \dots y_{m-1}, & \dots, & z_{m,1} &= y_1, \\
 z_{1,2} &= y_{1+m} \dots y_{2m}, & z_{2,2} &= y_{1+m} \dots y_{2m-1}, & \dots, & z_{m,2} &= y_{1+m}, \\
 &\vdots & &\vdots & & &\vdots \\
 z_{1,n} &= y_{1+(n-1)m} \dots y_{nm}, & z_{2,n} &= y_{1+(n-1)m} \dots y_{nm-1}, & \dots, & z_{m,n} &= y_{1+(n-1)m}.
 \end{aligned}
 \tag{3.14}$$

LEMMA 3.3. $\{z_{i,j}\}$ is a basis of G_{nm} .

Let $\bar{\tau}_r$ be the automorphism on G_{nm} that corresponds to τ_r which is the image of the braid generator σ_r of B_n under the Cohen map. When there is no danger of confusion, we will still denote the automorphism $\bar{\tau}_r$ by τ_r .

Using Lemma 2.4 in Section 2 of our work and [6, Theorem 3.1, page 172], we easily get the following theorem.

THEOREM 3.4. *For $1 \leq r \leq n - 1$ and $1 \leq i \leq m$, the action of τ_r on the basis $\{z_{i,j}\}$ of G_{nm} is given by*

- (1) $z_{i,r} \rightarrow z_{1,r} z_{i,r+1} z_{1,r}^{-1}$,
- (2) $z_{i,r+1} \rightarrow z_{i,r}$,
- (3) $z_{i,j} \rightarrow z_{i,j}$, $1 \leq j \leq n$ ($j \neq r, r + 1$).

Let $\phi(z_{i,j}) = t^k$ for $1 \leq i \leq m$ and $1 \leq j \leq n$. Let $D_{i,j} = \phi(\partial/\partial z_{i,j})$. Now to find the linear representation

$$B_n \longrightarrow B_{nm} \longrightarrow GL(nm, \mathbb{Z})[t^{\pm 1}], \tag{3.15}$$

we determine the Jacobian matrix of the image of the braid generator σ_r under Cohen map, namely the automorphism τ_r on the group G_{nm} . But first, we give an order to the generators of G_{nm} as follows:

$$z_{1,1}, z_{1,2}, \dots, z_{1,n}, z_{2,1}, z_{2,2}, \dots, z_{2,n}, \dots, z_{m,1}, z_{m,2}, \dots, z_{m,n}. \tag{3.16}$$

Then we define the Jacobian matrix as follows:

$$J(\tau_r) = \begin{pmatrix} D_{1,1}(\tau_r(z_{1,1})) & \cdots & D_{m,n}(\tau_r(z_{1,1})) \\ \vdots & & \vdots \\ D_{1,1}(\tau_r(z_{m,n})) & \cdots & D_{m,n}(\tau_r(z_{m,n})) \end{pmatrix}. \tag{3.17}$$

We now prove our main theorem.

THEOREM 3.5. *The linear representation obtained by composing the Cohen representation with Wada’s representation has a subrepresentation isomorphic to the Burau representation of B_n , and the quotient is isomorphic to the direct sum of $m - 1$ copies of the standard representation of B_n after changing the parameter t to t^k in the definitions of the Burau and standard representations. More precisely,*

$$\sigma_r \longrightarrow \begin{pmatrix} \beta_n(t^k)(\sigma_r) & 0 & \cdots & 0 \\ & \gamma_n(t^k)(\sigma_r) & & \vdots \\ & & \ddots & 0 \\ & & & \gamma_n(t^k)(\sigma_r) \end{pmatrix}. \tag{3.18}$$

Proof. Using Definition 2.3 for free derivatives and Theorem 3.4, we get for $1 \leq i \leq m$

$$D_{1,r}(\tau_r(z_{i,r})) = 1 - t^k, \quad D_{i,r+1}(\tau_r(z_{i,r})) = t^k. \tag{3.19}$$

- [6] M. N. Abdulrahim, "On the composition of the Burau representation and the natural map $B_n \rightarrow B_{nk}$," *Journal of Algebra and Its Applications*, vol. 2, no. 2, pp. 169–175, 2003.
- [7] J. S. Birman, *Braids, Links, and Mapping Class Groups*, Annals of Mathematics Studies, no. 82, Princeton University Press, Princeton, NJ, USA, 1974.

Mohammad N. Abdulrahim: Department of Mathematics, Beirut Arab University, P.O. Box 11-5020, Beirut 11072809, Lebanon
Email address: mna@bau.edu.lb

Nibal H. Kassem: Department of Mathematics, Beirut Arab University, P.O. Box 11-5020, Beirut 11072809, Lebanon
Email address: nibal_rose@hotmail.com



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

