

## GENERALIZATIONS OF INEQUALITIES OF LITTLEWOOD AND PALEY

LOU ZENGJIAN

Mathematics Department  
Qufu Normal University  
Qufu Shandong 273165  
P. R. China

(Received July 30, 1991 and in revised form December 13, 1991)

**ABSTRACT.** For a function  $f$ , holomorphic in the open unit ball  $B_n$  in  $C^n$ , with  $f(0) = 0$ , we prove(I) If  $0 < s < 2$  and  $s < p < \infty$  Then

$$\|f\|_p^p < C \int_0^1 \int_{\partial B_n} |f(\rho \xi)|^{p-s} |Rf(\rho \xi)|^s (\log 1/\rho)^{s-1} \rho^{-1} d\sigma(\xi) d\rho$$

(II) If  $2 < s < p < \infty$  Then

$$\int_0^1 \int_{\partial B_n} |f(\rho \xi)|^{p-s} |Rf(\rho \xi)|^s (\log 1/\rho)^{s-1} \rho^{-1} d\sigma(\xi) d\rho < C \|f\|_p^p$$

where  $Rf$  is the radial derivative of  $f$ , generalizing the known cases  $p = s$  ([1]) and  $p = s$ ,  $n = 1$  ([2]).**KEY WORDS AND PHRASES.** Radial derivative, slice function.1991 **AMS SUBJECT CLASSIFICATION CODES.** 32A10.

## 1. INTRODUCTION

Let  $C^n$  denote the  $n$ -dimensional vector space over  $C$ , let  $B_n$  denote the open unit ball in  $C^n$  with boundary  $\partial B^n$  and let  $\sigma$  denote the rotation-invariant positive measure on  $\partial B_n$  for which  $\sigma(\partial B_n) = 1$ .Throughout this paper, we assume that  $f$  is holomorphic in  $B_n$  with  $f(0) = 0$ , and  $Rf(z) = \sum_{\alpha \geq 0} \alpha |a_\alpha| z^\alpha$  is the radial derivative of  $f(z) = \sum_{\alpha \geq 0} a_\alpha z^\alpha$ .For  $0 < p < \infty$  and  $0 < s < \infty$ , we set

$$M_p^s(r, f) = \int_{\partial B_n} |f(r\xi)|^{p-s} d\sigma(\xi)$$

$$\|f\|_p = \sup_{0 < r < 1} M_p(r, f) \text{ and}$$

$$G_{p,s} \|f\| = \int_0^1 \int_{\partial B_n} |f(\rho \xi)|^{p-s} |Rf(\rho \xi)|^s (\log 1/\rho)^{s-1} \rho^{-1} d\sigma(\xi) d\rho$$

In [1, Theorem 4 and Theorem 7] J. H. Shi generalizes the inequalities of Littlewood and Paley of one complex variable ([2]) to the unit ball  $B_n$ . That is

THEOREM A (1) Let  $0 < p < 2$ . Then

$$\|f\|_p^p < C G_{p,p}[f] \quad (1)$$

(2) Let  $2 < p < \infty$ . Then

$$G_{p,p}[f] < C \|f\|_p^p \quad (2)$$

In this notes, we generalize these results, namely, we prove the following

THEOREM (I) Let  $0 < s < 2$  and  $s < p < \infty$ . Then

$$\|f\|_p^p < C G_{p,s}[f] \quad (3)$$

(II) Let  $2 < s < p < \infty$ . Then

$$G_{p,s}[f] < C \|f\|_p^p \quad (4)$$

Throughout this paper  $C$  denotes a positive constant depending only on  $p$  and  $s$ . The magnitude of  $C$  may vary from occurrence to occurrence even in the proof of the same theorem.

## 2. PROOF OF THE THEOREM.

For the proof of the Theorem we need the following

LEMMA. For  $0 < p < \infty$ . Then

$$\|f\|_p^p = p^s G_{p,s}[f] \quad (5)$$

PROOF. For  $\zeta \in \partial B_n$  the slice functions are defined by  $f_\zeta(\lambda) = f(\lambda \zeta)$ ,  $\lambda \in B_1$ . Then  $Rf(\lambda \zeta) = \lambda f'_\zeta(\lambda)$ .

By the Hardy\_Stein identity for one complex variable ([3]) we have

$$\begin{aligned} M_p^s(r, f_\zeta) &= (p^s/2\pi) \int_0^r \int_0^{2\pi} |f_\zeta(\rho e^{i\theta})|^{p-s} |f'_\zeta(\rho e^{i\theta})|^s \log(r/\rho) \rho d\rho d\theta \\ &= (p^s/2\pi) \int_0^r \int_0^{2\pi} |f(\rho \zeta e^{i\theta})|^{p-s} |Rf(\rho \zeta e^{i\theta})|^s \rho^{-1} \log(r/\rho) d\theta d\rho \end{aligned}$$

Integrating with respect to  $d\sigma(\zeta)$ , using the Fubini theorem and the formular

$$\int_{\partial B_n} g(\zeta) d\sigma(\zeta) = (1/2\pi) \int_{\partial B_n} d\sigma(\zeta) \int_0^{2\pi} g(e^{i\theta} \zeta) d\theta, \quad g \in L^1(\sigma).$$

(see [4, P.15]), we have

$$M_p^s(r, f) = p^s \int_0^r \int_{\partial B_n} |f(\rho \zeta)|^{p-s} |Rf(\rho \zeta)|^s \rho^{-1} \log(r/\rho) d\sigma(\zeta) d\rho \quad (6)$$

By letting  $r \rightarrow 1$  in (6), we obtain (5).

We also need the following fact whose easy proof (by Holder's inequality) we omit.

For a fixed  $p$ ,  $\log G_{p,s}[f]$  is a convex function of  $s$  ( $0 < s < \infty$ ). That is, if  $0 < s_1 < s < s_2 < \infty$  then

$$G_{p, s}[f] < G_{p, s_1}[f]^t G_{p, s_2}[f]^{1-t} \quad (7)$$

Where  $t = (s_2 - s) / (s_2 - s_1)$ .

We now turn to the proof of the Theorem

(I) Case 1.  $s < p < 2$ . Set  $t = (2 - p) / (2 - s)$

$$\begin{aligned} \|f\|_p^p &< C G_{p, p}[f] && \text{( by (1) )} \\ &< C G_{p, s}[f]^t G_{p, 2}[f]^{1-t} && \text{( by (7) )} \\ &< C G_{p, s}[f]^t \|f\|_p^{p(1-t)} && \text{( by (5) )} \end{aligned}$$

so that

$$\|f\|_p^p < C G_{p, s}[f]$$

Case 2.  $s < 2 < p$ . Set  $t = (p - 2) / (p - s)$

$$\begin{aligned} \|f\|_p^p &= C G_{p, 2}[f] && \text{( by (5) )} \\ &< C G_{p, s}[f]^t G_{p, p}[f]^{1-t} && \text{( by (7) )} \\ &< C G_{p, s}[f]^t \|f\|_p^{p(1-t)} && \text{( by (2) )} \end{aligned}$$

so that

$$\|f\|_p^p < C G_{p, s}[f]$$

This gives (3).

(II) Set  $t = (p - s) / (p - 2)$




$$\begin{aligned} G_{p, s}[f] &< G_{p, 2}[f]^t G_{p, p}[f]^{1-t} && \text{( by (7) )} \\ &< C \|f\|_p^{pt} G_{p, p}[f]^{1-t} && \text{( by (6) )} \\ &< C \|f\|_p^{pt} \|f\|_p^{p(1-t)} && \text{( by (2) )} \\ &= C \|f\|_p^p \end{aligned}$$

This gives (4).

**ACKNOWLEDGEMENT.** The author is grateful to Professor Li Qingyao for her encouragement. He also express his thanks to the referee and the Editor Dr. Lokenath Debnath for their suggestions and cooperation.

#### REFERENCES

1. SHI, J. H. Hadamard product of holomorphic functions on unit ball, Chinese Annals of Math. 9 (4) (1988), 49 \_ 59.
2. LITTLEWOOD, J. E. and PALEY, R. E. A. C. Theorems on fourier series and power series, Proc. London Math. Soc. 42 (1936), 52 \_ 89.
3. STEIN, P. On a theorem of Riesz, J. London Math. Soc. 8 (1933), 242 \_ 247.
4. RIDIN, W. Function theory in the unit ball of  $C^n$ , Springer \_ Verlag, New York, 1980.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

