

Research Article

The Local and Global Existence of Solutions for a Generalized Camassa-Holm Equation

Nan Li, Shaoyong Lai, Shuang Li, and Meng Wu

Department of Applied Mathematics, Southwestern University of Finance and Economics,
Chengdu 610074, China

Correspondence should be addressed to Shaoyong Lai, laishaoy@swufe.edu.cn

Received 13 October 2011; Accepted 13 January 2012

Academic Editor: Sergey Piskarev

Copyright © 2012 Nan Li et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

A nonlinear generalization of the Camassa-Holm equation is investigated. By making use of the pseudoparabolic regularization technique, its local well posedness in Sobolev space $H^s(\mathbb{R})$ with $s > 3/2$ is established via a limiting procedure. Provided that the initial value u_0 satisfies the sign condition and $u_0 \in H^s(\mathbb{R})$ ($s > 3/2$), it is shown that there exists a unique global solution for the equation in space $C([0, \infty); H^s(\mathbb{R})) \cap C^1([0, \infty); H^{s-1}(\mathbb{R}))$.

1. Introduction

Camassa and Holm [1] employed the Hamiltonian method to derive a completely integrable shallow water wave model

$$u_t - u_{txx} + 2ku_x + 3uu_x = 2u_x u_{xx} + uu_{xxx}, \quad (1.1)$$

which was alternatively established as a water wave equation in [2–4]. Equation (1.1) also models wave current interaction [5], while Dai [6] derived it as a model in elasticity (see [7]). In addition, it was pointed out in Lakshmanan [8] that the Camassa-Holm equation (1.1) could be relevant to the modeling of tsunami waves (see Constantin and Johnson [9]).

After the birth of the Camassa-Holm equation (1.1), many works have been carried out to probe its dynamic properties. For $k = 0$, (1.1) has travelling wave solutions of the form $ce^{-|x-ct|}$, called peakons, which describes an essential feature of the travelling waves of largest amplitude (see [10–14]). For $k > 0$, its solitary waves are stable solitons [15]. It is shown in [16–18] that the inverse spectral or scattering approach is a powerful tool to handle the Camassa-Holm equation and analyze its dynamics. It is worthwhile to mention

that (1.1) gives rise to geodesic flow of a certain invariant metric on the Bott-Virasoro group [19–21], and this geometric illustration leads to a proof that the least action principle holds. Xin and Zhang [22] proved the global existence of the weak solution in the energy space $H^1(R)$ without any sign conditions on the initial value, and the uniqueness of this weak solution is obtained under some assumptions on the solution [23]. Coclite et al. [24] extended the analysis presented in [22, 23] and obtained many useful dynamic properties to other partial differential equations (see [25–28] for an alternative approach). Li and Olver [29] established the local well posedness in the Sobolev space $H^s(R)$ with $s > 3/2$ for (1.1) and gave conditions on the initial data that lead to finite time blowup of certain solutions. It is shown in Constantin and Escher [30] that the blowup occurs in the form of breaking waves, namely, the solution remains bounded but its slope becomes unbounded in finite time. For other methods to handle the problems relating to various dynamic properties of the Camassa-Holm equation and other shallow water equations, the reader is referred to [31–39] and the references therein.

Motivated by the work in Hakkaev and Kirchev [33] to investigate the generalization forms of the Camassa-Holm equation with high-order nonlinear terms, we study the following generalized Camassa-Holm equation:

$$u_t - u_{txx} + ku^m u_x + (m+3)u^{m+1} u_x = (m+2)u^m u_x u_{xx} + u^{m+1} u_{xxx}, \quad (1.2)$$

where $m \geq 0$ is a natural number and $k \geq 0$. Obviously, (1.2) reduces to (1.1) if we set $m = 0$. As the Camassa-Holm equation (1.1) has been discussed by many mathematicians, we let the natural number $m \geq 1$ in this paper.

The objective of this paper is to study (1.2). Its local well posedness of solutions in the Sobolev space $H^s(R)$ with $s > 3/2$ is developed by using the pseudoparabolic regularization method. Provided that $(1 - \partial_x^2)u_0 + k/2(m+1) \geq 0$ and $u_0 \in H^s$ ($s > 3/2$), the existence and uniqueness of the global solutions are established in space $C([0, \infty); H^s(R)) \cap C^1([0, \infty); H^{s-1}(R))$. It should be mentioned that the existence and uniqueness of global strong solutions for the nonlinear generalized Camassa-Holm models like (1.2) have never been investigated in the literatures.

2. Main Results

The space of all infinitely differentiable functions $\phi(t, x)$ with compact support in $[0, +\infty) \times R$ is denoted by C_0^∞ . $L^p = L^p(R)$ ($1 \leq p < +\infty$) is the space of all measurable functions h such that $\|h\|_{L^p}^p = \int_R |h(t, x)|^p dx < \infty$. We define $L^\infty = L^\infty(R)$ with the standard norm $\|h\|_{L^\infty} = \inf_{m(e)=0} \sup_{x \in R \setminus e} |h(t, x)|$. For any real number s , $H^s = H^s(R)$ denotes the Sobolev space with the norm defined by

$$\|h\|_{H^s} = \left(\int_R (1 + |\xi|^2)^s |\widehat{h}(t, \xi)|^2 d\xi \right)^{1/2} < \infty, \quad (2.1)$$

where $\widehat{h}(t, \xi) = \int_R e^{-ix\xi} h(t, x) dx$.

For $T > 0$ and nonnegative number s , $C([0, T]; H^s(R))$ denotes the Frechet space of all continuous H^s -valued functions on $[0, T]$. We set $\Lambda = (1 - \partial_x^2)^{1/2}$. For simplicity, throughout this paper, we let c denote any positive constant which is independent of parameter ε .

We consider the Cauchy problem of (1.2), which has the equivalent form

$$\begin{aligned} u_t - u_{txx} &= -\frac{k}{m+1} \left(u^{m+1} \right)_x - \frac{m+3}{m+2} \left(u^{m+2} \right)_x + \frac{1}{m+2} \partial_x^3 \left(u^{m+2} \right) \\ &\quad - (m+1) \partial_x \left(u^m u_x^2 \right) + u^m u_x u_{xx}, \quad k \geq 0, \quad m \geq 1, \\ u(0, x) &= u_0(x). \end{aligned} \tag{2.2}$$

Now, we give our main results for problem (2.2).

Theorem 2.1. *Suppose that the initial function $u_0(x)$ belongs to the Sobolev space $H^s(\mathbb{R})$ with $s > 3/2$. Then there is a $T > 0$, which depends on $\|u_0\|_{H^s}$, such that there exists a unique solution $u(t, x)$ of the problem (2.2) and*

$$u(t, x) \in C([0, T]; H^s(\mathbb{R})) \cap C^1([0, T]; H^{s-1}(\mathbb{R})). \tag{2.3}$$

Theorem 2.2. *Let $u_0(x) \in H^s$, $s > 3/2$ and $(1 - \partial_x^2)u_0 + k/2(m+1) \geq 0$ for all $x \in \mathbb{R}$. Then problem (2.2) has a unique solution satisfying that*

$$u(t, x) \in C([0, \infty); H^s(\mathbb{R})) \cap C^1([0, \infty); H^{s-1}(\mathbb{R})). \tag{2.4}$$

3. Local Well-Posedness

In order to prove Theorem 2.1, we consider the associated regularized problem

$$\begin{aligned} u_t - u_{txx} + \varepsilon u_{txxxx} &= -\frac{k}{m+1} \left(u^{m+1} \right)_x - \frac{m+3}{m+2} \left(u^{m+2} \right)_x + \frac{1}{m+2} \partial_x^3 \left(u^{m+2} \right) \\ &\quad - (m+1) \partial_x \left(u^m u_x^2 \right) + u^m u_x u_{xx}, \\ u(0, x) &= u_0(x), \end{aligned} \tag{3.1}$$

where the parameter ε satisfies $0 < \varepsilon < 1/4$.

Lemma 3.1. *Let r and q be real numbers such that $-r < q \leq r$. Then*

$$\begin{aligned} \|uv\|_{H^q} &\leq c \|u\|_{H^r} \|v\|_{H^q}, \quad \text{if } r > \frac{1}{2}, \\ \|uv\|_{H^{r+q-1/2}} &\leq c \|u\|_{H^r} \|v\|_{H^q}, \quad \text{if } r < \frac{1}{2}. \end{aligned} \tag{3.2}$$

This lemma can be found in [34, 40].

Lemma 3.2. *Let $u_0(x) \in H^s(\mathbb{R})$ with $s > 3/2$. Then the Cauchy problem (3.1) has a unique solution $u(t, x) \in C([0, T]; H^s(\mathbb{R}))$ where $T > 0$ depends on $\|u_0\|_{H^s(\mathbb{R})}$. If $s \geq 2$, the solution $u \in C([0, +\infty); H^s)$ exists for all time.*

Proof. Assuming that $D = (1 - \partial_x^2 + \varepsilon \partial_x^4)^{-1}$, we know that $D : H^s \rightarrow H^{s+4}$ is a bounded linear operator. Applying the operator D on both sides of the first equation of system (3.1) and then integrating the resultant equation with respect to t over the interval $(0, t)$ lead to

$$u(t, x) = u_0(x) + \int_0^t D \left[-\frac{k}{m+1} (u^{m+1})_x - \frac{m+3}{m+2} (u^{m+2})_x + \frac{1}{m+2} \partial_x^3 (u^{m+2}) - (m+1) \partial_x (u^m u_x^2) + u^m u_x u_{xx} \right] dt. \quad (3.3)$$

Suppose that both u and v are in the closed ball $B_{M_0}(0)$ of radius M_0 about the zero function in $C([0, T]; H^s(R))$ and A is the operator in the right-hand side of (3.3). For any fixed $t \in [0, T]$, we get the following:

$$\begin{aligned} & \left\| \int_0^t D \left[-\frac{k}{m+1} (u^{m+1})_x - \frac{m+3}{m+2} (u^{m+2})_x + \frac{1}{m+2} \partial_x^3 (u^{m+2}) - (m+1) \partial_x (u^m u_x^2) + u^m u_x u_{xx} \right] dt \right. \\ & \quad \left. - \int_0^t D \left[-\frac{k}{m+1} (v^{m+1})_x - \frac{m+3}{m+2} (v^{m+2})_x + \frac{1}{m+2} \partial_x^3 (v^{m+2}) - (m+1) \partial_x (v^m v_x^2) + v^m v_x v_{xx} \right] dt \right\|_{H^s} \\ & \leq TC_1 \left(\sup_{0 \leq t \leq T} \|u^{m+1} - v^{m+1}\|_{H^s} + \sup_{0 \leq t \leq T} \|u^{m+2} - v^{m+2}\|_{H^s} \right. \\ & \quad \left. + \sup_{0 \leq t \leq T} \|D \partial_x [u^m u_x^2 - v^m v_x^2]\|_{H^s} + \sup_{0 \leq t \leq T} \|D [u^m u_x u_{xx} - v^m v_x v_{xx}]\|_{H^s} \right), \end{aligned} \quad (3.4)$$

where C_1 may depend on ε . The algebraic property of $H^{s_0}(R)$ with $s_0 > 1/2$ derives

$$\begin{aligned} \|u^{m+2} - v^{m+2}\|_{H^s} &= \|(u-v)(u^{m+1} + u^m v + \dots + u v^m + v^{m+1})\|_{H^s} \\ &\leq \|u-v\|_{H^s} \sum_{j=0}^{m+1} \|u\|_{H^s}^{m+1-j} \|v\|_{H^s}^j \\ &\leq M_0^{m+1} \|u-v\|_{H^s}, \end{aligned} \quad (3.5)$$

$$\begin{aligned} \|u^{m+1} - v^{m+1}\|_{H^s} &\leq M_0^m \|u-v\|_{H^s}, \\ \|D \partial_x (u^m u_x^2 - v^m v_x^2)\|_{H^s} &\leq \|D \partial_x [u^m (u_x^2 - v_x^2)]\|_{H^s} + \|D \partial_x [v_x^2 (u^m - v^m)]\|_{H^s} \\ &\leq C \left(\|u^m (u_x^2 - v_x^2)\|_{H^{s-1}} + \|v_x^2 (u^m - v^m)\|_{H^{s-1}} \right) \\ &\leq CM_0^{m+1} \|u-v\|_{H^s}. \end{aligned} \quad (3.6)$$

Using the first inequality of Lemma 3.1, we have

$$\begin{aligned}
 \|D[u^m u_x u_{xx} - v^m v_x v_{xx}]\|_{H^s} &= \left\| \frac{1}{2} D[u^m (u_x^2)_x - v^m (v_x^2)_x] \right\|_{H^s} \\
 &\leq \frac{1}{2} \left(\|D[u^m (u_x^2 - v_x^2)_x]\|_{H^s} + \|D[(v_x^2)_x (u^m - v^m)]\|_{H^s} \right) \\
 &\leq C \left(\|u^m (u_x^2 - v_x^2)_x\|_{H^{s-2}} + \|(v_x^2)_x (u^m - v^m)\|_{H^{s-2}} \right) \\
 &\leq C \left(\|u^m\|_{H^s} \|u_x^2 - v_x^2\|_{H^{s-1}} + \|v_x^2\|_{H^{s-1}} \|u^m - v^m\|_{H^s} \right) \\
 &\leq CM_0^{m+1} \|u - v\|_{H^s},
 \end{aligned} \tag{3.7}$$

where C may depend on ε . From (3.5)–(3.7), we obtain that

$$\|Au - Av\|_{H^s} \leq \theta \|u - v\|_{H^s}, \tag{3.8}$$

where $\theta = TC_2(M_0^m + M_0^{m+1})$ and C_2 is independent of $0 < t < T$. Choosing T sufficiently small such that $\theta < 1$, we know that A is a contraction. Applying the above inequality yields that

$$\|Au\|_{H^s} \leq \|u_0\|_{H^s} + \theta \|u\|_{H^s}. \tag{3.9}$$

Choosing T sufficiently small such that $\theta M_0 + \|u_0\|_{H^s} < M_0$, we deduce that A maps $B_{M_0}(0)$ to itself. It follows from the contraction-mapping principle that the mapping A has a unique fixed-point u in $B_{M_0}(0)$.

For $s \geq 2$, using the first equation of system (3.1) derives

$$\frac{d}{dt} \int_{\mathbb{R}} \left(u^2 + u_x^2 + \varepsilon u_{xx}^2 \right) dx = 0, \tag{3.10}$$

from which we have the conservation law

$$\int_{\mathbb{R}} \left(u^2 + u_x^2 + \varepsilon u_{xx}^2 \right) dx = \int_{\mathbb{R}} \left(u_0^2 + u_{0x}^2 + \varepsilon u_{0xx}^2 \right) dx. \tag{3.11}$$

The proof of the global existence result is a routine argument by using (3.11) (see Xin and Zhang [22]). □

Lemma 3.3 (Kato and Ponce [41]). *If $r \geq 0$, then $H^r \cap L^\infty$ is an algebra. Moreover*

$$\|uv\|_r \leq c(\|u\|_{L^\infty} \|v\|_r + \|u\|_r \|v\|_{L^\infty}), \tag{3.12}$$

where c is a constant depending only on r .

Lemma 3.4 (Kato and Ponce [41]). *Let $r > 0$. If $u \in H^r \cap W^{1,\infty}$ and $v \in H^{r-1} \cap L^\infty$, then*

$$\|[\Lambda^r, u]v\|_{L^2} \leq c \left(\|\partial_x u\|_{L^\infty} \|\Lambda^{r-1} v\|_{L^2} + \|\Lambda^r u\|_{L^2} \|v\|_{L^\infty} \right). \quad (3.13)$$

Lemma 3.5. *Let $s \geq 2$, and the function $u(t, x)$ is a solution of problem (3.1) and the initial data $u_0(x) \in H^s$. Then the following inequality holds:*

$$\begin{aligned} \|u\|_{H^1}^2 &\leq \int_{\mathbb{R}} \left(u^2 + u_x^2 + \varepsilon u_{xx}^2 \right) dx \\ &= \int_{\mathbb{R}} \left(u_0^2 + u_{0x}^2 + \varepsilon u_{0xx}^2 \right) dx. \end{aligned} \quad (3.14)$$

For $q \in (0, s-1]$, there is a constant c independent of ε such that

$$\begin{aligned} \int_{\mathbb{R}} \left(\Lambda^{q+1} u \right)^2 dx &\leq \int_{\mathbb{R}} \left[\left(\Lambda^{q+1} u_0 \right)^2 + \varepsilon \left(\Lambda^q u_{0xx} \right)^2 \right] dx \\ &+ c \int_0^t \|u\|_{H^{q+1}}^2 \left(\left(\|u\|_{L^\infty}^{m-1} + \|u\|_{L^\infty}^m \right) \|u_x\|_{L^\infty} + \|u\|_{L^\infty}^{m-1} \|u_x\|_{L^\infty}^2 \right) d\tau. \end{aligned} \quad (3.15)$$

For $q \in [0, s-1]$, there is a constant c independent of ε such that

$$(1 - 2\varepsilon) \|u_t\|_{H^q} \leq c \|u\|_{H^{q+1}} \left(\left(\|u\|_{L^\infty}^{m-1} + \|u\|_{L^\infty}^m \right) \|u\|_{H^1} + \|u\|_{L^\infty}^m \|u_x\|_{L^\infty} + \|u\|_{L^\infty}^{m-1} \|u_x\|_{L^\infty}^2 \right). \quad (3.16)$$

Proof. The inequality $\|u\|_{H^1}^2 \leq \int_{\mathbb{R}} (u^2 + u_x^2) dx$ and (3.11) derives (3.14).

Using $\partial_x^2 = -\Lambda^2 + 1$ and the Parseval equality gives rise to

$$\begin{aligned} \int_{\mathbb{R}} \Lambda^q u \Lambda^q \partial_x^3 (u^{m+2}) dx &= -(m+2) \int_{\mathbb{R}} \left(\Lambda^{q+1} u \right) \Lambda^{q+1} \left(u^{m+1} u_x \right) dx \\ &+ (m+2) \int_{\mathbb{R}} \left(\Lambda^q u \right) \Lambda^q \left(u^{m+1} u_x \right) dx. \end{aligned} \quad (3.17)$$

For $q \in (0, s-1]$, applying $(\Lambda^q u) \Lambda^q$ to both sides of the first equation of system (3.1) and integrating with respect to x by parts, we have the identity

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}} \left((\Lambda^q u)^2 + (\Lambda^q u_x)^2 + \varepsilon (\Lambda^q u_{xx})^2 \right) dx \\ &= -\frac{k}{m+1} \int_{\mathbb{R}} (\Lambda^q u) \Lambda^q \left(u^{m+1} \right)_x dx - (m+2) \int_{\mathbb{R}} (\Lambda^q u) \Lambda^q \left(u^{m+1} u_x \right) dx \\ &\quad - \int_{\mathbb{R}} \left(\Lambda^{q+1} u \right) \Lambda^{q+1} \left(u^{m+1} u_x \right) dx + (m+1) \int_{\mathbb{R}} (\Lambda^q u_x) \Lambda^q \left(u^m u_x^2 \right) dx \\ &\quad + \int_{\mathbb{R}} \Lambda^q u \Lambda^q \left(u^m u_x u_{xx} \right) dx. \end{aligned} \quad (3.18)$$

We will estimate the terms on the right-hand side of (3.18) separately. For the second term, by using the Cauchy-Schwartz inequality and Lemmas 3.3 and 3.4, we have

$$\begin{aligned}
\left| \int_{\mathbb{R}} (\Lambda^q u) \Lambda^q (u^{m+1} u_x) dx \right| &= \left| \int_{\mathbb{R}} (\Lambda^q u) \left[\Lambda^q (u^{m+1} u_x) - u^{m+1} \Lambda^q u_x \right] dx + \int_{\mathbb{R}} (\Lambda^q u) u^{m+1} \Lambda^q u_x dx \right| \\
&\leq c \|u\|_{H^q} \left((m+1) \|u\|_{L^\infty}^m \|u_x\|_{L^\infty} \|u\|_{H^q} + \|u_x\|_{L^\infty} \|u\|_{L^\infty}^m \|u\|_{H^q} \right) \\
&\quad + \frac{m+1}{2} \|u\|_{L^\infty}^m \|u_x\|_{L^\infty} \|\Lambda^q u\|_{L^2}^2 \\
&\leq c \|u\|_{H^q}^2 \|u\|_{L^\infty}^m \|u_x\|_{L^\infty}.
\end{aligned} \tag{3.19}$$

Similarly, for the first term in (3.18), we have

$$\left| \int_{\mathbb{R}} (\Lambda^q u) \Lambda^q (u^m u_x) dx \right| \leq c \|u\|_{H^q}^2 \|u\|_{L^\infty}^{m-1} \|u_x\|_{L^\infty}. \tag{3.20}$$

Using the above estimate to the third term yields that

$$\left| \int_{\mathbb{R}} (\Lambda^{q+1} u) \Lambda^{q+1} (u^{m+1} u_x) dx \right| \leq c \|u\|_{H^{q+1}}^2 \|u\|_{L^\infty}^m \|u_x\|_{L^\infty}. \tag{3.21}$$

For the fourth term, using the Cauchy-Schwartz inequality and Lemma 3.3, we obtain that

$$\begin{aligned}
\left| \int_{\mathbb{R}} (\Lambda^q u_x) \Lambda^q (u^m u_x^2) dx \right| &\leq \|\Lambda^q u_x\|_{L^2} \left\| \Lambda^q (u^m u_x^2) \right\|_{L^2} \\
&\leq c \|u\|_{H^{q+1}} \left(\|u^m u_x\|_{L^\infty} \|u_x\|_{H^q} + \|u_x\|_{L^\infty} \|u^m u_x\|_{H^q} \right) \\
&\leq c \|u\|_{H^{q+1}}^2 \|u_x\|_{L^\infty} \|u\|_{L^\infty}^m,
\end{aligned} \tag{3.22}$$

in which we have used $\|u^m u_x\|_{H^q} \leq c \|(u^{m+1})_x\|_{H^q} \leq c \|u\|_{L^\infty}^m \|u\|_{H^{q+1}}$.

For the last term in (3.18), using $u^m (u_x^2)_x = (u^m u_x^2)_x - (u^m)_x u_x^2$ results in

$$\begin{aligned}
\left| \int_{\mathbb{R}} (\Lambda^q u) \Lambda^q (u^m u_x u_{xx}) dx \right| &\leq \left| \int_{\mathbb{R}} \Lambda^q u_x \Lambda^q (u^m u_x^2) dx \right| + \int_{\mathbb{R}} \Lambda^q u \Lambda^q \left[(u^m)_x u_x^2 \right] dx \\
&= K_1 + K_2.
\end{aligned} \tag{3.23}$$

For K_1 , it follows from (3.22) that

$$K_1 \leq c \|u\|_{H^{q+1}}^2 \|u_x\|_{L^\infty} \|u\|_{L^\infty}^m. \tag{3.24}$$

For K_2 , applying Lemma 3.3 derives

$$\begin{aligned} K_2 &\leq c \|u\|_{H^q} \left\| (u^m)_x u_x^2 \right\|_{H^q} \\ &\leq c \|u\|_{H^q} \left(\|(u^m)_x\|_{L^\infty} \|u_x^2\|_{H^q} + \|(u^m)_x\|_{H^q} \|u_x^2\|_{L^\infty} \right) \\ &\leq c \|u\|_{H^{q+1}}^2 \left(\|u\|_{L^\infty}^{m-1} \|u_x\|_{L^\infty}^2 \right). \end{aligned} \quad (3.25)$$

It follows from (3.19)–(3.25) that there exists a constant c depending only on m such that

$$\frac{1}{2} \frac{d}{dt} \int_R \left[(\Lambda^q u)^2 + (\Lambda^q u_x)^2 + \varepsilon (\Lambda^q u_{xx})^2 \right] dx \leq c \|u\|_{H^{q+1}}^2 \left(\|u_x\|_{L^\infty} \|u\|_{L^\infty}^m + \|u\|_{L^\infty}^{m-1} \|u_x\|_{L^\infty}^2 \right). \quad (3.26)$$

Integrating both sides of the above inequality with respect to t results in (3.15).

To estimate the norm of u_t , we apply the operator $(1 - \partial_x^2)^{-1}$ to both sides of the first equation of system (3.1) to obtain the equation

$$\begin{aligned} (1 - \varepsilon) u_t - \varepsilon u_{txx} &= (1 - \partial_x^2)^{-1} \left[-\varepsilon u_t - k u^m u_x - \frac{m+3}{m+2} (u^{m+2})_x \right. \\ &\quad \left. + \frac{1}{m+2} \partial_x^3 (u^{m+2}) - (m+1) \partial_x (u^m u_x^2) + u^m u_x u_{xx} \right]. \end{aligned} \quad (3.27)$$

Applying $(\Lambda^q u_t) \Lambda^q$ to both sides of (3.27) for $q \in [0, s-1]$ gives rise to

$$\begin{aligned} (1 - \varepsilon) \int_R (\Lambda^q u_t)^2 dx + \varepsilon \int_R (\Lambda^q u_{xt})^2 dx \\ = \int_R (\Lambda^q u_t) \Lambda^{q-2} \left[-\varepsilon u_t + \partial_x \left(-\frac{k}{m+1} u^{m+1} - \frac{m+3}{m+2} u^{m+2} + \frac{1}{m+2} \partial_x^2 u^{m+2} - (m+1) u^m u_x^2 \right) \right. \\ \left. + u^m u_x u_{xx} \right] d\tau. \end{aligned} \quad (3.28)$$

For the right hand of (3.28), we have

$$\begin{aligned} \left| \int_R (\Lambda^q u_t) \Lambda^{q-2} (-\varepsilon u_t) dx \right| &\leq \varepsilon \|u_t\|_{H^q}^2, \\ \left| \int_R (\Lambda^q u_t) (1 - \partial_x^2)^{-1} \Lambda^q \partial_x \left(-\frac{k}{m+1} u^{m+1} - \frac{m+3}{m+2} u^{m+2} - (m+1) u^m u_x^2 \right) dx \right| \\ &\leq c \|u_t\|_{H^q} \left(\int_R (1 + \xi^2)^{q-1} \left[\int_R \left[-\frac{k}{m+1} \widehat{u^m}(\xi - \eta) \widehat{u}(\eta) - \frac{m+3}{m+2} \widehat{u^{m+1}}(\xi - \eta) \widehat{u}(\eta) \right. \right. \right. \\ &\quad \left. \left. \left. - (m+1) \widehat{u^m u_x}(\xi - \eta) \widehat{u_x}(\eta) \right] d\eta \right]^2 \right)^{1/2} \\ &\leq c \|u_t\|_{H^q} \|u\|_{H^1} \|u\|_{H^{q+1}} \left(\|u\|_{L^\infty}^{m-1} + \|u\|_{L^\infty}^m \right). \end{aligned} \quad (3.29)$$

Since

$$\begin{aligned} \int (\Lambda^q u_t) (1 - \partial_x^2)^{-1} \Lambda^q \partial_x^2 (u^{m+1} u_x) dx &= - \int (\Lambda^q u_t) \Lambda^q (u^{m+1} u_x) dx \\ &+ \int (\Lambda^q u_t) (1 - \partial_x^2)^{-1} \Lambda^q (u^{m+1} u_x) dx, \end{aligned} \quad (3.30)$$

using Lemma 3.3, $\|u^{m+1} u_x\|_{H^q} \leq c \|(u^{m+2})_x\|_{H^q} \leq c(m+2) \|u\|_{L^\infty}^{m+1} \|u\|_{H^{q+1}}$ and $\|u\|_{L^\infty} \leq \|u\|_{H^1}$, we have

$$\begin{aligned} \left| \int (\Lambda^q u_t) \Lambda^q (u^{m+1} u_x) dx \right| &\leq c \|u_t\|_{H^q} \|u^{m+1} u_x\|_{H^q} \\ &\leq c \|u_t\|_{H^q} \|u\|_{L^\infty}^m \|u\|_{H^1} \|u\|_{H^{q+1}}, \\ \left| \int (\Lambda^q u_t) (1 - \partial_x^2)^{-1} \Lambda^q (u^{m+1} u_x) dx \right| &\leq c \|u_t\|_{H^q} \|u\|_{L^\infty}^m \|u\|_{H^1} \|u\|_{H^{q+1}}. \end{aligned} \quad (3.31)$$

Using the Cauchy-Schwartz inequality and Lemmas 3.1 and 3.3 yields that

$$\begin{aligned} \left| \int_R (\Lambda^q u_t) (1 - \partial_x^2)^{-1} \Lambda^q (u^m u_x u_{xx}) dx \right| &\leq c \|u_t\|_{H^q} \|u^m u_x u_{xx}\|_{H^{q-2}} \\ &\leq c \|u_t\|_{H^q} \|u^m (u_x^2)_x\|_{H^{q-2}} \\ &\leq c \|u_t\|_{H^q} \left\| \left[u^m (u_x^2) \right]_x - (u^m)_x u_x^2 \right\|_{H^{q-2}} \\ &\leq c \|u_t\|_{H^q} \left(\|u^m u_x^2\|_{H^{q-1}} + \|(u^m)_x u_x^2\|_{H^{q-2}} \right) \\ &\leq c \|u_t\|_{H^q} \left(\|u^m u_x^2\|_{H^q} + \|(u^m)_x u_x^2\|_{H^q} \right) \\ &\leq c \|u_t\|_{H^q} \|u\|_{H^{q+1}} \left(\|u\|_{L^\infty}^m \|u_x\|_{L^\infty} + \|u\|_{L^\infty}^{m-1} \|u_x\|_{L^\infty}^2 \right), \end{aligned} \quad (3.32)$$

in which we have used (3.25).

Applying (3.29)–(3.32) into (3.28) yields the inequality

$$\begin{aligned} (1 - 2\varepsilon) \|u_t\|_{H^q} &\leq c \|u\|_{H^{q+1}} \left(\left(\|u\|_{L^\infty}^{m-1} + \|u\|_{L^\infty}^m \right) \|u\|_{H^1} \right. \\ &\quad \left. + \|u\|_{L^\infty}^m \|u_x\|_{L^\infty} + \|u\|_{L^\infty}^{m-1} \|u_x\|_{L^\infty}^2 \right) \end{aligned} \quad (3.33)$$

for a constant $c > 0$. This completes the proof of Lemma 3.5. □

Remark 3.6. In fact, letting $\varepsilon = 0$ in problem (3.1), (3.14), (3.15), and (3.16) are still valid.

Setting $\phi_\varepsilon(x) = \varepsilon^{-1/4}\phi(\varepsilon^{-1/4}x)$ with $0 < \varepsilon < 1/4$ and $u_{\varepsilon 0} = \phi_\varepsilon \star u_0$, we know that $u_{\varepsilon 0} \in C^\infty$ for any $u_0 \in H^s$, $s > 0$. From Lemma 3.2, it derives that the Cauchy problem

$$\begin{aligned} u_t - u_{txx} + \varepsilon u_{txxxx} &= -\frac{k}{m+1} \left(u^{m+1} \right)_x - \frac{m+3}{m+2} \left(u^{m+2} \right)_x + \frac{1}{m+2} \partial_x^3 \left(u^{m+2} \right) \\ &\quad - (m+1) \partial_x \left(u^m u_x^2 \right) + u^m u_x u_{xx}, \\ u(0, x) &= u_{\varepsilon 0}(x), \quad x \in R \end{aligned} \quad (3.34)$$

has a unique solution $u_\varepsilon(t, x) \in C^\infty([0, \infty); H^\infty)$.

Furthermore, we have the following.

Lemma 3.7. *For $s > 0$, $u_0 \in H^s$, it holds that*

$$\|u_{\varepsilon 0x}\|_{L^\infty} \leq c \|u_{0x}\|_{L^\infty}, \quad (3.35)$$

$$\|u_{\varepsilon 0}\|_{H^q} \leq c \quad \text{if } q \leq s, \quad (3.36)$$

$$\|u_{\varepsilon 0}\|_{H^q} \leq c \varepsilon^{(s-q)/4} \quad \text{if } q > s, \quad (3.37)$$

$$\|u_{\varepsilon 0} - u_0\|_{H^q} \leq c \varepsilon^{(s-q)/4} \quad \text{if } q \leq s, \quad (3.38)$$

$$\|u_{\varepsilon 0} - u_0\|_{H^s} = o(1), \quad (3.39)$$

where c is a constant independent of ε .

The proof of Lemma 3.7 can be found in [38].

Remark 3.8. For $s \geq 1$, using $\|u_\varepsilon\|_{L^\infty} \leq c \|u_\varepsilon\|_{H^{1/2+}} \leq c \|u_\varepsilon\|_{H^1}$, $\|u_\varepsilon\|_{H^1}^2 \leq c \int_{\mathbb{R}} (u_\varepsilon^2 + u_{\varepsilon x}^2) dx$, (3.14), (3.36), and (3.37), we know that

$$\begin{aligned} \|u_\varepsilon\|_{L^\infty}^2 &\leq c \|u_\varepsilon\|_{H^1} \leq c \int_{\mathbb{R}} \left(u_{\varepsilon 0}^2 + u_{\varepsilon 0x}^2 + \varepsilon u_{\varepsilon 0xx}^2 \right) dx \\ &\leq c \left(\|u_{\varepsilon 0}\|_{H^1}^2 + \varepsilon \|u_{\varepsilon 0}\|_{H^2}^2 \right) \\ &\leq c \left(c + c\varepsilon \times \varepsilon^{(s-2)/2} \right) \\ &\leq c_0, \end{aligned} \quad (3.40)$$

where c_0 is independent of ε .

Lemma 3.9. *If $u_0(x) \in H^s(\mathbb{R})$ with $s \geq 1$ such that $\|u_{0x}\|_{L^\infty} < \infty$. Let $u_{\varepsilon 0}$ be defined as in system (3.34). Then there exist two positive constants T and c , which are independent of ε , such that the solution u_ε of problem (3.34) satisfies $\|u_{\varepsilon x}\|_{L^\infty} \leq c$ for any $t \in [0, T)$.*

Proof. Using notation $u = u_\varepsilon$ and differentiating both sides of the first equation of problem (3.34) or (3.27) with respect to x give rise to

$$\begin{aligned} & (1 - \varepsilon)u_{tx} - \varepsilon u_{txxx} + \frac{1}{m+2} \partial^2 (u^{m+2}) - \left(m + \frac{1}{2}\right) (u^m u_x^2) \\ &= \frac{k}{m+1} u^{m+1} + u^{m+2} - \Lambda^{-2} \left[\varepsilon u_{tx} + \frac{k}{m+1} u^{m+1} + u^{m+2} + \left(m + \frac{1}{2}\right) (u^m u_x^2) \right. \\ & \quad \left. + \frac{1}{2} \partial_x [(u^m)_x u_x^2] \right]. \end{aligned} \tag{3.41}$$

Letting $p > 0$ be an integer and multiplying the above equation by $(u_x)^{2p+1}$ and then integrating the resulting equation with respect to x yield the equality

$$\begin{aligned} & \frac{1 - \varepsilon}{2p+2} \frac{d}{dt} \int_R (u_x)^{2p+2} dx - \varepsilon \int_R (u_x)^{2p+1} u_{txxx} dx + \frac{p-m}{2p+2} \int_R (u_x)^{2p+3} u^m dx \\ &= \int_R (u_x)^{2p+1} \left(\frac{k}{m+1} u^{m+1} + u^{m+2} \right) dx \\ & \quad - \int_R (u_x)^{2p+1} \Lambda^{-2} \left[\varepsilon u_{tx} + \frac{k}{m+1} u^{m+1} + u^{m+2} + \left(m + \frac{1}{2}\right) (u^m u_x^2) + \frac{1}{2} \partial_x [(u^m)_x u_x^2] \right] dx. \end{aligned} \tag{3.42}$$

Applying the Hölder's inequality yields that

$$\begin{aligned} & \frac{1 - \varepsilon}{2p+2} \frac{d}{dt} \int_R (u_x)^{2p+2} dx \leq \left\{ \varepsilon \left(\int_R |u_{txxx}|^{2p+2} dx \right)^{1/(2p+2)} + \left(\int_R |u^{m+1}|^{2p+2} dx \right)^{1/(2p+2)} \right. \\ & \quad \left. + \left(\int_R |u^{m+2}|^{2p+2} dx \right)^{1/(2p+2)} + \left(\int_R |G|^{2p+2} dx \right)^{1/(2p+2)} \right\} \\ & \quad \times \left(\int_R |u_x|^{2p+2} dx \right)^{(2p+1)/(2p+2)} + \left| \frac{p-m}{2p+2} \right| \|u_x\|_{L^\infty} \|u\|_{L^\infty}^m \int_R |u_x|^{2p+2} dx \end{aligned} \tag{3.43}$$

or

$$\begin{aligned} & (1 - \varepsilon) \frac{d}{dt} \left(\int_R (u_x)^{2p+2} dx \right)^{1/(2p+2)} \leq \left\{ \varepsilon \left(\int_R |u_{txxx}|^{2p+2} dx \right)^{1/(2p+2)} + \left(\int_R |u^{m+1}|^{2p+2} dx \right)^{1/(2p+2)} \right. \\ & \quad \left. + \left(\int_R |u^{m+2}|^{2p+2} dx \right)^{1/(2p+2)} + \left(\int_R |G|^{2p+2} dx \right)^{1/(2p+2)} \right\} \\ & \quad + \left| \frac{p-m}{2p+2} \right| \|u_x\|_{L^\infty} \|u\|_{L^\infty}^m \left(\int_R |u_x|^{2p+2} dx \right)^{1/(2p+2)}, \end{aligned} \tag{3.44}$$

where

$$G = \Lambda^{-2} \left[\varepsilon u_{tx} + \frac{k}{m+1} u^{m+1} + u^{m+2} + \left(m + \frac{1}{2} \right) (u^m u_x^2) + \frac{1}{2} \partial_x [(u^m)_x u_x^2] \right]. \quad (3.45)$$

Since $\|f\|_{L^p} \rightarrow \|f\|_{L^\infty}$ as $p \rightarrow \infty$ for any $f \in L^\infty \cap L^2$, integrating both sides of (3.44) with respect to t and taking the limit as $p \rightarrow \infty$ result in the estimate

$$(1 - \varepsilon) \|u_x\|_{L^\infty} \leq (1 - \varepsilon) \|u_{0x}\|_{L^\infty} + \int_0^t \left[\varepsilon \|u_{txxx}\|_{L^\infty} + c \left(\|u\|_{L^\infty}^{m+1} + \|u\|_{L^\infty}^{m+2} + \|G\|_{L^\infty} \right) + \frac{1}{2} \|u\|_{L^\infty}^m \|u_x\|_{L^\infty}^2 \right] d\tau. \quad (3.46)$$

Using the algebraic property of $H^{s_0}(R)$ with $s_0 > 1/2$ and (3.40) yields that

$$\|u\|_{L^\infty}^{m+2} \leq c \|u\|_{H^1}^{m+2} \leq c, \quad (3.47)$$

$$\begin{aligned} \|G\|_{L^\infty} &\leq c \|G\|_{H^{1/2+}} \\ &= c \left\| \Lambda^{-2} \left[\varepsilon u_{tx} + u^{m+1} + u^{m+2} + \left(m + \frac{1}{2} \right) (u^m u_x^2) + \frac{1}{2} \partial_x [(u^m)_x u_x^2] \right] \right\|_{H^{1/2+}} \\ &\leq c \left(\left\| \Lambda^{-2} u_{xt} \right\|_{H^{1/2+}} + \left\| \Lambda^{-2} (u^m u_x^2) \right\|_{H^{1/2+}} + \left\| \Lambda^{-2} \partial_x [(u^m)_x u_x^2] \right\|_{H^{1/2+}} \right) + c \\ &\leq c \left(\|u_t\|_{L^2} + \left\| u^m u_x^2 \right\|_{H^0} + \left\| (u^m)_x u_x^2 \right\|_{H^0} \right) + c \\ &\leq c \left(\|u_t\|_{L^2} + \|u\|_{L^\infty}^m \|u_x\|_{L^\infty} \|u\|_{H^1} + \|u_x\|_{L^\infty}^2 \|u\|_{L^\infty}^{m-1} \|u\|_{H^1} \right) + c \\ &\leq c \left(\|u_t\|_{L^2} + \|u_x\|_{L^\infty} + \|u_x\|_{L^\infty}^2 \right) + c \\ &\leq c \left(1 + \|u_x\|_{L^\infty}^2 \right), \end{aligned} \quad (3.48)$$

where we have used (3.16) and (3.40). Using (3.48), we have

$$\int_0^t \|G\|_{L^\infty} d\tau \leq c \int_0^t \left(1 + \|u_x\|_{L^\infty}^2 \right) d\tau, \quad (3.49)$$

where c is a constant independent of ε . Moreover, for any fixed $r \in (1/2, 1)$, there exists a constant c_r such that $\|u_{txxx}\|_{L^\infty} \leq c_r \|u_{txxx}\|_{H^r} \leq c_r \|u_t\|_{H^{r+3}}$. Using (3.16) and (3.40) yields that

$$\|u_{txxx}\|_{L^\infty} \leq c \|u\|_{H^{r+4}} \left(1 + \|u_x\|_{L^\infty}^2 \right). \quad (3.50)$$

Making use of the Gronwall's inequality to (3.15) with $q = r + 3$, $u = u_\varepsilon$ and (3.40) gives rise to

$$\|u\|_{H^{r+4}}^2 \leq \left(\int_{\mathbb{R}} (\Lambda^{r+4} u_0)^2 + \varepsilon (\Lambda^{r+3} u_{0xx})^2 \right) \exp \left[c \int_0^t \left(1 + \|u_x\|_{L^\infty}^2 \right) d\tau \right]. \quad (3.51)$$

From (3.36), (3.37), (3.50), and (3.51), one has

$$\|u_{txxx}\|_{L^\infty} \leq c\varepsilon^{(s-r-4)/4} \left(1 + \|u_x\|_{L^\infty}^2\right) \exp \left[c \int_0^t \left(1 + \|u_x\|_{L^\infty}^2\right) d\tau \right]. \quad (3.52)$$

For $\varepsilon < 1/4$, it follows from (3.46), (3.49), and (3.52) that

$$\begin{aligned} \|u_x\|_{L^\infty} &\leq \|u_{0x}\|_{L^\infty} \\ &+ c \int_0^t \left[\varepsilon^{(s-r)/4} \left(1 + \|u_x\|_{L^\infty}^2\right) \exp \left(c \int_0^\tau \left(1 + \|u_x\|_{L^\infty}^2\right) d\zeta \right) + 1 + \|u_x\|_{L^\infty}^2 \right] d\tau. \end{aligned} \quad (3.53)$$

It follows from the contraction mapping principle that there is a $T > 0$ such that the equation

$$\begin{aligned} \|W\|_{L^\infty} &= \|u_{0x}\|_{L^\infty} \\ &+ c \int_0^t \left[\left(1 + \|W\|_{L^\infty}^2\right) \exp \left(c \int_0^\tau \left(1 + \|W\|_{L^\infty}^2\right) d\zeta \right) + 1 + \|W\|_{L^\infty}^2 \right] d\tau \end{aligned} \quad (3.54)$$

has a unique solution $W \in C[0, T]$. Using the Theorem presented at page 51 in Li and Olver [29] or Theorem II in section I.1 presented in [42] yields that there are constants $T > 0$ and $c > 0$, which are independent of ε , such that $\|u_x\|_{L^\infty} \leq W(t)$ for arbitrary $t \in [0, T]$, which leads to the conclusion of Lemma 3.9. \square

Lemma 3.10 (Li and Olver [29]). *If u and f are functions in $H^{q+1} \cap \{\|u_x\|_{L^\infty} < \infty\}$, then*

$$\left| \int_{\mathbb{R}} \Lambda^q u \Lambda^q (uf)_x dx \right| \leq \begin{cases} c_q \|f\|_{H^{q+1}} \|u\|_{H^q}^2, & q \in \left(\frac{1}{2}, 1\right], \\ c_q \left(\|f\|_{H^{q+1}} \|u\|_{H^q} \|u\|_{L^\infty} \right. \\ \quad \left. + \|f_x\|_{L^\infty} \|u\|_{H^q}^2 + \|f\|_{H^q} \|u\|_{H^q} \|u_x\|_{L^\infty} \right), & q \in (0, \infty). \end{cases} \quad (3.55)$$

Lemma 3.11. *For $u, v \in H^s(\mathbb{R})$ with $s > 3/2$, $w = u - v$, $q > 1/2$, and a natural number n , it holds that*

$$\left| \int_{\mathbb{R}} \Lambda^s w \Lambda^s \left(u^{n+1} - v^{n+1} \right)_x dx \right| \leq c \left(\|w\|_{H^s} \|w\|_{H^q} \|v\|_{H^{s+1}} + \|w\|_{H^s}^2 \right). \quad (3.56)$$

The proof of this Lemma can be found in [38].

Lemma 3.12. For problem (3.34), $s > 3/2$ and $u_0 \in H^s(\mathbb{R})$, there exist two positive constants c and M , which are independent of ε , such that the following inequalities hold for any sufficiently small ε and $t \in [0, T]$

$$\|u_\varepsilon\|_{H^s} \leq M e^{ct}, \quad (3.57)$$

$$\|u_\varepsilon\|_{H^{s+k_1}} \leq \varepsilon^{-k_1/4} M e^{ct}, \quad k_1 > 0, \quad (3.58)$$

$$\|u_{\varepsilon t}\|_{H^{s+k_1}} \leq \varepsilon^{-(k_1+1)/4} M e^{ct}, \quad k_1 > -1. \quad (3.59)$$

Proof. If $s > 3/2$, $u_0 \in H^s$, we obtain that

$$\begin{aligned} u_0 &\in H^{s_1} \quad \text{with } 1 \leq s_1 \leq \frac{3}{2}, \\ \|u_{0x}\|_{L^\infty} &\leq c \|u_{0x}\|_{H^{1/2+}} \leq c \|u_0\|_{H^s} \leq c. \end{aligned} \quad (3.60)$$

From Lemma 3.9, we know that there exist two constants T and c (both independent of ε) such that

$$\|u_{\varepsilon x}\|_{L^\infty} \leq c \quad \text{for any } t \in [0, T]. \quad (3.61)$$

Applying the inequality (3.15) with $q + 1 = s$ and the bounded property of solution u (see (3.40) and (3.60)), we have

$$\begin{aligned} \int_{\mathbb{R}} (\Lambda^s u_\varepsilon)^2 dx &\leq \int_{\mathbb{R}} \left[(\Lambda^s u_{\varepsilon 0})^2 + \varepsilon (\Lambda^{s-1} u_{\varepsilon 0xx})^2 \right] dx + c \int_0^t \|u_\varepsilon\|_{H^s}^2 d\tau, \\ &= A + c \int_0^t \|u_\varepsilon\|_{H^s}^2 d\tau, \end{aligned} \quad (3.62)$$

where

$$\begin{aligned} A &= \int_{\mathbb{R}} \left[(\Lambda^s u_{\varepsilon 0})^2 + \varepsilon (\Lambda^{s-1} u_{\varepsilon 0xx})^2 \right] dx \leq \|u_{\varepsilon 0}\|_{H^s}^2 + \|u_{\varepsilon 0}\|_{H^{s+1}}^2 \\ &\leq c + c\varepsilon\varepsilon^{-1/2} \leq 2c, \end{aligned} \quad (3.63)$$

in which we have used (3.36) and (3.37).

From (3.61) and (3.62) and using the Gronwall's inequality, we get the following:

$$\|u_\varepsilon\|_{H^s} \leq 2c e^{ct}, \quad (3.64)$$

from which we know that (3.57) holds.

In a similar manner, for $q + 1 = s + k_1$ and $k_1 > 0$, applying (3.40) and (3.60) to (3.15), we have

$$\|u_\varepsilon\|_{H^{s+k_1}}^2 \leq \left(c\varepsilon^{-k_1/2} + c\varepsilon^{-(k_1+1)/2} \varepsilon \right) + c \int_0^t \|u_\varepsilon\|_{H^{s+k_1}}^2 d\tau, \quad (3.65)$$

which results in (3.58) by using Gronwall's inequality.

From (3.16), for $q = s + k_1$, we have

$$(1 - 2\varepsilon)\|u_{\varepsilon t}\|_{H^{s+k_1}} \leq c\|u_{\varepsilon}\|_{H^{s+k_1+1}}, \quad (3.66)$$

which leads to (3.59) by (3.58). \square

Lemma 3.13. *If $1/2 < q < \min\{1, s - 1\}$ and $s > 3/2$, then for any functions w, f defined on R , it holds that*

$$\left| \int_R \Lambda^q w \Lambda^{q-2} (wf)_x dx \right| \leq c\|w\|_{H^q}^2 \|f\|_{H^q}, \quad (3.67)$$

$$\left| \int_R \Lambda^q w \Lambda^{q-2} (w_x f_x)_x dx \right| \leq c\|w\|_{H^q}^2 \|f\|_{H^s}. \quad (3.68)$$

The proof of this lemma can be found in [38].

Our next step is to demonstrate that u_{ε} is a Cauchy sequence. Let u_{ε} and u_{δ} be solutions of problem (3.34), corresponding to the parameters ε and δ , respectively, with $0 < \varepsilon < \delta < 1/4$, and let $w = u_{\varepsilon} - u_{\delta}$. Then w satisfies the problem

$$\begin{aligned} & (1 - \varepsilon)w_t - \varepsilon w_{xxt} + (\delta - \varepsilon)(u_{\delta t} + u_{\delta xxt}) \\ &= (1 - \partial_x^2)^{-1} \left[-\varepsilon w_t + (\delta - \varepsilon)u_{\delta t} - \frac{k}{m+1} \partial_x (u_{\varepsilon}^{m+1} - u_{\delta}^{m+1}) - \partial_x (u_{\varepsilon}^{m+2} - u_{\delta}^{m+2}) \right. \\ & \quad \left. - \partial_x \left[\partial_x (u_{\varepsilon}^{m+1}) \partial_x w + \partial_x (u_{\varepsilon}^{m+1} - u_{\delta}^{m+1}) \partial_x u_{\delta} \right] \right. \\ & \quad \left. + [u_{\varepsilon}^m u_{\varepsilon x} u_{\varepsilon xx} - u_{\delta}^m u_{\delta x} u_{\delta xx}] \right] - \frac{1}{m+2} \partial_x (u_{\varepsilon}^{m+2} - u_{\delta}^{m+2}), \end{aligned} \quad (3.69)$$

$$w(x, 0) = w_0(x) = u_{\varepsilon 0}(x) - u_{\delta 0}(x). \quad (3.70)$$

Lemma 3.14. *For $s > 3/2$, $u_0 \in H^s(R)$, there exists $T > 0$ such that the solution u_{ε} of (3.34) is a Cauchy sequence in $C([0, T]; H^s(R)) \cap C^1([0, T]; H^{s-1}(R))$.*

Proof. For q with $1/2 < q < \min\{1, s - 1\}$, multiplying both sides of (3.69) by $\Lambda^q w \Lambda^q$ and then integrating with respect to x give rise to

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_R \left[(1 - \varepsilon)(\Lambda^q w)^2 + \varepsilon(\Lambda^q w_x)^2 \right] dx \\ &= (\varepsilon - \delta) \int_R (\Lambda^q w) [(\Lambda^q u_{\delta t}) + (\Lambda^q u_{\delta xxt})] dx \\ & \quad - \varepsilon \int_R \Lambda^q w \Lambda^{q-2} w_t dx + (\delta - \varepsilon) \int_R \Lambda^q w \Lambda^{q-2} u_{\delta t} dx \\ & \quad - \frac{1}{m+2} \int_R (\Lambda^q w) \Lambda^q (u_{\varepsilon}^{m+2} - u_{\delta}^{m+2})_x dx \\ & \quad - \frac{k}{m+1} \int_R \Lambda^q w \Lambda^{q-2} (u_{\varepsilon}^{m+1} - u_{\delta}^{m+1})_x dx \end{aligned}$$

$$\begin{aligned}
 & - \int_R \Lambda^q \omega \Lambda^{q-2} \left(u_\varepsilon^{m+2} - u_\delta^{m+2} \right)_x dx \\
 & - \int_R \Lambda^q \omega \Lambda^{q-2} \left[\partial_x \left(u_\varepsilon^{m+1} \right) \partial_x \omega \right]_x dx \\
 & - \int_R \Lambda^q \omega \Lambda^{q-2} \left[\partial_x \left(u_\varepsilon^{m+1} - u_\delta^{m+1} \right) \partial_x u_\delta \right]_x dx \\
 & + \int_R \Lambda^q \omega \Lambda^{q-2} \left[u_\varepsilon^m u_{\varepsilon x} u_{\varepsilon x x} - u_\delta^m u_{\delta x} u_{\delta x x} \right] dx.
 \end{aligned} \tag{3.71}$$

It follows from the Schwarz inequality that

$$\begin{aligned}
 & \frac{d}{dt} \int \left[(1 - \varepsilon) (\Lambda^q \omega)^2 + \varepsilon (\Lambda^q \omega_x)^2 \right] dx \\
 & \leq c \left\{ \|\Lambda^q \omega\|_{L^2} \left[(\delta - \varepsilon) (\|\Lambda^q u_{\delta t}\|_{L^2} + \|\Lambda^q u_{\delta x x t}\|_{L^2}) \right. \right. \\
 & \quad \left. \left. + \varepsilon \|\Lambda^{q-2} \omega_t\|_{L^2} + (\delta - \varepsilon) \|\Lambda^{q-2} u_{\delta t}\|_{L^2} \right] \right. \\
 & \quad + \left| \int_R \Lambda^q \omega \Lambda^q \left(u_\varepsilon^{m+2} - u_\delta^{m+2} \right)_x dx \right| \left| \int \Lambda^q \omega \Lambda^{q-2} \left(u_\varepsilon^{m+1} - u_\delta^{m+1} \right)_x dx \right| \\
 & \quad + \left| \int \Lambda^q \omega \Lambda^{q-2} \left(u_\varepsilon^{m+2} - u_\delta^{m+2} \right)_x dx \right| + \left| \int_R \Lambda^q \omega \Lambda^{q-2} \left[\partial_x \left(u_\varepsilon^{m+1} \right) \partial_x \omega \right]_x dx \right| \\
 & \quad + \left| \int_R \Lambda^q \omega \Lambda^{q-2} \left[\partial_x \left(u_\varepsilon^{m+1} - u_\delta^{m+1} \right) \partial_x u_\delta \right]_x dx \right| \\
 & \quad \left. + \left| \int_R \Lambda^q \omega \Lambda^{q-2} \left[u_\varepsilon^m u_{\varepsilon x} u_{\varepsilon x x} - u_\delta^m u_{\delta x} u_{\delta x x} \right] dx \right| \right\}.
 \end{aligned} \tag{3.72}$$

Using the first inequality in Lemma 3.10, we have

$$\begin{aligned}
 & \left| \int_R \Lambda^q \omega \Lambda^q \left(u_\varepsilon^{m+2} - u_\delta^{m+2} \right)_x dx \right| = \left| \int_R \Lambda^q \omega \Lambda^q \left(\omega g_{m+1} \right)_x dx \right| \\
 & \leq c \|\omega\|_{H^q}^2 \|g_{m+1}\|_{H^{q+1}},
 \end{aligned} \tag{3.73}$$

where $g_{m+1} = \sum_{j=0}^{m+1} u_\varepsilon^{m+1-j} u_\delta^j$. For the last three terms in (3.72), using Lemmas 3.1 and 3.13, $1/2 < q < \min\{1, s - 1\}$, $s > 3/2$, the algebra property of H^{s_0} with $s_0 > 1/2$ and (3.40), we have

$$\left| \int_R \Lambda^q \omega \Lambda^{q-2} \left(\partial_x \left(u_\varepsilon^{m+1} \right) \partial_x \omega \right)_x dx \right| \leq c \|\omega\|_{H^q}^2 \|u_\varepsilon\|_{H^s}^{m+1}, \tag{3.74}$$

$$\begin{aligned}
 & \left| \int_R \Lambda^q \omega \Lambda^{q-2} \left(\partial_x \left(u_\varepsilon^{m+1} - u_\delta^{m+1} \right) \partial_x u_\delta \right)_x dx \right| \\
 & \leq c \|\omega\|_{H^q} \|u_\delta\|_{H^s} \left\| u_\varepsilon^{m+1} - u_\delta^{m+1} \right\|_{H^q} \\
 & \leq c \|\omega\|_{H^q}^2 \|u_\delta\|_{H^s},
 \end{aligned}$$

$$\begin{aligned}
 & \left| \int_{\mathbb{R}} \Lambda^q w \Lambda^{q-2} [u_\varepsilon^m u_{\varepsilon x} u_{\varepsilon x x} - u_\delta^m u_{\delta x} u_{\delta x x}] dx \right| \\
 & \leq c \|w\|_{H^q} \left\| (u_\varepsilon^m - u_\delta^m) (u_{\varepsilon x}^2)_x + u_\delta^m [u_{\varepsilon x}^2 - u_{\delta x}^2]_x \right\|_{H^{q-2}} \\
 & \leq c \|w\|_{H^q} \left(\left\| (u_\varepsilon^m - u_\delta^m) (u_{\varepsilon x}^2)_x \right\|_{H^{q-1}} + \left\| u_\delta^m [u_{\varepsilon x}^2 - u_{\delta x}^2]_x \right\|_{H^{q-2}} \right) \\
 & \leq c \|w\|_{H^q} \left(\|u_\varepsilon^m - u_\delta^m\|_{H^q} \left\| (u_{\varepsilon x}^2)_x \right\|_{H^{q-1}} + \|u_\delta^m\|_{H^s} \left\| [u_{\varepsilon x}^2 - u_{\delta x}^2]_x \right\|_{H^{q-2}} \right) \\
 & \leq c \|w\|_{H^q} \left(\|w\|_{H^q} \|g_{m-1}\|_{H^q} \|u\|_{H^s}^2 + \|u_\delta^m\|_{H^s} \|u_{\varepsilon x} + u_{\delta x}\|_{H^q} \|w\|_{H^q} \right) \\
 & \leq c \|w\|_{H^q}^2 \left(\|g_{m-1}\|_{H^q} \|u\|_{H^s}^2 + \|u_\delta^m\|_{H^s} \|u_{\varepsilon x} + u_{\delta x}\|_{H^q} \right).
 \end{aligned} \tag{3.75}$$

Using (3.67), we derives that the inequality

$$\begin{aligned}
 \left| \int_{\mathbb{R}} \Lambda^q w \Lambda^{q-2} (u_\varepsilon^{m+2} - u_\delta^{m+2})_x dx \right| &= \left| \int_{\mathbb{R}} \Lambda^q w \Lambda^{q-2} (w g_{m+1})_x dx \right| \\
 &\leq c \|g_{m+1}\|_{H^q} \|w\|_{H^q}^2
 \end{aligned} \tag{3.76}$$

holds for some constant c , where $g_{m+1} = \sum_{j=0}^{m+1} u_\varepsilon^{m+1-j} u_\delta^j$. Using the algebra property of H^q with $q > 1/2$, $q + 1 < s$ and Lemma 3.12, we have $\|g_m\|_{H^{q+1}} \leq c$ for $t \in (0, \tilde{T}]$. Then it follows from (3.57)–(3.59) and (3.73)–(3.76) that there is a constant c depending on \tilde{T} such that the estimate

$$\frac{d}{dt} \int_{\mathbb{R}} \left[(1 - \varepsilon) (\Lambda^q w)^2 + \varepsilon (\Lambda^q w_x)^2 \right] dx \leq c \left(\delta^\gamma \|w\|_{H^q} + \|w\|_{H^q}^2 \right) \tag{3.77}$$

holds for any $t \in [0, \tilde{T})$, where $\gamma = 1$ if $s \geq 3 + q$ and $\gamma = (1 + s - q)/4$ if $s < 3 + q$. Integrating (3.77) with respect to t , one obtains the estimate

$$\begin{aligned}
 \frac{1}{2} \|w\|_{H^q}^2 &= \frac{1}{2} \int_{\mathbb{R}} (\Lambda^q w)^2 dx \\
 &\leq \int_{\mathbb{R}} \left[(1 - \varepsilon) (\Lambda^q w)^2 + \varepsilon (\Lambda^q w)^2 \right] dx \\
 &\leq \int_{\mathbb{R}} \left[(\Lambda^q w_0)^2 + \varepsilon (\Lambda^q w_{0x})^2 \right] dx + c \int_0^t \left(\delta^\gamma \|w\|_{H^q} + \|w\|_{H^q}^2 \right) d\tau.
 \end{aligned} \tag{3.78}$$

Applying the Gronswall inequality, (3.37) and (3.39) yields that

$$\|u\|_{H^q} \leq c \delta^{(s-q)/4} e^{ct} + \delta^\gamma (e^{ct} - 1), \tag{3.79}$$

for any $t \in [0, \tilde{T})$.

Multiplying both sides of (3.69) by $\Lambda^s w \Lambda^s$ and integrating the resultant equation with respect to x , one obtains that

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}} \left[(1 - \varepsilon) (\Lambda^s w)^2 + \varepsilon (\Lambda^s w_x)^2 \right] dx \\
&= (\varepsilon - \delta) \int_{\mathbb{R}} (\Lambda^s w) [(\Lambda^s u_{\delta t}) + (\Lambda^s u_{\delta x x t})] dx \\
&\quad - \varepsilon \int_{\mathbb{R}} \Lambda^s w \Lambda^{s-2} w_t dx + (\delta - \varepsilon) \int_{\mathbb{R}} \Lambda^s w \Lambda^{s-2} u_{\delta t} dx \\
&\quad - \frac{k}{m+1} \int_{\mathbb{R}} (\Lambda^s w) \Lambda^s (u_\varepsilon^{m+1} - u_\delta^{m+1})_x dx \\
&\quad - \frac{1}{m+2} \int_{\mathbb{R}} (\Lambda^s w) \Lambda^s (u_\varepsilon^{m+2} - u_\delta^{m+2})_x dx \\
&\quad - \int_{\mathbb{R}} \Lambda^s w \Lambda^{s-2} (u_\varepsilon^{m+2} - u_\delta^{m+2})_x dx \\
&\quad - \int_{\mathbb{R}} \Lambda^s w \Lambda^{s-2} [\partial_x (u_\varepsilon^{m+1}) \partial_x w]_x dx \\
&\quad - \int_{\mathbb{R}} \Lambda^s w \Lambda^{s-2} [\partial_x (u_\varepsilon^{m+1} - u_\delta^{m+1}) \partial_x u_\delta]_x dx \\
&\quad + \int_{\mathbb{R}} \Lambda^s w \Lambda^{s-2} [u_\varepsilon^m u_{\varepsilon x} u_{\varepsilon x x} - u_\delta^m u_{\delta x} u_{\delta x x}] dx.
\end{aligned} \tag{3.80}$$

From Lemma 3.13, we have

$$\left| \int_{\mathbb{R}} \Lambda^s w \Lambda^{s-2} (u_\varepsilon^{m+2} - u_\delta^{m+2})_x dx \right| \leq c_3 \|g_{m+1}\|_{H^s} \|w\|_{H^s}^2. \tag{3.81}$$

From Lemma 3.11, it holds that

$$\left| \int_{\mathbb{R}} \Lambda^s w \Lambda^s (u_\varepsilon^{m+2} - u_\delta^{m+2})_x dx \right| \leq c (\|w\|_{H^s} \|w\|_{H^q} \|u_\delta\|_{H^{s+1}} + \|w\|_{H^s}^2). \tag{3.82}$$

Using the Cauchy-Schwartz inequality and the algebra property of H^{s_0} with $s_0 > 1/2$, for $s > 3/2$, we have

$$\begin{aligned}
\left| \int_{\mathbb{R}} \Lambda^s w \Lambda^{s-2} [\partial_x (u_\varepsilon^{m+1}) \partial_x w]_x dx \right| &= \left| \int_{\mathbb{R}} \Lambda^q w \Lambda^{s-2} [\partial_x (u_\varepsilon^{m+1}) \partial_x w]_x dx \right| \\
&\leq c \|\Lambda^s w\|_{L^2} \left\| \Lambda^{s-2} [\partial_x (u_\varepsilon^{m+1}) \partial_x w]_x \right\|_{L^2} \\
&\leq c \|w\|_{H^q} \left\| \partial_x (u_\varepsilon^{m+1}) \partial_x w \right\|_{H^{s-1}} \\
&\leq c \|u_\varepsilon\|_{H^s}^{m+1} \|w\|_{H^s}^2 \\
\left| \int_{\mathbb{R}} \Lambda^s w \Lambda^{s-2} [\partial_x (u_\varepsilon^{m+1} - u_\delta^{m+1}) \partial_x u_\delta]_x dx \right| &\leq c \|w\|_{H^s} \left\| \Lambda^{s-2} [\partial_x (u_\varepsilon^{m+1} - u_\delta^{m+1}) \partial_x u_\delta]_x \right\|_{L^2} \\
&\leq c \|u_\delta\|_{H^s} \|g_m\|_{H^s} \|w\|_{H^s}^2,
\end{aligned} \tag{3.83}$$

$$\begin{aligned}
 & \left| \int_{\mathbb{R}} \Lambda^s \omega \Lambda^{s-2} [u_\varepsilon^m u_{\varepsilon x} u_{\varepsilon x x} - u_\delta^m u_{\delta x} u_{\delta x x}] dx \right| \\
 & \leq c \|\omega\|_{H^s} \left(\left\| (u_\varepsilon^m - u_\delta^m) (u_{\varepsilon x}^2)_x \right\|_{H^{s-2}} + \left\| u_\delta^m [u_{\varepsilon x}^2 - u_{\delta x}^2]_x \right\|_{H^{s-2}} \right) \\
 & \leq c \|\omega\|_{H^s} \left(\|u_\varepsilon^m - u_\delta^m\|_{H^s} \left\| (u_{\varepsilon x}^2)_x \right\|_{H^{s-2}} + \|u_\delta^m\|_{H^s} \left\| [u_{\varepsilon x}^2 - u_{\delta x}^2]_x \right\|_{H^{s-2}} \right) \\
 & \leq c \|\omega\|_{H^s}^2,
 \end{aligned} \tag{3.84}$$

in which we have used Lemma 3.1 and the bounded property of $\|u_\varepsilon\|_{H^s}$ and $\|u_\delta\|_{H^s}$ (see Lemma 3.12). It follows from (3.80)–(3.84) and (3.57)–(3.59) and (3.79) that there exists a constant c depending on m such that

$$\begin{aligned}
 & \frac{d}{dt} \int_{\mathbb{R}} \left[(1 - \varepsilon) (\Lambda^s \omega)^2 + \varepsilon (\Lambda^s \omega_x)^2 \right] dx \\
 & \leq 2\delta \left(\|u_{\delta t}\|_{H^s} + \|u_{\delta x x t}\|_{H^s} + \left\| \Lambda^{s-2} \omega_t \right\|_{L^2} + \left\| \Lambda^{s-2} u_{\delta t} \right\| \right) \|\omega\|_{H^s} \\
 & \quad + c \left(\|\omega\|_{H^s}^2 + \|\omega\|_{H^q} \|\omega\|_{H^s} \|u_\delta\|_{H^{s+1}} \right) \\
 & \leq c \left(\delta^{\gamma_1} \|\omega\|_{H^s} + \|\omega\|_{H^s}^2 \right),
 \end{aligned} \tag{3.85}$$

where $\gamma_1 = \min(1/4, (s - q - 1)/4) > 0$. Integrating (3.85) with respect to t leads to the estimate

$$\begin{aligned}
 \frac{1}{2} \|\omega\|_{H^s}^2 & \leq \int_{\mathbb{R}} \left[(1 - \varepsilon) (\Lambda^s \omega)^2 + \varepsilon (\Lambda^s \omega_x)^2 \right] dx \\
 & \leq \int_{\mathbb{R}} \left[(\Lambda^s \omega_0)^2 + \varepsilon (\Lambda^s \omega_{0x})^2 \right] dx + c \int_0^t \left(\delta^{\gamma_1} \|\omega\|_{H^s} + \|\omega\|_{H^s}^2 \right) d\tau.
 \end{aligned} \tag{3.86}$$

It follows from the Gronwall inequality and (3.86) that

$$\begin{aligned}
 \|\omega\|_{H^s} & \leq \left(2 \int_{\mathbb{R}} \left[(\Lambda^s \omega_0)^2 + \varepsilon (\Lambda^s \omega_{0x})^2 \right] dx \right)^{1/2} e^{ct} + \delta^{\gamma_1} (e^{ct} - 1) \\
 & \leq c_1 \left(\|\omega_0\|_{H^s} + \delta^{3/4} \right) e^{ct} + \delta^{\gamma_1} (e^{ct} - 1),
 \end{aligned} \tag{3.87}$$

where c_1 is independent of ε and δ .

Then (3.39) and the above inequality show that

$$\|\omega\|_{H^s} \longrightarrow 0 \quad \text{as } \varepsilon \longrightarrow 0, \delta \longrightarrow 0. \tag{3.88}$$

Next, we consider the convergence of the sequence $\{u_{\varepsilon t}\}$. Multiplying both sides of (3.69) by $\Lambda^{s-1}w_t\Lambda^{s-1}$ and integrating the resultant equation with respect to x , we obtain

$$\begin{aligned}
(1-\varepsilon)\|w_t\|_{H^{s-1}}^2 &+ \frac{1}{m+2} \int_R \left(\Lambda^{s-1}w_t\right)\Lambda^{s-1}\left(u_\varepsilon^{m+2} - u_\delta^{m+2}\right)_x dx \\
&+ \int_R \left[-\varepsilon\left(\Lambda^{s-1}w_t\right)\left(\Lambda^{s-1}w_{xxt}\right) + (\delta-\varepsilon)\left(\Lambda^{s-1}w_t\right)\Lambda^{s-1}\left(u_{\delta t} + u_{\delta xxt}\right)\right] dx \\
&= \int_R \left(\Lambda^{s-1}w_t\right)\Lambda^{s-3} \left[-\varepsilon w_t + (\delta-\varepsilon)u_{\delta t} - \frac{k}{m+1}\partial_x\left(u_\varepsilon^{m+1} - u_\delta^{m+1}\right) - \partial_x\left(u_\varepsilon^{m+2} - u_\delta^{m+2}\right)\right. \\
&\quad \left.- \partial_x\left[\partial_x\left(u_\varepsilon^{m+1}\right)\partial_x w + \partial_x\left(u_\varepsilon^{m+1} - u_\delta^{m+1}\right)\partial_x u_\delta\right]\right. \\
&\quad \left.+ \left[u_\varepsilon^m u_{\varepsilon x} u_{\varepsilon xx} - u_\delta^m u_{\delta x} u_{\delta xx}\right]\right] dx.
\end{aligned} \tag{3.89}$$

It follows from (3.57)–(3.60) and the Schwartz inequality that there is a constant c depending on \tilde{T} and m such that

$$(1-\varepsilon)\|w_t\|_{H^{s-1}}^2 \leq c\left(\delta^{1/2} + \|w\|_{H^s} + \|w\|_{s-1}\right)\|w_t\|_{H^{s-1}} + \varepsilon\|w_t\|_{H^{s-1}}^2. \tag{3.90}$$

Hence,

$$\begin{aligned}
\frac{1}{2}\|w_t\|_{H^{s-1}}^2 &\leq (1-2\varepsilon)\|w_t\|_{H^{s-1}}^2 \\
&\leq c\left(\delta^{1/2} + \|w\|_{H^s} + \|w\|_{H^{s-1}}\right)\|w_t\|_{H^{s-1}},
\end{aligned} \tag{3.91}$$

which results in

$$\frac{1}{2}\|w_t\|_{H^{s-1}} \leq c\left(\delta^{1/2} + \|w\|_{H^s} + \|w\|_{H^{s-1}}\right). \tag{3.92}$$

It follows from (3.79) and (3.88) that $w_t \rightarrow 0$ as $\varepsilon, \delta \rightarrow 0$ in the H^{s-1} norm. This implies that u_ε is a Cauchy sequence in the spaces $C([0, T]; H^s(R))$ and $C([0, T]; H^{s-1}(R))$, respectively. The proof is completed. \square

Proof of Theorem 2.1. We consider the problem

$$\begin{aligned}
(1-\varepsilon)u_t - \varepsilon u_{txx} &= \left(1 - \partial_x^2\right)^{-1} \left[-\frac{k}{m+1}\left(u^{m+1}\right)_x - \frac{m+3}{m+2}\left(u^{m+2}\right)_x + \frac{1}{m+2}\partial_x^3\left(u^{m+2}\right)\right. \\
&\quad \left.- (m+1)\partial_x\left(u^m u_x^2\right) + u^m u_x u_{xx}\right],
\end{aligned} \tag{3.93}$$

$$u(0, x) = u_{\varepsilon 0}(x).$$

Letting $u(t, x)$ be the limit of the sequence u_ε and taking the limit in problem (3.93) as $\varepsilon \rightarrow 0$, from Lemma 3.14, it is easy to see that u is a solution of the problem

$$u_t = \left(1 - \partial_x^2\right)^{-1} \left[-\frac{k}{m+1} \left(u^{m+1}\right)_x - \frac{m+3}{m+2} \left(u^{m+2}\right)_x + \frac{1}{m+2} \partial_x^3 \left(u^{m+2}\right) - (m+1) \partial_x \left(u^m u_x^2\right) + u^m u_x u_{xx} \right], \quad (3.94)$$

$$u(0, x) = u_0(x),$$

and hence u is a solution of problem (3.94) in the sense of distribution. In particular, if $s \geq 4$, u is also a classical solution. Let u and v be two solutions of (3.94) corresponding to the same initial data u_0 such that $u, v \in C([0, T]; H^s(\mathbb{R}))$. Then $w = u - v$ satisfies the Cauchy problem

$$w_t = \left(1 - \partial_x^2\right)^{-1} \left\{ \partial_x \left[-\frac{k}{m+1} w g_m - \frac{m+3}{m+2} w g_{m+1} + \frac{1}{m+2} \partial_x^2 (w g_{m+1}) - \partial_x \left(u^{m+1}\right) \partial_x w - \partial_x \left(u^{m+1} - v^{m+1}\right) \partial_x v \right] + u^m u_x u_{xx} - v^m v_x v_{xx} \right\}, \quad (3.95)$$

$$w(0, x) = 0.$$

For any $1/2 < q < \min\{1, s-1\}$, applying the operator $\Lambda^q w \Lambda^q$ to both sides of equation (3.95) and integrating the resultant equation with respect to x , we obtain the equality

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|w\|_{H^q}^2 &= \int_{\mathbb{R}} (\Lambda^q w) \Lambda^{q-2} \left\{ \partial_x \left[-\frac{k}{m+1} w g_m - \frac{m+3}{m+2} w g_{m+1} + \frac{1}{m+2} \partial_x^2 (w g_{m+1}) - \partial_x \left(u^{m+1}\right) \partial_x w - \partial_x \left(u^{m+1} - v^{m+1}\right) \partial_x v \right] + u^m u_x u_{xx} - v^m v_x v_{xx} \right\} dx. \end{aligned} \quad (3.96)$$

By the similar estimates presented in Lemma 3.14, we have

$$\frac{d}{dt} \|w\|_{H^q}^2 \leq \tilde{c} \|w\|_{H^q}^2. \quad (3.97)$$

Using the Gronwall inequality leads to the conclusion that

$$\|w\|_{H^q} \leq 0 \times e^{\tilde{c}t} = 0 \quad (3.98)$$

for $t \in [0, \tilde{T})$. This completes the proof. \square

4. Global Existence of Strong Solutions

We study the differential equation

$$\begin{aligned} p_t &= u^{m+1}(t, p), \quad t \in [0, T], \\ p(0, x) &= x. \end{aligned} \quad (4.1)$$

Motivated by the Lagrangian viewpoint in fluid mechanics, by which one looks at the motion of individual fluid particles (see [43]), we state the following Lemma.

Lemma 4.1. *Let $u_0 \in H^s$, $s \geq 3$ and let $T > 0$ be the maximal existence time of the solution to problem (2.2). Then problem (4.1) has a unique solution $p \in C^1([0, T] \times R)$. Moreover, the map $p(t, \cdot)$ is an increasing diffeomorphism of R with $p_x(t, x) > 0$ for $(t, x) \in [0, T] \times R$.*

Proof. From Theorem 2.1, we have $u(t, x) \in C([0, T]; H^s(R)) \cap C^1([0, T]; H^{s-1}(R))$ and $H^s(R) \in C^1(R)$, where the Sobolev imbedding theorem is used. Thus, we conclude that both functions $u(t, x)$ and $u_x(t, x)$ are bounded, Lipschitz in space and C^1 in time. Using the existence and uniqueness theorem of ordinary differential equations derives that problem (4.1) has a unique solution $p \in C^1([0, T] \times R)$.

Differentiating (4.1) with respect to x yields that

$$\begin{aligned} \frac{d}{dt} p_x &= (m+1)u^m u_x(t, p) p_x, \quad t \in [0, T], \quad b \neq 0, \\ p_x(0, x) &= 1, \end{aligned} \quad (4.2)$$

which leads to

$$p_x(t, x) = \exp\left(\int_0^t (m+1)u^m u_x(\tau, p(\tau, x)) d\tau\right). \quad (4.3)$$

For every $T' < T$, using the Sobolev imbedding theorem yields that

$$\sup_{(\tau, x) \in [0, T'] \times R} |u_x(\tau, x)| < \infty. \quad (4.4)$$

It is inferred that there exists a constant $K_0 > 0$ such that $p_x(t, x) \geq e^{-K_0 t}$ for $(t, x) \in [0, T] \times R$. It completes the proof. \square

The next Lemma is reminiscent of a strong invariance property of the Camassa-Holm equation (the conservation of momentum [44, 45]).

Lemma 4.2. *Let $u_0 \in H^s$ with $s \geq 3$, and let $T > 0$ be the maximal existence time of the problem (2.2), it holds that*

$$y(t, p(t, x)) p_x^2(t, x) = y_0(x) e^{\int_0^t m u^m u_x d\tau}, \quad (4.5)$$

where $(t, x) \in [0, T] \times R$ and $y := u - u_{xx} + k/2(m+1)$.

Proof. We have

$$\begin{aligned}
 \frac{d}{dt} \left[y(t, p(t, x)) p_x^2(t, x) \right] &= y_t p_x^2 + 2y p_x p_{xt} + y_x p_t p_x^2 \\
 &= y_t p_x^2 + 2y(m+1)u^m u_x p_x^2 + u^{m+1} y_x p_x^2 \\
 &= \left[y_t + k u^m u_x + (m+2)u^m u_x y + y_x u^{m+1} \right] p_x^2 + m u^m u_x y p_x^2 \\
 &= \left[u_t - u_{txx} + k u^m u_x + (m+2)u^m u_x (u - u_{xx}) + u^{m+1} (u_x - u_{xxx}) \right] p_x^2 \\
 &\quad + m u^m u_x y p_x^2 \\
 &= m u^m u_x y p_x^2.
 \end{aligned} \tag{4.6}$$

Using $p_x(0, x) = 1$ and solving the above equation, we complete the proof of this lemma. \square

Lemma 4.3. *If $u_0 \in H^s$, $s \geq 3/2$, such that $(1 - \partial_x^2)u_0 + k/2(m+1) \geq 0$, then the solution of problem (2.2) satisfies the following:*

$$\|u_x\|_{L^\infty} \leq \|u\|_{L^\infty} + \frac{k}{2(m+1)} \leq c. \tag{4.7}$$

Proof. Using $u_0 - u_{0xx} + k/2(m+1) \geq 0$, it follows from Lemma 4.2 that $u - u_{xx} + k/2(m+1) \geq 0$. Letting $Y_1 = u - u_{xx}$, we have

$$u = \frac{1}{2} e^{-x} \int_{-\infty}^x e^\eta Y_1(t, \eta) d\eta + \frac{1}{2} e^x \int_x^\infty e^{-\eta} Y_1(t, \eta) d\eta, \tag{4.8}$$

from which we obtain that

$$\begin{aligned}
 \partial_x u(t, x) &= -\frac{1}{2} \left(e^{-x} \int_{-\infty}^x e^\eta Y_1(t, \eta) d\eta + e^x \int_x^\infty e^{-\eta} Y_1(t, \eta) d\eta \right) + e^x \int_x^\infty e^{-\eta} Y_1(t, \eta) d\eta \\
 &= -u(t, x) + e^x \int_x^\infty e^{-\eta} Y_1(t, \eta) d\eta \\
 &= -u(t, x) + e^x \int_x^\infty e^{-\eta} \left(Y_1(t, \eta) + \frac{k}{2(m+1)} \right) d\eta - \frac{k}{2(m+1)} e^x \int_x^\infty e^{-\eta} d\eta \\
 &= -u(t, x) + e^x \int_x^\infty e^{-\eta} (y(t, \eta)) d\eta - \frac{k}{2(m+1)} \\
 &\geq -u(t, x) - \frac{k}{2(m+1)}.
 \end{aligned} \tag{4.9}$$

On the other hand, we have

$$\begin{aligned}
 \partial_x u(t, x) &= \frac{1}{2} \left(e^{-x} \int_{-\infty}^x e^\eta Y_1(t, \eta) d\eta + e^x \int_x^\infty e^{-\eta} Y_1(t, \eta) d\eta \right) - e^{-x} \int_{-\infty}^x e^\eta Y_1(t, \eta) d\eta \\
 &= u(t, x) - e^{-x} \int_{-\infty}^x e^\eta Y_1(t, \eta) d\eta
 \end{aligned}$$

$$\begin{aligned}
&= u(t, x) - e^{-x} \int_{-\infty}^x e^{\eta} \left(Y_1(t, \eta) + \frac{k}{2(m+1)} \right) d\eta + \frac{k}{2(m+1)} e^{-x} \int_{-\infty}^x e^{\eta} d\eta \\
&= u(t, x) - e^{-x} \int_{-\infty}^x e^{\eta} y(t, \eta) d\eta + \frac{k}{2(m+1)} \\
&\leq u(t, x) + \frac{k}{2(m+1)}.
\end{aligned} \tag{4.10}$$

The inequalities (3.40), (4.9), and (4.10) derive that (4.7) is valid. \square

Proof of Theorem 2.2. Noting Remarks 3.6 and 3.8, $\|u\|_{H^1} \leq c$ and taking $q + 1 = s$ in inequality (3.15), we have

$$\|u\|_{H^s}^2 \leq \|u_0\|_{H^s}^2 + c \int_0^t \|u\|_{H^s}^2 (\|u_x\|_{L^\infty} + \|u_x\|_{L^\infty}^2) d\tau, \tag{4.11}$$

from which we obtain that

$$\|u\|_{H^s} \leq \|u_0\|_{H^s} e^{c \int_0^t (\|u_x\|_{L^\infty} + \|u_x\|_{L^\infty}^2) d\tau}. \tag{4.12}$$

Applying Lemma 4.3 derives

$$\|u\|_{H^s} \leq \|u_0\|_{H^s} e^{(c+c^2)t}. \tag{4.13}$$

From Theorem 2.1 and (4.13), we know that the result of Theorem 2.2 holds. \square

Acknowledgment

This work is supported by the Applied and Basic Project of Sichuan Province (2012JY0020).

References

- [1] R. Camassa and D. D. Holm, "An integrable shallow water equation with peaked solitons," *Physical Review Letters*, vol. 71, no. 11, pp. 1661–1664, 1993.
- [2] A. Constantin and D. Lannes, "The hydrodynamical relevance of the Camassa-Holm and Degasperis-Procesi equations," *Archive for Rational Mechanics and Analysis*, vol. 192, no. 1, pp. 165–186, 2009.
- [3] R. S. Johnson, "Camassa-Holm, Korteweg-de Vries and related models for water waves," *Journal of Fluid Mechanics*, vol. 455, no. 1, pp. 63–82, 2002.
- [4] D. Ionescu-Kruse, "Variational derivation of the Camassa-Holm shallow water equation," *Journal of Nonlinear Mathematical Physics*, vol. 14, no. 3, pp. 303–312, 2007.
- [5] R. S. Johnson, "The Camassa-Holm equation for water waves moving over a shear flow," *Japan Society of Fluid Mechanics. Fluid Dynamics Research. An International Journal*, vol. 33, no. 1-2, pp. 97–111, 2003.
- [6] H.-H. Dai, "Model equations for nonlinear dispersive waves in a compressible Mooney-Rivlin rod," *Acta Mechanica*, vol. 127, no. 1–4, pp. 193–207, 1998.
- [7] A. Constantin and W. A. Strauss, "Stability of a class of solitary waves in compressible elastic rods," *Physics Letters. A*, vol. 270, no. 3–4, pp. 140–148, 2000.
- [8] M. Lakshmanan, "Integrable nonlinear wave equations and possible connections to tsunami dynamics," in *Tsunami and Nonlinear Waves*, A. Kundu, Ed., pp. 31–49, Springer, Berlin, Germany, 2007.

- [9] A. Constantin and R. S. Johnson, "Propagation of very long water waves, with vorticity, over variable depth, with applications to tsunamis," *Fluid Dynamics Research*, vol. 40, no. 3, pp. 175–211, 2008.
- [10] A. Constantin, "The trajectories of particles in Stokes waves," *Inventiones Mathematicae*, vol. 166, no. 3, pp. 523–535, 2006.
- [11] A. Constantin and J. Escher, "Particle trajectories in solitary water waves," *Bulletin of American Mathematical Society*, vol. 44, no. 3, pp. 423–431, 2007.
- [12] A. Constantin and W. A. Strauss, "Stability of peakons," *Communications on Pure and Applied Mathematics*, vol. 53, no. 5, pp. 603–610, 2000.
- [13] A. Constantin and J. Escher, "Analyticity of periodic traveling free surface water waves with vorticity," *Annals of Mathematics*, vol. 173, no. 1, pp. 559–568, 2011.
- [14] K. El Dika and L. Molinet, "Stability of multipeakons," *Annales de l'Institut Henri Poincaré. Analyse Non Linéaire*, vol. 26, no. 4, pp. 1517–1532, 2009.
- [15] A. Constantin and W. A. Strauss, "Stability of the Camassa-Holm solitons," *Journal of Nonlinear Science*, vol. 12, no. 4, pp. 415–422, 2002.
- [16] A. Constantin and H. P. McKean, "A shallow water equation on the circle," *Communications on Pure and Applied Mathematics*, vol. 52, no. 8, pp. 949–982, 1999.
- [17] A. Constantin, "On the inverse spectral problem for the Camassa-Holm equation," *Journal of Functional Analysis*, vol. 155, no. 2, pp. 352–363, 1998.
- [18] A. Constantin, V. S. Gerdjikov, and R. I. Ivanov, "Inverse scattering transform for the Camassa-Holm equation," *Inverse Problems*, vol. 22, no. 6, pp. 2197–2207, 2006.
- [19] H. P. McKean, "Integrable systems and algebraic curves," in *Global Analysis*, vol. 755 of *Lecture Notes in Mathematics*, pp. 83–200, Springer, Berlin, Germany, 1979.
- [20] A. Constantin, T. Kappeler, B. Kolev, and P. Topalov, "On geodesic exponential maps of the Virasoro group," *Annals of Global Analysis and Geometry*, vol. 31, no. 2, pp. 155–180, 2007.
- [21] G. Misiulek, "A shallow water equation as a geodesic flow on the Bott-Virasoro group," *Journal of Geometry and Physics*, vol. 24, no. 3, pp. 203–208, 1998.
- [22] Z. Xin and P. Zhang, "On the weak solutions to a shallow water equation," *Communications on Pure and Applied Mathematics*, vol. 53, no. 11, pp. 1411–1433, 2000.
- [23] Z. Xin and P. Zhang, "On the uniqueness and large time behavior of the weak solutions to a shallow water equation," *Communications in Partial Differential Equations*, vol. 27, no. 9-10, pp. 1815–1844, 2002.
- [24] G. M. Coclite, H. Holden, and K. H. Karlsen, "Global weak solutions to a generalized hyperelastic-rod wave equation," *SIAM Journal on Mathematical Analysis*, vol. 37, no. 4, pp. 1044–1069, 2005.
- [25] A. Bressan and A. Constantin, "Global conservative solutions of the Camassa-Holm equation," *Archive for Rational Mechanics and Analysis*, vol. 183, no. 2, pp. 215–239, 2007.
- [26] A. Bressan and A. Constantin, "Global dissipative solutions of the Camassa-Holm equation," *Analysis and Applications*, vol. 5, no. 1, pp. 1–27, 2007.
- [27] H. Holden and X. Raynaud, "Dissipative solutions for the Camassa-Holm equation," *Discrete and Continuous Dynamical Systems*, vol. 24, no. 4, pp. 1047–1112, 2009.
- [28] H. Holden and X. Raynaud, "Global conservative solutions of the Camassa-Holm equation—a Lagrangian point of view," *Communications in Partial Differential Equations*, vol. 32, no. 10–12, pp. 1511–1549, 2007.
- [29] Y. A. Li and P. J. Olver, "Well-posedness and blow-up solutions for an integrable nonlinearly dispersive model wave equation," *Journal of Differential Equations*, vol. 162, no. 1, pp. 27–63, 2000.
- [30] A. Constantin and J. Escher, "Wave breaking for nonlinear nonlocal shallow water equations," *Acta Mathematica*, vol. 181, no. 2, pp. 229–243, 1998.
- [31] R. Beals, D. H. Sattinger, and J. Szmigielski, "Multi-peakons and a theorem of Stieltjes," *Inverse Problems*, vol. 15, no. 1, pp. L1–L4, 1999.
- [32] D. Henry, "Persistence properties for a family of nonlinear partial differential equations," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 70, no. 4, pp. 1565–1573, 2009.
- [33] S. Hakkaev and K. Kirchev, "Local well-posedness and orbital stability of solitary wave solutions for the generalized Camassa-Holm equation," *Communications in Partial Differential Equations*, vol. 30, no. 4–6, pp. 761–781, 2005.
- [34] G. Rodríguez-Blanco, "On the Cauchy problem for the Camassa-Holm equation," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 46, no. 3, pp. 309–327, 2001.
- [35] Z. Yin, "On the blow-up scenario for the generalized Camassa-Holm equation," *Communications in Partial Differential Equations*, vol. 29, no. 5–6, pp. 867–877, 2004.
- [36] Z. Yin, "On the Cauchy problem for an integrable equation with peakon solutions," *Illinois Journal of Mathematics*, vol. 47, no. 3, pp. 649–666, 2003.

- [37] Y. Zhou, "Wave breaking for a periodic shallow water equation," *Journal of Mathematical Analysis and Applications*, vol. 290, no. 2, pp. 591–604, 2004.
- [38] S. Lai and Y. Wu, "The local well-posedness and existence of weak solutions for a generalized Camassa-Holm equation," *Journal of Differential Equations*, vol. 248, no. 8, pp. 2038–2063, 2010.
- [39] S. Y. Wu and Z. Yin, "Global existence and blow-up phenomena for the weakly dissipative Camassa-Holm equation," *Journal of Differential Equations*, vol. 246, no. 11, pp. 4309–4321, 2009.
- [40] T. Kato, "Quasi-linear equations of evolution, with applications to partial differential equations," in *Spectral Theory and Differential Equations*, vol. 448 of *Lecture notes in Mathematics*, pp. 25–70, Springer, Berlin, 1975.
- [41] T. Kato and G. Ponce, "Commutator estimates and the Euler and Navier-Stokes equations," *Communications on Pure and Applied Mathematics*, vol. 41, no. 7, pp. 891–907, 1988.
- [42] W. Walter, *Differential and Integral Inequalities*, Springer-Verlag, New York, NY, USA, 1970.
- [43] B. Kolev, "Lie groups and mechanics: an introduction," *Journal of Nonlinear Mathematical Physics*, vol. 11, no. 4, pp. 480–498, 2004.
- [44] A. Constantin, "Existence of permanent and breaking waves for a shallow water equation: a geometric approach," *Annales de l'Institut Fourier*, vol. 50, no. 2, pp. 321–362, 2000.
- [45] B. Kolev, "Poisson brackets in hydrodynamics," *Discrete and Continuous Dynamical Systems*, vol. 19, no. 3, pp. 555–574, 2007.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

