

## Research Article

# Powers of Convex-Cyclic Operators

Fernando León-Saavedra<sup>1</sup> and María del Pilar Romero-de la Rosa<sup>2</sup>

<sup>1</sup> Department of Mathematics, University of Cádiz, Campus de Jerez de la Frontera, Cádiz, 11405 Jerez de la Frontera, Spain

<sup>2</sup> IES Sofía, Doctor Marañón 13, Cádiz, 11407 Jerez de la Frontera, Spain

Correspondence should be addressed to María del Pilar Romero-de la Rosa; pilar.romero@uca.es

Received 21 January 2014; Revised 15 April 2014; Accepted 16 April 2014; Published 8 May 2014

Academic Editor: Manuel Maestre

Copyright © 2014 F. León-Saavedra and M. P. Romero-de la Rosa. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

A bounded operator  $T$  on a Banach space  $X$  is convex cyclic if there exists a vector  $x$  such that the convex hull generated by the orbit  $\{T^n x\}_{n \geq 0}$  is dense in  $X$ . In this note we study some questions concerned with convex-cyclic operators. We provide an example of a convex-cyclic operator  $T$  such that the power  $T^n$  fails to be convex cyclic. Using this result we solve three questions posed by Rezaei (2013).

## 1. Introduction and Main Results

Throughout this paper we denote by  $L(X)$  the algebra of all bounded linear operators on a real or complex infinite dimensional Banach space  $X$ . An operator  $T \in L(X)$  is said to be cyclic if there exists a vector  $x \in X$  (later called cyclic vector for  $T$ ) such that the linear span of the orbit

$$\text{linear span}(\{T^n x : n \in \mathbb{N}\}) \quad (1)$$

is dense in  $X$ . If the orbit  $\text{Orb}(T, x) := \{T^n x : n \in \mathbb{N}\}$  is dense itself, without the help of the linear span, then  $T$  is called hypercyclic and  $x$  is called hypercyclic for  $T$ . In the midway stand several notions studied by different authors. For instance, the operator  $T$  is said to be supercyclic if the projective orbit is dense in  $X$ . We refer to the books [1, 2] and references therein for further information on hypercyclic operators.

When we sometimes abusively say that a polynomial  $p(z)$  is a convex polynomial, what we really mean is that  $p(z) = t_0 + t_1 z + t_2 z^2 + \dots + t_n z^n$ ,  $t_i \in \mathbb{R}$ ,  $i = 0, \dots, n$ , and  $t_0 + t_1 + \dots + t_n = 1$ . We will focus our attention on the notion of convex cyclicity introduced by Rezaei in [3]. An operator  $T$  is said to be convex cyclic if there exists a vector  $x \in X$  such that the real convex hull of the orbit (denoted by  $\text{co}(\text{Orb}(T, x))$ )

$$\text{co}(\text{Orb}(T, x)) = \{p(T)x : p \text{ convex polynomial}\} \quad (2)$$

is dense in  $X$ .

In [3] are characterized the convex-cyclic matrices in finite dimension, and the author develops the main properties in the infinite dimensional setting.

A result by Ansari [4] states that if  $T$  is a hypercyclic operator then  $T^n$  is also hypercyclic; this fact is not true for cyclic operators. In this paper we show that Ansari's result fails also for convex-cyclic operators, solving a question posed in [3].

Another result proved by Bourdon and Feldman on hypercyclic operators says that if the orbit of a vector is somewhere dense, then it is dense (see [5]). From our previous counterexample we can construct a non-convex-cyclic operator  $T$  such that the  $\text{co}(\text{Orb}(T, x))$  has nonempty interior. That is, Bourdon and Feldman's result is not true in the convex-cyclic setting. Finally we can construct a convex-cyclic operator  $T$  such that  $T$  is not weakly hypercyclic; that is, its orbit is not dense in the weak operator topology. The later examples solve Questions 5.5 and 5.6 in [3].

## 2. Powers of a Convex-Cyclic Operator

The first example of hypercyclic operator on Banach spaces was discovered by Rolewicz (see [6]). Throughout this section  $\mathcal{B} = \ell_p$ ,  $1 \leq p < \infty$  or  $c_0$  of complex valued sequences. Rolewicz's operator  $\mu B$  with  $|\mu| > 1$  is defined on  $\mathcal{B}$  by

$$\mu B(x_0, x_1, \dots, x_n, \dots) = \mu(x_1, x_2, \dots, x_n, \dots), \quad (3)$$

where  $B$  denotes the backward shift operator.

**Lemma 1.** Set  $\alpha = e^{2\pi i/3}$  and  $r_0 > 1$ . For any  $z_0 \in \mathbb{C} \setminus \{0\}$  there exist  $k_0 \geq 0$  and a sequence of polynomials  $p_k(z)$  such that

- (1)  $p_k(z) = (t_{1,k} + t_{2,k}z + t_{3,k}z^2)z^k$  for all  $k \geq k_0$ ;
- (2)  $t_{i,k} \in [0, 1]$  and  $t_{1,k} + t_{2,k} + t_{3,k} = 1, i \in \{1, 2, 3\}$  and  $k \geq k_0$ ;
- (3)  $p_k(r_0\alpha) = z_0, k \geq k_0$ .

*Proof.* Let us denote by  $\mathcal{T}$  the triangle with vertices  $\{1, r_0\alpha, r_0^2\alpha^2\}$ . Since  $|r_0\alpha| > 1$  and  $0 \in \mathcal{T}$ , there exists  $k_0$  such that  $z_k = z_0/(r_0\alpha)^k \in \mathcal{T}$  for all  $k \geq k_0$ . Then, there exist barycentric coordinates  $t_{i,k} \in [0, 1] i = 1, 2, 3$  satisfying  $t_{1,k} + t_{2,k}r_0\alpha + t_{3,k}(r_0\alpha)^2 = z_k$  and  $t_{1,k} + t_{2,k} + t_{3,k} = 1$ . Then, the polynomials  $p_k(z) = (t_{1,k} + t_{2,k}z + t_{3,k}z^2)z^k$ , for all  $k \geq k_0$ , yield the desired result.  $\square$

**Lemma 2.** Let  $p_k$  be a sequence of polynomials satisfying Conditions (1)–(3) of Lemma 1. Then, there exists a  $G_\delta$  dense subset  $Z \subset \mathcal{B}$  of vectors such that  $\{p_k(\mu B)x_0\}_{k \geq k_0}$  is dense in  $\mathcal{B}$  for all  $x_0 \in Z$ .

*Proof.* We will use some hypercyclicity criterion version for sequence of operators (see [2, Theorem 3.24]); that is, we will show the existence of two dense subsets  $X$  and  $Y$  and a sequence of mappings  $S_k$  such that

- (i)  $\lim_k p_k(\mu B)x = 0 \forall x \in X$ ;
- (ii)  $p_k(\mu B)S_k y = y \forall y \in Y$ ;
- (iii)  $\lim_k S_k y = 0 \forall y \in Y$ .

Let us consider the subsets

$$X = \text{span} \{ \text{Ker}(\mu B - \lambda I) : |\lambda| < 1 \}, \tag{4}$$

$$Y = \{ \text{Ker}(\mu B - \lambda I) : \lambda \in \mathbb{R}, 1 < \lambda < |\mu| \},$$

which are dense in  $\mathcal{B}$  (see [2, Example 3.2, page 70]).

If  $x \in \text{Ker}(\mu B - \lambda I)$  with  $|\lambda| < 1$ , then

$$p_k(\mu B)x = (t_{1,k}(\mu B)^k + t_{2,k}(\mu B)^{k+1} + t_{3,k}(\mu B)^{k+2})x$$

$$= (t_{1,k}I + t_{2,k}(\mu B) + t_{3,k}(\mu B)^2)(\mu B)^k x \tag{5}$$

$$\leq \text{const}|\lambda|^k,$$

which goes to zero when  $k \rightarrow \infty$ ; therefore, Condition (i) is fulfilled.

Denoting by  $q_k(z) = t_{1,k} + t_{2,k}z + t_{3,k}z^2$ , since  $t_{1,k}, t_{2,k}, t_{3,k}$  are barycentric coordinates of a triangle, then  $q_k(\lambda)$  lies in the degenerate triangle with vertices  $\{1, \lambda, \lambda^2\}$ , in particular  $q_k(\lambda) \geq 1$ .

Let us take  $y \in \text{Ker}(\mu B - \lambda I)$  with  $\lambda \in \mathbb{R}$  and  $1 < \lambda < |\mu|$ , and let us define the mapping  $S_k$  on  $y$  as

$$S_k y = \frac{1}{\lambda^k q_k(\lambda)} y \tag{6}$$

and we extend linearly  $S_k$  on  $Y$ . Clearly  $S_k y \rightarrow 0$  as  $k \rightarrow \infty$  for all  $y \in Y$  and  $p_k(\mu B)S_k y = y$  for all  $y \in Y$ . Thus, by the hypercyclicity criterion there exists a  $G_\delta$  dense subset of vectors  $x_0 \in \mathcal{B}$  such that  $\{p_k(\mu B)x_0\}_{k \geq k_0}$  is dense in  $\mathcal{B}$ .  $\square$

Now, let us prove the main result of this section, which solves Question 5.6 in [3].

**Theorem 3.** The operator  $T = r_0\alpha I_{\mathbb{C}} \oplus \mu B$  is convex cyclic on  $\mathbb{C} \oplus \mathcal{B}$ ; however  $T^3$  is not.

*Proof.* If  $p$  is a polynomial, then  $p(T) = p(r_0\alpha) \oplus p(\mu B)$ . Let us observe that the first coordinate of the powers of  $(T^3)^n$  are only real numbers. Take  $x = \sum_{n=0}^{\infty} x_0 e_0 \in \mathbb{C} \oplus \mathcal{B}$ . If  $f^*$  is the projection on the first coordinate,

$$\{f^*(\text{co}(\text{Orb}(T^3, x)))\} = tx_0, \forall t \geq r_0 \tag{7}$$

which is not dense in  $\mathbb{C}$ . Therefore,  $T^3$  is not convex cyclic.

Now, let us prove that  $T$  is a convex-cyclic operator using a direct application of the Baire category theorem (see, for instance, [2, Theorem 1.57]). Thus  $T$  is convex cyclic if for any nonempty open subsets  $U, V \subset \mathbb{C} \oplus \mathcal{B}$ , there exists a convex polynomial  $p(z)$  such that  $p(T)(U) \cap V \neq \emptyset$ .

Indeed, let  $U = G_1 \times W_1$  and  $V = G_2 \times W_2$  open subsets of  $\mathbb{C} \oplus \mathcal{B}$ , where  $G_i \subset \mathbb{C}$  and  $W_i \subset \mathcal{B}, i = 1, 2$ , are nonempty open subsets. Let  $z_1 \in G_1$  and  $z_2 \in G_2$  with  $z_1 z_2 \neq 0$ . Set  $z_0 = z_2/z_1$  and  $p_k(z)$  the sequence of polynomials which guarantees Lemma 1. Hence we have  $p_k(r_0\alpha) = z_2/z_1$  and therefore  $p_k(r_0\alpha)z_1 = z_2$  (this fact will imply that  $p_k(T)$  acting on  $G_1$  will intersect  $G_2$ ). Now we apply Lemma 2 and we obtain a  $G_\delta$  dense subset  $Z \subset \mathcal{B}$  of hypercyclic vectors for the sequence  $\{p_k(\mu B)\}$ . Thus there exist  $x_0 \in W_1$  and a subsequence  $\{n_k\}$  such that  $p_{n_k}(\mu B)x_0 \in W_2$ . Therefore  $p_{n_k}(T)(U) \cap V \neq \emptyset$ , which yields the desired result.  $\square$

*Remark 4.* If we take  $\alpha = e^{2\pi i/n}$  with  $n \geq 4$ , using similar arguments as in Theorem 3, we can show that  $T = r_0\alpha \oplus \mu B$  is convex cyclic on  $\mathbb{C} \oplus \mathcal{B}$  but  $T^n$  is not (if  $n = 4$  the operator  $T$  is convex cyclic but  $T^2$  is not).

*Remark 5.* If we consider the Rolewicz operator on real spaces  $\ell^p, 1 \leq p < \infty$  or  $c_0$ , then we can get that the operator  $T = -r_0 \oplus \mu B (r_0 > 1)$  is convex cyclic on  $\mathbb{R} \oplus \ell^p, 1 \leq p < \infty$  or  $\mathbb{R} \oplus c_0$ , but clearly  $T^2$  is not. Lemma 1 can be adapted clearly to the real case. Now it is well known that if we consider Rolewicz's operator on real spaces  $\ell^p, 1 \leq p < \infty$  or  $c_0$ , then its complexification can be identified with the same operator on the corresponding spaces of complex sequences. With some slight modification in the proof of Corollary 2.51 in [2] we can obtain that Lemma 2 continues being true on real spaces. The rest of the proof is straightforward.

*Remark 6.* Another difference between hypercyclic operators and convex-cyclic operators is the following: hypercyclic operators are invariant under unimodular multiplications (see [7]); that is, if  $T$  is hypercyclic, then  $\lambda T$  is also with  $|\lambda| = 1$ . However this is not true for convex-cyclic operators; the previous counterexample  $T = r_0\alpha I_{\mathbb{C}} \oplus \mu B$  is convex cyclic; however  $\bar{\alpha}T$  is not.

Now, let  $\alpha = e^{2\pi i/3}$ . Let us consider the operator  $T = \alpha I_{\mathbb{C}} \oplus \mu B$ , that is, the same operator provided by Theorem 3 but without the multiplier factor  $r_0$ . Then, an easy check shows

that the set  $S = \{p_k(\alpha) : p_k(z) \text{ convex polynomial}\}$  is contained in the unit disk. Moreover, the set  $S$  has nonempty interior in  $\mathbb{C}$ ; for instance,  $S$  contains the triangle  $\mathcal{T}$  with vertices  $\{1, \alpha, \alpha^2\}$ . Using the arguments of Theorem 3 we can find a vector  $x_0 \in \mathcal{B}$  such that the convex orbit

$$\{p(T)(1 \oplus x_0) : p \text{ convex polynomial}\} \quad (8)$$

is dense in  $\mathcal{T} \oplus \mathcal{B}$ . Therefore the convex orbit has nonempty interior. However the operator  $T$  is not convex cyclic. This solves Question 5.5 in Rezaei's paper.

**Proposition 7.** *Bourdon and Feldman's result fails for convex-cyclic operators.*

The adjoint of the operator  $T$  in Theorem 3 has an eigenvalue; therefore,  $T$  cannot be weakly hypercyclic. This solves Question 5.4 in [3].

## Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

## Acknowledgments

The authors are indebted to the referees for alerting them to some details that they had to take care of in the proof of the first version. Fernando León-Saavedra and María del Pilar Romero de la Rosa were partially supported by Junta de Andalucía FQM-257.

## References

- [1] F. Bayart and E. Matheron, *Dynamics of Linear Operators*, vol. 179 of *Cambridge Tracts in Mathematics*, Cambridge University Press, Cambridge, UK, 2009.
- [2] K.-G. Grosse-Erdmann and A. P. Manguillot, *Linear Chaos. Universitext*, Springer, London, UK, 2011.
- [3] H. Rezaei, "On the convex hull generated by orbit of operators," *Linear Algebra and Its Applications*, vol. 438, no. 11, pp. 4190–4203, 2013.
- [4] S. I. Ansari, "Hypercyclic and cyclic vectors," *Journal of Functional Analysis*, vol. 128, no. 2, pp. 374–383, 1995.
- [5] P. S. Bourdon and N. S. Feldman, "Somewhere dense orbits are everywhere dense," *Indiana University Mathematics Journal*, vol. 52, no. 3, pp. 811–819, 2003.
- [6] S. Rolewicz, "On orbits of elements," *Studia Mathematica*, vol. 32, pp. 17–22, 1969.
- [7] F. León-Saavedra and V. Müller, "Rotations of hypercyclic and supercyclic operators," *Integral Equations and Operator Theory*, vol. 50, no. 3, pp. 385–391, 2004.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

