

## Research Article

# Algorithms of Common Solutions for Generalized Mixed Equilibria, Variational Inclusions, and Constrained Convex Minimization

Lu-Chuan Ceng<sup>1</sup> and Suliman Al-Homidan<sup>2</sup>

<sup>1</sup> Department of Mathematics, Shanghai Normal University and Scientific Computing Key Laboratory of Shanghai Universities, Shanghai 200234, China

<sup>2</sup> Department of Mathematics and Statistics, King Fahd University of Petroleum and Minerals, Dhahran, Saudi Arabia

Correspondence should be addressed to Lu-Chuan Ceng; [zenglc@hotmail.com](mailto:zenglc@hotmail.com)

Received 3 November 2013; Accepted 12 November 2013; Published 23 January 2014

Academic Editor: Qamrul Hasan Ansari

Copyright © 2014 L.-C. Ceng and S. Al-Homidan. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We introduce new implicit and explicit iterative algorithms for finding a common element of the set of solutions of the minimization problem for a convex and continuously Fréchet differentiable functional, the set of solutions of a finite family of generalized mixed equilibrium problems, and the set of solutions of a finite family of variational inclusions in a real Hilbert space. Under suitable control conditions, we prove that the sequences generated by the proposed algorithms converge strongly to a common element of three sets, which is the unique solution of a variational inequality defined over the intersection of three sets.

## 1. Introduction

Let  $C$  be a nonempty closed convex subset of a real Hilbert space  $H$  and let  $P_C$  be the metric projection of  $H$  onto  $C$ . Let  $S : C \rightarrow C$  be a self-mapping on  $C$ . We denote by  $\text{Fix}(S)$  the set of fixed points of  $S$  and by  $\mathbf{R}$  the set of all real numbers. A mapping  $A : C \rightarrow H$  is called  $L$ -Lipschitz continuous if there exists a constant  $L \geq 0$  such that

$$\|Ax - Ay\| \leq L \|x - y\|, \quad \forall x, y \in C. \quad (1)$$

In particular, if  $L = 1$ , then  $A$  is called a nonexpansive mapping [1]; if  $L \in [0, 1)$ , then  $A$  is called a contraction.

A mapping  $V$  is called strongly positive on  $H$  if there exists a constant  $\mu > 0$  such that

$$\langle Vx, x \rangle \geq \mu \|x\|^2, \quad \forall x \in H. \quad (2)$$

Let  $A : C \rightarrow H$  be a nonlinear mapping on  $C$ . We consider the following variational inequality problem (VIP): find a point  $x \in C$  such that

$$\langle Ax, y - x \rangle \geq 0, \quad \forall y \in C. \quad (3)$$

The solution set of VIP (3) is denoted by  $\text{VI}(C, A)$ .

The VIP (3) was first discussed by Lions [2]. There are many applications of VIP (3) in various fields; see, for example, [3–6]. It is well known that if  $A$  is a strongly monotone and Lipschitz continuous mapping on  $C$ , then VIP (3) has a unique solution. In 1976, Korpelevič [7] proposed an iterative algorithm for solving the VIP (3) in Euclidean space  $\mathbf{R}^n$ :

$$\begin{aligned} y_n &= P_C(x_n - \tau Ax_n), \\ x_{n+1} &= P_C(x_n - \tau Ay_n), \quad \forall n \geq 0, \end{aligned} \quad (4)$$

with  $\tau > 0$  a given number, which is known as the extragradient method (see also [8]). The literature on the VIP is vast and Korpelevič's extragradient method has received great attention given by many authors, who improved it in various ways; see, for example, [9–24] and references therein, to name but a few.

Let  $\varphi : C \rightarrow \mathbf{R}$  be a real-valued function,  $A : H \rightarrow H$  a nonlinear mapping, and  $\Theta : C \times C \rightarrow \mathbf{R}$  a bifunction. In 2008, Peng and Yao [12] introduced the following generalized

mixed equilibrium problem (GMEP) of finding  $x \in C$  such that

$$\Theta(x, y) + \varphi(y) - \varphi(x) + \langle Ax, y - x \rangle \geq 0, \quad \forall y \in C. \quad (5)$$

We denote the set of solutions of GMEP (5) by  $\text{GMEP}(\Theta, \varphi, A)$ . The GMEP (5) is very general in the sense that it includes, as special cases, optimization problems, variational inequalities, minimax problems, and Nash equilibrium problems in noncooperative games. The GMEP is further considered and studied; see, for example, [11, 14, 23, 25–28]. If  $\varphi = 0$  and  $A = 0$ , then GMEP (5) reduces to the equilibrium problem (EP) which is to find  $x \in C$  such that

$$\Theta(x, y) \geq 0, \quad \forall y \in C. \quad (6)$$

It is considered and studied in [29]. The set of solutions of EP is denoted by  $\text{EP}(\Theta)$ . It is worth mentioning that the EP is a unified model of several problems, namely, variational inequality problems, optimization problems, saddle point problems, complementarity problems, fixed point problems, Nash equilibrium problems, and so forth.

Throughout this paper, it is assumed as in [12] that  $\Theta : C \times C \rightarrow \mathbf{R}$  is a bifunction satisfying conditions (A1)–(A4) and  $\varphi : C \rightarrow \mathbf{R}$  is a lower semicontinuous and convex function with restriction (B1) or (B2), where

- (A1)  $\Theta(x, x) = 0$  for all  $x \in C$ ;
- (A2)  $\Theta$  is monotone; that is,  $\Theta(x, y) + \Theta(y, x) \leq 0$  for any  $x, y \in C$ ;
- (A3)  $\Theta$  is upper-hemicontinuous; that is, for each  $x, y, z \in C$ ,

$$\limsup_{t \rightarrow 0^+} \Theta(tz + (1-t)x, y) \leq \Theta(x, y); \quad (7)$$

- (A4)  $\Theta(x, \cdot)$  is convex and lower semicontinuous for each  $x \in C$ ;
- (B1) for each  $x \in H$  and  $r > 0$ , there exists a bounded subset  $D_x \subset C$  and  $y_x \in C$  such that, for any  $z \in C \setminus D_x$ ,

$$\Theta(z, y_x) + \varphi(y_x) - \varphi(z) + \frac{1}{r} \langle y_x - z, z - x \rangle < 0; \quad (8)$$

- (B2)  $C$  is a bounded set.

Next we list some elementary results for the MEP.

**Proposition 1** (see [26]). *Assume that  $\Theta : C \times C \rightarrow \mathbf{R}$  satisfies (A1)–(A4) and let  $\varphi : C \rightarrow \mathbf{R}$  be a proper lower semicontinuous and convex function. Assume that either (B1) or (B2) holds. For  $r > 0$  and  $x \in H$ , define a mapping  $T_r^{(\Theta, \varphi)} : H \rightarrow C$  as follows:*

$$T_r^{(\Theta, \varphi)}(x) = \left\{ z \in C : \Theta(z, y) + \varphi(y) - \varphi(z) + \frac{1}{r} \langle y - z, z - x \rangle \geq 0, \forall y \in C \right\}, \quad (9)$$

for all  $x \in H$ . Then the following hold:

(i) for each  $x \in H$ ,  $T_r^{(\Theta, \varphi)}(x)$  is nonempty and single-valued;

(ii)  $T_r^{(\Theta, \varphi)}$  is firmly nonexpansive; that is, for any  $x, y \in H$ ,

$$\|T_r^{(\Theta, \varphi)}x - T_r^{(\Theta, \varphi)}y\|^2 \leq \langle T_r^{(\Theta, \varphi)}x - T_r^{(\Theta, \varphi)}y, x - y \rangle; \quad (10)$$

(iii)  $\text{Fix}(T_r^{(\Theta, \varphi)}) = \text{MEP}(\Theta, \varphi)$ ;

(iv)  $\text{MEP}(\Theta, \varphi)$  is closed and convex;

(v)  $\|T_s^{(\Theta, \varphi)}x - T_t^{(\Theta, \varphi)}x\|^2 \leq ((s-t)/s) \langle T_s^{(\Theta, \varphi)}x - T_t^{(\Theta, \varphi)}x, T_s^{(\Theta, \varphi)}x - x \rangle$  for all  $s, t > 0$  and  $x \in H$ .

Let  $\lambda_{n,1}, \lambda_{n,2}, \dots, \lambda_{n,N} \in (0, 1]$ ,  $n \geq 1$ . Given the nonexpansive mappings  $S_1, S_2, \dots, S_N$  on  $H$ , for each  $n \geq 1$ , the mappings  $U_{n,1}, U_{n,2}, \dots, U_{n,N}$  are defined by

$$\begin{aligned} U_{n,1} &= \lambda_{n,1}S_1 + (1 - \lambda_{n,1})I, \\ U_{n,2} &= \lambda_{n,2}S_nU_{n,1} + (1 - \lambda_{n,2})I, \\ U_{n,n-1} &= \lambda_{n-1}T_{n-1}U_{n,n} + (1 - \lambda_{n-1})I, \\ &\vdots \\ U_{n,N-1} &= \lambda_{n,N-1}S_{N-1}U_{n,N-2} + (1 - \lambda_{n,N-1})I, \\ W_n &:= U_{n,N} = \lambda_{n,N}S_NU_{n,N-1} + (1 - \lambda_{n,N})I. \end{aligned} \quad (11)$$

The  $W_n$  is called the  $W$ -mapping generated by  $S_1, \dots, S_N$  and  $\lambda_{n,1}, \lambda_{n,2}, \dots, \lambda_{n,N}$ . Note that the nonexpansivity of  $S_i$  implies the nonexpansivity of  $W_n$ .

In 2012, combining the hybrid steepest-descent method in [30] and hybrid viscosity approximation method in [31], Ceng et al. [27] proposed and analyzed the following hybrid iterative method for finding a common element of the set of solutions of GMEP (5) and the set of fixed points of a finite family of nonexpansive mappings  $\{S_i\}_{i=1}^N$ .

**Theorem CGY** (see [27, Theorem 3.1]). *Let  $C$  be a nonempty closed convex subset of a real Hilbert space  $H$ . Let  $\Theta : C \times C \rightarrow \mathbf{R}$  be a bifunction satisfying assumptions (A1)–(A4) and  $\varphi : C \rightarrow \mathbf{R}$  a lower semicontinuous and convex function with restriction (B1) or (B2). Let the mapping  $A : H \rightarrow H$  be  $\delta$ -inverse-strongly monotone and  $\{S_i\}_{i=1}^N$  a finite family of nonexpansive mappings on  $H$  such that  $\bigcap_{i=1}^N \text{Fix}(S_i) \cap \text{GMEP}(\Theta, \varphi, A) \neq \emptyset$ . Let  $F : H \rightarrow H$  be a  $\kappa$ -Lipschitzian and  $\eta$ -strongly monotone operator with constants  $\kappa, \eta > 0$  and  $V : H \rightarrow H$  a  $\rho$ -Lipschitzian mapping with constant  $\rho \geq 0$ . Let  $0 < \mu < 2\eta/\kappa^2$  and  $0 \leq \gamma\rho < \tau$ , where  $\tau = 1 - \sqrt{1 - \mu(2\eta - \mu\kappa^2)}$ . Suppose  $\{\alpha_n\}$  and  $\{\beta_n\}$  are two sequences in  $(0, 1)$ ,  $\{\gamma_n\}$  is a sequence in  $(0, 2\delta]$ , and  $\{\lambda_{n,i}\}_{i=1}^N$  is a sequence in  $[a, b]$  with  $0 < a \leq b < 1$ . For every  $n \geq 1$ , let  $W_n$  be the  $W$ -mapping generated by  $S_1, \dots, S_N$  and  $\lambda_{n,1}, \lambda_{n,2}, \dots, \lambda_{n,N}$ .*



- (C1)  $\lim_{n \rightarrow \infty} \alpha_n = 0$  and  $\sum_{n=1}^{\infty} \alpha_n = \infty$ ;
- (C2)  $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$ .

Then the sequence  $\{x_n\}$  converges strongly to  $x^* \in \Omega$ , where  $x^* = P_{\Omega}(\gamma f(x^*) + (I - V)x^*)$  is a unique solution of the VIP:

$$\langle (\gamma f - V)x^*, y - x^* \rangle \leq 0, \quad \forall y \in \Omega. \tag{17}$$

Let  $f : C \rightarrow \mathbf{R}$  be a convex and continuously Fréchet differentiable functional. Consider the convex minimization problem (CMP) of minimizing  $f$  over the constraint set  $C$

$$\min_{x \in C} f(x) \tag{18}$$

(assuming the existence of minimizers). We denote by  $\Gamma$  the set of minimizers of CMP (18). It is well known that the gradient-projection algorithm (GPA) generates a sequence  $\{x_n\}$  determined by the gradient  $\nabla f$  and the metric projection  $P_C$ :

$$x_{n+1} := P_C(x_n - \lambda \nabla f(x_n)), \quad \forall n \geq 0, \tag{19}$$

or more generally,

$$x_{n+1} := P_C(x_n - \lambda_n \nabla f(x_n)), \quad \forall n \geq 0, \tag{20}$$

where, in both (19) and (20), the initial guess  $x_0$  is taken from  $C$  arbitrarily and the parameters  $\lambda$  or  $\lambda_n$  are positive real numbers. The convergence of algorithms (19) and (20) depends on the behavior of the gradient  $\nabla f$ . As a matter of fact, it is known that if  $\nabla f$  is  $\alpha$ -strongly monotone and  $L$ -Lipschitz continuous, then, for  $0 < \lambda < 2\alpha/L^2$ , the operator  $P_C(I - \lambda \nabla f)$  is a contraction; hence, the sequence  $\{x_n\}$  defined by the GPA (19) converges in norm to the unique solution of CMP (18). More generally, if the sequence  $\{\lambda_n\}$  is chosen to satisfy the property

$$0 < \liminf_{n \rightarrow \infty} \lambda_n \leq \limsup_{n \rightarrow \infty} \lambda_n < \frac{2\alpha}{L^2}, \tag{21}$$

then the sequence  $\{x_n\}$  defined by the GPA (20) converges in norm to the unique minimizer of CMP (18). If the gradient  $\nabla f$  is only assumed to be Lipschitz continuous, then  $\{x_n\}$  can only be weakly convergent if  $H$  is infinite-dimensional (a counterexample is given in Section 5 of Xu [40]).

Since the Lipschitz continuity of the gradient  $\nabla f$  implies that it is actually  $(1/L)$ -inverse-strongly monotone (ism) [41], its complement can be an averaged mapping (i.e., it can be expressed as a proper convex combination of the identity mapping and a nonexpansive mapping). Consequently, the GPA can be rewritten as the composite of a projection and an averaged mapping, which is again an averaged mapping. This shows that averaged mappings play an important role in the GPA. Recently, Xu [40] used averaged mappings to study the convergence analysis of the GPA, which is hence an operator-oriented approach.

Motivated and inspired by the above facts, we in this paper introduce new implicit and explicit iterative algorithms for finding a common element of the set of solutions of the CMP (18) for a convex functional  $f : C \rightarrow \mathbf{R}$  with  $L$ -Lipschitz continuous gradient  $\nabla f$ , the set of solutions of a

finite family of GMEPs, and the set of solutions of a finite family of variational inclusions for maximal monotone and inverse-strong monotone mappings in a real Hilbert space. Under mild control conditions, we prove that the sequences generated by the proposed algorithms converge strongly to a common element of three sets, which is the unique solution of a variational inequality defined over the intersection of three sets. Our iterative algorithms are based on Korpelevich's extragradient method, hybrid steepest-descent method in [30], viscosity approximation method, and averaged mapping approach to the GPA in [40]. The results obtained in this paper improve and extend the corresponding results announced by many others.

## 2. Preliminaries

Throughout this paper, we assume that  $H$  is a real Hilbert space whose inner product and norm are denoted by  $\langle \cdot, \cdot \rangle$  and  $\| \cdot \|$ , respectively. Let  $C$  be a nonempty closed convex subset of  $H$ . We write  $x_n \rightharpoonup x$  to indicate that the sequence  $\{x_n\}$  converges weakly to  $x$  and  $x_n \rightarrow x$  to indicate that the sequence  $\{x_n\}$  converges strongly to  $x$ . Moreover, we use  $\omega_w(x_n)$  to denote the weak  $\omega$ -limit set of the sequence  $\{x_n\}$ ; that is,

$$\omega_w(x_n) := \{x \in H : x_{n_i} \rightharpoonup x \text{ for some subsequence } \{x_{n_i}\} \text{ of } \{x_n\}\}. \tag{22}$$

Recall that a mapping  $A : C \rightarrow H$  is called

- (i) monotone if

$$\langle Ax - Ay, x - y \rangle \geq 0, \quad \forall x, y \in C; \tag{23}$$

- (ii)  $\eta$ -strongly monotone if there exists a constant  $\eta > 0$  such that

$$\langle Ax - Ay, x - y \rangle \geq \eta \|x - y\|^2, \quad \forall x, y \in C; \tag{24}$$

- (iii)  $\alpha$ -inverse-strongly monotone if there exists a constant  $\alpha > 0$  such that

$$\langle Ax - Ay, x - y \rangle \geq \alpha \|Ax - Ay\|^2, \quad \forall x, y \in C. \tag{25}$$

It is obvious that if  $A$  is  $\alpha$ -inverse-strongly monotone, then  $A$  is monotone and  $(1/\alpha)$ -Lipschitz continuous.

The metric (or nearest point) projection from  $H$  onto  $C$  is the mapping  $P_C : H \rightarrow C$  which assigns to each point  $x \in H$  the unique point  $P_C x \in C$  satisfying the property

$$\|x - P_C x\| = \inf_{y \in C} \|x - y\| =: d(x, C). \tag{26}$$

Some important properties of projections are gathered in the following proposition.



**Proposition 2.** For given  $x \in H$  and  $z \in C$ ,

- (i)  $z = P_C x \Leftrightarrow \langle x - z, y - z \rangle \leq 0$ , for all  $y \in C$ ;
- (ii)  $z = P_C x \Leftrightarrow \|x - z\|^2 \leq \|x - y\|^2 - \|y - z\|^2$ , for all  $y \in C$ ;
- (iii)  $\langle P_C x - P_C y, x - y \rangle \geq \|P_C x - P_C y\|^2$ , for all  $y \in H$ .

Consequently,  $P_C$  is nonexpansive and monotone.

If  $A$  is an  $\alpha$ -inverse-strongly monotone mapping of  $C$  into  $H$ , then it is obvious that  $A$  is  $(1/\alpha)$ -Lipschitz continuous. We also have that, for all  $u, v \in C$  and  $\lambda > 0$ ,

$$\begin{aligned} \|(I - \lambda A)u - (I - \lambda A)v\|^2 &= \|(u - v) - \lambda(Au - Av)\|^2 \\ &= \|u - v\|^2 - 2\lambda \langle Au - Av, u - v \rangle \\ &\quad + \lambda^2 \|Au - Av\|^2 \\ &\leq \|u - v\|^2 + \lambda(\lambda - 2\alpha) \\ &\quad \times \|Au - Av\|^2. \end{aligned} \tag{27}$$

So, if  $\lambda \leq 2\alpha$ , then  $I - \lambda A$  is a nonexpansive mapping from  $C$  to  $H$ .

**Definition 3.** A mapping  $T : H \rightarrow H$  is said to be

- (a) nonexpansive [1] if

$$\|Tx - Ty\| \leq \|x - y\|, \quad \forall x, y \in H; \tag{28}$$

- (b) firmly nonexpansive if  $2T - I$  is nonexpansive or, equivalently, if  $T$  is 1-inverse-strongly monotone (1-ism),

$$\langle x - y, Tx - Ty \rangle \geq \|Tx - Ty\|^2, \quad \forall x, y \in H; \tag{29}$$

alternatively,  $T$  is firmly nonexpansive if and only if  $T$  can be expressed as

$$T = \frac{1}{2}(I + S), \tag{30}$$

where  $S : H \rightarrow H$  is nonexpansive; projections are firmly nonexpansive.

It can be easily seen that if  $T$  is nonexpansive, then  $I - T$  is monotone. It is also easy to see that a projection  $P_C$  is 1-ism. Inverse-strongly monotone (also referred to as co-coercive) operators have been applied widely in solving practical problems in various fields.

**Definition 4.** A mapping  $T : H \rightarrow H$  is said to be an averaged mapping if it can be written as the average of the identity  $I$  and a nonexpansive mapping; that is,

$$T \equiv (1 - \alpha)I + \alpha S, \tag{31}$$

where  $\alpha \in (0, 1)$  and  $S : H \rightarrow H$  is nonexpansive. More precisely, when the last equality holds, we say that  $T$  is  $\alpha$ -averaged. Thus, firmly nonexpansive mappings (in particular, projections) are  $(1/2)$ -averaged mappings.

**Proposition 5** (see [42]). Let  $T : H \rightarrow H$  be a given mapping.

- (i)  $T$  is nonexpansive if and only if the complement  $I - T$  is  $(1/2)$ -ism.
- (ii) If  $T$  is  $\nu$ -ism, then, for  $\gamma > 0$ ,  $\gamma T$  is  $(\nu/\gamma)$ -ism.
- (iii)  $T$  is averaged if and only if the complement  $I - T$  is  $\nu$ -ism for some  $\nu > 1/2$ . Indeed, for  $\alpha \in (0, 1)$ ,  $T$  is  $\alpha$ -averaged if and only if  $I - T$  is  $(1/2\alpha)$ -ism.

**Proposition 6** (see [42, 43]). Let  $S, T, V : H \rightarrow H$  be given operators.

- (i) If  $T = (1 - \alpha)S + \alpha V$  for some  $\alpha \in (0, 1)$  and if  $S$  is averaged and  $V$  is nonexpansive, then  $T$  is averaged.
- (ii)  $T$  is firmly nonexpansive if and only if the complement  $I - T$  is firmly nonexpansive.
- (iii) If  $T = (1 - \alpha)S + \alpha V$  for some  $\alpha \in (0, 1)$  and if  $S$  is firmly nonexpansive and  $V$  is nonexpansive, then  $T$  is averaged.
- (iv) The composite of finitely many averaged mappings is averaged. That is, if each of the mappings  $\{T_i\}_{i=1}^N$  is averaged, then so is the composite  $T_1 \cdots T_N$ . In particular, if  $T_1$  is  $\alpha_1$ -averaged and  $T_2$  is  $\alpha_2$ -averaged, where  $\alpha_1, \alpha_2 \in (0, 1)$ , then the composite  $T_1 T_2$  is  $\alpha$ -averaged, where  $\alpha = \alpha_1 + \alpha_2 - \alpha_1 \alpha_2$ .

- (v) If the mappings  $\{T_i\}_{i=1}^N$  are averaged and have a common fixed point, then

$$\bigcap_{i=1}^N \text{Fix}(T_i) = \text{Fix}(T_1 \cdots T_N). \tag{32}$$

The notation  $\text{Fix}(T)$  denotes the set of all fixed points of the mapping  $T$ ; that is,  $\text{Fix}(T) = \{x \in H : Tx = x\}$ .

We need some facts and tools in a real Hilbert space  $H$  which are listed as lemmas below.

**Lemma 7.** Let  $X$  be a real inner product space. Then the following inequality holds:

$$\|x + y\|^2 \leq \|x\|^2 + 2 \langle y, x + y \rangle, \quad \forall x, y \in X. \tag{33}$$

**Lemma 8.** Let  $A : C \rightarrow H$  be a monotone mapping. In the context of the variational inequality problem the characterization of the projection (see Proposition 2(i)) implies

$$u \in \text{VI}(C, A) \iff u = P_C(u - \lambda Au), \quad \text{for some } \lambda > 0. \tag{34}$$

**Lemma 9** (see [44, Demiclosedness principle]). Let  $C$  be a nonempty closed convex subset of a real Hilbert space  $H$ . Let  $T$  be a nonexpansive self-mapping on  $C$  with  $\text{Fix}(T) \neq \emptyset$ . Then  $I - T$  is demiclosed. That is, whenever  $\{x_n\}$  is a sequence in  $C$  weakly converging to some  $x \in C$  and the sequence  $\{(I - T)x_n\}$  strongly converges to some  $y$ , it follows that  $(I - T)x = y$ . Here  $I$  is the identity operator of  $H$ .

**Lemma 10** (see [45]). Let  $\{s_n\}$  be a sequence of nonnegative numbers satisfying the conditions

$$s_{n+1} \leq (1 - \alpha_n)s_n + \alpha_n\beta_n, \quad \forall n \geq 1, \quad (35)$$

where  $\{\alpha_n\}$  and  $\{\beta_n\}$  are sequences of real numbers such that

(i)  $\{\alpha_n\} \subset [0, 1]$  and  $\sum_{n=1}^{\infty} \alpha_n = \infty$  or, equivalently,

$$\prod_{n=1}^{\infty} (1 - \alpha_n) := \lim_{n \rightarrow \infty} \prod_{k=1}^n (1 - \alpha_k) = 0; \quad (36)$$

(ii)  $\limsup_{n \rightarrow \infty} \beta_n \leq 0$ , or  $\sum_{n=1}^{\infty} |\alpha_n\beta_n| < \infty$ .

Then  $\lim_{n \rightarrow \infty} s_n = 0$ .

**Lemma 11** (see [46]). Let  $\{x_n\}$  and  $\{z_n\}$  be bounded sequences in a Banach space  $X$  and  $\{\beta_n\}$  a sequence in  $[0, 1]$  with

$$0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1. \quad (37)$$

Suppose that  $x_{n+1} = (1 - \beta_n)z_n + \beta_nx_n$  for each  $n \geq 1$  and

$$\limsup_{n \rightarrow \infty} (\|z_{n+1} - z_n\| - \|x_{n+1} - x_n\|) \leq 0. \quad (38)$$

Then  $\lim_{n \rightarrow \infty} \|z_n - x_n\| = 0$ .

The following lemma can be easily proven and, therefore, we omit the proof.

**Lemma 12.** Let  $V : H \rightarrow H$  be an  $l$ -Lipschitzian mapping with constant  $l \geq 0$ , and let  $F : H \rightarrow H$  be a  $\kappa$ -Lipschitzian and  $\eta$ -strongly monotone operator with positive constants  $\kappa, \eta > 0$ . Then, for  $0 \leq \gamma l < \mu\eta$ ,

$$\langle (\mu F - \gamma V)x - (\mu F - \gamma V)y, x - y \rangle \geq (\mu\eta - \gamma l) \|x - y\|^2, \quad \forall x, y \in H. \quad (39)$$

That is,  $\mu F - \gamma V$  is strongly monotone with constant  $\mu\eta - \gamma l$ .

Let  $C$  be a nonempty closed convex subset of a real Hilbert space  $H$ . We introduce some notations. Let  $\lambda$  be a number in  $(0, 1]$  and let  $\mu > 0$ . Associating with a nonexpansive mapping  $T : C \rightarrow H$ , we define the mapping  $T^\lambda : C \rightarrow H$  by

$$T^\lambda x := Tx - \lambda\mu F(Tx), \quad \forall x \in C, \quad (40)$$

where  $F : H \rightarrow H$  is an operator such that, for some positive constants  $\kappa, \eta > 0$ ,  $F$  is  $\kappa$ -Lipschitzian and  $\eta$ -strongly monotone on  $H$ ; that is,  $F$  satisfies the conditions:

$$\|Fx - Fy\| \leq \kappa \|x - y\|, \quad \langle Fx - Fy, x - y \rangle \geq \eta \|x - y\|^2, \quad (41)$$

for all  $x, y \in H$ .

**Lemma 13** (see [45, Lemma 3.1]).  $T^\lambda$  is a contraction provided  $0 < \mu < 2\eta/\kappa^2$ ; that is,

$$\|T^\lambda x - T^\lambda y\| \leq (1 - \lambda\tau) \|x - y\|, \quad \forall x, y \in C, \quad (42)$$

where  $\tau = 1 - \sqrt{1 - \mu(2\eta - \mu\kappa^2)} \in (0, 1]$ .

Recall that a set-valued mapping  $R : D(R) \subset H \rightarrow 2^H$  is called monotone if for all  $x, y \in D(R)$ ,  $f \in R(x)$  and  $g \in R(y)$  imply

$$\langle f - g, x - y \rangle \geq 0. \quad (43)$$

A set-valued mapping  $R$  is called maximal monotone if  $R$  is monotone and  $(I + \lambda R)D(R) = H$  for each  $\lambda > 0$ , where  $I$  is the identity mapping of  $H$ . We denote by  $G(R)$  the graph of  $R$ . It is known that a monotone mapping  $R$  is maximal if and only if, for  $(x, f) \in H \times H$ ,  $\langle f - g, x - y \rangle \geq 0$  for every  $(y, g) \in G(R)$  implies  $f \in R(x)$ .

Let  $A : C \rightarrow H$  be a monotone,  $k$ -Lipschitz continuous mapping and let  $N_C v$  be the normal cone to  $C$  at  $v \in C$ ; that is,

$$N_C v = \{w \in H : \langle v - u, w \rangle \geq 0, \forall u \in C\}. \quad (44)$$

Define

$$Tv = \begin{cases} Av + N_C v, & \text{if } v \in C, \\ \emptyset, & \text{if } v \notin C. \end{cases} \quad (45)$$

Then,  $T$  is maximal monotone and  $0 \in Tv$  if and only if  $v \in VI(C, A)$ ; see [32].

Assume that  $R : D(R) \subset H \rightarrow 2^H$  is a maximal monotone mapping. Then, for  $\lambda > 0$ , associated with  $R$ , the resolvent operator  $J_{R,\lambda}$  can be defined as

$$J_{R,\lambda} x = (I + \lambda R)^{-1} x, \quad \forall x \in H. \quad (46)$$

In terms of Huang [33] (see also [34]), the following property holds for the resolvent operator  $J_{R,\lambda} : H \rightarrow \overline{D(R)}$ .

**Lemma 14.**  $J_{R,\lambda}$  is single-valued and firmly nonexpansive; that is,

$$\langle J_{R,\lambda} x - J_{R,\lambda} y, x - y \rangle \geq \|J_{R,\lambda} x - J_{R,\lambda} y\|^2, \quad \forall x, y \in H. \quad (47)$$

Consequently,  $J_{R,\lambda}$  is nonexpansive and monotone.

**Lemma 15** (see [39]). Let  $R$  be a maximal monotone mapping with  $D(R) = C$ . Then, for any given  $\lambda > 0$ ,  $u \in C$  is a solution of problem (14) if and only if  $u \in C$  satisfies

$$u = J_{R,\lambda} (u - \lambda Bu). \quad (48)$$

**Lemma 16** (see [34]). Let  $R$  be a maximal monotone mapping with  $D(R) = C$  and let  $B : C \rightarrow H$  be a strongly monotone, continuous, and single-valued mapping. Then, for each  $z \in H$ , the equation  $z \in (B + \lambda R)x$  has a unique solution  $x_\lambda$  for  $\lambda > 0$ .

**Lemma 17** (see [39]). Let  $R$  be a maximal monotone mapping with  $D(R) = C$  and let  $B : C \rightarrow H$  be a monotone, continuous, and single-valued mapping. Then  $(I + \lambda(R + B))C = H$  for each  $\lambda > 0$ . In this case,  $R + B$  is maximal monotone.

### 3. Implicit Iterative Algorithm and Its Convergence Criteria

We now state and prove the first main result of this paper.

**Theorem 18.** *Let  $C$  be a nonempty closed convex subset of a real Hilbert space  $H$ . Let  $f : C \rightarrow \mathbf{R}$  be a convex functional with  $L$ -Lipschitz continuous gradient  $\nabla f$ . Let  $M, N$  be two integers. Let  $\Theta_k$  be a bifunction from  $C \times C$  to  $\mathbf{R}$  satisfying (A1)–(A4) and let  $\varphi_k : C \rightarrow \mathbf{R} \cup \{+\infty\}$  be a proper lower semicontinuous and convex function, where  $k \in \{1, 2, \dots, M\}$ . Let  $R_i : C \rightarrow 2^H$  be a maximal monotone mapping and let  $A_k : H \rightarrow H$  and  $B_i : C \rightarrow H$  be  $\mu_k$ -inverse-strongly monotone and  $\eta_i$ -inverse-strongly monotone, respectively, where  $k \in \{1, 2, \dots, M\}$ ,  $i \in \{1, 2, \dots, N\}$ . Let  $F : H \rightarrow H$  be a  $\kappa$ -Lipschitzian and  $\eta$ -strongly monotone operator with positive constants  $\kappa, \eta > 0$ . Let  $V : H \rightarrow H$  be an  $l$ -Lipschitzian mapping with constant  $l \geq 0$ . Let  $0 < \mu < 2\eta/\kappa^2$  and  $0 \leq \gamma l < \tau$ , where  $\tau = 1 - \sqrt{1 - \mu(2\eta - \mu\kappa^2)}$ . Assume that  $\Omega := \bigcap_{k=1}^M \text{GMEP}(\Theta_k, \varphi_k, A_k) \cap \bigcap_{i=1}^N I(B_i, R_i) \cap \Gamma \neq \emptyset$  and that either (B1) or (B2) holds. Let  $\{x_n\}$  be a sequence generated by*

$$\begin{aligned} u_n &= T_{r_{M,n}}^{(\Theta_M, \varphi_M)} (I - r_{M,n} A_M) T_{r_{M-1,n}}^{(\Theta_{M-1}, \varphi_{M-1})} \\ &\quad \times (I - r_{M-1,n} A_{M-1}) \cdots T_{r_{1,n}}^{(\Theta_1, \varphi_1)} (I - r_{1,n} A_1) x_n, \\ v_n &= J_{R_N, \lambda_{N,n}} (I - \lambda_{N,n} B_N) J_{R_{N-1}, \lambda_{N-1,n}} \\ &\quad \times (I - \lambda_{N-1,n} B_{N-1}) \cdots J_{R_1, \lambda_{1,n}} (I - \lambda_{1,n} B_1) u_n, \\ x_n &= s_n \gamma V x_n + (I - s_n \mu F) T_n v_n, \\ &\quad \forall n \geq 1, \end{aligned} \tag{49}$$

where  $P_C(I - \lambda_n \nabla f) = s_n I + (1 - s_n) T_n$  (here  $T_n$  is nonexpansive and  $s_n = (2 - \lambda_n L)/4 \in (0, 1/2)$  for each  $\lambda_n \in (0, 2/L)$ ). Assume that the following conditions hold:

- (i)  $s_n \in (0, 1/2)$  for each  $\lambda_n \in (0, 2/L)$ ,  $\lim_{n \rightarrow \infty} s_n = 0$  ( $\Leftrightarrow \lim_{n \rightarrow \infty} \lambda_n = 2/L$ );
- (ii)  $\{\lambda_{i,n}\} \subset [a_i, b_i] \subset (0, 2\eta_i)$ , for all  $i \in \{1, 2, \dots, N\}$ ;
- (iii)  $\{r_{k,n}\} \subset [e_k, f_k] \subset (0, 2\mu_k)$ , for all  $k \in \{1, 2, \dots, M\}$ .

Then  $\{x_n\}$  converges strongly as  $\lambda_n \rightarrow 2/L$  ( $\Leftrightarrow s_n \rightarrow 0$ ) to a point  $q \in \Omega$ , which is a unique solution of the VIP:

$$\langle (\mu F - \gamma V) q, p - q \rangle \geq 0, \quad \forall p \in \Omega. \tag{50}$$

Equivalently,  $q = P_\Omega(I - \mu F + \gamma V)q$ .

*Proof.* First of all, let us show that the sequence  $\{x_n\}$  is well defined. Indeed, since  $\nabla f$  is  $L$ -Lipschitzian, it follows that  $\nabla f$  is  $1/L$ -ism; see [41]. By Proposition 5(ii) we know that, for  $\lambda > 0$ ,  $\lambda \nabla f$  is  $(1/\lambda L)$ -ism. So by Proposition 5(iii) we deduce that  $I - \lambda \nabla f$  is  $(\lambda L/2)$ -averaged. Now since the projection  $P_C$  is  $(1/2)$ -averaged, it is easy to see from Proposition 6(iv) that the composite  $P_C(I - \lambda \nabla f)$  is  $(2 + \lambda L)/4$ -averaged for  $\lambda \in (0, 2/L)$ . Hence, we obtain that, for each  $n \geq 1$ ,  $P_C(I - \lambda_n \nabla f)$

is  $((2 + \lambda_n L)/4)$ -averaged for each  $\lambda_n \in (0, 2/L)$ . Therefore, we can write

$$P_C(I - \lambda_n \nabla f) = \frac{2 - \lambda_n L}{4} I + \frac{2 + \lambda_n L}{4} T_n = s_n I + (1 - s_n) T_n, \tag{51}$$

where  $T_n$  is nonexpansive and  $s_n := s_n(\lambda_n) = (2 - \lambda_n L)/4 \in (0, 1/2)$  for each  $\lambda_n \in (0, 2/L)$ . It is clear that

$$\lambda_n \rightarrow \frac{2}{L} \iff s_n \rightarrow 0. \tag{52}$$

Put

$$\begin{aligned} \Delta_n^k &= T_{r_{k,n}}^{(\Theta_k, \varphi_k)} (I - r_{k,n} A_k) T_{r_{k-1,n}}^{(\Theta_{k-1}, \varphi_{k-1})} \\ &\quad \times (I - r_{k-1,n} A_{k-1}) \cdots T_{r_{1,n}}^{(\Theta_1, \varphi_1)} (I - r_{1,n} A_1) x_n, \end{aligned} \tag{53}$$

for all  $k \in \{1, 2, \dots, M\}$  and  $n \geq 1$ ,

$$\begin{aligned} \Lambda_n^i &= J_{R_i, \lambda_{i,n}} (I - \lambda_{i,n} B_i) J_{R_{i-1}, \lambda_{i-1,n}} \\ &\quad \times (I - \lambda_{i-1,n} B_{i-1}) \cdots J_{R_1, \lambda_{1,n}} (I - \lambda_{1,n} B_1), \end{aligned} \tag{54}$$

for all  $i \in \{1, 2, \dots, N\}$  and  $n \geq 1$ , and  $\Delta_n^0 = \Lambda_n^0 = I$ , where  $I$  is the identity mapping on  $H$ . Then we have that  $u_n = \Delta_n^M x_n$  and  $v_n = \Lambda_n^N u_n$ .

Consider the following mapping  $G_n$  on  $H$  defined by

$$G_n x = s_n \gamma V x + (I - s_n \mu F) T_n \Lambda_n^N \Delta_n^M x, \quad \forall x \in H, \quad n \geq 1, \tag{55}$$

where  $s_n = (2 - \lambda_n L)/4 \in (0, 1/2)$  for each  $\lambda_n \in (0, 2/L)$ . By Proposition 1(ii) and Lemma 13 we obtain from (27) that for all  $x, y \in H$

$$\begin{aligned} &\|G_n x - G_n y\| \\ &\leq s_n \gamma \|Vx - Vy\| + \|(I - s_n \mu F) T_n \Lambda_n^N \Delta_n^M x \\ &\quad - (I - s_n \mu F) T_n \Lambda_n^N \Delta_n^M y\| \\ &\leq s_n \gamma l \|x - y\| + (1 - s_n \tau) \|\Lambda_n^N \Delta_n^M x - \Lambda_n^N \Delta_n^M y\| \\ &\leq s_n \gamma l \|x - y\| + (1 - s_n \tau) \|(I - \lambda_{N,n} B_N) \Lambda_n^{N-1} \Delta_n^M x \\ &\quad - (I - \lambda_{N,n} B_N) \Lambda_n^{N-1} \Delta_n^M y\| \\ &\leq s_n \gamma l \|x - y\| + (1 - s_n \tau) \|\Lambda_n^{N-1} \Delta_n^M x - \Lambda_n^{N-1} \Delta_n^M y\| \\ &\vdots \\ &\leq s_n \gamma l \|x - y\| + (1 - s_n \tau) \|\Lambda_n^0 \Delta_n^M x - \Lambda_n^0 \Delta_n^M y\| \end{aligned}$$

$$\begin{aligned}
&= s_n \gamma l \|x - y\| + (1 - s_n \tau) \|\Delta_n^M x - \Delta_n^M y\| \\
&\leq s_n \gamma l \|x - y\| + (1 - s_n \tau) \|(I - r_{M,n} A_M) \Delta_n^{M-1} x \\
&\quad - (I - r_{M,n} A_M) \Delta_n^{M-1} y\| \\
&\leq s_n \gamma l \|x - y\| + (1 - s_n \tau) \|\Delta_n^{M-1} x - \Delta_n^{M-1} y\| \\
&\vdots \\
&\leq s_n \gamma l \|x - y\| + (1 - s_n \tau) \|\Delta_n^0 x - \Delta_n^0 y\| \\
&= s_n \gamma l \|x - y\| + (1 - s_n \tau) \|x - y\| \\
&= (1 - s_n (\tau - \gamma l)) \|x - y\|.
\end{aligned} \tag{56}$$

Since  $0 < 1 - s_n (\tau - \gamma l) < 1$ ,  $G_n : H \rightarrow H$  is a contraction. Therefore, by the Banach contraction principle,  $G_n$  has a unique fixed point  $x_n \in H$ , which uniquely solves the fixed point equation:

$$x_n = s_n \gamma V x_n + (I - s_n \mu F) T_n \Lambda_n^N \Delta_n^M x_n. \tag{57}$$

This shows that the sequence  $\{x_n\}$  is defined well.

Note that  $0 \leq \gamma l < \tau$  and  $\mu \eta \geq \tau \Leftrightarrow \kappa \geq \eta$ . Hence, by Lemma 12 we know that

$$\begin{aligned}
\langle (\mu F - \gamma V) x - (\mu F - \gamma V) y, x - y \rangle &\geq (\mu \eta - \gamma l) \|x - y\|^2, \\
\forall x, y \in H.
\end{aligned} \tag{58}$$

That is,  $\mu F - \gamma V$  is strongly monotone for  $0 \leq \gamma l < \tau \leq \mu \eta$ . Moreover, it is clear that  $\mu F - \gamma V$  is Lipschitz continuous. So the VIP (50) has only one solution. Below we use  $q \in \Omega$  to denote the unique solution of the VIP (50).

Now, let us show that  $\{x_n\}$  is bounded. In fact, take  $p \in \Omega$  arbitrarily. Then from (27) and Proposition 1(ii) we have

$$\begin{aligned}
\|u_n - p\| &= \left\| T_{r_{M,n}}^{(\Theta_M, \varphi_M)} (I - r_{M,n} A_M) \Delta_n^{M-1} x_n \right. \\
&\quad \left. - T_{r_{M,n}}^{(\Theta_M, \varphi_M)} (I - r_{M,n} A_M) \Delta_n^{M-1} p \right\| \\
&\leq \|(I - r_{M,n} A_M) \Delta_n^{M-1} x_n \\
&\quad - (I - r_{M,n} A_M) \Delta_n^{M-1} p\| \\
&\leq \|\Delta_n^{M-1} x_n - \Delta_n^{M-1} p\| \\
&\vdots \\
&\leq \|\Delta_n^0 x_n - \Delta_n^0 p\| \\
&= \|x_n - p\|.
\end{aligned} \tag{59}$$

Similarly, we have

$$\begin{aligned}
\|v_n - p\| &= \left\| J_{R_N, \lambda_{N,n}} (I - \lambda_{N,n} B_N) \Lambda_n^{N-1} u_n \right. \\
&\quad \left. - J_{R_N, \lambda_{N,n}} (I - \lambda_{N,n} B_N) \Lambda_n^{N-1} p \right\| \\
&\leq \|(I - \lambda_{N,n} B_N) \Lambda_n^{N-1} u_n - (I - \lambda_{N,n} B_N) \Lambda_n^{N-1} p\| \\
&\leq \|\Lambda_n^{N-1} u_n - \Lambda_n^{N-1} p\| \\
&\vdots \\
&\leq \|\Lambda_n^0 u_n - \Lambda_n^0 p\| \\
&= \|u_n - p\|.
\end{aligned} \tag{60}$$

Combining (59) and (60), we have

$$\|v_n - p\| \leq \|x_n - p\|. \tag{61}$$

Since

$$p = P_C (I - \lambda_n \nabla f) p = s_n p + (1 - s_n) T_n p, \quad \forall \lambda_n \in \left(0, \frac{2}{L}\right), \tag{62}$$

where  $s_n := s_n(\lambda_n) = (2 - \lambda_n L)/4 \in (0, 1/2)$ , it is clear that  $T_n p = p$  for each  $\lambda_n \in (0, 2/L)$ . Thus, utilizing Lemma 13 and the nonexpansivity of  $T_n$ , we obtain from (61) that

$$\begin{aligned}
\|x_n - p\| &= \|s_n (\gamma V x_n - \mu F p) + (I - s_n \mu F) T_n v_n \\
&\quad - (I - s_n \mu F) T_n p\| \\
&\leq \|(I - s_n \mu F) T_n v_n - (I - s_n \mu F) T_n p\| \\
&\quad + s_n \|\gamma V x_n - \mu F p\| \\
&\leq (1 - s_n \tau) \|v_n - p\| \\
&\quad + s_n (\gamma \|V x_n - V p\| + \|\gamma V p - \mu F p\|) \\
&\leq (1 - s_n \tau) \|x_n - p\| \\
&\quad + s_n (\gamma l \|x_n - p\| + \|\gamma V p - \mu F p\|) \\
&= (1 - s_n (\tau - \gamma l)) \|x_n - p\| + s_n \|\gamma V p - \mu F p\|.
\end{aligned} \tag{63}$$

This implies that  $\|x_n - p\| \leq \|\gamma V p - \mu F p\| / (\tau - \gamma l)$ . Hence,  $\{x_n\}$  is bounded. So, according to (59) and (61) we know that  $\{u_n\}$ ,  $\{v_n\}$ ,  $\{T_n v_n\}$ ,  $\{V x_n\}$ , and  $\{F T_n v_n\}$  are bounded.

Next let us show that  $\|u_n - x_n\| \rightarrow 0$ ,  $\|v_n - u_n\| \rightarrow 0$ , and  $\|x_n - T_n x_n\| \rightarrow 0$  as  $n \rightarrow \infty$ .



Indeed, from (27) it follows that for all  $i \in \{1, 2, \dots, N\}$  and  $k \in \{1, 2, \dots, M\}$

$$\begin{aligned}
 \|v_n - p\|^2 &= \|\Lambda_n^N u_n - p\|^2 \\
 &\leq \|\Lambda_n^i u_n - p\|^2 \\
 &= \|J_{R_i, \lambda_{i,n}}(I - \lambda_{i,n} B_i) \Lambda_n^{i-1} u_n \\
 &\quad - J_{R_i, \lambda_{i,n}}(I - \lambda_{i,n} B_i) p\|^2 \\
 &\leq \|(I - \lambda_{i,n} B_i) \Lambda_n^{i-1} u_n - (I - \lambda_{i,n} B_i) p\|^2 \\
 &\leq \|\Lambda_n^{i-1} u_n - p\|^2 + \lambda_{i,n} (\lambda_{i,n} - 2\eta_i) \\
 &\quad \times \|B_i \Lambda_n^{i-1} u_n - B_i p\|^2 \\
 &\leq \|u_n - p\|^2 + \lambda_{i,n} (\lambda_{i,n} - 2\eta_i) \\
 &\quad \times \|B_i \Lambda_n^{i-1} u_n - B_i p\|^2 \\
 &\leq \|x_n - p\|^2 + \lambda_{i,n} (\lambda_{i,n} - 2\eta_i) \\
 &\quad \times \|B_i \Lambda_n^{i-1} u_n - B_i p\|^2, \\
 \|u_n - p\|^2 &= \|\Delta_n^M x_n - p\|^2 \\
 &\leq \|\Delta_n^k x_n - p\|^2 \\
 &= \|T_{r_{k,n}}^{(\Theta_k, \varphi_k)}(I - r_{k,n} A_k) \Delta_n^{k-1} x_n \\
 &\quad - T_{r_{k,n}}^{(\Theta_k, \varphi_k)}(I - r_{k,n} A_k) p\|^2 \\
 &\leq \|(I - r_{k,n} A_k) \Delta_n^{k-1} x_n - (I - r_{k,n} A_k) p\|^2 \\
 &\leq \|\Delta_n^{k-1} x_n - p\|^2 + r_{k,n} (r_{k,n} - 2\mu_k) \\
 &\quad \times \|A_k \Delta_n^{k-1} x_n - A_k p\|^2 \\
 &\leq \|x_n - p\|^2 + r_{k,n} (r_{k,n} - 2\mu_k) \\
 &\quad \times \|A_k \Delta_n^{k-1} x_n - A_k p\|^2.
 \end{aligned} \tag{64}$$

Thus, utilizing Lemma 7, from (49) and (64) we have

$$\begin{aligned}
 \|x_n - p\|^2 &= \|s_n (\gamma V x_n - \mu F p) + (I - s_n \mu F) T_n v_n \\
 &\quad - (I - s_n \mu F) T_n p\|^2 \\
 &\leq \|(I - s_n \mu F) T_n v_n - (I - s_n \mu F) T_n p\|^2 \\
 &\quad + 2s_n \langle \gamma V x_n - \mu F p, x_n - p \rangle \\
 &\leq (1 - s_n \tau)^2 \|v_n - p\|^2 + 2s_n \langle \gamma V x_n - \mu F p, x_n - p \rangle
 \end{aligned}$$

$$\begin{aligned}
 &= (1 - s_n \tau)^2 \|v_n - p\|^2 + 2s_n \gamma \langle V x_n - V p, x_n - p \rangle \\
 &\quad + 2s_n \langle \gamma V p - \mu F p, x_n - p \rangle \\
 &\leq (1 - s_n \tau)^2 \|v_n - p\|^2 + 2s_n \gamma l \|x_n - p\|^2 \\
 &\quad + 2s_n \|\gamma V p - \mu F p\| \|x_n - p\| \\
 &\leq (1 - s_n \tau)^2 [\|u_n - p\|^2 + \lambda_{i,n} (\lambda_{i,n} - 2\eta_i) \\
 &\quad \times \|B_i \Lambda_n^{i-1} u_n - B_i p\|^2] \\
 &\quad + 2s_n \gamma l \|x_n - p\|^2 + 2s_n \|\gamma V p - \mu F p\| \|x_n - p\| \\
 &\leq (1 - s_n \tau)^2 [\|x_n - p\|^2 + r_{k,n} (r_{k,n} - 2\mu_k) \\
 &\quad \times \|A_k \Delta_n^{k-1} x_n - A_k p\|^2 \\
 &\quad + \lambda_{i,n} (\lambda_{i,n} - 2\eta_i) \|B_i \Lambda_n^{i-1} u_n - B_i p\|^2] \\
 &\quad + 2s_n \gamma l \|x_n - p\|^2 + 2s_n \|\gamma V p - \mu F p\| \|x_n - p\| \\
 &= [1 - 2s_n (\tau - \gamma l) + s_n^2 \tau^2] \|x_n - p\|^2 \\
 &\quad - (1 - s_n \tau)^2 [r_{k,n} (2\mu_k - r_{k,n}) \|A_k \Delta_n^{k-1} x_n - A_k p\|^2 \\
 &\quad + \lambda_{i,n} (2\eta_i - \lambda_{i,n}) \|B_i \Lambda_n^{i-1} u_n - B_i p\|^2] \\
 &\quad + 2s_n \|\gamma V p - \mu F p\| \|x_n - p\| \\
 &\leq \|x_n - p\|^2 + s_n^2 \tau^2 \|x_n - p\|^2 - (1 - s_n \tau)^2 \\
 &\quad \times [r_{k,n} (2\mu_k - r_{k,n}) \|A_k \Delta_n^{k-1} x_n - A_k p\|^2 \\
 &\quad + \lambda_{i,n} (2\eta_i - \lambda_{i,n}) \|B_i \Lambda_n^{i-1} u_n - B_i p\|^2] \\
 &\quad + 2s_n \|\gamma V p - \mu F p\| \|x_n - p\|,
 \end{aligned} \tag{65}$$

which implies that

$$\begin{aligned}
 &(1 - s_n \tau)^2 [r_{k,n} (2\mu_k - r_{k,n}) \|A_k \Delta_n^{k-1} x_n - A_k p\|^2 \\
 &\quad + \lambda_{i,n} (2\eta_i - \lambda_{i,n}) \|B_i \Lambda_n^{i-1} u_n - B_i p\|^2] \\
 &\leq s_n^2 \tau^2 \|x_n - p\|^2 + 2s_n \|\gamma V p - \mu F p\| \|x_n - p\|.
 \end{aligned} \tag{66}$$

Since  $\{\lambda_{i,n}\} \subset [a_i, b_i] \subset (0, 2\eta_i)$  and  $\{r_{k,n}\} \subset [e_k, f_k] \subset (0, 2\mu_k)$  for all  $i \in \{1, 2, \dots, N\}$  and  $k \in \{1, 2, \dots, M\}$ , from  $s_n \rightarrow 0$  we conclude immediately that

$$\begin{aligned}
 \lim_{n \rightarrow \infty} \|A_k \Delta_n^{k-1} x_n - A_k p\| &= 0, \\
 \lim_{n \rightarrow \infty} \|B_i \Lambda_n^{i-1} u_n - B_i p\| &= 0,
 \end{aligned} \tag{67}$$

for all  $k \in \{1, 2, \dots, M\}$  and  $i \in \{1, 2, \dots, N\}$ .

Furthermore, by Proposition 1(ii) we obtain that for each  $k \in \{1, 2, \dots, M\}$

$$\begin{aligned} & \|\Delta_n^k x_n - p\|^2 \\ &= \left\| T_{r_{k,n}}^{(\Theta_k, \varphi_k)} (I - r_{k,n} A_k) \Delta_n^{k-1} x_n \right. \\ & \quad \left. - T_{r_{k,n}}^{(\Theta_k, \varphi_k)} (I - r_{k,n} A_k) p \right\|^2 \\ &\leq \langle (I - r_{k,n} A_k) \Delta_n^{k-1} x_n - (I - r_{k,n} A_k) p, \Delta_n^k x_n - p \rangle \\ &= \frac{1}{2} \left( \|(I - r_{k,n} A_k) \Delta_n^{k-1} x_n - (I - r_{k,n} A_k) p\|^2 \right. \\ & \quad + \|\Delta_n^k x_n - p\|^2 - \|(I - r_{k,n} A_k) \Delta_n^{k-1} x_n \\ & \quad \quad - (I - r_{k,n} A_k) p \\ & \quad \quad \left. - (\Delta_n^k x_n - p)\|^2 \right) \\ &\leq \frac{1}{2} \left( \|\Delta_n^{k-1} x_n - p\|^2 + \|\Delta_n^k x_n - p\|^2 \right. \\ & \quad \left. - \|\Delta_n^{k-1} x_n - \Delta_n^k x_n - r_{k,n} (A_k \Delta_n^{k-1} x_n - A_k p)\|^2 \right), \end{aligned} \tag{68}$$

which implies that

$$\begin{aligned} & \|\Delta_n^k x_n - p\|^2 \\ &\leq \|\Delta_n^{k-1} x_n - p\|^2 \\ & \quad - \|\Delta_n^{k-1} x_n - \Delta_n^k x_n - r_{k,n} (A_k \Delta_n^{k-1} x_n - A_k p)\|^2 \\ &= \|\Delta_n^{k-1} x_n - p\|^2 - \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\|^2 \\ & \quad - r_{k,n}^2 \|A_k \Delta_n^{k-1} x_n - A_k p\|^2 \\ & \quad + 2r_{k,n} \langle \Delta_n^{k-1} x_n - \Delta_n^k x_n, A_k \Delta_n^{k-1} x_n - A_k p \rangle \\ &\leq \|\Delta_n^{k-1} x_n - p\|^2 - \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\|^2 \\ & \quad + 2r_{k,n} \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\| \|A_k \Delta_n^{k-1} x_n - A_k p\| \\ &\leq \|x_n - p\|^2 - \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\|^2 \\ & \quad + 2r_{k,n} \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\| \\ & \quad \times \|A_k \Delta_n^{k-1} x_n - A_k p\|. \end{aligned} \tag{69}$$

Also, by Lemma 14, we obtain that for each  $i \in \{1, 2, \dots, N\}$

$$\begin{aligned} & \|\Lambda_n^i u_n - p\|^2 \\ &= \|J_{R_i, \lambda_{i,n}} (I - \lambda_{i,n} B_i) \Lambda_n^{i-1} u_n - J_{R_i, \lambda_{i,n}} (I - \lambda_{i,n} B_i) p\|^2 \\ &\leq \langle (I - \lambda_{i,n} B_i) \Lambda_n^{i-1} u_n - (I - \lambda_{i,n} B_i) p, \Lambda_n^i u_n - p \rangle \end{aligned}$$

$$\begin{aligned} &= \frac{1}{2} \left( \|(I - \lambda_{i,n} B_i) \Lambda_n^{i-1} u_n - (I - \lambda_{i,n} B_i) p\|^2 \right. \\ & \quad + \|\Lambda_n^i u_n - p\|^2 - \|(I - \lambda_{i,n} B_i) \Lambda_n^{i-1} u_n \\ & \quad \quad - (I - \lambda_{i,n} B_i) p \\ & \quad \quad \left. - (\Lambda_n^i u_n - p)\|^2 \right) \\ &\leq \frac{1}{2} \left( \|\Lambda_n^{i-1} u_n - p\|^2 + \|\Lambda_n^i u_n - p\|^2 \right. \\ & \quad \left. - \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n - \lambda_{i,n} (B_i \Lambda_n^{i-1} u_n - B_i p)\|^2 \right) \\ &\leq \frac{1}{2} \left( \|u_n - p\|^2 + \|\Lambda_n^i u_n - p\|^2 \right. \\ & \quad \left. - \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n - \lambda_{i,n} (B_i \Lambda_n^{i-1} u_n - B_i p)\|^2 \right), \end{aligned} \tag{70}$$

which implies

$$\begin{aligned} \|\Lambda_n^i u_n - p\|^2 &\leq \|u_n - p\|^2 - \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n \\ & \quad - \lambda_{i,n} (B_i \Lambda_n^{i-1} u_n - B_i p)\|^2 \\ &= \|u_n - p\|^2 - \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\|^2 \\ & \quad - \lambda_{i,n}^2 \|B_i \Lambda_n^{i-1} u_n - B_i p\|^2 \\ & \quad + 2\lambda_{i,n} \langle \Lambda_n^{i-1} u_n - \Lambda_n^i u_n, \\ & \quad \quad B_i \Lambda_n^{i-1} u_n - B_i p \rangle \\ &\leq \|u_n - p\|^2 - \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\|^2 \\ & \quad + 2\lambda_{i,n} \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\| \\ & \quad \times \|B_i \Lambda_n^{i-1} u_n - B_i p\|. \end{aligned} \tag{71}$$

Thus, utilizing Lemma 7, from (49), (69), and (71) we have

$$\begin{aligned} & \|x_n - p\|^2 \\ &= \|s_n (\gamma V x_n - \mu F p) + (I - s_n \mu F) T_n v_n \\ & \quad - (I - s_n \mu F) T_n p\|^2 \\ &\leq \|(I - s_n \mu F) T_n v_n - (I - s_n \mu F) T_n p\|^2 \\ & \quad + 2s_n \langle \gamma V x_n - \mu F p, x_n - p \rangle \\ &\leq (1 - s_n \tau)^2 \|v_n - p\|^2 + 2s_n \gamma l \|x_n - p\|^2 \\ & \quad + 2s_n \|\gamma V p - \mu F p\| \|x_n - p\| \end{aligned}$$

$$\begin{aligned}
 &= (1 - s_n \tau)^2 \|\Lambda_n^N u_n - p\|^2 + 2s_n \gamma l \|x_n - p\|^2 \\
 &\quad + 2s_n \|\gamma Vp - \mu Fp\| \|x_n - p\| \\
 &\leq (1 - s_n \tau)^2 \|\Lambda_n^i u_n - p\|^2 + 2s_n \gamma l \|x_n - p\|^2 \\
 &\quad + 2s_n \|\gamma Vp - \mu Fp\| \|x_n - p\| \\
 &\leq (1 - s_n \tau)^2 \left[ \|u_n - p\|^2 - \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\|^2 \right. \\
 &\quad \left. + 2\lambda_{i,n} \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\| \right. \\
 &\quad \left. \times \|B_i \Lambda_n^{i-1} u_n - B_i p\| \right] \\
 &\quad + 2s_n \gamma l \|x_n - p\|^2 + 2s_n \|\gamma Vp - \mu Fp\| \|x_n - p\| \\
 &= (1 - s_n \tau)^2 \left[ \|\Delta_n^M x_n - p\|^2 - \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\|^2 \right. \\
 &\quad \left. + 2\lambda_{i,n} \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\| \right. \\
 &\quad \left. \times \|B_i \Lambda_n^{i-1} u_n - B_i p\| \right] \\
 &\quad + 2s_n \gamma l \|x_n - p\|^2 + 2s_n \|\gamma Vp - \mu Fp\| \|x_n - p\| \\
 &\leq (1 - s_n \tau)^2 \left[ \|\Delta_n^k x_n - p\|^2 - \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\|^2 \right. \\
 &\quad \left. + 2\lambda_{i,n} \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\| \right. \\
 &\quad \left. \times \|B_i \Lambda_n^{i-1} u_n - B_i p\| \right] \\
 &\quad + 2s_n \gamma l \|x_n - p\|^2 + 2s_n \|\gamma Vp - \mu Fp\| \|x_n - p\| \\
 &\leq (1 - s_n \tau)^2 \left[ \|x_n - p\|^2 - \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\|^2 \right. \\
 &\quad \left. + 2r_{k,n} \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\| \right. \\
 &\quad \left. \times \|A_k \Delta_n^{k-1} x_n - A_k p\| \right. \\
 &\quad \left. - \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\|^2 \right. \\
 &\quad \left. + 2\lambda_{i,n} \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\| \right. \\
 &\quad \left. \times \|B_i \Lambda_n^{i-1} u_n - B_i p\| \right] \\
 &\quad + 2s_n \gamma l \|x_n - p\|^2 \\
 &\quad + 2s_n \|\gamma Vp - \mu Fp\| \|x_n - p\| \\
 &\leq (1 - 2s_n (\tau - \gamma l) + s_n^2 \tau^2) \|x_n - p\|^2 \\
 &\quad - (1 - s_n \tau)^2 \left( \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\|^2 \right. \\
 &\quad \left. + \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\|^2 \right) \\
 &\quad + 2r_{k,n} \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\| \|A_k \Delta_n^{k-1} x_n - A_k p\| \\
 &\quad + 2\lambda_{i,n} \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\| \|B_i \Lambda_n^{i-1} u_n - B_i p\| \\
 &\quad + 2s_n \|\gamma Vp - \mu Fp\| \|x_n - p\|. \tag{72}
 \end{aligned}$$

It immediately follows that

$$\begin{aligned}
 &(1 - s_n \tau)^2 \left( \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\|^2 + \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\|^2 \right) \\
 &\leq s_n^2 \tau^2 \|x_n - p\|^2 + 2r_{k,n} \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\| \\
 &\quad \times \|A_k \Delta_n^{k-1} x_n - A_k p\| + 2\lambda_{i,n} \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\| \\
 &\quad \times \|B_i \Lambda_n^{i-1} u_n - B_i p\| + 2s_n \|\gamma Vp - \mu Fp\| \|x_n - p\|. \tag{73}
 \end{aligned}$$

Since  $\{\lambda_{i,n}\} \subset [a_i, b_i] \subset (0, 2\eta_i)$  and  $\{r_{k,n}\} \subset [e_k, f_k] \subset (0, 2\mu_k)$  for all  $i \in \{1, 2, \dots, N\}$  and  $k \in \{1, 2, \dots, M\}$ , from (67) and  $s_n \rightarrow 0$  we deduce that

$$\lim_{n \rightarrow \infty} \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\| = 0, \quad \lim_{n \rightarrow \infty} \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\| = 0, \tag{74}$$

for all  $k \in \{1, 2, \dots, M\}$  and  $i \in \{1, 2, \dots, N\}$ . Hence, we get

$$\begin{aligned}
 \|x_n - u_n\| &= \|\Delta_n^0 x_n - \Delta_n^M x_n\| \\
 &\leq \|\Delta_n^0 x_n - \Delta_n^1 x_n\| + \|\Delta_n^1 x_n - \Delta_n^2 x_n\| \\
 &\quad + \dots + \|\Delta_n^{M-1} x_n - \Delta_n^M x_n\| \rightarrow 0 \\
 &\quad \text{as } n \rightarrow \infty, \tag{75}
 \end{aligned}$$

$$\begin{aligned}
 \|u_n - v_n\| &= \|\Lambda_n^0 u_n - \Lambda_n^N u_n\| \\
 &\leq \|\Lambda_n^0 u_n - \Lambda_n^1 u_n\| + \|\Lambda_n^1 u_n - \Lambda_n^2 u_n\| \\
 &\quad + \dots + \|\Lambda_n^{N-1} u_n - \Lambda_n^N u_n\| \rightarrow 0 \\
 &\quad \text{as } n \rightarrow \infty. \tag{76}
 \end{aligned}$$

So, taking into account that  $\|x_n - v_n\| \leq \|x_n - u_n\| + \|u_n - v_n\|$ , we have

$$\lim_{n \rightarrow \infty} \|x_n - v_n\| = 0. \tag{77}$$

Thus, from (77) and  $s_n \rightarrow 0$  we have

$$\begin{aligned}
 \|v_n - T_n v_n\| &\leq \|v_n - x_n\| + \|x_n - T_n v_n\| \\
 &= \|v_n - x_n\| + s_n \|\gamma Vx_n - \mu FT_n v_n\| \rightarrow 0 \\
 &\quad \text{as } n \rightarrow \infty. \tag{78}
 \end{aligned}$$

Now we show that  $\|x_n - T_n x_n\| \rightarrow 0$  as  $n \rightarrow \infty$ . In fact, from the nonexpansivity of  $T_n$ , we have

$$\begin{aligned} \|x_n - T_n x_n\| &\leq \|x_n - v_n\| + \|v_n - T_n v_n\| \\ &\quad + \|T_n v_n - T_n x_n\| \\ &\leq 2 \|x_n - v_n\| + \|v_n - T_n v_n\|. \end{aligned} \quad (79)$$

By (77) and (78), we get

$$\lim_{n \rightarrow \infty} \|x_n - T_n x_n\| = 0. \quad (80)$$

From (78) it is easy to see that

$$\lim_{n \rightarrow \infty} \|x_n - T_n v_n\| = 0. \quad (81)$$

Observe that

$$\begin{aligned} &\|P_C(I - \lambda_n \nabla f) v_n - v_n\| \\ &= \|s_n v_n + (1 - s_n) T_n v_n - v_n\| \\ &= (1 - s_n) \|T_n v_n - v_n\| \\ &\leq \|T_n v_n - v_n\|, \end{aligned} \quad (82)$$

where  $s_n = (2 - \lambda_n L)/4 \in (0, 1/2)$  for each  $\lambda_n \in (0, 2/L)$ . Hence, we have

$$\begin{aligned} &\|P_C\left(I - \frac{2}{L} \nabla f\right) v_n - v_n\| \\ &\leq \|P_C\left(I - \frac{2}{L} \nabla f\right) v_n - P_C(I - \lambda_n \nabla f) v_n\| \\ &\quad + \|P_C(I - \lambda_n \nabla f) v_n - v_n\| \\ &\leq \left\| \left(I - \frac{2}{L} \nabla f\right) v_n - (I - \lambda_n \nabla f) v_n \right\| \\ &\quad + \|P_C(I - \lambda_n \nabla f) v_n - v_n\| \\ &\leq \left(\frac{2}{L} - \lambda_n\right) \|\nabla f(v_n)\| + \|T_n v_n - v_n\|. \end{aligned} \quad (83)$$

From the boundedness of  $\{v_n\}$ ,  $s_n \rightarrow 0$  ( $\Leftrightarrow \lambda_n \rightarrow 2/L$ ) and  $\|T_n v_n - v_n\| \rightarrow 0$  (due to (78)), it follows that

$$\lim_{n \rightarrow \infty} \|v_n - P_C\left(I - \frac{2}{L} \nabla f\right) v_n\| = 0. \quad (84)$$

Further, we show that  $\omega_w(x_n) \subset \Omega$ . Indeed, since  $\{x_n\}$  is bounded, there exists a subsequence  $\{x_{n_i}\}$  of  $\{x_n\}$  which converges weakly to some  $w$ . Note that  $\lim_{n \rightarrow \infty} \|x_n - u_n\| = 0$  (due to (75)). Hence,  $x_{n_i} \rightharpoonup w$ . Since  $C$  is closed and convex,  $C$  is weakly closed. So, we have  $w \in C$ . From (74) and (75), we have that  $\Delta_{n_i}^k x_{n_i} \rightharpoonup w$ ,  $\Lambda_{n_i}^m u_{n_i} \rightharpoonup w$ ,  $u_{n_i} \rightharpoonup w$ ,  $v_{n_i} \rightharpoonup w$ , where  $k \in \{1, 2, \dots, M\}$ ,  $m \in \{1, 2, \dots, N\}$ . First, we prove that  $w \in \cap_{m=1}^N I(B_m, R_m)$ . As a matter of fact, since  $B_m$  is  $\eta_m$ -inverse-strongly monotone,  $B_m$  is a monotone and Lipschitz continuous mapping. It follows from Lemma 17 that  $R_m + B_m$  is maximal monotone. Let  $(v, g) \in G(R_m + B_m)$ ; that is,

$g - B_m v \in R_m v$ . Again, since  $\Lambda_n^m u_n = J_{R_m, \lambda_{m,n}}(I - \lambda_{m,n} B_m) \Lambda_n^{m-1} u_n$ ,  $n \geq 1$ ,  $m \in \{1, 2, \dots, N\}$ , we have

$$\Lambda_n^{m-1} u_n - \lambda_{m,n} B_m \Lambda_n^{m-1} u_n \in (I + \lambda_{m,n} R_m) \Lambda_n^m u_n; \quad (85)$$

that is,

$$\frac{1}{\lambda_{m,n}} (\Lambda_n^{m-1} u_n - \Lambda_n^m u_n - \lambda_{m,n} B_m \Lambda_n^{m-1} u_n) \in R_m \Lambda_n^m u_n. \quad (86)$$

In terms of the monotonicity of  $R_m$ , we get

$$\begin{aligned} &\left\langle v - \Lambda_n^m u_n, g - B_m v - \frac{1}{\lambda_{m,n}} \right. \\ &\quad \left. \times (\Lambda_n^{m-1} u_n - \Lambda_n^m u_n - \lambda_{m,n} B_m \Lambda_n^{m-1} u_n) \right\rangle \geq 0 \end{aligned} \quad (87)$$

and hence

$$\begin{aligned} &\langle v - \Lambda_n^m u_n, g \rangle \\ &\geq \left\langle v - \Lambda_n^m u_n, B_m v + \frac{1}{\lambda_{m,n}} \right. \\ &\quad \left. \times (\Lambda_n^{m-1} u_n - \Lambda_n^m u_n - \lambda_{m,n} B_m \Lambda_n^{m-1} u_n) \right\rangle \\ &= \left\langle v - \Lambda_n^m u_n, B_m v - B_m \Lambda_n^m u_n + B_m \Lambda_n^m u_n \right. \\ &\quad \left. - B_m \Lambda_n^{m-1} u_n + \frac{1}{\lambda_{m,n}} (\Lambda_n^{m-1} u_n - \Lambda_n^m u_n) \right\rangle \\ &\geq \langle v - \Lambda_n^m u_n, B_m \Lambda_n^m u_n - B_m \Lambda_n^{m-1} u_n \rangle \\ &\quad + \left\langle v - \Lambda_n^m u_n, \frac{1}{\lambda_{m,n}} (\Lambda_n^{m-1} u_n - \Lambda_n^m u_n) \right\rangle. \end{aligned} \quad (88)$$

In particular,

$$\begin{aligned} &\langle v - \Lambda_{n_i}^m u_{n_i}, g \rangle \\ &\geq \langle v - \Lambda_{n_i}^m u_{n_i}, B_m \Lambda_{n_i}^m u_{n_i} - B_m \Lambda_{n_i}^{m-1} u_{n_i} \rangle \\ &\quad + \left\langle v - \Lambda_{n_i}^m u_{n_i}, \frac{1}{\lambda_{m,n_i}} (\Lambda_{n_i}^{m-1} u_{n_i} - \Lambda_{n_i}^m u_{n_i}) \right\rangle. \end{aligned} \quad (89)$$

Since  $\|\Lambda_n^m u_n - \Lambda_n^{m-1} u_n\| \rightarrow 0$  (due to (74)) and  $\|B_m \Lambda_n^m u_n - B_m \Lambda_n^{m-1} u_n\| \rightarrow 0$  (due to the Lipschitz continuity of  $B_m$ ), we conclude from  $\Lambda_{n_i}^m u_{n_i} \rightharpoonup w$  and condition (ii) that

$$\lim_{i \rightarrow \infty} \langle v - \Lambda_{n_i}^m u_{n_i}, g \rangle = \langle v - w, g \rangle \geq 0. \quad (90)$$

It follows from the maximal monotonicity of  $B_m + R_m$  that  $0 \in (R_m + B_m)w$ ; that is,  $w \in I(B_m, R_m)$ . Therefore,  $w \in \cap_{m=1}^N I(B_m, R_m)$ . Next we prove that

$w \in \cap_{k=1}^M \text{GMEP}(\Theta_k, \varphi_k, A_k)$ . Since  $\Delta_n^k x_n = T_{r_{k,n}}^{(\Theta_k, \varphi_k)}(I - r_{k,n} A_k) \Delta_n^{k-1} x_n, n \geq 1, k \in \{1, 2, \dots, M\}$ , we have

$$\begin{aligned} & \Theta_k(\Delta_n^k x_n, y) + \varphi_k(y) - \varphi_k(\Delta_n^k x_n) \\ & + \langle A_k \Delta_n^{k-1} x_n, y - \Delta_n^k x_n \rangle \\ & + \frac{1}{r_{k,n}} \langle y - \Delta_n^k x_n, \Delta_n^k x_n - \Delta_n^{k-1} x_n \rangle \geq 0. \end{aligned} \tag{91}$$

By (A2), we have

$$\begin{aligned} & \varphi_k(y) - \varphi_k(\Delta_n^k x_n) + \langle A_k \Delta_n^{k-1} x_n, y - \Delta_n^k x_n \rangle \\ & + \frac{1}{r_{k,n}} \langle y - \Delta_n^k x_n, \Delta_n^k x_n - \Delta_n^{k-1} x_n \rangle \geq \Theta_k(y, \Delta_n^k x_n). \end{aligned} \tag{92}$$

Let  $z_t = ty + (1-t)w$  for all  $t \in (0, 1]$  and  $y \in C$ . This implies that  $z_t \in C$ . Then, we have

$$\begin{aligned} & \langle z_t - \Delta_n^k x_n, A_k z_t \rangle \\ & \geq \varphi_k(\Delta_n^k x_n) - \varphi_k(z_t) + \langle z_t - \Delta_n^k x_n, A_k z_t \rangle \\ & \quad - \langle z_t - \Delta_n^k x_n, A_k \Delta_n^{k-1} x_n \rangle \\ & \quad - \left\langle z_t - \Delta_n^k x_n, \frac{\Delta_n^k x_n - \Delta_n^{k-1} x_n}{r_{k,n}} \right\rangle + \Theta_k(z_t, \Delta_n^k x_n) \\ & = \varphi_k(\Delta_n^k x_n) - \varphi_k(z_t) \\ & \quad + \langle z_t - \Delta_n^k x_n, A_k z_t - A_k \Delta_n^k x_n \rangle \\ & \quad + \langle z_t - \Delta_n^k x_n, A_k \Delta_n^k x_n - A_k \Delta_n^{k-1} x_n \rangle \\ & \quad - \left\langle z_t - \Delta_n^k x_n, \frac{\Delta_n^k x_n - \Delta_n^{k-1} x_n}{r_{k,n}} \right\rangle + \Theta_k(z_t, \Delta_n^k x_n). \end{aligned} \tag{93}$$

By (74), we have  $\|A_k \Delta_n^k x_n - A_k \Delta_n^{k-1} x_n\| \rightarrow 0$  (due to the Lipschitz continuity of  $A_k$ ). Furthermore, by the monotonicity of  $A_k$ , we obtain  $\langle z_t - \Delta_n^k x_n, A_k z_t - A_k \Delta_n^k x_n \rangle \geq 0$ . Then, by (A4) we obtain

$$\langle z_t - w, A_k z_t \rangle \geq \varphi_k(w) - \varphi_k(z_t) + \Theta_k(z_t, w). \tag{94}$$

Utilizing (A1), (A4), and (94), we obtain

$$\begin{aligned} 0 & = \Theta_k(z_t, z_t) + \varphi_k(z_t) - \varphi_k(z_t) \\ & \leq t \Theta_k(z_t, y) + (1-t) \Theta_k(z_t, w) + t \varphi_k(y) \\ & \quad + (1-t) \varphi_k(w) - \varphi_k(z_t) \\ & \leq t [\Theta_k(z_t, y) + \varphi_k(y) - \varphi_k(z_t)] \\ & \quad + (1-t) \langle z_t - w, A_k z_t \rangle \\ & = t [\Theta_k(z_t, y) + \varphi_k(y) - \varphi_k(z_t)] \\ & \quad + (1-t) t \langle y - w, A_k z_t \rangle, \end{aligned} \tag{95}$$

and hence

$$0 \leq \Theta_k(z_t, y) + \varphi_k(y) - \varphi_k(z_t) + (1-t) \langle y - w, A_k z_t \rangle. \tag{96}$$

Letting  $t \rightarrow 0$ , we have, for each  $y \in C$ ,

$$0 \leq \Theta_k(w, y) + \varphi_k(y) - \varphi_k(w) + \langle y - w, A_k w \rangle. \tag{97}$$

This implies that  $w \in \text{GMEP}(\Theta_k, \varphi_k, A_k)$  and hence  $w \in \cap_{k=1}^M \text{GMEP}(\Theta_k, \varphi_k, A_k)$ . Further, let us show that  $w \in \Gamma$ . As a matter of fact, from (84),  $v_{n_i} \rightarrow w$ , and Lemma 9, we conclude that

$$w = P_C \left( I - \frac{2}{L} \nabla f \right) w. \tag{98}$$

So,  $w \in \text{VI}(C, \nabla f) = \Gamma$ . Therefore,  $w \in \cap_{k=1}^M \text{GMEP}(\Theta_k, \varphi_k, A_k) \cap \cap_{i=1}^N I(B_i, R_i) \cap \Gamma =: \Omega$ . This shows that  $\omega_w(x_n) \subset \Omega$ .

Finally, let us show that  $x_n \rightarrow q$  as  $n \rightarrow \infty$  where  $q$  is the unique solution of the VIP (50). Indeed, we note that, for  $w \in \Omega$  with  $x_{n_i} \rightarrow w$ ,

$$\begin{aligned} x_n - w & = s_n (\gamma V x_n - \mu F w) + (I - s_n \mu F) T_n v_n \\ & \quad - (I - s_n \mu F) w. \end{aligned} \tag{99}$$

By (61) and Lemma 13, we obtain that

$$\begin{aligned} \|x_n - w\|^2 & = s_n \langle \gamma V x_n - \mu F w, x_n - w \rangle \\ & \quad + \langle (I - s_n \mu F) T_n v_n - (I - s_n \mu F) w, x_n - w \rangle \\ & = s_n \langle \gamma V x_n - \mu F w, x_n - w \rangle \\ & \quad + \|(I - s_n \mu F) T_n v_n - (I - s_n \mu F) w\| \|x_n - w\| \\ & \leq s_n \langle \gamma V x_n - \mu F w, x_n - w \rangle \\ & \quad + (1 - s_n \tau) \|v_n - w\| \|x_n - w\| \\ & \leq s_n \langle \gamma V x_n - \mu F w, x_n - w \rangle + (1 - s_n \tau) \|x_n - w\|^2. \end{aligned} \tag{100}$$

Hence, it follows that

$$\begin{aligned} \|x_n - w\|^2 & \leq \frac{1}{\tau} \langle \gamma V x_n - \mu F w, x_n - w \rangle \\ & = \frac{1}{\tau} (\gamma \langle V x_n - V w, x_n - w \rangle \\ & \quad + \langle \gamma V w - \mu F w, x_n - w \rangle) \\ & \leq \frac{1}{\tau} (\gamma l \|x_n - w\|^2 + \langle \gamma V w - \mu F w, x_n - w \rangle), \end{aligned} \tag{101}$$

which hence leads to

$$\|x_n - w\|^2 \leq \frac{\langle \gamma V w - \mu F w, x_n - w \rangle}{\tau - \gamma l}. \tag{102}$$



In particular, we have

$$\|x_{n_i} - w\|^2 \leq \frac{\langle \gamma Vw - \mu Fw, x_{n_i} - w \rangle}{\tau - \gamma l}. \tag{103}$$

Since  $x_{n_i} \rightarrow w$ , it follows from (103) that  $x_{n_i} \rightarrow w$  as  $i \rightarrow \infty$ .

Now we show that  $w$  solves the VIP (50). Since  $x_n = s_n \gamma Vx_n + (I - s_n \mu F)T_n v_n$ , we have

$$(\mu F - \gamma V)x_n = -\frac{1}{s_n} ((I - s_n \mu F)x_n - (I - s_n \mu F)T_n v_n). \tag{104}$$

It follows that, for each  $p \in \Omega$ ,

$$\begin{aligned} & \langle (\mu F - \gamma V)x_n, x_n - p \rangle \\ &= -\frac{1}{s_n} \langle (I - s_n \mu F)x_n - (I - s_n \mu F)T_n v_n, x_n - p \rangle \\ &= -\frac{1}{s_n} \langle (I - s_n \mu F)x_n - (I - s_n \mu F)T_n v_n, x_n - p \rangle \\ &= -\frac{1}{s_n} \langle (I - s_n \mu F)x_n - (I - s_n \mu F)T_n \Lambda_n^N \Delta_n^M x_n, x_n - p \rangle \\ &= -\frac{1}{s_n} \langle (I - T_n \Lambda_n^N \Delta_n^M)x_n - (I - T_n \Lambda_n^N \Delta_n^M)p, x_n - p \rangle \\ &\quad + \langle \mu Fx_n - \mu FT_n \Lambda_n^N \Delta_n^M x_n, x_n - p \rangle \\ &\leq \langle \mu Fx_n - \mu FT_n \Lambda_n^N \Delta_n^M x_n, x_n - p \rangle, \end{aligned} \tag{105}$$

since  $I - T_n \Lambda_n^N \Delta_n^M$  is monotone (i.e.,  $\langle (I - T_n \Lambda_n^N \Delta_n^M)x - (I - T_n \Lambda_n^N \Delta_n^M)y, x - y \rangle \geq 0$  for all  $x, y \in H$ ). This is due to the nonexpansivity of  $T_n \Lambda_n^N \Delta_n^M$ . Since  $\|x_n - T_n v_n\| = \|(I - T_n \Lambda_n^N \Delta_n^M)x_n\| \rightarrow 0$  as  $n \rightarrow \infty$ , by replacing  $n$  in (105) with  $n_i$  and letting  $i \rightarrow \infty$ , we get

$$\begin{aligned} & \langle (\mu F - \gamma V)w, w - p \rangle \\ &= \lim_{i \rightarrow \infty} \langle (\mu F - \gamma V)x_{n_i}, x_{n_i} - p \rangle \\ &\leq \lim_{i \rightarrow \infty} \langle \mu Fx_{n_i} - \mu FT_{n_i} v_{n_i}, x_{n_i} - p \rangle = 0. \end{aligned} \tag{106}$$

That is,  $w \in \Omega$  is a solution of VIP (50).

Finally, in terms of the uniqueness of solutions of VIP (50), we deduce that  $w = q$  and  $x_{n_i} \rightarrow q$  as  $n \rightarrow \infty$ . So, every weak convergence subsequence of  $\{x_n\}$  converges strongly to the unique solution  $q$  of VIP (50). Therefore,  $\{x_n\}$  converges strongly to the unique solution  $q$  of VIP (50). In addition, the VIP (50) can be rewritten as

$$\langle (I - \mu F + \gamma V)q - q, q - p \rangle \geq 0, \quad \forall p \in \Omega. \tag{107}$$

By Proposition 2(i), this is equivalent to the fixed point equation:

$$P_\Omega (I - \mu F + \gamma V)q = q. \tag{108}$$

This completes the proof.  $\square$

**Corollary 19.** Let  $C$  be a nonempty closed convex subset of a real Hilbert space  $H$ . Let  $f : C \rightarrow \mathbf{R}$  be a convex functional with  $L$ -Lipschitz continuous gradient  $\nabla f$ . Let  $\Theta$  be a bifunction from  $C \times C$  to  $\mathbf{R}$  satisfying (A1)–(A4) and let  $\varphi : C \rightarrow \mathbf{R} \cup \{+\infty\}$  be a proper lower semicontinuous and convex function. Let  $R_i : C \rightarrow 2^H$  be a maximal monotone mapping and let  $A : H \rightarrow H$  and  $B_i : C \rightarrow H$  be  $\zeta$ -inverse-strongly monotone and  $\eta_i$ -inverse-strongly monotone, respectively, for  $i = 1, 2$ . Let  $F : H \rightarrow H$  be a  $\kappa$ -Lipschitzian and  $\eta$ -strongly monotone operator with positive constants  $\kappa, \eta > 0$ . Let  $V : H \rightarrow H$  be an  $l$ -Lipschitzian mapping with constant  $l \geq 0$ . Let  $0 < \mu < 2\eta/\kappa^2$  and  $0 \leq \gamma l < \tau$ , where  $\tau = 1 - \sqrt{1 - \mu(2\eta - \mu\kappa^2)}$ . Assume that  $\Omega := \text{GMEP}(\Theta, \varphi, A) \cap I(B_1, R_1) \cap I(B_2, R_2) \cap \Gamma \neq \emptyset$  and that either (B1) or (B2) holds. Let  $\{x_n\}$  be a sequence generated by

$$\begin{aligned} & \Theta(u_n, y) + \varphi(y) - \varphi(u_n) + \langle Ax_n, y - u_n \rangle \\ & \quad + \frac{1}{r_n} \langle y - u_n, u_n - x_n \rangle \geq 0, \quad \forall y \in C, \end{aligned} \tag{109}$$

$$v_n = J_{R_2, \lambda_{2,n}} (I - \lambda_{2,n} B_2) J_{R_1, \lambda_{1,n}} (I - \lambda_{1,n} B_1) u_n,$$

$$x_n = s_n \gamma Vx_n + (I - s_n \mu F)T_n v_n, \quad \forall n \geq 1,$$

where  $P_C(I - \lambda_n \nabla f) = s_n I + (1 - s_n)T_n$  (here  $T_n$  is nonexpansive and  $s_n = (2 - \lambda_n L)/4 \in (0, 1/2)$  for each  $\lambda_n \in (0, 2/L)$ ). Assume that the following conditions hold:

- (i)  $s_n \in (0, 1/2)$  for each  $\lambda_n \in (0, 2/L)$ ,  $\lim_{n \rightarrow \infty} s_n = 0$  ( $\Leftrightarrow \lim_{n \rightarrow \infty} \lambda_n = 2/L$ );
- (ii)  $\{\lambda_{i,n}\} \subset [a_i, b_i] \subset (0, 2\eta_i)$  for  $i = 1, 2$ ;
- (iii)  $\{r_n\} \subset [e, f] \subset (0, 2\zeta)$ .

Then  $\{x_n\}$  converges strongly as  $\lambda_n \rightarrow 2/L$  ( $\Leftrightarrow s_n \rightarrow 0$ ) to a point  $q \in \Omega$ , which is a unique solution of the VIP:

$$\langle (\mu F - \gamma V)q, p - q \rangle \geq 0, \quad \forall p \in \Omega. \tag{110}$$

**Corollary 20.** Let  $C$  be a nonempty closed convex subset of a real Hilbert space  $H$ . Let  $f : C \rightarrow \mathbf{R}$  be a convex functional with  $L$ -Lipschitz continuous gradient  $\nabla f$ . Let  $\Theta$  be a bifunction from  $C \times C$  to  $\mathbf{R}$  satisfying (A1)–(A4) and let  $\varphi : C \rightarrow \mathbf{R} \cup \{+\infty\}$  be a proper lower semicontinuous and convex function. Let  $R : C \rightarrow 2^H$  be a maximal monotone mapping and let  $A : H \rightarrow H$  and  $B : C \rightarrow H$  be  $\zeta$ -inverse-strongly monotone and  $\xi$ -inverse-strongly monotone, respectively. Let  $F : H \rightarrow H$  be a  $\kappa$ -Lipschitzian and  $\eta$ -strongly monotone operator with positive constants  $\kappa, \eta > 0$ . Let  $V : H \rightarrow H$  be an  $l$ -Lipschitzian mapping with constant  $l \geq 0$ . Let  $0 < \mu < 2\eta/\kappa^2$  and  $0 \leq \gamma l < \tau$ , where  $\tau = 1 - \sqrt{1 - \mu(2\eta - \mu\kappa^2)}$ . Assume that  $\Omega := \text{GMEP}(\Theta, \varphi, A) \cap I(B, R) \cap \Gamma \neq \emptyset$  and that either (B1) or (B2) holds. Let  $\{x_n\}$  be a sequence generated by

$$\begin{aligned} & \Theta(u_n, y) + \varphi(y) - \varphi(u_n) + \langle Ax_n, y - u_n \rangle \\ & \quad + \frac{1}{r_n} \langle y - u_n, u_n - x_n \rangle \geq 0, \quad \forall y \in C, \end{aligned} \tag{111}$$

$$x_n = s_n \gamma Vx_n + (I - s_n \mu F)T_n J_{R, \rho_n} (u_n - \rho_n B u_n),$$

$$\forall n \geq 1,$$

where  $P_C(I - \lambda_n \nabla f) = s_n I + (1 - s_n)T_n$  (here  $T_n$  is nonexpansive and  $s_n = (2 - \lambda_n L)/4 \in (0, 1/2)$  for each  $\lambda_n \in (0, 2/L)$ ). Assume that the following conditions hold:

- (i)  $s_n \in (0, 1/2)$  for each  $\lambda_n \in (0, 2/L)$ ,  $\lim_{n \rightarrow \infty} s_n = 0$  ( $\Leftrightarrow \lim_{n \rightarrow \infty} \lambda_n = 2/L$ );
- (ii)  $\{\rho_n\} \subset [a, b] \subset (0, 2\xi)$ ;
- (iii)  $\{r_n\} \subset [e, f] \subset (0, 2\zeta)$ .

Then  $\{x_n\}$  converges strongly as  $\lambda_n \rightarrow 2/L$  ( $\Leftrightarrow s_n \rightarrow 0$ ) to a point  $q \in \Omega$ , which is a unique solution of the VIP:

$$\langle (\mu F - \gamma V)q, p - q \rangle \geq 0, \quad \forall p \in \Omega. \quad (112)$$

#### 4. Explicit Iterative Algorithm and Its Convergence Criteria

We next state and prove the second main result of this paper.

**Theorem 21.** Let  $C$  be a nonempty closed convex subset of a real Hilbert space  $H$ . Let  $f : C \rightarrow \mathbf{R}$  be a convex functional with  $L$ -Lipschitz continuous gradient  $\nabla f$ . Let  $M, N$  be two integers. Let  $\Theta_k$  be a bifunction from  $C \times C$  to  $\mathbf{R}$  satisfying (A1)–(A4) and let  $\varphi_k : C \rightarrow \mathbf{R} \cup \{+\infty\}$  be a proper lower semicontinuous and convex function, where  $k \in \{1, 2, \dots, M\}$ . Let  $R_i : C \rightarrow 2^H$  be a maximal monotone mapping and let  $A_k : H \rightarrow H$  and  $B_i : C \rightarrow H$  be  $\mu_k$ -inverse-strongly monotone and  $\eta_i$ -inverse-strongly monotone, respectively, where  $k \in \{1, 2, \dots, M\}$ ,  $i \in \{1, 2, \dots, N\}$ . Let  $F : H \rightarrow H$  be a  $\kappa$ -Lipschitzian and  $\eta$ -strongly monotone operator with positive constants  $\kappa, \eta > 0$ . Let  $V : H \rightarrow H$  be an  $l$ -Lipschitzian mapping with constant  $l \geq 0$ . Let  $0 < \mu < 2\eta/\kappa^2$  and  $0 \leq \gamma l < \tau$ , where  $\tau = 1 - \sqrt{1 - \mu(2\eta - \mu\kappa^2)}$ . Assume that  $\Omega := \bigcap_{k=1}^M \text{GMEP}(\Theta_k, \varphi_k, A_k) \cap \bigcap_{i=1}^N I(B_i, R_i) \cap \Gamma \neq \emptyset$  and that either (B1) or (B2) holds. For arbitrarily given  $x_1 \in H$ , let  $\{x_n\}$  be a sequence generated by

$$\begin{aligned} u_n &= T_{r_{M,n}}^{(\Theta_M, \varphi_M)} (I - r_{M,n} A_M) T_{r_{M-1,n}}^{(\Theta_{M-1}, \varphi_{M-1})} \\ &\quad \times (I - r_{M-1,n} A_{M-1}) \cdots T_{r_{1,n}}^{(\Theta_1, \varphi_1)} (I - r_{1,n} A_1) x_n, \\ v_n &= J_{R_N, \lambda_{N,n}} (I - \lambda_{N,n} B_N) J_{R_{N-1}, \lambda_{N-1,n}} \\ &\quad \times (I - \lambda_{N-1,n} A_{N-1}) \cdots J_{R_1, \lambda_{1,n}} (I - \lambda_{1,n} B_1) u_n, \end{aligned} \quad (113)$$

$$\begin{aligned} x_{n+1} &= s_n \gamma V x_n + \beta_n x_n + ((1 - \beta_n) I - s_n \mu F) T_n v_n, \\ &\quad \forall n \geq 1, \end{aligned}$$

where  $P_C(I - \lambda_n \nabla f) = s_n I + (1 - s_n)T_n$  (here  $T_n$  is nonexpansive and  $s_n = (2 - \lambda_n L)/4 \in (0, 1/2)$  for each  $\lambda_n \in (0, 2/L)$ ). Assume that the following conditions hold:

- (i)  $s_n \in (0, 1/2)$  for each  $\lambda_n \in (0, 2/L)$ , and  $\lim_{n \rightarrow \infty} s_n = 0$  ( $\Leftrightarrow \lim_{n \rightarrow \infty} \lambda_n = 2/L$ );
- (ii)  $\{\beta_n\} \subset (0, 1)$  and  $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$ ;
- (iii)  $\{\lambda_{i,n}\} \subset [a_i, b_i] \subset (0, 2\eta_i)$  and  $\lim_{n \rightarrow \infty} |\lambda_{i,n+1} - \lambda_{i,n}| = 0$  for all  $i \in \{1, 2, \dots, N\}$ ;
- (iv)  $\{r_{k,n}\} \subset [e_k, f_k] \subset (0, 2\mu_k)$  and  $\lim_{n \rightarrow \infty} |r_{k,n+1} - r_{k,n}| = 0$  for all  $k \in \{1, 2, \dots, M\}$ .

Then  $\{x_n\}$  converges strongly as  $\lambda_n \rightarrow 2/L$  ( $\Leftrightarrow s_n \rightarrow 0$ ) to a point  $q \in \Omega$ , which is a unique solution of VIP (50).

*Proof.* First of all, repeating the same arguments as in Theorem 18, we can write

$$P_C(I - \lambda_n \nabla f) = \frac{2 - \lambda_n L}{4} I + \frac{2 + \lambda_n L}{4} T_n = s_n I + (1 - s_n) T_n, \quad (114)$$

where  $T_n$  is nonexpansive and  $s_n := s_n(\lambda_n) = (2 - \lambda_n L)/4 \in (0, 1/2)$  for each  $\lambda_n \in (0, 2/L)$ . It is clear that

$$\lambda_n \rightarrow \frac{2}{L} \iff s_n \rightarrow 0. \quad (115)$$

Put

$$\begin{aligned} \Delta_n^k &= T_{r_{k,n}}^{(\Theta_k, \varphi_k)} (I - r_{k,n} A_k) T_{r_{k-1,n}}^{(\Theta_{k-1}, \varphi_{k-1})} \\ &\quad \times (I - r_{k-1,n} A_{k-1}) \cdots T_{r_{1,n}}^{(\Theta_1, \varphi_1)} (I - r_{1,n} A_1) x_n \end{aligned} \quad (116)$$

for all  $k \in \{1, 2, \dots, M\}$  and  $n \geq 1$ ,

$$\begin{aligned} \Lambda_n^i &= J_{R_i, \lambda_{i,n}} (I - \lambda_{i,n} B_i) J_{R_{i-1}, \lambda_{i-1,n}} \\ &\quad \times (I - \lambda_{i-1,n} B_{i-1}) \cdots J_{R_1, \lambda_{1,n}} (I - \lambda_{1,n} B_1) \end{aligned} \quad (117)$$

for all  $i \in \{1, 2, \dots, N\}$  and  $n \geq 1$ , and  $\Delta_n^0 = \Lambda_n^0 = I$ , where  $I$  is the identity mapping on  $H$ . Then we have that  $u_n = \Delta_n^M x_n$  and  $v_n = \Lambda_n^N u_n$ . In addition, taking into consideration conditions (i) and (ii), we may assume, without loss of generality, that  $s_n \leq 1 - \beta_n$  for all  $n \geq 1$ .

We divide the remainder of the proof into several steps.

*Step 1.* Let us show that  $\|x_n - p\| \leq \max\{\|x_1 - p\|, \|\gamma V p - \mu F p\|/(\tau - \gamma l)\}$  for all  $n \geq 1$  and  $p \in \Omega$ . Indeed, take  $p \in \Omega$  arbitrarily. Repeating the same arguments as those of (59)–(61) in the proof of Theorem 18, we obtain

$$\begin{aligned} \|u_n - p\| &\leq \|x_n - p\|, \\ \|v_n - p\| &\leq \|u_n - p\|, \\ \|v_n - p\| &\leq \|x_n - p\|. \end{aligned} \quad (118)$$

Then from (118),  $T_n p = p$ , and Lemma 13, we have

$$\begin{aligned} \|x_{n+1} - p\| &= \|s_n (\gamma V x_n - \mu F p) + \beta_n (x_n - p) \\ &\quad + ((1 - \beta_n) I - s_n \mu F) T_n v_n \\ &\quad - ((1 - \beta_n) I - s_n \mu F) T_n p\| \end{aligned}$$

$$\begin{aligned}
 &\leq s_n \|\gamma Vx_n - \mu Fp\| + \beta_n \|x_n - p\| \\
 &\quad + (1 - \beta_n) \left\| \left( I - \frac{s_n}{1 - \beta_n} \mu F \right) T_n v_n \right. \\
 &\quad \quad \left. - \left( I - \frac{s_n}{1 - \beta_n} \mu F \right) T_n p \right\| \\
 &\leq s_n (\|\gamma Vx_n - \gamma Vp\| + \|\gamma Vp - \mu Fp\|) \\
 &\quad + \beta_n \|x_n - p\| + (1 - \beta_n) \left( 1 - \frac{s_n \tau}{1 - \beta_n} \right) \|v_n - p\| \\
 &\leq s_n (\|\gamma Vx_n - \gamma Vp\| + \|\gamma Vp - \mu Fp\|) \\
 &\quad + \beta_n \|x_n - p\| + (1 - \beta_n - s_n \tau) \|x_n - p\| \\
 &\leq s_n \gamma l \|x_n - p\| + s_n \|\gamma Vp - \mu Fp\| \\
 &\quad + (1 - s_n \tau) \|x_n - p\| \\
 &= (1 - s_n (\tau - \gamma l)) \|x_n - p\| \\
 &\quad + s_n (\tau - \gamma l) \frac{\|\gamma Vp - \mu Fp\|}{\tau - \gamma l} \\
 &\leq \max \left\{ \|x_n - p\|, \frac{\|\gamma Vp - \mu Fp\|}{\tau - \gamma l} \right\}. \tag{119}
 \end{aligned}$$

By induction, we have

$$\|x_n - p\| \leq \max \left\{ \|x_1 - p\|, \frac{\|\gamma Vp - \mu Fp\|}{\tau - \gamma l} \right\}, \quad \forall n \geq 1. \tag{120}$$

Hence,  $\{x_n\}$  is bounded. According to (118),  $\{u_n\}$ ,  $\{v_n\}$ ,  $\{T_n v_n\}$ ,  $\{Vx_n\}$ , and  $\{FT_n v_n\}$  are also bounded.

*Step 2.* Let us show that  $\|x_{n+1} - x_n\| \rightarrow 0$  as  $n \rightarrow \infty$ . To this end, define

$$x_{n+1} = \beta_n x_n + (1 - \beta_n) z_n, \quad \forall n \geq 1. \tag{121}$$

Observe that, from the definition of  $z_n$ ,

$$\begin{aligned}
 z_{n+1} - z_n &= \frac{x_{n+2} - \beta_{n+1} x_{n+1}}{1 - \beta_{n+1}} - \frac{x_{n+1} - \beta_n x_n}{1 - \beta_n} \\
 &= \frac{s_{n+1} \gamma Vx_{n+1} + ((1 - \beta_{n+1}) I - s_{n+1} \mu F) T_{n+1} v_{n+1}}{1 - \beta_{n+1}} \\
 &\quad - \frac{s_n \gamma Vx_n + ((1 - \beta_n) I - s_n \mu F) T_n v_n}{1 - \beta_n}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{s_{n+1}}{1 - \beta_{n+1}} \gamma Vx_{n+1} - \frac{s_n}{1 - \beta_n} \gamma Vx_n + T_{n+1} v_{n+1} \\
 &\quad - T_n v_n + \frac{s_n}{1 - \beta_n} \mu FT_n v_n - \frac{s_{n+1}}{1 - \beta_{n+1}} \mu FT_{n+1} v_{n+1} \\
 &= \frac{s_{n+1}}{1 - \beta_{n+1}} (\gamma Vx_{n+1} - \mu FT_{n+1} v_{n+1}) \\
 &\quad + \frac{s_n}{1 - \beta_n} (\mu FT_n v_n - \gamma Vx_n) + T_{n+1} v_{n+1} - T_n v_n. \tag{122}
 \end{aligned}$$

Thus, it follows that

$$\begin{aligned}
 \|z_{n+1} - z_n\| &\leq \frac{s_{n+1}}{1 - \beta_{n+1}} (\gamma \|Vx_{n+1}\| + \mu \|FT_{n+1} v_{n+1}\|) \\
 &\quad + \frac{s_n}{1 - \beta_n} (\mu \|FT_n v_n\| + \gamma \|Vx_n\|) \\
 &\quad + \|T_{n+1} v_{n+1} - T_n v_n\|. \tag{123}
 \end{aligned}$$

On the other hand, since  $\nabla f$  is  $(1/L)$ -ism,  $P_C(I - \lambda_n \nabla f)$  is nonexpansive for  $\lambda_n \in (0, 2/L)$ . So, it follows that, for any given  $p \in \Omega$ ,

$$\begin{aligned}
 \|P_C(I - \lambda_{n+1} \nabla f) v_n\| &\leq \|P_C(I - \lambda_{n+1} \nabla f) v_n - p\| + \|p\| \\
 &= \|P_C(I - \lambda_{n+1} \nabla f) v_n \\
 &\quad - P_C(I - \lambda_{n+1} \nabla f) p\| + \|p\| \\
 &\leq \|v_n - p\| + \|p\| \\
 &\leq \|v_n\| + 2 \|p\|. \tag{124}
 \end{aligned}$$

This together with the boundedness of  $\{v_n\}$  implies that  $\{P_C(I - \lambda_{n+1} \nabla f) v_n\}$  is bounded. Also, observe that

$$\begin{aligned}
 &\|T_{n+1} v_n - T_n v_n\| \\
 &= \left\| \frac{4P_C(I - \lambda_{n+1} \nabla f) - (2 - \lambda_{n+1} L) I}{2 + \lambda_{n+1} L} v_n \right. \\
 &\quad \left. - \frac{4P_C(I - \lambda_n \nabla f) - (2 - \lambda_n L) I}{2 + \lambda_n L} v_n \right\| \\
 &\leq \left\| \frac{4P_C(I - \lambda_{n+1} \nabla f)}{2 + \lambda_{n+1} L} v_n - \frac{4P_C(I - \lambda_n \nabla f)}{2 + \lambda_n L} v_n \right\| \\
 &\quad + \left\| \frac{2 - \lambda_n L}{2 + \lambda_n L} v_n - \frac{2 - \lambda_{n+1} L}{2 + \lambda_{n+1} L} v_n \right\| \\
 &= \left\| (4(2 + \lambda_n L) P_C(I - \lambda_{n+1} \nabla f) v_n \right. \\
 &\quad - 4(2 + \lambda_{n+1} L) P_C(I - \lambda_n \nabla f) v_n) \\
 &\quad \left. \times ((2 + \lambda_{n+1} L)(2 + \lambda_n L))^{-1} \right\| \\
 &\quad + \frac{4L |\lambda_{n+1} - \lambda_n|}{(2 + \lambda_{n+1} L)(2 + \lambda_n L)} \|v_n\|
 \end{aligned}$$

$$\begin{aligned}
 &= \left\| (4L(\lambda_n - \lambda_{n+1})P_C(I - \lambda_{n+1}\nabla f)v_n + 4(2 + \lambda_{n+1}L) \right. \\
 &\quad \times (P_C(I - \lambda_{n+1}\nabla f)v_n - P_C(I - \lambda_n\nabla f)v_n)) \\
 &\quad \times ((2 + \lambda_{n+1}L)(2 + \lambda_nL))^{-1} \left. \right\| \\
 &\quad + \frac{4L|\lambda_{n+1} - \lambda_n|}{(2 + \lambda_{n+1}L)(2 + \lambda_nL)} \|v_n\| \\
 &\leq \frac{4L|\lambda_n - \lambda_{n+1}| \|P_C(I - \lambda_{n+1}\nabla f)v_n\|}{(2 + \lambda_{n+1}L)(2 + \lambda_nL)} \\
 &\quad + (4(2 + \lambda_{n+1}L) \|P_C(I - \lambda_{n+1}\nabla f)v_n \\
 &\quad \quad - P_C(I - \lambda_n\nabla f)v_n\|) \\
 &\quad \times ((2 + \lambda_{n+1}L)(2 + \lambda_nL))^{-1} \\
 &\quad + \frac{4L|\lambda_{n+1} - \lambda_n|}{(2 + \lambda_{n+1}L)(2 + \lambda_nL)} \|v_n\| \\
 &\leq |\lambda_{n+1} - \lambda_n| [L \|P_C(I - \lambda_{n+1}\nabla f)v_n\| \\
 &\quad + 4 \|\nabla f(v_n)\| + L \|v_n\|] \\
 &\leq \widetilde{M} |\lambda_{n+1} - \lambda_n|,
 \end{aligned} \tag{125}$$

where  $\sup_{n \geq 1} \{L \|P_C(I - \lambda_{n+1}\nabla f)v_n\| + 4 \|\nabla f(v_n)\| + L \|v_n\|\} \leq \widetilde{M}$  for some  $\widetilde{M} > 0$ . So, by (125), we have that

$$\begin{aligned}
 \|T_{n+1}v_{n+1} - T_n v_n\| &\leq \|T_{n+1}v_{n+1} - T_{n+1}v_n\| \\
 &\quad + \|T_{n+1}v_n - T_n v_n\| \\
 &\leq \|v_{n+1} - v_n\| + \widetilde{M} |\lambda_{n+1} - \lambda_n| \\
 &\leq \|v_{n+1} - v_n\| + \frac{4\widetilde{M}}{L} (s_{n+1} + s_n).
 \end{aligned} \tag{126}$$

Note that

$$\begin{aligned}
 \|v_{n+1} - v_n\| &= \|\Lambda_{n+1}^N u_{n+1} - \Lambda_n^N u_n\| \\
 &= \|J_{R_N, \lambda_{N, n+1}}(I - \lambda_{N, n+1} B_N) \Lambda_{n+1}^{N-1} u_{n+1} \\
 &\quad - J_{R_N, \lambda_{N, n}}(I - \lambda_{N, n} B_N) \Lambda_n^{N-1} u_n\| \\
 &\leq \|J_{R_N, \lambda_{N, n+1}}(I - \lambda_{N, n+1} B_N) \Lambda_{n+1}^{N-1} u_{n+1} \\
 &\quad - J_{R_N, \lambda_{N, n}}(I - \lambda_{N, n} B_N) \Lambda_{n+1}^{N-1} u_{n+1}\| \\
 &\quad + \|J_{R_N, \lambda_{N, n}}(I - \lambda_{N, n} B_N) \Lambda_{n+1}^{N-1} u_{n+1} \\
 &\quad - J_{R_N, \lambda_{N, n}}(I - \lambda_{N, n} B_N) \Lambda_n^{N-1} u_n\|
 \end{aligned}$$

$$\begin{aligned}
 &\leq \|(I - \lambda_{N, n+1} B_N) \Lambda_{n+1}^{N-1} u_{n+1} \\
 &\quad - (I - \lambda_{N, n} B_N) \Lambda_{n+1}^{N-1} u_{n+1}\| \\
 &\quad + \|(I - \lambda_{N, n} B_N) \Lambda_{n+1}^{N-1} u_{n+1} \\
 &\quad \quad - (I - \lambda_{N, n} B_N) \Lambda_n^{N-1} u_n\| \\
 &\leq |\lambda_{N, n+1} - \lambda_{N, n}| \|B_N \Lambda_{n+1}^{N-1} u_{n+1}\| \\
 &\quad + \|\Lambda_{n+1}^{N-1} u_{n+1} - \Lambda_n^{N-1} u_n\| \\
 &\leq |\lambda_{N, n+1} - \lambda_{N, n}| \|B_N \Lambda_{n+1}^{N-1} u_{n+1}\| \\
 &\quad + |\lambda_{N-1, n+1} - \lambda_{N-1, n}| \|B_{N-1} \Lambda_{n+1}^{N-2} u_{n+1}\| \\
 &\quad + \|\Lambda_{n+1}^{N-2} u_{n+1} - \Lambda_n^{N-2} u_n\| \\
 &\quad \vdots \\
 &\leq |\lambda_{N, n+1} - \lambda_{N, n}| \|B_N \Lambda_{n+1}^{N-1} u_{n+1}\| \\
 &\quad + |\lambda_{N-1, n+1} - \lambda_{N-1, n}| \|B_{N-1} \Lambda_{n+1}^{N-2} u_{n+1}\| \\
 &\quad + \dots + |\lambda_{1, n+1} - \lambda_{1, n}| \|B_1 \Lambda_{n+1}^0 u_{n+1}\| \\
 &\quad + \|\Lambda_{n+1}^0 u_{n+1} - \Lambda_n^0 u_n\| \\
 &\leq \widetilde{M}_0 \sum_{i=1}^N |\lambda_{i, n+1} - \lambda_{i, n}| + \|u_{n+1} - u_n\|,
 \end{aligned} \tag{127}$$

where  $\sup_{n \geq 1} \{\sum_{i=1}^N \|B_i \Lambda_{n+1}^{i-1} u_{n+1}\|\} \leq \widetilde{M}_0$  for some  $\widetilde{M}_0 > 0$ . Also, utilizing Proposition 1(v) we deduce that

$$\begin{aligned}
 \|u_{n+1} - u_n\| &= \|\Delta_{n+1}^M x_{n+1} - \Delta_n^M x_n\| \\
 &= \|T_{r_{M, n+1}}^{(\Theta_M, \varphi_M)}(I - r_{M, n+1} A_M) \Delta_{n+1}^{M-1} x_{n+1} \\
 &\quad - T_{r_{M, n}}^{(\Theta_M, \varphi_M)}(I - r_{M, n} A_M) \Delta_n^{M-1} x_n\| \\
 &\leq \|T_{r_{M, n+1}}^{(\Theta_M, \varphi_M)}(I - r_{M, n+1} A_M) \Delta_{n+1}^{M-1} x_{n+1} \\
 &\quad - T_{r_{M, n}}^{(\Theta_M, \varphi_M)}(I - r_{M, n} A_M) \Delta_{n+1}^{M-1} x_{n+1}\| \\
 &\quad + \|T_{r_{M, n}}^{(\Theta_M, \varphi_M)}(I - r_{M, n} A_M) \Delta_{n+1}^{M-1} x_{n+1} \\
 &\quad \quad - T_{r_{M, n}}^{(\Theta_M, \varphi_M)}(I - r_{M, n} A_M) \Delta_n^{M-1} x_n\| \\
 &\leq \|T_{r_{M, n+1}}^{(\Theta_M, \varphi_M)}(I - r_{M, n+1} A_M) \Delta_{n+1}^{M-1} x_{n+1} \\
 &\quad - T_{r_{M, n}}^{(\Theta_M, \varphi_M)}(I - r_{M, n+1} A_M) \Delta_{n+1}^{M-1} x_{n+1}\|
 \end{aligned}$$

$$\begin{aligned}
 &+ \left\| T_{r_{M,n}}^{(\Theta_M, \varphi_M)} (I - r_{M,n+1} A_M) \Delta_{n+1}^{M-1} x_{n+1} \right. \\
 &\quad \left. - T_{r_{M,n}}^{(\Theta_M, \varphi_M)} (I - r_{M,n} A_M) \Delta_{n+1}^{M-1} x_{n+1} \right\| \\
 &+ \left\| (I - r_{M,n} A_M) \Delta_{n+1}^{M-1} x_{n+1} - (I - r_{M,n} A_M) \Delta_n^{M-1} x_n \right\| \\
 &\leq \frac{|r_{M,n+1} - r_{M,n}|}{r_{M,n+1}} \left\| T_{r_{M,n+1}}^{(\Theta_M, \varphi_M)} (I - r_{M,n+1} A_M) \Delta_{n+1}^{M-1} x_{n+1} \right. \\
 &\quad \left. - (I - r_{M,n+1} A_M) \Delta_{n+1}^{M-1} x_{n+1} \right\| \\
 &+ |r_{M,n+1} - r_{M,n}| \left\| A_M \Delta_{n+1}^{M-1} x_{n+1} \right\| \\
 &+ \left\| \Delta_{n+1}^{M-1} x_{n+1} - \Delta_n^{M-1} x_n \right\| \\
 &= |r_{M,n+1} - r_{M,n}| \\
 &\quad \times \left[ \left\| A_M \Delta_{n+1}^{M-1} x_{n+1} \right\| \right. \\
 &\quad \left. + \frac{1}{r_{M,n+1}} \left\| T_{r_{M,n+1}}^{(\Theta_M, \varphi_M)} (I - r_{M,n+1} A_M) \Delta_{n+1}^{M-1} x_{n+1} \right. \right. \\
 &\quad \quad \left. \left. - (I - r_{M,n+1} A_M) \Delta_{n+1}^{M-1} x_{n+1} \right\| \right] \\
 &+ \left\| \Delta_{n+1}^{M-1} x_{n+1} - \Delta_n^{M-1} x_n \right\| \\
 &\vdots \\
 &\leq |r_{M,n+1} - r_{M,n}| \\
 &\quad \times \left[ \left\| A_M \Delta_{n+1}^{M-1} x_{n+1} \right\| \right. \\
 &\quad \left. + \frac{1}{r_{M,n+1}} \left\| T_{r_{M,n+1}}^{(\Theta_M, \varphi_M)} (I - r_{M,n+1} A_M) \Delta_{n+1}^{M-1} x_{n+1} \right. \right. \\
 &\quad \quad \left. \left. - (I - r_{M,n+1} A_M) \Delta_{n+1}^{M-1} x_{n+1} \right\| \right] \\
 &+ \dots + |r_{1,n+1} - r_{1,n}| \\
 &\quad \times \left[ \left\| A_1 \Delta_{n+1}^0 x_{n+1} \right\| \right. \\
 &\quad \left. + \frac{1}{r_{1,n+1}} \left\| T_{r_{1,n+1}}^{(\Theta_1, \varphi_1)} (I - r_{1,n+1} A_1) \Delta_{n+1}^0 x_{n+1} \right. \right. \\
 &\quad \quad \left. \left. - (I - r_{1,n+1} A_1) \Delta_{n+1}^0 x_{n+1} \right\| \right] \\
 &+ \left\| \Delta_{n+1}^0 x_{n+1} - \Delta_n^0 x_n \right\| \\
 &\leq \widetilde{M}_1 \sum_{k=1}^M |r_{k,n+1} - r_{k,n}| + \|x_{n+1} - x_n\|,
 \end{aligned}$$

(128)

where  $\widetilde{M}_1 > 0$  is a constant such that for each  $n \geq 1$

$$\begin{aligned}
 &\sum_{k=1}^M \left[ \left\| A_k \Delta_{n+1}^{k-1} x_{n+1} \right\| + \frac{1}{r_{k,n+1}} \right. \\
 &\quad \times \left\| T_{r_{k,n+1}}^{(\Theta_k, \varphi_k)} (I - r_{k,n+1} A_k) \Delta_{n+1}^{k-1} x_{n+1} \right. \\
 &\quad \quad \left. \left. - (I - r_{k,n+1} A_k) \Delta_{n+1}^{k-1} x_{n+1} \right\| \right] \leq \widetilde{M}_1.
 \end{aligned}$$

Combining (123)–(128), we get

$$\begin{aligned}
 &\|z_{n+1} - z_n\| - \|x_{n+1} - x_n\| \\
 &\leq \frac{s_{n+1}}{1 - \beta_{n+1}} (\gamma \|Vx_{n+1}\| + \mu \|FT_{n+1}v_{n+1}\|) \\
 &\quad + \frac{s_n}{1 - \beta_n} (\mu \|FT_n v_n\| + \gamma \|Vx_n\|) \\
 &\quad + \|T_{n+1}v_{n+1} - T_n v_n\| - \|x_{n+1} - x_n\| \\
 &\leq \frac{s_{n+1}}{1 - \beta_{n+1}} (\gamma \|Vx_{n+1}\| + \mu \|FT_{n+1}v_{n+1}\|) \\
 &\quad + \frac{s_n}{1 - \beta_n} (\mu \|FT_n v_n\| + \gamma \|Vx_n\|) \\
 &\quad + \|v_{n+1} - v_n\| + \frac{4\widetilde{M}}{L} (s_{n+1} + s_n) - \|x_{n+1} - x_n\| \\
 &\leq \frac{s_{n+1}}{1 - \beta_{n+1}} (\gamma \|Vx_{n+1}\| + \mu \|FT_{n+1}v_{n+1}\|) \\
 &\quad + \frac{s_n}{1 - \beta_n} (\mu \|FT_n v_n\| + \gamma \|Vx_n\|) \\
 &\quad + \widetilde{M}_0 \sum_{i=1}^N |\lambda_{i,n+1} - \lambda_{i,n}| + \|u_{n+1} - u_n\| \\
 &\quad + \frac{4\widetilde{M}}{L} (s_{n+1} + s_n) - \|x_{n+1} - x_n\| \\
 &\leq \frac{s_{n+1}}{1 - \beta_{n+1}} (\gamma \|Vx_{n+1}\| + \mu \|FT_{n+1}v_{n+1}\|) \\
 &\quad + \frac{s_n}{1 - \beta_n} (\mu \|FT_n v_n\| + \gamma \|Vx_n\|) \\
 &\quad + \widetilde{M}_0 \sum_{i=1}^N |\lambda_{i,n+1} - \lambda_{i,n}| + \widetilde{M}_1 \sum_{k=1}^M |r_{k,n+1} - r_{k,n}| \\
 &\quad + \|x_{n+1} - x_n\| + \frac{4\widetilde{M}}{L} (s_{n+1} + s_n) - \|x_{n+1} - x_n\| \\
 &= \frac{s_{n+1}}{1 - \beta_{n+1}} (\gamma \|Vx_{n+1}\| + \mu \|FT_{n+1}v_{n+1}\|) \\
 &\quad + \frac{s_n}{1 - \beta_n} (\mu \|FT_n v_n\| + \gamma \|Vx_n\|)
 \end{aligned}$$



$$\begin{aligned}
 & + \widetilde{M}_0 \sum_{i=1}^N |\lambda_{i,n+1} - \lambda_{i,n}| + \widetilde{M}_1 \sum_{k=1}^M |r_{k,n+1} - r_{k,n}| \\
 & + \frac{4\widetilde{M}}{L} (s_{n+1} + s_n).
 \end{aligned} \tag{130}$$

Thus, it follows from (130) and conditions (i)–(iv) that

$$\limsup_{n \rightarrow \infty} (\|z_{n+1} - z_n\| - \|x_{n+1} - x_n\|) \leq 0. \tag{131}$$

Hence, by Lemma 11 we have

$$\lim_{n \rightarrow \infty} \|z_n - x_n\| = 0. \tag{132}$$

Consequently,

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = \lim_{n \rightarrow \infty} (1 - \beta_n) \|z_n - x_n\| = 0, \tag{133}$$

and, by (126)–(128),

$$\begin{aligned}
 \lim_{n \rightarrow \infty} \|u_{n+1} - u_n\| & = 0, \\
 \lim_{n \rightarrow \infty} \|v_{n+1} - v_n\| & = 0, \\
 \lim_{n \rightarrow \infty} \|T_{n+1}v_{n+1} - T_nv_n\| & = 0.
 \end{aligned} \tag{134}$$

*Step 3.* Let us show that  $\|A_k \Delta_n^{k-1} x_n - A_k p\| \rightarrow 0$  and  $\|B_i \Lambda_n^{i-1} u_n - B_i p\| \rightarrow 0$  for all  $k \in \{1, 2, \dots, M\}$  and  $i \in \{1, 2, \dots, N\}$ .

Indeed, since

$$x_{n+1} = s_n \gamma Vx_n + \beta_n x_n + ((1 - \beta_n)I - s_n \mu F) T_n v_n, \tag{135}$$

we have

$$\begin{aligned}
 \|x_n - T_n v_n\| & \leq \|x_n - x_{n+1}\| + \|x_{n+1} - T_n v_n\| \\
 & \leq \|x_n - x_{n+1}\| + s_n \|\gamma Vx_n - \mu FT_n v_n\| \\
 & \quad + \beta_n \|x_n - T_n v_n\|;
 \end{aligned} \tag{136}$$

that is,

$$\begin{aligned}
 \|x_n - T_n v_n\| & \leq \frac{1}{1 - \beta_n} \|x_n - x_{n+1}\| \\
 & \quad + \frac{s_n}{1 - \beta_n} (\gamma \|Vx_n\| + \mu \|FT_n v_n\|).
 \end{aligned} \tag{137}$$

So, from  $s_n \rightarrow 0$ ,  $\|x_{n+1} - x_n\| \rightarrow 0$ , and condition (ii), it follows that

$$\lim_{n \rightarrow \infty} \|x_n - T_n v_n\| = 0. \tag{138}$$

Also, from (27) it follows that for all  $i \in \{1, 2, \dots, N\}$  and  $k \in \{1, 2, \dots, M\}$

$$\begin{aligned}
 \|v_n - p\|^2 & = \|\Lambda_n^N u_n - p\|^2 \\
 & \leq \|\Lambda_n^i u_n - p\|^2 \\
 & = \|J_{R_i, \lambda_{i,n}} (I - \lambda_{i,n} B_i) \Lambda_n^{i-1} u_n \\
 & \quad - J_{R_i, \lambda_{i,n}} (I - \lambda_{i,n} B_i) p\|^2 \\
 & \leq \|(I - \lambda_{i,n} B_i) \Lambda_n^{i-1} u_n \\
 & \quad - (I - \lambda_{i,n} B_i) p\|^2 \\
 & \leq \|\Lambda_n^{i-1} u_n - p\|^2 + \lambda_{i,n} (\lambda_{i,n} - 2\eta_i) \\
 & \quad \times \|B_i \Lambda_n^{i-1} u_n - B_i p\|^2 \\
 & \leq \|u_n - p\|^2 + \lambda_{i,n} (\lambda_{i,n} - 2\eta_i) \\
 & \quad \times \|B_i \Lambda_n^{i-1} u_n - B_i p\|^2 \\
 & \leq \|x_n - p\|^2 + \lambda_{i,n} (\lambda_{i,n} - 2\eta_i) \\
 & \quad \times \|B_i \Lambda_n^{i-1} u_n - B_i p\|^2,
 \end{aligned} \tag{139}$$

$$\begin{aligned}
 \|u_n - p\|^2 & = \|\Delta_n^M x_n - p\|^2 \\
 & \leq \|\Delta_n^k x_n - p\|^2 \\
 & = \|T_{r_{k,n}}^{(\Theta_k, \varphi_k)} (I - r_{k,n} A_k) \Delta_n^{k-1} x_n \\
 & \quad - T_{r_{k,n}}^{(\Theta_k, \varphi_k)} (I - r_{k,n} A_k) p\|^2 \\
 & \leq \|(I - r_{k,n} A_k) \Delta_n^{k-1} x_n - (I - r_{k,n} A_k) p\|^2 \\
 & \leq \|\Delta_n^{k-1} x_n - p\|^2 + r_{k,n} (r_{k,n} - 2\mu_k) \\
 & \quad \times \|A_k \Delta_n^{k-1} x_n - A_k p\|^2 \\
 & \leq \|x_n - p\|^2 + r_{k,n} (r_{k,n} - 2\mu_k) \\
 & \quad \times \|A_k \Delta_n^{k-1} x_n - A_k p\|^2.
 \end{aligned}$$

Furthermore, utilizing Lemma 7, we deduce from (113) that

$$\begin{aligned}
 \|x_{n+1} - p\|^2 & = \|s_n (\gamma Vx_n - \mu Fp) + \beta_n (x_n - T_n v_n) \\
 & \quad + (I - s_n \mu F) T_n v_n - (I - s_n \mu F) p\|^2 \\
 & \leq \|\beta_n (x_n - T_n v_n) + (I - s_n \mu F) T_n v_n - (I - s_n \mu F) p\|^2 \\
 & \quad + 2s_n \langle \gamma Vx_n - \mu Fp, x_{n+1} - p \rangle
 \end{aligned}$$

$$\begin{aligned}
 &\leq [\beta_n \|x_n - T_n v_n\| + \|(I - s_n \mu F) T_n v_n - (I - s_n \mu F) T_n p\|]^2 \\
 &\quad + 2s_n \|\gamma V x_n - \mu F p\| \|x_{n+1} - p\| \\
 &\leq [\beta_n \|x_n - T_n v_n\| + (1 - s_n \tau) \|v_n - p\|]^2 \\
 &\quad + 2s_n \|\gamma V x_n - \mu F p\| \|x_{n+1} - p\| \\
 &= (1 - s_n \tau)^2 \|v_n - p\|^2 + \beta_n^2 \|x_n - T_n v_n\|^2 \\
 &\quad + 2(1 - s_n \tau) \beta_n \|v_n - p\| \|x_n - T_n v_n\| \\
 &\quad + 2s_n \|\gamma V x_n - \mu F p\| \|x_{n+1} - p\|.
 \end{aligned} \tag{140}$$

From (139)–(140), it follows that

$$\begin{aligned}
 \|x_{n+1} - p\|^2 &\leq \|v_n - p\|^2 + \beta_n^2 \|x_n - T_n v_n\|^2 \\
 &\quad + 2(1 - s_n \tau) \beta_n \|v_n - p\| \|x_n - T_n v_n\| \\
 &\quad + 2s_n \|\gamma V x_n - \mu F p\| \|x_{n+1} - p\| \\
 &\leq \|u_n - p\|^2 + \lambda_{i,n} (\lambda_{i,n} - 2\eta_i) \\
 &\quad \times \|B_i \Lambda_n^{i-1} u_n - B_i p\|^2 + \beta_n^2 \|x_n - T_n v_n\|^2 \\
 &\quad + 2(1 - s_n \tau) \beta_n \|v_n - p\| \|x_n - T_n v_n\| \\
 &\quad + 2s_n \|\gamma V x_n - \mu F p\| \|x_{n+1} - p\| \\
 &\leq \|x_n - p\|^2 + r_{k,n} (r_{k,n} - 2\mu_k) \\
 &\quad \times \|A_k \Delta_n^{k-1} x_n - A_k p\|^2 \\
 &\quad + \lambda_{i,n} (\lambda_{i,n} - 2\eta_i) \|B_i \Lambda_n^{i-1} u_n - B_i p\|^2 \\
 &\quad + \beta_n^2 \|x_n - T_n v_n\|^2 \\
 &\quad + 2(1 - s_n \tau) \beta_n \|v_n - p\| \|x_n - T_n v_n\| \\
 &\quad + 2s_n \|\gamma V x_n - \mu F p\| \|x_{n+1} - p\|,
 \end{aligned} \tag{141}$$

and so

$$\begin{aligned}
 &r_{k,n} (2\mu_k - r_{k,n}) \|A_k \Delta_n^{k-1} x_n - A_k p\|^2 \\
 &\quad + \lambda_{i,n} (2\eta_i - \lambda_{i,n}) \|B_i \Lambda_n^{i-1} u_n - B_i p\|^2 \\
 &\leq \|x_n - p\|^2 - \|x_{n+1} - p\|^2 + \beta_n^2 \|x_n - T_n v_n\|^2 \\
 &\quad + 2(1 - s_n \tau) \beta_n \|v_n - p\| \|x_n - T_n v_n\| \\
 &\quad + 2s_n \|\gamma V x_n - \mu F p\| \|x_{n+1} - p\| \\
 &\leq (\|x_n - p\| + \|x_{n+1} - p\|) \\
 &\quad \times \|x_n - x_{n+1}\| + \beta_n^2 \|x_n - T_n v_n\|^2 \\
 &\quad + 2(1 - s_n \tau) \beta_n \|v_n - p\| \|x_n - T_n v_n\| \\
 &\quad + 2s_n \|\gamma V x_n - \mu F p\| \|x_{n+1} - p\|.
 \end{aligned} \tag{142}$$

Since  $\{\lambda_{i,n}\} \subset [a_i, b_i] \subset (0, 2\eta_i)$  and  $\{r_{k,n}\} \subset [e_k, f_k] \subset (0, 2\mu_k)$  for all  $i \in \{1, 2, \dots, N\}$  and  $k \in \{1, 2, \dots, M\}$ , by (133), (138), and (142) we conclude immediately that

$$\begin{aligned}
 \lim_{n \rightarrow \infty} \|A_k \Delta_n^{k-1} x_n - A_k p\| &= 0, \\
 \lim_{n \rightarrow \infty} \|B_i \Lambda_n^{i-1} u_n - B_i p\| &= 0,
 \end{aligned} \tag{143}$$

for all  $k \in \{1, 2, \dots, M\}$  and  $i \in \{1, 2, \dots, N\}$ .

*Step 4.* Let us show that  $\|x_n - u_n\| \rightarrow 0$ ,  $\|u_n - v_n\| \rightarrow 0$ , and  $\|v_n - T_n v_n\| \rightarrow 0$  as  $n \rightarrow \infty$ .

Indeed, by Proposition 1(iii) we obtain that for each  $k \in \{1, 2, \dots, M\}$

$$\begin{aligned}
 &\|\Delta_n^k x_n - p\|^2 \\
 &= \|T_{r_{k,n}}^{(\Theta_k, \varphi_k)} (I - r_{k,n} A_k) \Delta_n^{k-1} x_n - T_{r_{k,n}}^{(\Theta_k, \varphi_k)} (I - r_{k,n} A_k) p\|^2 \\
 &\leq \langle (I - r_{k,n} A_k) \Delta_n^{k-1} x_n - (I - r_{k,n} A_k) p, \Delta_n^k x_n - p \rangle \\
 &= \frac{1}{2} \left( \|(I - r_{k,n} A_k) \Delta_n^{k-1} x_n - (I - r_{k,n} A_k) p\|^2 \right. \\
 &\quad \left. + \|\Delta_n^k x_n - p\|^2 - \|(I - r_{k,n} A_k) \Delta_n^{k-1} x_n \right. \\
 &\quad \left. - (I - r_{k,n} A_k) p \right. \\
 &\quad \left. - (\Delta_n^k x_n - p) \right\|^2 \\
 &\leq \frac{1}{2} \left( \|\Delta_n^{k-1} x_n - p\|^2 + \|\Delta_n^k x_n - p\|^2 \right. \\
 &\quad \left. - \|\Delta_n^{k-1} x_n - \Delta_n^k x_n - r_{k,n} (A_k \Delta_n^{k-1} x_n - A_k p)\|^2 \right),
 \end{aligned} \tag{144}$$

which implies that

$$\begin{aligned}
 \|\Delta_n^k x_n - p\|^2 &\leq \|\Delta_n^{k-1} x_n - p\|^2 \\
 &\quad - \|\Delta_n^{k-1} x_n - \Delta_n^k x_n \\
 &\quad \quad - r_{k,n} (A_k \Delta_n^{k-1} x_n - A_k p)\|^2 \\
 &= \|\Delta_n^{k-1} x_n - p\|^2 - \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\|^2 \\
 &\quad - r_{k,n}^2 \|A_k \Delta_n^{k-1} x_n - A_k p\|^2 \\
 &\quad + 2r_{k,n} \langle \Delta_n^{k-1} x_n - \Delta_n^k x_n, A_k \Delta_n^{k-1} x_n - A_k p \rangle \\
 &\leq \|\Delta_n^{k-1} x_n - p\|^2 - \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\|^2 \\
 &\quad + 2r_{k,n} \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\| \|A_k \Delta_n^{k-1} x_n - A_k p\| \\
 &\leq \|x_n - p\|^2 - \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\|^2 \\
 &\quad + 2r_{k,n} \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\| \|A_k \Delta_n^{k-1} x_n - A_k p\|.
 \end{aligned} \tag{145}$$

Also, by Lemma 14, we obtain that for each  $i \in \{1, 2, \dots, N\}$

$$\begin{aligned}
 & \|\Lambda_n^i u_n - p\|^2 \\
 &= \|J_{R_i, \lambda_{i,n}}(I - \lambda_{i,n} B_i) \Lambda_n^{i-1} u_n - J_{R_i, \lambda_{i,n}}(I - \lambda_{i,n} B_i) p\|^2 \\
 &\leq \langle (I - \lambda_{i,n} B_i) \Lambda_n^{i-1} u_n - (I - \lambda_{i,n} B_i) p, \Lambda_n^i u_n - p \rangle \\
 &= \frac{1}{2} \left( \|(I - \lambda_{i,n} B_i) \Lambda_n^{i-1} u_n - (I - \lambda_{i,n} B_i) p\|^2 \right. \\
 &\quad \left. + \|\Lambda_n^i u_n - p\|^2 - \|(I - \lambda_{i,n} B_i) \Lambda_n^{i-1} u_n \right. \\
 &\quad \left. - (I - \lambda_{i,n} B_i) p \right. \\
 &\quad \left. - (\Lambda_n^i u_n - p)\|^2 \right) \\
 &\leq \frac{1}{2} \left( \|\Lambda_n^{i-1} u_n - p\|^2 + \|\Lambda_n^i u_n - p\|^2 \right. \\
 &\quad \left. - \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n - \lambda_{i,n} (B_i \Lambda_n^{i-1} u_n - B_i p)\|^2 \right) \\
 &\leq \frac{1}{2} \left( \|u_n - p\|^2 + \|\Lambda_n^i u_n - p\|^2 \right. \\
 &\quad \left. - \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n - \lambda_{i,n} (B_i \Lambda_n^{i-1} u_n - B_i p)\|^2 \right), \tag{146}
 \end{aligned}$$

which implies

$$\begin{aligned}
 & \|\Lambda_n^i u_n - p\|^2 \leq \|u_n - p\|^2 \\
 &\quad - \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n - \lambda_{i,n} (B_i \Lambda_n^{i-1} u_n - B_i p)\|^2 \\
 &= \|u_n - p\|^2 - \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\|^2 \\
 &\quad - \lambda_{i,n}^2 \|B_i \Lambda_n^{i-1} u_n - B_i p\|^2 \\
 &\quad + 2\lambda_{i,n} \langle \Lambda_n^{i-1} u_n - \Lambda_n^i u_n, B_i \Lambda_n^{i-1} u_n - B_i p \rangle \\
 &\leq \|u_n - p\|^2 - \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\|^2 \\
 &\quad + 2\lambda_{i,n} \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\| \|B_i \Lambda_n^{i-1} u_n - B_i p\|. \tag{147}
 \end{aligned}$$

Thus, from (140), (145), and (147), we have

$$\begin{aligned}
 & \|x_{n+1} - p\|^2 \\
 &\leq (1 - s_n \tau)^2 \|v_n - p\|^2 + \beta_n^2 \|x_n - T_n v_n\|^2 \\
 &\quad + 2(1 - s_n \tau) \beta_n \|v_n - p\| \|x_n - T_n v_n\| \\
 &\quad + 2s_n \|\gamma V x_n - \mu F p\| \|x_{n+1} - p\| \\
 &= (1 - s_n \tau)^2 \|\Lambda_n^N u_n - p\|^2 + \beta_n^2 \|x_n - T_n v_n\|^2 \\
 &\quad + 2(1 - s_n \tau) \beta_n \|v_n - p\| \|x_n - T_n v_n\| \\
 &\quad + 2s_n \|\gamma V x_n - \mu F p\| \|x_{n+1} - p\|
 \end{aligned}$$

$$\begin{aligned}
 & \leq (1 - s_n \tau)^2 \|\Lambda_n^i u_n - p\|^2 + \beta_n^2 \|x_n - T_n v_n\|^2 \\
 &\quad + 2(1 - s_n \tau) \beta_n \|v_n - p\| \|x_n - T_n v_n\| \\
 &\quad + 2s_n \|\gamma V x_n - \mu F p\| \|x_{n+1} - p\| \\
 &\leq (1 - s_n \tau)^2 \left[ \|u_n - p\|^2 - \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\|^2 \right. \\
 &\quad \left. + 2\lambda_{i,n} \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\| \right. \\
 &\quad \left. \times \|B_i \Lambda_n^{i-1} u_n - B_i p\| \right] \\
 &\quad + \beta_n^2 \|x_n - T_n v_n\|^2 + 2(1 - s_n \tau) \beta_n \|v_n - p\| \\
 &\quad \times \|x_n - T_n v_n\| + 2s_n \|\gamma V x_n - \mu F p\| \|x_{n+1} - p\| \\
 &= (1 - s_n \tau)^2 \left[ \|\Delta_n^M x_n - p\|^2 - \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\|^2 \right. \\
 &\quad \left. + 2\lambda_{i,n} \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\| \right. \\
 &\quad \left. \times \|B_i \Lambda_n^{i-1} u_n - B_i p\| \right] \\
 &\quad + \beta_n^2 \|x_n - T_n v_n\|^2 + 2(1 - s_n \tau) \beta_n \|v_n - p\| \\
 &\quad \times \|x_n - T_n v_n\| + 2s_n \|\gamma V x_n - \mu F p\| \|x_{n+1} - p\| \\
 &\leq (1 - s_n \tau)^2 \left[ \|\Delta_n^k x_n - p\|^2 - \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\|^2 \right. \\
 &\quad \left. + 2\lambda_{i,n} \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\| \right. \\
 &\quad \left. \times \|B_i \Lambda_n^{i-1} u_n - B_i p\| \right] \\
 &\quad + \beta_n^2 \|x_n - T_n v_n\|^2 + 2(1 - s_n \tau) \beta_n \|v_n - p\| \\
 &\quad \times \|x_n - T_n v_n\| + 2s_n \|\gamma V x_n - \mu F p\| \|x_{n+1} - p\| \\
 &\leq (1 - s_n \tau)^2 \left[ \|x_n - p\|^2 - \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\|^2 \right. \\
 &\quad \left. + 2r_{k,n} \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\| \right. \\
 &\quad \left. \times \|A_k \Delta_n^{k-1} x_n - A_k p\| \right. \\
 &\quad \left. - \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\|^2 \right. \\
 &\quad \left. + 2\lambda_{i,n} \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\| \right. \\
 &\quad \left. \times \|B_i \Lambda_n^{i-1} u_n - B_i p\| \right] \\
 &\quad + \beta_n^2 \|x_n - T_n v_n\|^2 \\
 &\quad + 2(1 - s_n \tau) \beta_n \|v_n - p\| \|x_n - T_n v_n\| \\
 &\quad + 2s_n \|\gamma V x_n - \mu F p\| \|x_{n+1} - p\| \\
 &\leq \|x_n - p\|^2 - (1 - s_n \tau)^2 \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\|^2 \\
 &\quad + 2r_{k,n} \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\| \|A_k \Delta_n^{k-1} x_n - A_k p\|
 \end{aligned}$$

$$\begin{aligned}
 & - (1 - s_n \tau)^2 \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\|^2 \\
 & + 2\lambda_{i,n} \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\| \|B_i \Lambda_n^{i-1} u_n - B_i p\| \\
 & + \beta_n^2 \|x_n - T_n v_n\|^2 + 2(1 - s_n \tau) \beta_n \|v_n - p\| \\
 & \times \|x_n - T_n v_n\| + 2s_n \|\gamma V x_n - \mu F p\| \|x_{n+1} - p\|;
 \end{aligned} \tag{148}$$

that is,

$$\begin{aligned}
 & (1 - s_n \tau)^2 \left[ \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\|^2 + \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\|^2 \right] \\
 & \leq \|x_n - p\|^2 - \|x_{n+1} - p\|^2 \\
 & + 2r_{k,n} \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\| \|A_k \Delta_n^{k-1} x_n - A_k p\| \\
 & + 2\lambda_{i,n} \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\| \|B_i \Lambda_n^{i-1} u_n - B_i p\| \\
 & + \beta_n^2 \|x_n - T_n v_n\|^2 \\
 & + 2(1 - s_n \tau) \beta_n \|v_n - p\| \|x_n - T_n v_n\| \\
 & + 2s_n \|\gamma V x_n - \mu F p\| \|x_{n+1} - p\| \\
 & \leq (\|x_n - p\| + \|x_{n+1} - p\|) \|x_n - x_{n+1}\| \\
 & + 2r_{k,n} \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\| \|A_k \Delta_n^{k-1} x_n - A_k p\| \\
 & + 2\lambda_{i,n} \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\| \|B_i \Lambda_n^{i-1} u_n - B_i p\| \\
 & + \beta_n^2 \|x_n - T_n v_n\|^2 + 2(1 - s_n \tau) \beta_n \|v_n - p\| \\
 & \times \|x_n - T_n v_n\| + 2s_n \|\gamma V x_n - \mu F p\| \|x_{n+1} - p\|.
 \end{aligned} \tag{149}$$

So, from  $s_n \rightarrow 0$ , (133), (138), and (143), we immediately get

$$\lim_{n \rightarrow \infty} \|\Lambda_n^{i-1} u_n - \Lambda_n^i u_n\| = 0, \quad \lim_{n \rightarrow \infty} \|\Delta_n^{k-1} x_n - \Delta_n^k x_n\| = 0, \tag{150}$$

for all  $i \in \{1, 2, \dots, N\}$  and  $k \in \{1, 2, \dots, M\}$ . Note that

$$\begin{aligned}
 \|x_n - u_n\| & = \|\Delta_n^0 x_n - \Delta_n^M x_n\| \\
 & \leq \|\Delta_n^0 x_n - \Delta_n^1 x_n\| + \|\Delta_n^1 x_n - \Delta_n^2 x_n\| \\
 & \quad + \dots + \|\Delta_n^{M-1} x_n - \Delta_n^M x_n\|, \\
 \|u_n - v_n\| & = \|\Lambda_n^0 u_n - \Lambda_n^N u_n\| \\
 & \leq \|\Lambda_n^0 u_n - \Lambda_n^1 u_n\| + \|\Lambda_n^1 u_n - \Lambda_n^2 u_n\| \\
 & \quad + \dots + \|\Lambda_n^{N-1} u_n - \Lambda_n^N u_n\|.
 \end{aligned} \tag{151}$$

Thus, from (150) we have

$$\lim_{n \rightarrow \infty} \|x_n - u_n\| = 0, \quad \lim_{n \rightarrow \infty} \|u_n - v_n\| = 0. \tag{152}$$

It is easy to see that as  $n \rightarrow \infty$

$$\|x_n - v_n\| \leq \|x_n - u_n\| + \|u_n - v_n\| \rightarrow 0. \tag{153}$$

Also, observe that

$$\|T_n v_n - v_n\| \leq \|T_n v_n - x_n\| + \|x_n - v_n\|. \tag{154}$$

Hence, we have from (138)

$$\lim_{n \rightarrow \infty} \|T_n v_n - v_n\| = 0. \tag{155}$$

*Step 5.* Let us show that  $\limsup_{n \rightarrow \infty} \langle (\mu F - \gamma V)q, q - x_n \rangle \leq 0$ , where  $q \in \Omega$  is the same as in Theorem 18; that is,  $q \in \Omega$  is a unique solution of VIP (50). To show this inequality, we choose a subsequence  $\{x_{n_i}\}$  of  $\{x_n\}$  such that

$$\begin{aligned}
 & \limsup_{n \rightarrow \infty} \langle (\mu F - \gamma V)q, q - x_n \rangle \\
 & = \lim_{i \rightarrow \infty} \langle (\mu F - \gamma V)q, q - x_{n_i} \rangle.
 \end{aligned} \tag{156}$$

Since  $\{x_n\}$  is bounded, there exists a subsequence  $\{x_{n_j}\}$  of  $\{x_{n_i}\}$  which converges weakly to  $w$ . Without loss of generality, we may assume that  $x_{n_i} \rightharpoonup w$ . From Step 4, we have that  $\Delta_{n_i}^k x_{n_i} \rightharpoonup w$ ,  $\Lambda_{n_i}^m u_{n_i} \rightharpoonup w$ ,  $u_{n_i} \rightharpoonup w$ , and  $v_{n_i} \rightharpoonup w$ , where  $k \in \{1, 2, \dots, M\}$ ,  $m \in \{1, 2, \dots, N\}$ . Since  $v_n - T_n v_n \rightarrow 0$  by Step 4, by the same arguments as in the proof of Theorem 18, we get  $w \in \Omega$ . Since  $q = P_\Omega(I - \mu F + \gamma V)q$ , it follows that

$$\begin{aligned}
 \limsup_{n \rightarrow \infty} \langle (\mu F - \gamma V)q, q - x_n \rangle & = \lim_{i \rightarrow \infty} \langle (\mu F - \gamma V)q, q - x_{n_i} \rangle \\
 & = \langle (\mu F - \gamma V)q, q - w \rangle \leq 0.
 \end{aligned} \tag{157}$$

*Step 6.* Let us show that  $\lim_{n \rightarrow \infty} \|x_n - q\| = 0$ , where  $q \in \Omega$  is the same as in Theorem 18; that is,  $q \in \Omega$  is a unique solution of VIP (50). From (113), we know that

$$\begin{aligned}
 x_{n+1} - q & = s_n (\gamma V x_n - \mu F q) + \beta_n (T_n v_n - q) \\
 & \quad + ((1 - \beta_n)I - s_n \mu F) T_n v_n \\
 & \quad - ((1 - \beta_n)I - s_n \mu F) q.
 \end{aligned} \tag{158}$$

Applying Lemmas 7 and 13 and noticing  $T_n q = q$  and  $\|v_n - q\| \leq \|x_n - q\|$  for all  $n \geq 1$ , we have

$$\begin{aligned}
 & \|x_{n+1} - q\|^2 \\
 & \leq \|\beta_n (T_n v_n - q) + ((1 - \beta_n)I - s_n \mu F) T_n v_n \\
 & \quad - ((1 - \beta_n)I - s_n \mu F) q\|^2 \\
 & \quad + 2s_n \langle \gamma V x_n - \mu F q, x_{n+1} - q \rangle
 \end{aligned}$$

$$\begin{aligned}
 &\leq [\beta_n \|T_n v_n - q\| + \|((1 - \beta_n)I - s_n \mu F) T_n v_n \\
 &\quad - ((1 - \beta_n)I - s_n \mu F) q\|]^2 \\
 &\quad + 2s_n \langle \gamma V x_n - \gamma V q, x_{n+1} - q \rangle \\
 &\quad + 2s_n \langle \gamma V q - \mu F q, x_{n+1} - q \rangle \\
 &= \left[ \beta_n \|T_n v_n - q\| + (1 - \beta_n) \right. \\
 &\quad \times \left\| \left( I - \frac{s_n}{1 - \beta_n} \mu F \right) T_n v_n - \left( I - \frac{s_n}{1 - \beta_n} \mu F \right) T_n q \right\|^2 \\
 &\quad + 2s_n \langle \gamma V x_n - \gamma V q, x_{n+1} - q \rangle \\
 &\quad + 2s_n \langle \gamma V q - \mu F q, x_{n+1} - q \rangle \\
 &\leq \left[ \beta_n \|v_n - q\| + (1 - \beta_n) \left( 1 - \frac{s_n \tau}{1 - \beta_n} \right) \|v_n - q\| \right]^2 \\
 &\quad + 2s_n \langle \gamma V x_n - \gamma V q, x_{n+1} - q \rangle \\
 &\quad + 2s_n \langle \gamma V q - \mu F q, x_{n+1} - q \rangle \\
 &= [\beta_n \|v_n - q\| + (1 - \beta_n - s_n \tau) \|v_n - q\|]^2 \\
 &\quad + 2s_n \langle \gamma V x_n - \gamma V q, x_{n+1} - q \rangle \\
 &\quad + 2s_n \langle \gamma V q - \mu F q, x_{n+1} - q \rangle \\
 &\leq [\beta_n \|x_n - q\| + (1 - \beta_n - s_n \tau) \|x_n - q\|]^2 \\
 &\quad + 2s_n \langle \gamma V x_n - \gamma V q, x_{n+1} - q \rangle \\
 &\quad + 2s_n \langle \gamma V q - \mu F q, x_{n+1} - q \rangle \\
 &\leq (1 - s_n \tau)^2 \|x_n - q\|^2 + 2s_n \gamma l \|x_n - q\| \\
 &\quad \times \|x_{n+1} - q\| + 2s_n \langle \gamma V q - \mu F q, x_{n+1} - q \rangle \\
 &\leq (1 - s_n \tau)^2 \|x_n - q\|^2 + s_n \gamma l (\|x_n - q\|^2 + \|x_{n+1} - q\|^2) \\
 &\quad + 2s_n \langle \gamma V q - \mu F q, x_{n+1} - q \rangle.
 \end{aligned} \tag{159}$$

This implies that

$$\begin{aligned}
 &\|x_{n+1} - q\|^2 \\
 &\leq \frac{1 - 2\tau s_n + \tau^2 s_n^2 + s_n \gamma l}{1 - s_n \gamma l} \|x_n - q\|^2 \\
 &\quad + \frac{2s_n}{1 - s_n \gamma l} \langle \gamma V q - \mu F q, x_{n+1} - q \rangle \\
 &= \left( 1 - \frac{2(\tau - \gamma l) s_n}{1 - s_n \gamma l} \right) \|x_n - q\|^2 + \frac{\tau^2 s_n^2}{1 - s_n \gamma l} \|x_n - q\|^2 \\
 &\quad + \frac{2s_n}{1 - s_n \gamma l} \langle \gamma V q - \mu F q, x_{n+1} - q \rangle
 \end{aligned}$$

$$\begin{aligned}
 &\leq \left( 1 - \frac{2(\tau - \gamma l) s_n}{1 - s_n \gamma l} \right) \|x_n - q\|^2 + \frac{2(\tau - \gamma l) s_n}{1 - s_n \gamma l} \\
 &\quad \times \left( \frac{\tau^2 s_n}{2(\tau - \gamma l)} \widetilde{M}_2 \right. \\
 &\quad \left. + \frac{1}{\tau - \gamma l} \langle \mu F q - \gamma V q, q - x_{n+1} \rangle \right) \\
 &= (1 - \sigma_n) \|x_n - q\|^2 + \sigma_n \delta_n,
 \end{aligned} \tag{160}$$

where  $\widetilde{M}_2 = \sup_{n \geq 1} \|x_n - q\|^2$ ,  $\sigma_n = (2(\tau - \gamma l)/(1 - s_n \gamma l))s_n$ , and

$$\delta_n = \frac{\tau^2 s_n}{2(\tau - \gamma l)} \widetilde{M}_2 + \frac{1}{\tau - \gamma l} \langle \mu F q - \gamma V q, q - x_{n+1} \rangle. \tag{161}$$

From condition (i) and Step 5, it is easy to see that  $\sigma_n \rightarrow 0$ ,  $\sum_{n=0}^{\infty} \sigma_n = \infty$  and  $\limsup_{n \rightarrow \infty} \delta_n \leq 0$ . Hence, by Lemma 10, we conclude that  $x_n \rightarrow q$  as  $n \rightarrow \infty$ . This completes the proof.  $\square$

**Corollary 22.** *Let  $C$  be a nonempty closed convex subset of a real Hilbert space  $H$ . Let  $f : C \rightarrow \mathbf{R}$  be a convex functional with  $L$ -Lipschitz continuous gradient  $\nabla f$ . Let  $\Theta$  be a bifunction from  $C \times C$  to  $\mathbf{R}$  satisfying (A1)–(A4) and let  $\varphi : C \rightarrow \mathbf{R} \cup \{+\infty\}$  be a proper lower semicontinuous and convex function. Let  $R_i : C \rightarrow 2^H$  be a maximal monotone mapping and let  $A : H \rightarrow H$  and  $B_i : C \rightarrow H$  be  $\zeta$ -inverse-strongly monotone and  $\eta_i$ -inverse-strongly monotone, respectively, for  $i = 1, 2$ . Let  $F : H \rightarrow H$  be a  $\kappa$ -Lipschitzian and  $\eta$ -strongly monotone operator with positive constants  $\kappa, \eta > 0$ . Let  $V : H \rightarrow H$  be an  $l$ -Lipschitzian mapping with constant  $l \geq 0$ . Let  $0 < \mu < 2\eta/\kappa^2$  and  $0 \leq \gamma l < \tau$ , where  $\tau = 1 - \sqrt{1 - \mu(2\eta - \mu\kappa^2)}$ . Assume that  $\Omega := \text{GMEP}(\Theta, \varphi, A) \cap I(B_1, R_1) \cap I(B_2, R_2) \cap \Gamma \neq \emptyset$  and that either (B1) or (B2) holds. For arbitrarily given  $x_1 \in H$ , let  $\{x_n\}$  be a sequence generated by*

$$\begin{aligned}
 &\Theta(u_n, y) + \varphi(y) - \varphi(u_n) + \langle Ax_n, y - u_n \rangle \\
 &\quad + \frac{1}{r_n} \langle y - u_n, u_n - x_n \rangle \geq 0, \quad \forall y \in C, \\
 &v_n = J_{R_2, \lambda_{2,n}} (I - \lambda_{2,n} B_2) J_{R_1, \lambda_{1,n}} (I - \lambda_{1,n} B_1) u_n, \\
 &x_{n+1} = s_n \gamma V x_n + \beta_n x_n + ((1 - \beta_n)I - s_n \mu F) T_n v_n,
 \end{aligned} \tag{162}$$

$\forall n \geq 1$ ,

where  $P_C(I - \lambda_n \nabla f) = s_n I + (1 - s_n) T_n$  (here  $T_n$  is nonexpansive and  $s_n = (2 - \lambda_n L)/4 \in (0, 1/2)$  for each  $\lambda_n \in (0, 2/L)$ ). Assume that the following conditions hold:

- (i)  $s_n \in (0, 1/2)$  for each  $\lambda_n \in (0, 2/L)$ , and  $\lim_{n \rightarrow \infty} s_n = 0$  ( $\Leftrightarrow \lim_{n \rightarrow \infty} \lambda_n = 2/L$ );
- (ii)  $\{\beta_n\} \subset (0, 1)$  and  $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$ ;
- (iii)  $\{\lambda_{i,n}\} \subset [a_i, b_i] \subset (0, 2\eta_i)$  and  $\lim_{n \rightarrow \infty} |\lambda_{i,n+1} - \lambda_{i,n}| = 0$  for  $i = 1, 2$ ;
- (iv)  $\{r_n\} \subset [e, f] \subset (0, 2\zeta)$  and  $\lim_{n \rightarrow \infty} |r_{n+1} - r_n| = 0$ .



Then  $\{x_n\}$  converges strongly as  $\lambda_n \rightarrow 2/L (\Leftrightarrow s_n \rightarrow 0)$  to a point  $q \in \Omega$ , which is a unique solution of VIP (110).

**Corollary 23.** Let  $C$  be a nonempty closed convex subset of a real Hilbert space  $H$ . Let  $f : C \rightarrow \mathbf{R}$  be a convex functional with  $L$ -Lipschitz continuous gradient  $\nabla f$ . Let  $\Theta$  be a bifunction from  $C \times C$  to  $\mathbf{R}$  satisfying (A1)–(A4) and let  $\varphi : C \rightarrow \mathbf{R} \cup \{+\infty\}$  be a proper lower semicontinuous and convex function. Let  $R : C \rightarrow 2^H$  be a maximal monotone mapping and let  $A : H \rightarrow H$  and  $B : C \rightarrow H$  be  $\zeta$ -inverse-strongly monotone and  $\xi$ -inverse-strongly monotone, respectively. Let  $F : H \rightarrow H$  be a  $\kappa$ -Lipschitzian and  $\eta$ -strongly monotone operator with positive constants  $\kappa, \eta > 0$ . Let  $V : H \rightarrow H$  be an  $l$ -Lipschitzian mapping with constant  $l \geq 0$ . Let  $0 < \mu < 2\eta/\kappa^2$  and  $0 \leq \gamma l < \tau$ , where  $\tau = 1 - \sqrt{1 - \mu(2\eta - \mu\kappa^2)}$ . Assume that  $\Omega := \text{GMEP}(\Theta, \varphi, A) \cap I(B, R) \cap \Gamma \neq \emptyset$  and that either (B1) or (B2) holds. For arbitrarily given  $x_1 \in H$ , let  $\{x_n\}$  be a sequence generated by

$$\begin{aligned} & \Theta(u_n, y) + \varphi(y) - \varphi(u_n) + \langle Ax_n, y - u_n \rangle \\ & + \frac{1}{r_n} \langle y - u_n, u_n - x_n \rangle \geq 0, \quad \forall y \in C, \\ x_{n+1} & = s_n \gamma Vx_n + \beta_n x_n \\ & + ((1 - \beta_n)I - s_n \mu F)T_n J_{R, \rho_n}(u_n - \rho_n B u_n), \quad \forall n \geq 1, \end{aligned} \tag{163}$$

where  $P_C(I - \lambda_n \nabla f) = s_n I + (1 - s_n)T_n$  (here  $T_n$  is nonexpansive and  $s_n = (2 - \lambda_n L)/4 \in (0, 1/2)$  for each  $\lambda_n \in (0, 2/L)$ ). Assume that the following conditions hold:

- (i)  $s_n \in (0, 1/2)$  for each  $\lambda_n \in (0, 2/L)$ , and  $\lim_{n \rightarrow \infty} s_n = 0 (\Leftrightarrow \lim_{n \rightarrow \infty} \lambda_n = 2/L)$ ;
- (ii)  $\{\beta_n\} \subset (0, 1)$  and  $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$ ;
- (iii)  $\{\rho_n\} \subset [a, b] \subset (0, 2\xi)$  and  $\lim_{n \rightarrow \infty} |\rho_{n+1} - \rho_n| = 0$ ;
- (iv)  $\{r_n\} \subset [e, f] \subset (0, 2\zeta)$  and  $\lim_{n \rightarrow \infty} |r_{n+1} - r_n| = 0$ .

Then  $\{x_n\}$  converges strongly as  $\lambda_n \rightarrow 2/L (\Leftrightarrow s_n \rightarrow 0)$  to a point  $q \in \Omega$ , which is a unique solution of VIP (112).

**Conflict of Interests**

The authors declare that there is no conflict of interests regarding the publication of this paper.

**Acknowledgments**

In this research, the first author was supported by King Fahd University of Petroleum and Minerals Project no. IN111015; he also is grateful to the university for providing excellent research facilities. This research was partially supported by the National Science Foundation of China (11071169), Innovation Program of Shanghai Municipal Education Commission (09ZZ133), and Ph.D. Program Foundation of Ministry of Education of China (20123127110002).

**References**

- [1] F. E. Browder and W. V. Petryshyn, “Construction of fixed points of nonlinear mappings in Hilbert space,” *Journal of Mathematical Analysis and Applications*, vol. 20, pp. 197–228, 1967.
- [2] J. L. Lions, *Quelques Méthodes de Résolution des Problèmes aux Limites Non Lineaires*, Dunod, Paris, France, 1969.
- [3] R. Glowinski, *Numerical Methods for Nonlinear Variational Problems*, Springer, New York, NY, USA, 1984.
- [4] W. Takahashi, *Nonlinear Functional Analysis*, Yokohama Publishers, Yokohama, Japan, 2000.
- [5] J. T. Oden, *Quantitative Methods on Nonlinear Mechanics*, Prentice-Hall, Englewood Cliffs, NJ, USA, 1986.
- [6] E. Zeidler, *Nonlinear Functional Analysis and Its Applications*, Springer, New York, NY, USA, 1985.
- [7] G. M. Korpelevič, “An extragradient method for finding saddle points and for other problems,” *Èkonomika i Matematicheskie Metody*, vol. 12, no. 4, pp. 747–756, 1976.
- [8] F. Facchinei and J.-S. Pang, *Finite-Dimensional Variational Inequalities and Complementarity Problems*, vol. 1-2, Springer, New York, NY, USA, 2003.
- [9] N. Nadezhkina and W. Takahashi, “Weak convergence theorem by an extragradient method for nonexpansive mappings and monotone mappings,” *Journal of Optimization Theory and Applications*, vol. 128, no. 1, pp. 191–201, 2006.
- [10] L.-C. Zeng and J.-C. Yao, “Strong convergence theorem by an extragradient method for fixed point problems and variational inequality problems,” *Taiwanese Journal of Mathematics*, vol. 10, no. 5, pp. 1293–1303, 2006.
- [11] Y. Yao, Y. J. Cho, and Y.-C. Liou, “Algorithms of common solutions for variational inclusions, mixed equilibrium problems and fixed point problems,” *European Journal of Operational Research*, vol. 212, no. 2, pp. 242–250, 2011.
- [12] J.-W. Peng and J.-C. Yao, “A new hybrid-extragradient method for generalized mixed equilibrium problems, fixed point problems and variational inequality problems,” *Taiwanese Journal of Mathematics*, vol. 12, no. 6, pp. 1401–1432, 2008.
- [13] L.-C. Ceng, C.-Y. Wang, and J.-C. Yao, “Strong convergence theorems by a relaxed extragradient method for a general system of variational inequalities,” *Mathematical Methods of Operations Research*, vol. 67, no. 3, pp. 375–390, 2008.
- [14] L.-C. Ceng, Q. H. Ansari, and S. Schaible, “Hybrid extragradient-like methods for generalized mixed equilibrium problems, systems of generalized equilibrium problems and optimization problems,” *Journal of Global Optimization*, vol. 53, no. 1, pp. 69–96, 2012.
- [15] L.-C. Ceng, Q. H. Ansari, M. M. Wong, and J.-C. Yao, “Mann type hybrid extragradient method for variational inequalities, variational inclusions and fixed point problems,” *Fixed Point Theory*, vol. 13, no. 2, pp. 403–422, 2012.
- [16] L.-C. Ceng, Q. H. Ansari, and J.-C. Yao, “An extragradient method for solving split feasibility and fixed point problems,” *Computers & Mathematics with Applications*, vol. 64, no. 4, pp. 633–642, 2012.
- [17] L.-C. Ceng, Q. H. Ansari, and J.-C. Yao, “Relaxed extragradient methods for finding minimum-norm solutions of the split feasibility problem,” *Nonlinear Analysis: Theory, Methods & Applications*, vol. 75, no. 4, pp. 2116–2125, 2012.
- [18] L.-C. Ceng, N. Hadjisavvas, and N.-C. Wong, “Strong convergence theorem by a hybrid extragradient-like approximation

- method for variational inequalities and fixed point problems,” *Journal of Global Optimization*, vol. 46, no. 4, pp. 635–646, 2010.
- [19] L. C. Ceng, M. Teboulle, and J. C. Yao, “Weak convergence of an iterative method for pseudomonotone variational inequalities and fixed-point problems,” *Journal of Optimization Theory and Applications*, vol. 146, no. 1, pp. 19–31, 2010.
- [20] L.-C. Ceng, Q. H. Ansari, and J.-C. Yao, “Relaxed extragradient iterative methods for variational inequalities,” *Applied Mathematics and Computation*, vol. 218, no. 3, pp. 1112–1123, 2011.
- [21] L.-C. Ceng, Q. H. Ansari, N.-C. Wong, and J.-C. Yao, “An extragradient-like approximation method for variational inequalities and fixed point problems,” *Fixed Point Theory and Applications*, vol. 2011, article 22s, 18 pages, 2011.
- [22] L.-C. Ceng, S.-M. Guu, and J.-C. Yao, “Finding common solutions of a variational inequality, a general system of variational inequalities, and a fixed-point problem via a hybrid extragradient method,” *Fixed Point Theory and Applications*, Article ID 626159, 22 pages, 2011.
- [23] L.-C. Ceng and A. Petruşel, “Relaxed extragradient-like method for general system of generalized mixed equilibria and fixed point problem,” *Taiwanese Journal of Mathematics*, vol. 16, no. 2, pp. 445–478, 2012.
- [24] Y. Yao, Y.-C. Liou, and S. M. Kang, “Approach to common elements of variational inequality problems and fixed point problems via a relaxed extragradient method,” *Computers & Mathematics with Applications*, vol. 59, no. 11, pp. 3472–3480, 2010.
- [25] L. C. Ceng, H.-Y. Hu, and M. M. Wong, “Strong and weak convergence theorems for generalized mixed equilibrium problem with perturbation and fixed pointed problem of infinitely many nonexpansive mappings,” *Taiwanese Journal of Mathematics*, vol. 15, no. 3, pp. 1341–1367, 2011.
- [26] L.-C. Ceng and J.-C. Yao, “A hybrid iterative scheme for mixed equilibrium problems and fixed point problems,” *Journal of Computational and Applied Mathematics*, vol. 214, no. 1, pp. 186–201, 2008.
- [27] L.-C. Ceng, S.-M. Guu, and J.-C. Yao, “Hybrid iterative method for finding common solutions of generalized mixed equilibrium and fixed point problems,” *Fixed Point Theory and Applications*, vol. 2012, article 92, 19 pages, 2012.
- [28] S. Takahashi and W. Takahashi, “Strong convergence theorem for a generalized equilibrium problem and a nonexpansive mapping in a Hilbert space,” *Nonlinear Analysis: Theory, Methods & Applications*, vol. 69, no. 3, pp. 1025–1033, 2008.
- [29] L. C. Ceng, A. Petruşel, and J. C. Yao, “Iterative approaches to solving equilibrium problems and fixed point problems of infinitely many nonexpansive mappings,” *Journal of Optimization Theory and Applications*, vol. 143, no. 1, pp. 37–58, 2009.
- [30] I. Yamada, “The hybrid steepest-descent method for the variational inequality problems over the intersection of the fixed-point sets of nonexpansive mappings,” in *Inherently Parallel Algorithms in Feasibility and Optimization and Their Applications*, D. Batnariu, Y. Censor, and S. Reich, Eds., pp. 473–504, North-Holland, Amsterdam, The Netherlands, 2001.
- [31] V. Colao, G. Marino, and H.-K. Xu, “An iterative method for finding common solutions of equilibrium and fixed point problems,” *Journal of Mathematical Analysis and Applications*, vol. 344, no. 1, pp. 340–352, 2008.
- [32] R. T. Rockafellar, “Monotone operators and the proximal point algorithm,” *SIAM Journal on Control and Optimization*, vol. 14, no. 5, pp. 877–898, 1976.
- [33] N.-J. Huang, “A new completely general class of variational inclusions with noncompact valued mappings,” *Computers & Mathematics with Applications*, vol. 35, no. 10, pp. 9–14, 1998.
- [34] L.-C. Zeng, S.-M. Guu, and J.-C. Yao, “Characterization of  $H$ -monotone operators with applications to variational inclusions,” *Computers & Mathematics with Applications*, vol. 50, no. 3-4, pp. 329–337, 2005.
- [35] L. C. Ceng, S. M. Guu, and J. C. Yao, “Weak convergence theorem by a modified extragradient method for variational inclusions, variational inequalities and fixed point problems,” *Journal of Nonlinear and Convex Analysis*, vol. 14, pp. 21–31, 2013.
- [36] Y.-P. Fang and N.-J. Huang, “ $H$ -accretive operators and resolvent operator technique for solving variational inclusions in Banach spaces,” *Applied Mathematics Letters*, vol. 17, no. 6, pp. 647–653, 2004.
- [37] S.-S. Zhang, J. H. W. Lee, and C. K. Chan, “Algorithms of common solutions to quasi variational inclusion and fixed point problems,” *Applied Mathematics and Mechanics (English Edition)*, vol. 29, no. 5, pp. 571–581, 2008.
- [38] J.-W. Peng, Y. Wang, D. S. Shyu, and J.-C. Yao, “Common solutions of an iterative scheme for variational inclusions, equilibrium problems, and fixed point problems,” *Journal of Inequalities and Applications*, vol. 2008, Article ID 720371, 15 pages, 2008.
- [39] L.-C. Ceng, Q. H. Ansari, M. M. Wong, and J.-C. Yao, “Mann type hybrid extragradient method for variational inequalities, variational inclusions and fixed point problems,” *Fixed Point Theory*, vol. 13, no. 2, pp. 403–422, 2012.
- [40] H.-K. Xu, “Averaged mappings and the gradient-projection algorithm,” *Journal of Optimization Theory and Applications*, vol. 150, no. 2, pp. 360–378, 2011.
- [41] J.-B. Baillon and G. Haddad, “Quelques propriétés des opérateurs angle-bornés et  $n$ -cycliquement monotones,” *Israel Journal of Mathematics*, vol. 26, no. 2, pp. 137–150, 1977.
- [42] C. Byrne, “A unified treatment of some iterative algorithms in signal processing and image reconstruction,” *Inverse Problems*, vol. 20, no. 1, pp. 103–120, 2004.
- [43] P. L. Combettes, “Solving monotone inclusions via compositions of nonexpansive averaged operators,” *Optimization*, vol. 53, no. 5-6, pp. 475–504, 2004.
- [44] K. Geobel and W. A. Kirk, *Topics on Metric Fixed-Point Theory*, Cambridge University Press, Cambridge, UK, 1990.
- [45] H. K. Xu and T. H. Kim, “Convergence of hybrid steepest-descent methods for variational inequalities,” *Journal of Optimization Theory and Applications*, vol. 119, no. 1, pp. 185–201, 2003.
- [46] T. Suzuki, “Strong convergence of Krasnoselskii and Mann’s type sequences for one-parameter nonexpansive semigroups without Bohnner integrals,” *Journal of Mathematical Analysis and Applications*, vol. 305, no. 1, pp. 227–239, 2005.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

