

Research Article

Antiageing Mechanisms of a Standardized Supercritical CO₂ Preparation of Black Jack (*Bidens pilosa* L.) in Human Fibroblasts and Skin Fragments

Gustavo Dieamant,¹ Maria Del Carmen V. Pereda,¹ Cecília Nogueira,¹ Samara Eberlin,¹ Gustavo Facchini,¹ Juliana Tibério Checon,² Camila Kappke Cesar,¹ Lilian Mussi,¹ Márcio Antonio Polezel,¹ Divino Martins-Oliveira Jr.,¹ and Luiz Claudio Di Stasi²

¹Chemyunion Química Ltda, Avenida Independência 1501, 18087-101 Sorocaba, SP, Brazil

²Laboratory of Phytomedicines, Pharmacology and Biotechnology (PhytoPharmaTech), Department of Pharmacology, Institute of Biosciences, Universidade Estadual Paulista (UNESP), 18618-000 Botucatu, SP, Brazil

Correspondence should be addressed to Luiz Claudio Di Stasi; ldistasi@ibb.unesp.br

Received 8 December 2014; Revised 23 February 2015; Accepted 15 March 2015

Academic Editor: Alessandra Guerrini

Copyright © 2015 Gustavo Dieamant et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The use of topical retinoids to treat skin disorders and ageing can induce local reactions, while oral retinoids are potent teratogens and produce several unwanted effects. This way, efforts to explore complementary care resources should be supported. Based on this, we evaluate the antiageing effects of a supercritical CO₂ extract from *Bidens pilosa* L. (BPE-CO₂A) containing a standardized multicomponent mixture of phytol, linolenic, palmitic, linoleic, and oleic acids. BPE-CO₂A was assessed for its effects on human dermal fibroblasts (TGF- β 1 and FGF levels using ELISA; collagen, elastin, and glycosaminoglycan by colorimetric assays, and mRNA expression of RXR, RAR, and EGFR by qRT-PCR) and human skin fragments (RAR, RXR, collagen, elastin, and glycosaminoglycan by immunohistochemical analysis). Levels of extracellular matrix elements, TGF- β 1 and FGF, and EGFR gene expression were significantly increased by BPE-CO₂A. The modulation of RXR and RAR was positively demonstrated after the treatment with BPE-CO₂A or phytol, a component of BPE-CO₂A. The effects produced by BPE-CO₂A were similar to or better than those produced by retinol and retinoic acid. The ability to stimulate extracellular matrix elements, increase growth factors, and modulate retinoid and retinoid receptors provides a basis for the development of preparation containing BPE-CO₂A as an antiageing/skin-repair agent.

1. Introduction

Retinoids are classically defined as compounds that exhibit vitamin A-like effects or bind to nuclear retinoid receptors, exerting their pharmacological effects on gene expression by activating retinoic acid receptors (RARs) or retinoid X receptors (RXRs). RARs are ligand-controlled transcription factors that function as heterodimers with RXRs to regulate cell growth, differentiation, survival, and death [1]. Activation of RAR and RXR has been associated with several diseases, and their ligands are widely used to treat skin disorders, mainly aging and photoaging, acne, and psoriasis, and have been used for cancer therapy and chemoprevention [2, 3].

The beneficial effects of retinoids on ageing skin are attributed to increased dermal function, mainly through the increased production of extracellular matrix components, reduced inflammatory response, and antioxidative properties. These effects are related to epidermal growth factor (EGF) and fibroblast growth factor (FGF), both acting as mediators of the retinoid response in the skin [4]. In addition, the beneficial effects of retinoids on ageing and photoageing include improved fine wrinkling, diminished tactile roughness, improved actinic keratosis, and reduced hyperpigmentation, which are histopathologically associated with epidermal hyperplasia, the compaction of the stratum corneum, the thickening of the granular layer, reduced melanocytic

hypertrophy, the restoration of cell polarity, increased angiogenesis, increased new collagen formation, and the normalization of the appearance of elastic tissue [5].

In contrast, the use of topical retinoids to treat skin disorders can induce local reactions including erythema, burning, dryness, desquamation, stinging, and photosensitivity reactions to ultraviolet radiation, whereas the use of oral retinoids is limited because these compounds are potent teratogens and produce several unwanted effects such as cheilitis, cutaneous photosensitivity, conjunctivitis, photophobia, and increased susceptibility to staphylococcal infections [4]. This way, efforts to explore alternative and complementary care resources should be supported [6], and the use of herbal preparations and natural products as a source of new active compounds for complementary and alternative therapies could be an important therapeutic strategy to obtain new products with retinoid-like effects and better management of skin diseases and ageing.

Black Jack (*Bidens pilosa* L. (Asteraceae)) is an annual and ruderal plant originating in South America and is also found in tropical and subtropical regions around the world [7]. This plant is a cosmopolitan herb with high distribution in disturbed areas and is widely used in the traditional medicine of different countries to treat internal and topical inflammatory processes, wounds, insect bites, fungal infections, diabetic ulcers, fever, malaria, inflammation, hepatitis, hemorrhoids, and cancer [7–11]. Recent review showed that *B. pilosa* has anti-inflammatory, immunomodulatory, antimalarial, antitumor, antioxidative, antiulcerogenic, antibacterial, hepatoprotective, and antihypertensive activities [7, 8]. Phytochemical studies indicated that approximately 200 different compounds have been identified and isolated in *B. pilosa*, and among them mainly flavonoids, polyacetylenes, terpenoids, phenylpropanoids, and hydrocarbons were reported [7, 8]. It has been demonstrated that nonpolar chemical constituents of *Bidens pilosa* such as docosahexaenoic acid, phytanic acid, phytol, α -tocopherol, ricinoleic acid, and esters of caffeic acid have similar retinoid chemical structures, and some of these compounds have been considered to be natural RXR ligands and modulators of the retinoic acid signalling pathway, directly influencing their transcriptional regulatory activity [12–16].

The study of new products from natural sources, mainly plant species, can be performed through several approaches; however, the supercritical carbon dioxide extraction and fractionation (SC-CO₂) of natural matter is one of the early and most studied applications with immediate advantages over traditional extraction techniques [17]. It is a flexible process because it allows for the continuous modulation of the solvent power/selectivity and elimination of polluting organic solvents, thereby eliminating the expensive postprocessing of extracts traditionally required for solvent elimination. Moreover, it is more economical and simple than other methods; it has diffusivity that reduces mass transfer limitations and a low surface tension, which allows for the penetration and wetting of pores smaller than those accessible with liquid solvents; it allows extraction at low to moderate temperatures, leaving no solvent residues; and it is environmentally acceptable [17, 18].

Based on the pharmacological properties of *B. pilosa*, such as antioxidative, anti-inflammatory and retinoid-like effects, and its phytochemical composition of mainly nonpolar compounds, the aim of this study was to evaluate a SC-CO₂ nonpolar extract from the aerial parts of *B. pilosa* to develop an herbal preparation with retinoid-like activity and to potentially use this preparation as an antiageing or skin-repair agent.

2. Materials and Methods

2.1. Plant Collection, Plant Extraction, and Gas Chromatography Analysis. *B. pilosa* was cultivated using organic agricultural methods as certified by Ecocert Brazil (Santa Rosa de Lima/Santa Catarina/Br) and submitted to taxonomic identification at Herbarium Irina Gemtchujnikov (Department of Botany, Institute of Biosciences, Universidade Estadual Paulista (UNESP), Botucatu, SP), where a voucher specimen was deposited. The aerial parts were collected, dehydrated in hothouse with air circulation and renewal, and triturated in an industrial mill. A supercritical extraction system (Autoclave Engineers) under the following conditions of 300 bar, 40°C, and CO₂ flux of 5 L/min was used to generate a nonpolar extract named BPE-CO₂A.

The extraction procedures of BPE-CO₂A by GC/MS analysis were made according to following conditions: 100 mg of BPE-CO₂-A extract with 5 mL of NaOH 0.5 M in methanol was heated in a water bath for 10 min at 37°C. After cooling, 5 mL of NH₄Cl was added in methanol and heated in a water bath for 5 min at 37°C. 1 mL of BPE-CO₂A was extracted with 5 mL hexane. After cooling, 1 μ L of sample was injected in GC equipment.

GC/MS analysis was performed according to the following conditions: the gas chromatography-mass spectroscopy analysis system consisted of a Gas Chromatograph Focus (Thermo Scientific) equipped with an automatic Triplus sampler and coupled to an ISQ 230ST (Thermo Scientific) mass selective detector. The GC was fitted with an OV-5MS fused silica capillary column (30 m \times 0.25 mm \times 0.24 μ m) and helium was used as the carrier gas at 1 mL/min. Injection was made in splitless mode with injection volume of 1.0 mL/min following a split mode (1:50). The injector temperature was 260°C and detector temperature was 200°C. Temperature of column was initially 35°C and then increased to 195°C at 25°C/min, to 205°C at 3°C/min, and finally to 230°C at 8°C/min. Xcalibur software, version 2.1 (Thermo Finigan), was used to acquire and process spectrometric data. Identification of compounds was based on the retention time using NIST08 libraries and comparing with mass spectrum from scientific literature.

2.2. Cell Culture. Primary human adult keratinocytes (Invitrogen) and normal human dermal fibroblasts (Cambrex/Lonza) were commercially obtained, grown in specific culture media (keratinocyte growth medium, Epilife-Invitrogen; fibroblast basal medium, Cambrex/Lonza), and cultured in a humidified environment at 37°C and 5% CO₂. After reaching confluence, the cells were trypsinised, and the total number of

cells was counted using a Neubauer camera. After counting, the cells were seeded in culture plates of 24 or 96 wells at densities of 2×10^5 cells/well and 1×10^4 cells/well, respectively.

The apolar extract of *Bidens pilosa* (BPE-CO₂A) obtained using supercritical fluid extraction was added to the cell cultures at concentrations of 0.05, 0.10, and 0.20 mg/mL. The selection of these doses was based on preliminary cytotoxicity tests using the XTT method (data not shown). As a positive control for comparison with retinoids, we used retinol and retinoic acid at a concentration of 10 mM, according to previous data [19, 20]. After 48-hour incubation, the supernatant and cell lysate were collected for the subsequent measurement of the proposed parameters.

2.3. Specimen Skin Collection and Sample Preparation. Human eyelids were surgically removed, dipped in 70% ethanol for 15 seconds, and rinsed 2 times with saline solution. The skin fragments were transferred to a petri dish containing culture medium RPMI-1640 supplemented with foetal bovine serum-Lonza, a combination of antibiotics containing gentamicin and amphotericin-B (gentamicin sulfate/amphotericin-B, Lonza), bovine insulin (Lonza), and growth factor (rhFGF b-r-human fibroblast growth factor-B, Lonza) for a maximum of 24 hours before treatment with BPE-CO₂A.

BPE-CO₂A was dissolved in culture medium at 0.5% (a commonly used concentration for retinol) and applied to the skin fragment homogeneously over the entire surface. As positive controls for comparison, we used retinol and retinoic acid, 30 μ M and 0.5%, respectively. The samples were kept in contact with the cultures for 48 hours in a humidified atmosphere at 37°C and 5% CO₂.

2.4. TGF- β and FGF Evaluation. Transforming growth factor beta (TGF- β) and fibroblast growth factor (FGF) were measured in fibroblast culture supernatant according to the instructions for the commercially obtained ELISA kits (R&D Systems), and the concentrations of both growth factors were calculated with reference to a standard curve generated by known concentrations.

2.5. Collagen, Elastin, and Glycosaminoglycan Evaluation. Collagen, elastin, and glycosaminoglycan were quantified in the supernatant of fibroblast cultures using commercial colorimetric kits (Bicolor). All extracellular matrix elements were quantified using a standard curve.

2.6. RXR, RAR, and EGFR Gene Expression. After human fibroblasts were incubated with predetermined concentrations of BPE-CO₂A for 48 hours, total RNA was extracted using TriReagent Solution (Applied Biosystems) and quantified using a Quant-iT RNA Assay Kit (Invitrogen) and a Qubit Fluorometer (Invitrogen). The tests were conducted in a StepOnePlus sequence detection system (Applied Biosystems).

The gene expression of RAR, RXR, and EGFR was evaluated using a commercially available kit (TaqMan RNA-to-CT 1-Step, Applied Biosystems) and probes (TaqMan Gene

Expression Assays: RAR: Hs0023097_m1; RXR: Hs01067635_m1; EGFR: Hs01075087_m1; B2M: Hs00984230_m1, Applied Biosystems). The B2M (beta-2-microglobulin) gene was used as a reference (endogenous control). The RT-PCR conditions were 48°C for 15 min for reverse transcription and 95°C for 10 minutes for the activation of the Ultra-Pure AmpliTaq Gold DNA Polymerase, followed by 40 cycles of 94°C for 15 seconds and 60°C for 1 minute for denaturation and annealing, respectively.

The relative amount of mRNA was calculated with the $2^{-\Delta\Delta CT}$ method. Gene expression was considered significant when the expression values were greater than 1.5 times compared to the control. For expression inhibition, values less than 0.5-fold were considered relevant.

2.7. Immunohistochemical Assessment in Human Skin Fragments. After 48 hours of incubation with concentrations of BPE-CO₂A and phytol, *ex vivo* skin fragments were fixed in 4% paraformaldehyde (pH 7.4) for 24 hours and cryoprotected in a 30% sucrose solution for 72 hours. Then, serial sections of 10 μ m were collected directly on silanised slides with a cryostat (Leica CN1850).

After the sections were collected, they were washed with PBS. Endogenous peroxidase activity and nonspecific sites were blocked as directed by the Histostain-SP Kit (Zymed Laboratories). Then, the primary anti-RAR, anti-RXR, anti-collagen, anti-elastin, and anti-glycosaminoglycan antibodies were diluted in PBS buffer pH 7.4 containing BSA (5%) and incubated overnight.

The incubation with secondary antibodies, the amplification of immunoreactivity using an avidin-biotin complex, the revelation of the staining using a hydrogen peroxide reaction, and the mounting of the slides were performed following the Histostain-SP Kit instructions.

2.8. Histological Sections and Haematoxylin-Eosin Staining. Skin samples were fixed in 10% buffered formalin. After fixation, the material was embedded in paraffin, and serial sections approximately 5 μ m thick were made and placed on glass slides. The sections were stained with hematoxylin and eosin (HE). The parameters evaluated were visual qualitative epidermal thickness, the number of viable keratinocytes (basal layer, granular layer, and spinous layer), and barrier condition using the integrity of the stratum corneum.

2.9. Ethical Aspects. This study was conducted in accordance with Brazilian law, based on the "Normative Resolution 196/96 CNS/MS," with the consent of the Committee of Ethics in Research of the School of Medical Sciences, State University of Campinas (UNICAMP). For these experiments, we used skin fragments obtained from blepharoplasty performed at the UNICAMP Hospital.

2.10. Statistical Analysis. A parametric method of analysis of variance (ANOVA) was used for statistical analysis followed by Tukey's multiple comparison test. In all groups studied, only those with *P* values less than 0.05 were considered statistically significant.




FIGURE 1: Total ion chromatogram of BPE-CO₂A extract. Identified compounds are described in Table 1.

3. Results

3.1. Chemical Analysis of BPE-CO₂A by GC/MS. In the total ion chromatogram of BPE-CO₂A (Figure 1) the presence of several compounds such as phytol (0.139%) and fatty acids such as palmitic (30%), oleic (27%), linoleic (24.3%), and linolenic (3.8%) acids was observed. In addition the following was detected in these plant extracts: thirteen alkanes (C₁₁, C₁₄, C₁₈, C₂₀, C₂₂, C₂₃, C₂₄, C₂₅, C₂₆, C₂₇, C₂₈, C₂₉, and C₃₀), two ethyl esters of fatty acids (ethyl hexadecanoate and ethyl 9,12-octadecadienoate), two sterols (stigmasterol and sitosterol), and six unidentified compounds (Table 1).

3.2. BPE-CO₂A Increases Extracellular Matrix Elements. To assess the retinoid-like activity of the *B. pilosa* extract, we analysed the synthesis of the collagen, elastin, and glycosaminoglycans in cultured human fibroblasts treated with various concentrations of *B. pilosa* extract, retinoic acid (10 μM), or retinol (10 μM). After treatment with BPE-CO₂A, collagen levels were elevated by 32.5% at 0.05 mg/mL, 23.3% at 0.1 mg/mL, and 21.6% at 0.2 mg/mL. For comparison, retinoic acid (10 μM) and retinol (10 μM) were able to increase collagen levels by 62.9% and 45.6%, respectively (Figure 2). BPE-CO₂A was also able to significantly increase the synthesis of elastin *in vitro*. At 0.2 mg/mL, the extract caused a 25.2% increase in elastin levels (Figure 2). As expected, retinoic acid (10 μM) and retinol (10 μM) increased elastin production by 66.7% and 16.7%, respectively. In contrast, BPE-CO₂A was unable to significantly alter the levels of glycosaminoglycans (data not shown). Corroborating with the *in vitro* results, BPE-CO₂A increased the immunofluorescent staining for collagen and elastin (Figures 2(b)(A) and 2(b)(B)). In addition, we also observed an increase in immunofluorescent staining for glycosaminoglycan when skin fragments were treated with BPE-CO₂A (Figure 2(b)(C)).

3.3. BPE-CO₂A Stimulates Growth Factors. In another set of experiments, we evaluated the effects of BPE-CO₂A on TGF-β1 and FGF levels in cultured human fibroblasts as well as

the modulation of epidermal growth factor receptor (EGFr). At the three concentrations tested, BPE-CO₂A significantly increased TGF-β1 levels, with its highest stimulatory effect (53.2%) at a concentration of 0.2 mg/mL (Figure 3). In the same assay, retinoic acid (10 μM) increased TGF-β1 synthesis by 101.3%, and retinol was ineffective (Figure 3). Similar to its TGF-β1 induction, BPE-CO₂A had a significant stimulatory effect on FGF levels (Figure 3) with an increase of 188.3% after the treatment of cultured human fibroblasts with BPE-CO₂A at 0.2 mg/mL. Retinoic acid (10 μM) and retinol (10 μM) also caused significant FGF increases, but these effects were lower than those produced by BPE-CO₂A (Figure 3). In addition, we evaluated the modulation of epidermal growth factor receptor (EGFr) gene expression. BPE-CO₂A produced a 2.6-, 2.3-, and 2-fold increase in EGFr gene expression at concentration of 0.05 mg/mL, 0.10 mg/mL, and 0.2 mg/mL, respectively. Retinoic acid increased EGFr mRNA levels by 52.3%, and retinol was not able to significantly increase the expression of EGFr (Figure 3).

3.4. BPE-CO₂A Modulates Retinoic Receptors. To evaluate whether the effects induced by BPE-CO₂A have similar effects to retinoic acid and retinol in the modulation of receptors RXR and RAR, we evaluated the ability of the extract to modulate the gene expression of these receptors in human fibroblasts (Figures 4 and 5). Gene expression of RXR increased 3.6-, 3.7-, and 1.6-fold in the group treated with BPE-CO₂A at the concentration of 0.05, 0.10, and 0.20 mg/mL, respectively, whereas retinoic acid and retinol increased gene expression 1.5-fold (Figure 4(a)). An increase in receptor staining in the immunohistochemical evaluation of RXR performed on human skin fragments corroborates the *in vitro* results (Figure 4(b)).

In contrast, the expression of RAR was not significantly changed after treatment with BPE-CO₂A, retinoic acid, or retinol (Figure 5(a)). However, when we evaluated RAR protein synthesis using immunohistochemistry, we found that treatment with retinoic acid markedly increased RAR staining, whereas treatment with BPE-CO₂A and retinol promoted only subtle stimulation (Figure 5(b)).

We also evaluated the effect of phytol, an acyclic isoprenoid compound present in BPE-CO₂A, on RAR and RXR production in skin fragments. Phytol (0.0002%) had similar effects to BPE-CO₂A, inducing RXR synthesis (Figure 6) and not altering RAR production (Figure 6). It is thus possible that the effect of BPE-CO₂A on retinoid receptors might be due to the presence of phytol.

3.5. BPE-CO₂A Improves Histological Characteristics. Haematoxylin-eosin staining of human skin fragments was performed after various treatments with retinoic acid, retinol, and BPE-CO₂A (Figure 7). A comparative analysis showed that the control group has a viable epidermal structure but a less cohesive and thinner stratum corneum than the treated groups. In this group, some vacuoles were observed, resulting in a less dense dermis. Retinoic acid exhibits its classical effects, such as the smooth peeling that improves the characteristics of the epidermal basal layer (viable cells),

TABLE 1: Chemical compounds identified in the BPE-CO₂ extract by gas chromatography and mass spectroscopy (GC/MS) according to chromatogram from Figure 1.

Retention time (min)	Compounds	Molecular formula
9.83	Undecane	C ₁₁
14.10	Tetradecane	C ₁₄
15.35	Unidentified	
16.59	Unidentified	
17.38	Unidentified	
18.80	Octadecane	C ₁₈
20.45	Hexadecanoic acid (palmitic acid)	C ₁₆ H ₃₂ O ₂
20.77	Ethyl hexadecanoate	C ₁₈ H ₃₆ O ₂
20.82	Eicosane	C ₂₀
21.93	3,7,11,15-Tetramethyl-2-hexadecen-1-ol (phytol)	C ₂₀ H ₄₀ O
22.08	9,12-Octadecadienoic acid (linoleic acid)	C ₁₈ H ₃₂ O ₂
22.13	9-Octadecenoic acid (oleic acid)	C ₁₈ H ₃₄ O ₂
22.31	Octadecanoic acid (stearic acid)	C ₁₈ H ₃₆ O ₂
22.36	Ethyl 9,12-octadecadienoate	C ₂₀ H ₃₆ O ₂
22.41	9,12,15-Octadecatrienoic acid (linolenic acid)	C ₁₈ H ₃₀ O ₂
22.61	Unidentified	
22.66	Docosane	C ₂₂
23.53	Tricosane	C ₂₃
24.37	Tetracosane	C ₂₄
25.16	Pentacosane	C ₂₅
25.83	Unidentified	
25.93	Hexacosane	C ₂₆
26.66	Heptacosane	C ₂₇
27.38	Octacosane	C ₂₈
28.05	Nonacosane	C ₂₉
28.75	Triacotane	C ₃₀
29.49	Unidentified	
31.35	Stigmasterol	C ₂₉ H ₄₈ O
31.98	Sitosterol	C ₂₉ H ₅₀ O

after a single application. The group treated with retinol had characteristics similar to those observed after treatment with retinoic acid, but with a smaller disruption of the horny layer, due to less aggressive behaviour towards retinoic acid and retinol. BPE-CO₂A, in turn, had a slight effect in increasing the thickness of the epidermal basal layer and dermis (Figure 7).

4. Discussion

The ageing of human skin is a complex biological phenomenon consisting of two components: intrinsic ageing (passage of time and individual genetic features) and extrinsic ageing caused by cumulative exposure to environmental factors such as ultraviolet radiation [21]. Intrinsic ageing is a slow, cumulative, progressive, and degradative process affecting mainly elastic fibre, while in extrinsic ageing the slow evolution can be enzymatically accelerated [22]. Both skin ageing processes are associated with structural and functional changes that occur in the dermal extracellular matrix, where

fibrillar collagens, elastic fibres, and glycosaminoglycans are necessary to confer tensile strength, resilience, and hydration [23]. In both intrinsic and extrinsic ageing, there is evidence for the degradation of fibrous extracellular matrix components including elastin, collagens, and the oligosaccharide fraction [23, 24]. Our results showed that BPE-CO₂A was able to significantly increase the synthesis of collagen and elastin *in vitro*. Although the *in vitro* stimulation promoted by BPE-CO₂A on extracellular matrix components is quantitatively lower than the effect exerted by retinoic acid, it is evident in the *ex vivo* experiments that BPE-CO₂A stimulates the production of collagen and elastin in the skin.

The fundamental mechanism for age-related collagen synthesis involves the transforming growth factor- β (TGF- β) signalling pathway, which stimulates collagen synthesis in cultured fibroblasts [25–28]. Moreover, it has been suggested that reactive oxygen species can activate different surface receptors, including those for epidermal growth factor, and this effect may be mediated in part by the mitogen-activated protein kinase (MAPK) pathway [22, 29].




FIGURE 2: Extracellular matrix elements evaluation. (a) Collagen and elastin production in human fibroblast cultures after 48 hours of incubation with BPE-CO₂A, retinoic acid, or retinol. (b) Collagen (A), elastin (B), and glycosaminoglycan (C) in human skin fragments labelled with anti-collagen (green), anti-elastin (green), and anti-glycosaminoglycan (green) antibodies, respectively, after a 48-hour incubation. The nuclei were counterstained with DAPI (blue) (40x magnification). The data represent the mean \pm SD of three individual experiments. * $P < 0.05$, ** $P < 0.01$.

The activation of several MAPK members, particularly c-Jun and c-Fos, is related to the transcription complex AP-1 that directly inhibits the production of procollagen by blocking the TGF- β receptor [30, 31]. Topical TGF- β treatment in wounds promotes collagen synthesis, increases tensile wound strength, stimulates granulation tissue formation, enhances the thickness of regenerate dermal tissue, and stabilizes the dermoepidermal junction [32, 33]. It has been related that the reduction of collagen in photoaged human skin occurs due to blocking the TGF- β receptor [30]. BPE-CO₂A, retinoic acid, and retinol enhanced the production of TGF- β in fibroblast cultures, and this effect was in synergy with the reduction in collagen production that we detected in our experiments. Our results clearly showed that the increase in collagen synthesis is closely related to the enhanced production of TGF- β .

In addition to TGF- β , fibroblast growth factor-1 has a wide activity spectrum, mainly increasing synthesis of matrix

macromolecules, notably the main dermal glycosaminoglycan, hyaluronic acid [34], which contributes to the wound-healing process by stimulating fibroblast proliferation and inhibits the expression of collagenase-1 in keratinocytes [35]. In our study, BPE-CO₂A exerted a potent stimulatory effect on FGF- β levels in cultured human fibroblasts that was higher than the effects produced by both retinoic acid and retinol.

Another representative member of the growth factor family is epidermal growth factor (EGF), which can bind to and activate members of the EGF receptor (EGFR) family, leading to the initiation of several MAPKs signal transduction cascades and playing an important role in many cellular functions, such as proliferation, apoptosis, migration, inflammation, and immunity [36–39]. In fact, EGF induces collagenase expression in human dermal fibroblasts with a consequent reduction in collagen synthesis [40, 41]. Our results showed that BPE-CO₂A was able to increase EGFR




FIGURE 3: Effects of BPE-CO₂A, retinoic acid, and retinol on the TGF-β1 (a) and FGF (b) levels and EGFR mRNA expression in human fibroblast cultures. The data represent the mean ± SD. * $P < 0.05$, ** $P < 0.01$.

gene expression (2.6-fold), although retinoic acid produced a smaller increase (53.6%), and retinol was not able to increase gene expression. Thus, it is clear that the protective effect of BPE-CO₂A on collagen and elastin degradation was closely related to the increase in the expression of growth factors, particularly TGF-β and FGF-β, and EGFR expression in human fibroblasts. This effect was similar to that produced by retinoic acid and was sometimes more marked than effects produced by both retinoic acid and retinol.

In a second set of experiments, we evaluated the effects of BPE-CO₂A on both the gene expression and synthesis of the retinoic acid and retinoid X receptors. BPE-CO₂A, retinoic acid, and retinol were not able to significantly alter the gene expression of RAR. However, retinoic acid markedly increased RAR protein synthesis as evidenced using immunohistochemistry, whereas BPE-CO₂A treatment produces a subtle stimulation, similar to that produced by retinol. In contrast, BPE-CO₂A treatment increased RXR gene expression significantly (3.6-fold compared to control) and to a greater extent than retinoic acid and retinol, both of which increased RXR gene expression approximately 1.5-fold. This effect was corroborated by an increase in RXR protein synthesis.

There is evidence that RAR and RXR mRNA levels decrease with age and that compounds able to increase

the expression of these genes have neuroprotective effects in ageing [42, 43]. Thus, expression of RXRs was reduced in healthy elderly men suggesting that RXRs reduction might be a common feature of physiological senescence [44]. In addition, a single dose of all-*trans* retinoic acid to old rats was found to increase RAR expression in the liver after 24 h, indicating that reversible alterations in retinoid receptors may also occur during ageing [44, 45]. This way, it is possible that the protective effects produced by BPE-CO₂A are also related to RXR mRNA expression.

Finally, BPE-CO₂A contains a mixture of phytol and several long chain fatty acids, mainly palmitic, oleic, linoleic, and linolenic acids. Phytol, a branched fatty alcohol, is a carbon side-chain of chlorophylls and their metabolites, such as phytanic acid. Phytol has been shown to be a natural ligand for RXR, able to mimic various effects of conjugated linolenic acids, which activate the peroxisome proliferator-activated receptor (PPAR) and the RXRs [46, 47]. There is evidence that PPAR-γ mRNA levels decrease with age [41] and that phytol is functional as a PPAR-α ligand, stimulating the expression of PPAR-α-target genes in intact cells [47]. Moreover, phytol metabolites are compelling candidates for physiological effectors because their RXR binding affinities and activation potencies match their micromolar circulating concentrations [46, 48]. In contrast, reduced concentrations







FIGURE 4: Effects of BPE-CO₂A, retinoic acid, and retinol on RXR mRNA expression in human fibroblast cultures (a) and RXR synthesis (brown precipitate) in human skin fragments labelled with anti-RXR antibodies (b) after a 48-hour incubation (40x magnification). The data represent the mean \pm SD. * $P < 0.05$, ** $P < 0.01$.

FIGURE 5: Effects of BPE-CO₂A, retinoic acid, and retinol on RAR mRNA expression in human fibroblasts (a) and RAR synthesis (brown precipitate) in human skin fragments labelled with anti-RXR antibodies (b) after a 48-hour incubation (40x magnification). The data represent the mean \pm SD. * $P < 0.05$, ** $P < 0.01$.

of long-chain fatty acids have been correlated with normal ageing and neurodegeneration [48], and some of these long-chain fatty acids are also RXR ligands or increase RXR gene expression [12–16, 49]. In our study, phytol was able to induce RXR synthesis but was not able to influence RAR production. Because phytol is exclusively of dietary origin, a *Bidens pilosa* extract containing phytol may represent an essential dietary product to control cellular metabolism through the RXR signalling pathways.

5. Conclusion

In conclusion, our results showed that BPE-CO₂A from *Bidens pilosa* enhanced extracellular matrix components, particularly collagen and elastin fibres through the maintenance of TGF- β , FGF- β , and EGF levels and, simultaneously, affected RXR gene expression and synthesis. It is an innovative active ingredient with several clinical applications and pharmacological activities that control intrinsic and extrinsic human skin ageing. This way, BPE-CO₂A from *Bidens pilosa*




FIGURE 6: Effects of phytol on RXR and RAR synthesis (brown precipitate) in human skin fragments labelled with anti-RXR and anti-RAR antibodies after a 48-hour incubation (40x magnification).




FIGURE 7: Photomicrographs of human skin fragments stained with haematoxylin-eosin and treated with retinoic acid, retinol, or BPE-CO₂A (40x magnification). Sc: stratum corneum, ep: epidermis, de: dermis, and va: vacuoles.

is an innovative active ingredient with potential clinical applications to control intrinsic and extrinsic human skin ageing and to develop an herbal preparation with retinoid-like activity useful as a skin-repair and antiageing agent.

Disclaimer

The content of this publication is solely the responsibility of the authors and does not necessarily represent the official views of the financial support agencies.

Conflict of Interests

The authors declare no conflict of interests.

Acknowledgments





Research in the Di Stasi Lab (PhytoPharmaTech) was supported by the São Paulo Research Foundation (FAPESP) and Chemyunion Quimica Ltda. Fellowships are as follows: LCDS from the National Council for Scientific and Technological Development (CNPq, Brazilian Ministry of Science and Technology).

References

- [1] A. le Maire, S. Álvarez, P. Shankaranarayanan, A. R. de Lera, W. Bourguet, and H. Gronemeyer, "Retinoid receptors and therapeutic applications of RAR/RXR modulators," *Current Topics in Medicinal Chemistry*, vol. 12, no. 6, pp. 505–527, 2012.
- [2] R. E. B. Watson, J. A. Ratnayaka, R. C. C. Brooke, S. Yee-Sit-Yu, P. Ancian, and C. E. M. Griffiths, "Retinoic acid receptor α expression and cutaneous ageing," *Mechanisms of Ageing and Development*, vol. 125, no. 7, pp. 465–473, 2004.
- [3] P. Germain, P. Chambon, G. Eichele et al., "International union of pharmacology. LXIII. Retinoid X receptors," *Pharmacological Reviews*, vol. 58, no. 4, pp. 760–772, 2006.
- [4] L. Rittié, G. L. Fisher, and C. E. M. Griffiths, "Anti-aging effects of retinoids and mechanisms of actions," in *Retinoids and Carotenoids in Dermatology*, A. Vahlquist and M. Duvic, Eds., pp. 77–102, Informa Healthcare, New York, NY, USA, 2007.
- [5] C. Stefanaki, A. Stratigos, and A. Katsambas, "Topical retinoids in the treatment of photoaging," *Journal of Cosmetic Dermatology*, vol. 4, pp. 130–134, 2005.
- [6] S. Knutsen-Larson, A. L. Dawson, C. A. Dunnick, and R. P. Dellavalle, "Acne vulgaris: pathogenesis, treatment, and needs assessment," *Dermatologic Clinics*, vol. 30, no. 1, pp. 99–106, 2012.
- [7] F. L. Silva, D. C. H. Fischer, J. F. Tavares, M. S. Silva, P. F. De Athayde-Filho, and J. M. Barbosa-Filho, "Compilation of secondary metabolites from *Bidens pilosa* L.," *Molecules*, vol. 16, no. 2, pp. 1070–1102, 2011.
- [8] A. P. Bartolome, I. M. Villaseñor, and W.-C. Yang, "*Bidens pilosa* L. (Asteraceae): botanical properties, traditional uses, phytochemistry, and pharmacology," *Evidence-Based Complementary and Alternative Medicine*, vol. 2013, Article ID 340215, 51 pages, 2013.
- [9] L. C. Di Stasi, C. A. Hiruma, E. M. Guimaraes, and C. M. Santos, "Medicinal plants popularly used in Brazilian Amazon," *Fitoterapia*, vol. 65, no. 6, pp. 529–540, 1994.
- [10] L. C. Di Stasi, G. P. Oliveira, M. A. Carvalhaes et al., "Medicinal plants popularly used in the Brazilian Tropical Atlantic Forest," *Fitoterapia*, vol. 73, no. 1, pp. 69–91, 2002.
- [11] B. Buaud, L. Esterle, C. Vaysse et al., "A high-fat diet induces lower expression of retinoid receptors and their target genes

- GAP-43/neuromodulin and RC3/neurogranin in the rat brain," *British Journal of Nutrition*, vol. 103, no. 12, pp. 1720–1729, 2010.
- [12] P. K. Lemotte, S. Keidel, and C. M. Apfel, "Phytanic acids is a retinoid X receptor ligand," *European Journal of Biochemistry*, vol. 236, no. 1, pp. 328–333, 1996.
- [13] J. Lengqvist, A. Mata de Urquiza, A.-C. Bergman et al., "Polyunsaturated fatty acids including docosahexaenoic and arachidonic acid bind to the retinoid X receptor α ligand-binding domain," *Molecular and Cellular Proteomics*, vol. 3, no. 7, pp. 692–703, 2004.
- [14] S. A. Moore, E. Hurt, E. Yoder, H. Sprecher, and A. A. Spector, "Docosahexaenoic acid synthesis in human skin fibroblasts involves peroxisomal retroconversion of tetracosahexaenoic acid," *Journal of Lipid Research*, vol. 36, no. 11, pp. 2433–2443, 1995.
- [15] F. Zapata-Gonzalez, F. Rueda, J. Petriz et al., "Human dendritic cell activities are modulated by the omega-3 fatty acid, docosahexaenoic acid, mainly through PPAR (gamma):RXR heterodimers: comparison with other polyunsaturated fatty acids," *Journal of Leukocyte Biology*, vol. 84, no. 4, pp. 1172–1182, 2008.
- [16] E. Reverchon and I. de Marco, "Supercritical fluid extraction and fractionation of natural matter," *Journal of Supercritical Fluids*, vol. 38, no. 2, pp. 146–166, 2006.
- [17] I. Gamliehl-Bonshtein, E. Korin, and S. Cohen, "Selective separation of cis-trans geometrical isomers of β -carotene via CO₂ supercritical fluid extraction," *Biotechnology and Bioengineering*, vol. 80, no. 2, pp. 169–174, 2002.
- [18] G. Jenkins, "Molecular mechanisms of skin ageing," *Mechanisms of Ageing and Development*, vol. 123, no. 7, pp. 801–810, 2002.
- [19] S. Kojima, H. Muramatsu, H. Amanuma, and T. Muramatsu, "Midkine enhances fibrinolytic activity of bovine endothelial cells," *The Journal of Biological Chemistry*, vol. 270, no. 16, pp. 9590–9596, 1995.
- [20] E. A. Bauer, J. L. Seltzer, and A. Z. Eisen, "Retinoic acid inhibition of collagenase and gelatinase expression in human skin fibroblast cultures. Evidence for a dual mechanism," *Journal of Investigative Dermatology*, vol. 81, no. 2, pp. 162–169, 1983.
- [21] M. Sárdy, "Role of matrix metalloproteinases in skin ageing," *Connective Tissue Research*, vol. 50, no. 2, pp. 132–138, 2009.
- [22] E. C. Naylor, R. E. B. Watson, and M. J. Sherratt, "Molecular aspects of skin ageing," *Maturitas*, vol. 69, no. 3, pp. 249–256, 2011.
- [23] A. J. Bailey, "Molecular mechanisms of ageing in connective tissues," *Mechanisms of Ageing and Development*, vol. 122, no. 7, pp. 735–755, 2001.
- [24] K.-A. Hwang, B.-R. Yi, and K.-C. Choi, "Molecular mechanisms and in vivo mouse models of skin aging associated with dermal matrix alterations," *Laboratory Animal Research*, vol. 27, no. 1, pp. 1–8, 2011.
- [25] R. A. Ignatz and J. Massague, "Transforming growth factor- β stimulates the expression of fibronectin and collagen and their incorporation into the extracellular matrix," *The Journal of Biological Chemistry*, vol. 261, no. 9, pp. 4337–4345, 1986.
- [26] M. Bitzer, G. von Gersdorff, D. Liang et al., "A mechanism of suppression of TGF- β /SMAD signaling by NF- κ B/RelA," *Genes and Development*, vol. 14, no. 2, pp. 187–197, 2000.
- [27] R. Raghov, A. E. Postlethwaite, J. Keski-Oja, H. L. Moses, and A. H. Kang, "Transforming growth factor β increases steady state levels of type I procollagen and fibronectin messenger RNAs posttranscriptionally in cultured human dermal fibroblasts," *Journal of Clinical Investigation*, vol. 79, no. 4, pp. 1285–1288, 1987.
- [28] K. K. Nelson and J. A. Melendez, "Mitochondrial redox control of matrix metalloproteinases," *Free Radical Biology and Medicine*, vol. 37, no. 6, pp. 768–784, 2004.
- [29] T. Quan, T. He, S. Kang, J. J. Voorhees, and G. J. Fisher, "Solar ultraviolet irradiation reduces collagen in photoaged human skin by blocking transforming growth factor- β type II receptor/Smad signaling," *The American Journal of Pathology*, vol. 165, no. 3, pp. 741–751, 2004.
- [30] M. Yaar and B. A. Gilchrist, "Photoageing: mechanism, prevention and therapy," *British Journal of Dermatology*, vol. 157, no. 5, pp. 874–887, 2007.
- [31] L. Tjäderhane, H. Palosaari, J. Wahlgren, M. Larmas, T. Sorsa, and T. Salo, "Human odontoblast culture method: the expression of collagen and matrix metalloproteinases (MMPs)," *Advances in Dental Research*, vol. 15, pp. 55–58, 2001.
- [32] R. Govinden and K. D. Bhoola, "Genealogy, expression, and cellular function of transforming growth factor- β ," *Pharmacology and Therapeutics*, vol. 98, no. 2, pp. 257–265, 2003.
- [33] K. Kuroda, A. Utani, Y. Hamasaki, and H. Shinkai, "Up-regulation of putative hyaluronan synthase mRNA by basic fibroblast growth factor and insulin-like growth factor-1 in human skin fibroblasts," *Journal of Dermatological Science*, vol. 26, no. 2, pp. 156–160, 2001.
- [34] B. K. Pilcher, J. Gaither-Ganim, W. C. Parks, and H. G. Welgus, "Cell type-specific inhibition of keratinocyte collagenase-1 expression by basic fibroblast growth factor and keratinocyte growth factor: a common receptor pathway," *Journal of Biological Chemistry*, vol. 272, no. 29, pp. 18147–18154, 1997.
- [35] M. R. Frey, R. S. Dize, K. L. Edelblum, and D. B. Polk, "p38 kinase regulates epidermal growth factor receptor downregulation and cellular migration," *The EMBO Journal*, vol. 25, no. 24, pp. 5683–5692, 2006.
- [36] S. Pastore, F. Mascia, V. Mariani, and G. Girolomoni, "The epidermal growth factor receptor system in skin repair and inflammation," *Journal of Investigative Dermatology*, vol. 128, no. 6, pp. 1365–1374, 2008.
- [37] S. A. Grando, J.-C. Bystry, A. I. Chernyavsky et al., "Apoptolysis: a novel mechanism of skin blistering in pemphigus vulgaris linking the apoptotic pathways to basal cell shrinkage and suprabasal acantholysis," *Experimental Dermatology*, vol. 18, no. 9, pp. 764–770, 2009.
- [38] C. H. Park and J. H. Chung, "Epidermal growth factor-induced matrix metalloproteinase-1 expression is negatively regulated by p38 MAPK in human skin fibroblasts," *Journal of Dermatological Science*, vol. 64, no. 2, pp. 134–141, 2011.
- [39] Y. Mimura, H. Ihn, M. Jinnin, Y. Asano, K. Yamane, and K. Tamaki, "Epidermal growth factor affects the synthesis and degradation of type I collagen in cultured human dermal fibroblasts," *Matrix Biology*, vol. 25, no. 4, pp. 202–212, 2006.
- [40] S. E. Moon, N. Bhagavathula, and J. Varani, "Keratinocyte stimulation of matrix metalloproteinase-1 production and proliferation in fibroblasts: regulation through mitogen-activated protein kinase signalling events," *British Journal of Cancer*, vol. 87, no. 4, pp. 457–464, 2002.
- [41] V. Enderlin, S. Alfons, V. Pallet et al., "Aging decreases the abundance of retinoic acid (RAR) and triiodothyronine (TR) nuclear receptor mRNA in rat brain: effect of the administration of retinoids," *FEBS Letters*, vol. 412, no. 3, pp. 629–632, 1997.

- [42] S. C. Dyal, G. J. Michael, and A. T. Michael-Titus, "Omega-3 fatty acids reverse age-related decreases in nuclear receptors and increase neurogenesis in old rats," *Journal of Neuroscience Research*, vol. 88, no. 10, pp. 2091–2102, 2010.
- [43] J. Brtko, E. Rock, P. Nezbedova et al., "Age-related change in the retinoid X receptor beta gene expression in peripheral blood mononuclear cells of healthy volunteers: effect of 13-*cis* retinoic acid supplementation," *Mechanisms of Ageing and Development*, vol. 128, no. 11-12, pp. 594–600, 2007.
- [44] V. Pallet, V. Azais-Braesco, V. Enderlin et al., "Aging decreases retinoic acid and triiodothyronine nuclear expression in rat liver: exogenous retinol and retinoic acid differentially modulate this decreased expression," *Mechanisms of Ageing and Development*, vol. 99, no. 2, pp. 123–136, 1997.
- [45] M. F. McCarty, "The chlorophyll metabolite phytanic acid is a natural retinoid—potential for treatment and prevention of diabetes," *Medical Hypotheses*, vol. 56, no. 2, pp. 217–219, 2001.
- [46] T. Goto, N. Takahashi, S. Kato et al., "Phytol directly activates peroxisome proliferator-activated receptor α (PPAR α) and regulates gene expression involved in lipid metabolism in PPAR α -expressing HepG2 hepatocytes," *Biochemical and Biophysical Research Communications*, vol. 337, no. 2, pp. 440–445, 2005.
- [47] S. Kitareewan, L. T. Burka, K. B. Tomer et al., "Phytol metabolites are circulating dietary factors that activate the nuclear receptor RXR," *Molecular Biology of the Cell*, vol. 7, no. 8, pp. 1153–1166, 1996.
- [48] C. I. F. Janssen and A. J. Kiliaan, "Long-chain polyunsaturated fatty acids (LCPUFA) from genesis to senescence: the influence of LCPUFA on neural development, aging, and neurodegeneration," *Progress in Lipid Research*, vol. 53, no. 1, pp. 1–17, 2014.
- [49] A. M. de Urquiza, S. Liu, M. Sjöberg et al., "Docosahexaenoic acid, a ligand for the retinoid X receptor in mouse brain," *Science*, vol. 290, no. 5499, pp. 2140–2144, 2000.



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

