

Research Article

Dynamical Behaviors of Stochastic Hopfield Neural Networks with Both Time-Varying and Continuously Distributed Delays

Qinghua Zhou,¹ Penglin Zhang,² and Li Wan³

¹ Department of Mathematics, Zhaoqing University, Zhaoqing 526061, China

² School of Remote Sensing and Information Engineering, Wuhan University, Wuhan 430072, China

³ School of Mathematics and Physics, Wuhan Textile University, Wuhan 430073, China

Correspondence should be addressed to Penglin Zhang; zpicy@263.com

Received 24 December 2012; Revised 19 March 2013; Accepted 20 March 2013

Academic Editor: Chuangxia Huang

Copyright © 2013 Qinghua Zhou et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper investigates dynamical behaviors of stochastic Hopfield neural networks with both time-varying and continuously distributed delays. By employing the Lyapunov functional theory and linear matrix inequality, some novel criteria on asymptotic stability, ultimate boundedness, and weak attractor are derived. Finally, an example is given to illustrate the correctness and effectiveness of our theoretical results.

1. Introduction

Hopfield neural networks [1] have been extensively studied in the past years and found many applications in different areas such as pattern recognition, associative memory, and combinatorial optimization. Such applications heavily depend on the dynamical behaviors such as stability, uniform boundedness, ultimate boundedness, attractor, bifurcation, and chaos. As it is well known, time delays are unavoidably encountered in the implementation of neural networks. Since time delays as a source of instability and bad performance always appear in many neural networks owing to the finite speed of information processing, the stability analysis for the delayed neural networks has received considerable attention. However, in these recent publications, most research on delayed neural networks has been restricted to simple cases of discrete delays. Since a neural network usually has a spatial nature due to the presence of an amount of parallel pathways of a variety of axon sizes and lengths, it is desired to model them by introducing distributed delays. Therefore, both discrete and distributed delays should be taken into account when modeling realistic neural networks [2, 3].

On the other hand, it has now been well recognized that stochastic disturbances are also ubiquitous owing to thermal noise in electronic implementations. Therefore, it

is important to understand how these disturbances affect the networks. Many results on stochastic neural networks have been reported in [4–24]. Some sufficient criteria on the stability of uncertain stochastic neural networks were derived in [4–7]. Almost sure exponential stability of stochastic neural networks was studied in [8–10]. In [11–16], mean square exponential stability and p th moment exponential stability of stochastic neural networks were discussed. The stability of stochastic impulsive neural networks was discussed in [17–19]. The stability of stochastic neural networks with the Markovian jumping parameters was investigated in [20–22]. The passivity analysis for stochastic neural networks was discussed in [23, 24]. These references mainly considered the stability of equilibrium point of stochastic delayed neural networks. What do we study to understand the asymptotic behaviors when the equilibrium point does not exist?

Except for the stability property, boundedness and attractor are also foundational concepts of dynamical systems. They play an important role in investigating the uniqueness of equilibrium, global asymptotic stability, global exponential stability, and the existence of periodic solution, its control, and synchronization [25]. Recently, ultimate boundedness and attractor of several classes of neural networks with time delays have been reported in [26–33]. Some sufficient criteria were derived in [26, 27], but these results hold only

under constant delays. Following, in [28], the globally robust ultimate boundedness of integrodifferential neural networks with uncertainties and varying delays were studied. After that, some sufficient criteria on the ultimate boundedness of neural networks with both varying and unbounded delays were derived in [29], but the concerned systems are deterministic ones. In [30, 31], a series of criteria on the boundedness, global exponential stability, and the existence of periodic solution for nonautonomous recurrent neural networks were established. In [32, 33], the ultimate boundedness and weak attractor of stochastic neural networks with time-varying delays were discussed. To the best of our knowledge, for stochastic neural networks with mixed time delays, there are few published results on the ultimate boundedness and weak attractor. Therefore, the arising questions about the ultimate boundedness, weak attractor, and asymptotic stability of the stochastic Hopfield neural networks with mixed time delays are important and meaningful.

The left paper is organized as follows. Some preliminaries are in Section 2, main results are presented in Section 3, a numerical example is given in Section 4, and conclusions are drawn in Section 5.

2. Preliminaries

Consider the following stochastic Hopfield neural networks with both time-varying and continuously distributed delays:

$$\begin{aligned} dx(t) = & \left[-Cx(t) + Af(x(t)) + Bf(x(t-\tau(t))) \right. \\ & \left. + D \int_{-\infty}^t K(t-s)g(x(s))ds + J \right] dt \\ & + [\sigma_1 x(t) + \sigma_2 x(t-\tau(t))] dw(t), \end{aligned} \quad (1)$$

in which $x = (x_1, \dots, x_n)^T$ is a state vector associated with the neurons; $C = \text{diag}(c_1, \dots, c_n)$, $c_i > 0$ represents the rate with which the i th unit will reset its potential to the resting state in isolation when being disconnected from the network and the external stochastic perturbation; $A = (a_{ij})_{n \times n}$, $B = (b_{ij})_{n \times n}$, and $D = (d_{ij})_{n \times n}$ represent the connection weight matrix, the delayed connection weight matrix, and the distributively delayed connection weight matrix, respectively; $J = (J_1, \dots, J_n)^T$, J_i denotes the external bias on the i th unit; $f(x(t)) = (f_1(x_1(t)), \dots, f_n(x_n(t)))^T$, $g(x(t)) = (g_1(x_1(t)), \dots, g_n(x_n(t)))^T$, f_j and g_j denote activation functions, $K(t) = \text{diag}(k_1(t), \dots, k_n(t))$, and the delay kernel $k_j(t)$ is a real-valued nonnegative continuous function defined on $[0, \infty)$; σ_1 and σ_2 are diffusion coefficient matrices; $w(t)$ is a one-dimensional Brownian motion or Wiener process, which is defined on a complete probability space (Ω, \mathcal{F}, P) with a natural filtration $\{\mathcal{F}_t\}_{t \geq 0}$ generated by $\{w(s) : 0 \leq s \leq t\}$; $\tau(t)$ is the transmission delay, and the initial conditions associated with system (1) are of the following forms: $x(t) = \xi(t)$, $-\infty < t \leq 0$, where $\xi(\cdot)$ is a \mathcal{F}_0 -measurable, bounded, and continuous R^n -valued random variable defined on $(-\infty, 0]$.

Throughout this paper, one always supposes that the following condition holds.

(A1) For $f(x(t))$ and $g(x(t))$ in (1), there are always constants $l_i^-, l_i^+, m_i^-,$ and m_i^+ such that

$$\begin{aligned} l_i^- & \leq \frac{f_i(x) - f_i(y)}{x - y} \leq l_i^+, \\ m_i^- & \leq \frac{g_i(x) - g_i(y)}{x - y} \leq m_i^+, \quad \forall x, y \in R. \end{aligned} \quad (2)$$

Moreover, there exist constant $\nu > 0$ and matrix $\bar{K}(\nu) = \text{diag}(\bar{k}_1(\nu), \dots, \bar{k}_n(\nu)) > 0$ such that

$$\int_0^\infty k_j(\theta) d\theta = 1, \quad \int_0^\infty k_j(\theta) e^{\nu\theta} d\theta = \bar{k}_j(\nu) < \infty. \quad (3)$$

Following, $A > 0$ (resp., $A \geq 0$) means that matrix A is a symmetric positive definite (resp., positive semi-definite). A^T and A^{-1} denote the transpose and inverse of the matrix A . $\lambda_{\max}(A)$ and $\lambda_{\min}(A)$ represent the maximum and minimum eigenvalues of matrix A , respectively.

Definition 1. System (1) is said to be stochastically ultimately bounded if, for any $\varepsilon \in (0, 1)$, there is a positive constant $C = C(\varepsilon)$ such that the solution $x(t)$ of system (1) satisfies

$$\limsup_{t \rightarrow \infty} P \{ \|x(t)\| \leq C \} \geq 1 - \varepsilon. \quad (4)$$

Lemma 2 (see [34]). *Let $Q(x) = Q^T(x)$, $R(x) = R^T(x)$ and $S(x)$ depends affinely on x . Then, linear matrix inequality*

$$\begin{pmatrix} Q(x) & S(x) \\ S^T(x) & R(x) \end{pmatrix} > 0 \quad (5)$$

is equivalent to

- (1) $R(x) > 0, Q(x) - S(x)R^{-1}(x)S^T(x) > 0,$
- (2) $Q(x) > 0, R(x) - S^T(x)Q^{-1}(x)S(x) > 0.$

3. Main Results

Theorem 3. *System (1) is stochastically ultimately bounded provided that $\tau(t)$ satisfies $0 \leq \tau(t) \leq \tau$, $\dot{\tau}(t) \leq \mu \leq 1$, and there exist some matrices $P > 0, Q_i \geq 0, U_i = \text{diag}(u_{i1}, \dots, u_{in}) \geq 0$ ($i = 1, 2, 3, 4$) such that the following linear matrix inequality holds:*

$$\begin{aligned} \Sigma = & \begin{pmatrix} \Sigma_{11} & 0 & PA + L_2 U_1 & PB & M_2 U_4 & PD & \sigma_1^T P \\ * & \Sigma_{22} & 0 & L_2 U_2 & 0 & 0 & \sigma_2^T P \\ * & * & \Sigma_{33} & 0 & 0 & 0 & 0 \\ * & * & * & \Sigma_{44} & 0 & 0 & 0 \\ * & * & * & * & \Sigma_{55} & 0 & 0 \\ * & * & * & * & * & -U_3 & 0 \\ * & * & * & * & * & * & -P \end{pmatrix} \\ & < 0, \end{aligned} \quad (6)$$

where $*$ denotes the corresponding symmetric terms,

$$\begin{aligned} \Sigma_{11} &= Q_1 + \tau Q_2 - PC - CP - 2L_1U_1 - 2M_1U_4, \\ \Sigma_{22} &= -(1 - \mu)Q_1 - 2L_1U_2, \quad \Sigma_{33} = Q_3 + \tau Q_4 - 2U_1, \\ \Sigma_{44} &= -(1 - \mu)Q_3 - 2U_2, \quad \Sigma_{55} = U_3\bar{K}(\nu) - 2U_4, \\ L_1 &= \text{diag}(l_1^+ l_1^+, \dots, l_n^+ l_n^+), \\ L_2 &= \text{diag}(l_1^- + l_1^+, \dots, l_n^- + l_n^+), \\ M_1 &= \text{diag}(m_1^- m_1^+, \dots, m_n^- m_n^+), \\ M_2 &= \text{diag}(m_1^- + m_1^+, \dots, m_n^- + m_n^+). \end{aligned} \tag{7}$$

Proof. The key of the proof is to prove that there exists a positive constant C_* , which is independent of the initial data, such that

$$\limsup_{t \rightarrow \infty} E\|x(t)\|^2 \leq C_*. \tag{8}$$

If (8) holds, then it follows from Chebyshev's inequality that for any $\varepsilon > 0$ and $C = \sqrt{C_*/\varepsilon}$,

$$\limsup_{t \rightarrow \infty} P\{\|x(t)\| > C\} \leq \frac{\limsup_{t \rightarrow \infty} E\|x(t)\|^2}{C^2} = \varepsilon, \tag{9}$$

which implies that (4) holds. Now, we begin to prove that (8) holds.

From $\Sigma < 0$ and Lemma 2, one may obtain

$$\begin{aligned} &\begin{pmatrix} \Sigma_{11} & 0 & PA + L_2U_1 & PB & M_2U_4 & PD \\ * & \Sigma_{22} & 0 & L_2U_2 & 0 & 0 \\ * & * & \Sigma_{33} & 0 & 0 & 0 \\ * & * & * & \Sigma_{44} & 0 & 0 \\ * & * & * & * & \Sigma_{55} & 0 \\ * & * & * & * & * & -U_3 \end{pmatrix} \\ &+ \begin{pmatrix} \sigma_1^T P \\ \sigma_2^T P \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} P^{-1} \begin{pmatrix} \sigma_1^T P \\ \sigma_2^T P \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}^T < 0. \end{aligned} \tag{10}$$

Hence, there exists a sufficiently small $\lambda \in (0, \nu)$ such that

$$\begin{aligned} \Gamma &= \begin{pmatrix} \Gamma_{11} & 0 & PA + L_2U_1 & PB & M_2U_4 & PD \\ * & \Gamma_{22} & 0 & L_2U_2 & 0 & 0 \\ * & * & \Gamma_{33} & 0 & 0 & 0 \\ * & * & * & \Gamma_{44} & 0 & 0 \\ * & * & * & * & \Gamma_{55} & 0 \\ * & * & * & * & * & -U_3 \end{pmatrix} \\ &+ \begin{pmatrix} \sigma_1^T P \\ \sigma_2^T P \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} P^{-1} \begin{pmatrix} \sigma_1^T P \\ \sigma_2^T P \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}^T < 0, \end{aligned} \tag{11}$$

where I is identity matrix,

$$\begin{aligned} \Gamma_{11} &= e^{\lambda\tau}Q_1 + \tau Q_2 - PC - CP - 2L_1U_1 \\ &\quad - 2M_1U_4 + 2\lambda P + 2\lambda I, \\ \Gamma_{22} &= \Sigma_{22} + \lambda I, \quad \Gamma_{33} = \lambda I + e^{\lambda\tau}Q_3 + \tau Q_4 - 2U_1, \\ \Gamma_{44} &= \Sigma_{44} + \lambda I, \quad \Gamma_{55} = \Sigma_{55} + \lambda I. \end{aligned} \tag{12}$$

Consider the Lyapunov-Krasovskii function as follows:

$$\begin{aligned} V(x(t), t) &= V_1(x(t), t) + V_2(x(t), t) \\ &\quad + V_3(x(t), t) + V_4(x(t), t), \end{aligned} \tag{13}$$

where

$$\begin{aligned} V_1(x(t), t) &= e^{\lambda t}x^T(t)Px(t), \\ V_2(x(t), t) &= \sum_{j=1}^n u_{3j} \int_0^\infty k_j(\theta) \int_{t-\theta}^t e^{\lambda(s+\theta)} g_j^2(x_j(s)) ds d\theta, \\ V_3(x(t), t) &= \int_{t-\tau(t)}^t e^{\lambda(s+\tau)} [x^T(s)Q_1x(s) + f^T(x(s))Q_3f(x(s))] ds, \\ V_4(x(t), t) &= \int_{t-\tau(t)}^t \int_s^t e^{\lambda\theta} [x^T(\theta)Q_2x(\theta) \\ &\quad + f^T(x(\theta))Q_4f(x(\theta))] d\theta ds. \end{aligned} \tag{14}$$

Then, it can be obtained by Ito's formula in [35] that

$$\begin{aligned} dV(x(t), t) &= e^{\lambda t}2x^T(t)P[\sigma_1x(t) + \sigma_2x(t - \tau(t))]dw(t) \\ &\quad + \left\{ \mathcal{L}V_1(x(t), t) \right. \\ &\quad \left. + \frac{\partial [V_2(x(t), t) + V_3(x(t), t) + V_4(x(t), t)]}{\partial t} \right\} dt, \end{aligned} \tag{15}$$

where

$$\begin{aligned} \frac{\partial V_2(x(t), t)}{\partial t} &= \sum_{j=1}^n u_{3j} \int_0^\infty k_j(\theta) [e^{\lambda(t+\theta)} g_j^2(x_j(t)) - e^{\lambda t} g_j^2(x_j(t - \theta))] d\theta \\ &\leq e^{\lambda t} \sum_{j=1}^n u_{3j} \bar{k}_j(\nu) g_j^2(x_j(t)) \end{aligned}$$

$$\begin{aligned}
& - e^{\lambda t} \sum_{j=1}^n u_{3j} \left[\int_0^{\infty} k_j(\theta) g_j(x_j(t-\theta)) d\theta \right]^2 \\
& = e^{\lambda t} g^T(x(t)) U_3 \bar{K}(v) g(x(t)) \\
& - e^{\lambda t} \left(\int_{-\infty}^t K(t-s) g(x(s)) ds \right)^T \\
& \times U_3 \left(\int_{-\infty}^t K(t-s) g(x(s)) ds \right), \\
\frac{\partial V_3(x(t), t)}{\partial t} & = e^{\lambda(t+\tau)} \left[x^T(t) Q_1 x(t) + f^T(x(t)) Q_3 f(x(t)) \right] \\
& - (1 - \dot{\tau}(t)) e^{\lambda(t-\tau(t)+\tau)} \\
& \times \left[x^T(t-\tau(t)) Q_1 x(t-\tau(t)) \right. \\
& \quad \left. + f^T(x(t-\tau(t))) Q_3 f(x(t-\tau(t))) \right] \\
& \leq e^{\lambda(t+\tau)} \left[x^T(t) Q_1 x(t) + f^T(x(t)) Q_3 f(x(t)) \right] \\
& - (1 - \mu) e^{\lambda t} \\
& \times \left[x^T(t-\tau(t)) Q_1 x(t-\tau(t)) \right. \\
& \quad \left. + f^T(x(t-\tau(t))) Q_3 f(x(t-\tau(t))) \right],
\end{aligned}$$

$$\begin{aligned}
\frac{\partial V_4(x(t), t)}{\partial t} & = e^{\lambda t} \tau(t) \left[x^T(t) Q_2 x(t) + f^T(x(t)) Q_4 f(x(t)) \right] \\
& - (1 - \dot{\tau}(t)) \\
& \times \int_{t-\tau(t)}^t e^{\lambda s} \left[x^T(s) Q_2 x(s) + f^T(x(s)) Q_4 f(x(s)) \right] ds \\
& \leq e^{\lambda t} \tau \left[x^T(t) Q_2 x(t) + f^T(x(t)) Q_4 f(x(t)) \right],
\end{aligned} \tag{16}$$

$$\begin{aligned}
\mathcal{L}V_1(x(t), t) & = \frac{\partial V_1(x(t), t)}{\partial t} + \frac{\partial V_1(x(t), t)}{\partial x} \\
& \times \left[-Cx(t) + Af(x(t)) + Bf(x(t-\tau(t))) \right. \\
& \quad \left. + D \int_{-\infty}^t K(t-s) g(x(s)) ds + J \right] \\
& + \text{trace} \left(\left[\sigma_1 x(t) + \sigma_2 x(t-\tau(t)) \right]^T \frac{\partial^2 V_1(x(t), t)}{\partial x^2} \right. \\
& \quad \left. \times \left[\sigma_1 x(t) + \sigma_2 x(t-\tau(t)) \right] \right)
\end{aligned}$$

$$\begin{aligned}
& = \lambda e^{\lambda t} x^T(t) Px(t) \\
& + e^{\lambda t} 2x^T(t) P \left[-Cx(t) + Af(x(t)) \right. \\
& \quad \left. + Bf(x(t-\tau(t))) \right. \\
& \quad \left. + D \int_{-\infty}^t K(t-s) g(x(s)) ds + J \right] \\
& + e^{\lambda t} [\sigma_1 x(t) + \sigma_2 x(t-\tau(t))]^T \\
& \times P [\sigma_1 x(t) + \sigma_2 x(t-\tau(t))] \\
& \leq \lambda e^{\lambda t} x^T(t) Px(t) \\
& + e^{\lambda t} 2x^T(t) P \left[-Cx(t) + Af(x(t)) \right. \\
& \quad \left. + Bf(x(t-\tau(t))) \right. \\
& \quad \left. + D \int_{-\infty}^t K(t-s) g(x(s)) ds \right] \\
& + e^{\lambda t} \left[\lambda x^T(t) Px(t) + \lambda^{-1} J^T P J \right] \\
& + e^{\lambda t} [\sigma_1 x(t) + \sigma_2 x(t-\tau(t))]^T \\
& \times P [\sigma_1 x(t) + \sigma_2 x(t-\tau(t))],
\end{aligned} \tag{17}$$

in which the following inequality is used:

$$\begin{aligned}
2x^T P J & \leq \lambda x^T P x + \lambda^{-1} J^T P^T P^{-1} P J \\
& = \lambda x^T P x + \lambda^{-1} J^T P J, \quad \text{for } P > 0, \lambda > 0.
\end{aligned} \tag{18}$$

On the other hand, it follows from (A1) that for $i = 1, \dots, n$,

$$\begin{aligned}
& [f_i(x_i(t)) - f_i(0) - l_i^+ x_i(t)] \\
& \times [f_i(x_i(t)) - f_i(0) - l_i^- x_i(t)] \leq 0, \\
& [f_i(x_i(t-\tau(t))) - f_i(0) - l_i^+ x_i(t-\tau(t))] \\
& \times [f_i(x_i(t-\tau(t))) - f_i(0) - l_i^- x_i(t-\tau(t))] \leq 0, \\
0 & \leq e^{\lambda t} \left\{ -2 \sum_{i=1}^n u_{1i} [f_i(x_i(t)) - f_i(0) - l_i^+ x_i(t)] \right. \\
& \quad \times [f_i(x_i(t)) - f_i(0) - l_i^- x_i(t)] \\
& \quad - 2 \sum_{i=1}^n u_{2i} [f_i(x_i(t-\tau(t))) \\
& \quad \left. - f_i(0) - l_i^+ x_i(t-\tau(t))] \right\}
\end{aligned} \tag{19}$$

$$\begin{aligned}
 & \times [f_i(x_i(t - \tau(t))) - f_i(0) \\
 & \quad - l_i^- x_i(t - \tau(t))] \Big\} \\
 = e^{\lambda t} & \left\{ -2 \sum_{i=1}^n u_{1i} [f_i(x_i(t)) - l_i^+ x_i(t)] \right. \\
 & \quad \times [f_i(x_i(t)) - l_i^- x_i(t)] \\
 & \quad - 2 \sum_{i=1}^n u_{2i} [f_i(x_i(t - \tau(t))) - l_i^+ x_i(t - \tau(t))] \\
 & \quad \times [f_i(x_i(t - \tau(t))) - l_i^- x_i(t - \tau(t))] \\
 & \quad - 2 \sum_{i=1}^n u_{1i} f_i^2(0) \\
 & \quad + 2 \sum_{i=1}^n u_{1i} f_i(0) [2f_i(x_i(t)) - (l_i^+ + l_i^-) x_i(t)] \\
 & \quad - 2 \sum_{i=1}^n u_{2i} f_i^2(0) \\
 & \quad \left. + 2 \sum_{i=1}^n u_{2i} f_i(0) [2f_i(x_i(t - \tau(t))) \right. \\
 & \quad \quad \left. - (l_i^+ + l_i^-) x_i(t - \tau(t))] \Big\} \\
 \leq e^{\lambda t} & \left\{ -2 \sum_{i=1}^n u_{1i} [f_i(x_i(t)) - l_i^+ x_i(t)] \right. \\
 & \quad \times [f_i(x_i(t)) - l_i^- x_i(t)] \\
 & \quad - 2 \sum_{i=1}^n u_{2i} [f_i(x_i(t - \tau(t))) - l_i^+ x_i(t - \tau(t))] \\
 & \quad \times [f_i(x_i(t - \tau(t))) - l_i^- x_i(t - \tau(t))] \\
 & \quad + \sum_{i=1}^n [|4u_{1i} f_i(0) f_i(x_i(t))| \\
 & \quad \quad + |2u_{1i} f_i(0) (l_i^+ + l_i^-) x_i(t)|] \\
 & \quad + \sum_{i=1}^n [|4u_{2i} f_i(0) f_i(x_i(t - \tau(t)))| \\
 & \quad \quad + |2u_{2i} f_i(0) (l_i^+ + l_i^-) x_i(t - \tau(t))|] \Big\}
 \end{aligned}$$

$$\begin{aligned}
 & \leq e^{\lambda t} \left\{ -2 \sum_{i=1}^n u_{1i} [f_i(x_i(t)) - l_i^+ x_i(t)] \right. \\
 & \quad \times [f_i(x_i(t)) - l_i^- x_i(t)] \\
 & \quad - 2 \sum_{i=1}^n u_{2i} [f_i(x_i(t - \tau(t))) - l_i^+ x_i(t - \tau(t))] \\
 & \quad \times [f_i(x_i(t - \tau(t))) - l_i^- x_i(t - \tau(t))] \\
 & \quad + \sum_{i=1}^n [\lambda f_i^2(x_i(t)) + 4\lambda^{-1} f_i^2(0) u_{1i}^2 \\
 & \quad \quad + \lambda x_i^2(t) + \lambda^{-1} f_i^2(0) u_{1i}^2 (l_i^+ + l_i^-)^2] \\
 & \quad + \sum_{i=1}^n [\lambda f_i^2(x_i(t - \tau(t))) + 4\lambda^{-1} f_i^2(0) u_{2i}^2 \\
 & \quad \quad + \lambda x_i^2(t - \tau(t)) \\
 & \quad \quad \left. + \lambda^{-1} f_i^2(0) u_{2i}^2 (l_i^+ + l_i^-)^2] \Big\}. \tag{20}
 \end{aligned}$$

Similarly, one may obtain

$$\begin{aligned}
 0 \leq e^{\lambda t} & \left\{ -2 \sum_{i=1}^n u_{4i} [g_i(x_i(t)) - g_i(0) - m_i^+ x_i(t)] \right. \\
 & \quad \times [g_i(x_i(t)) - g_i(0) - m_i^- x_i(t)] \Big\} \\
 \leq e^{\lambda t} & \left\{ -2 \sum_{i=1}^n u_{4i} [g_i(x_i(t)) - m_i^+ x_i(t)] \right. \\
 & \quad \times [g_i(x_i(t)) - m_i^- x_i(t)] \\
 & \quad + \sum_{i=1}^n [\lambda g_i^2(x_i(t)) + 4\lambda^{-1} g_i^2(0) u_{4i}^2 \\
 & \quad \quad + \lambda x_i^2(t) + \lambda^{-1} g_i^2(0) u_{4i}^2 (m_i^+ + m_i^-)^2] \Big\}. \tag{21}
 \end{aligned}$$

Therefore, from (15)–(21), it follows that

$$\begin{aligned}
 dV(x(t), t) & \leq e^{\lambda t} 2x^T(t) P [\sigma_1 x(t) + \sigma_2 x(t - \tau(t))] dw(t) \\
 & \quad + e^{\lambda t} \eta^T(t) \Gamma \eta(t) dt + e^{\lambda t} C_1 dt \\
 & \leq e^{\lambda t} 2x^T(t) P [\sigma_1 x(t) + \sigma_2 x(t - \tau(t))] dw(t) \\
 & \quad + e^{\lambda t} C_1 dt, \tag{22}
 \end{aligned}$$

where

$$\eta(t) = \left(x^T(t), x^T(t - \tau(t)), f^T(x(t)), f^T(x(t - \tau(t))), \right. \\ \left. g^T(x(t)), \left(\int_{-\infty}^t K(t-s)g(x(s))ds \right)^T \right)^T, \tag{23}$$

$$C_1 = \lambda^{-1} J^T P J \\ + \sum_{i=1}^n \left[4\lambda^{-1} f_i^2(0) u_{1i}^2 + \lambda^{-1} f_i^2(0) u_{1i}^2 (l_i^+ + l_i^-)^2 \right. \\ \left. + 4\lambda^{-1} f_i^2(0) u_{2i}^2 \right. \\ \left. + \lambda^{-1} f_i^2(0) u_{2i}^2 (l_i^+ + l_i^-)^2 \right. \\ \left. + 4\lambda^{-1} g_i^2(0) u_{4i}^2 \right. \\ \left. + \lambda^{-1} g_i^2(0) u_{4i}^2 (m_i^+ + m_i^-)^2 \right]. \tag{24}$$

Thus, one may obtain that

$$V(x(t), t) \leq V(x(0), 0) + e^{\lambda t} \lambda^{-1} C_1 \\ + \int_0^t 2x^T(s) P [\sigma_1 x(s) + \sigma_2 x(s - \tau(s))] dw(s), \tag{25}$$

$$E\|x(t)\|^2 \leq \frac{e^{-\lambda t} EV(x(0), 0) + \lambda^{-1} C_1}{\lambda_{\min}(P)} \\ = \frac{e^{-\lambda t} EV(x(0), 0)}{\lambda_{\min}(P)} + C_*, \tag{26}$$

where $C_* = \lambda^{-1} C_1 / \lambda_{\min}(P)$. Equation (26) implies that (8) holds. The proof is completed. \square

Theorem 3 shows that there exists $t_0 > 0$ such that for any $t \geq t_0$, $P\{\|x(t)\| \leq C\} \geq 1 - \delta$. Let B_C be denoted by

$$B_C = \{x \in R^n \mid \|x(t)\| \leq C, t \geq t_0\}. \tag{27}$$

Clearly, B_C is closed, bounded, and invariant. Moreover,

$$\limsup_{t \rightarrow \infty} \inf_{y \in B_C} \|x(t) - y\| = 0 \tag{28}$$

with no less than probability $1 - \delta$, which means that B_C attracts the solutions infinitely many times with no less than probability $1 - \delta$; so we may say that B_C is a weak attractor for the solutions.

Theorem 4. Suppose that all conditions of Theorem 3 hold, then there exists a weak attractor B_C for the solutions of system (1).

Theorem 5. Suppose that all conditions of Theorem 3 hold and $f(0) = g(0) = J = 0$; then, zero solution of system (1) is mean square exponential stability and almost sure exponential stability.

Proof. If $f(0) = g(0) = J = 0$, then $C_1 = C_* = 0$. By (25) and the semimartingale convergence theorem used in [35], zero solution of system (1) is almost sure exponential stability. It follows from (26) that zero solution of system (1) is mean square exponential stability. \square

Remark 6. If one takes $Q_2 = Q_4 = 0$ in Theorem 3, then it is not required that $\mu \leq 1$. Furthermore, If one takes $Q_1 = Q_2 = Q_3 = Q_4 = 0$, then $\tau(t)$ may be nondifferentiable or the boundedness of $\dot{\tau}(t)$ is unknown.

Remark 7. Assumption (A1) is less conservative than that in [32] since the constants $l_i^-, l_i^+, m_i^-,$ and m_i^+ are allowed to be positive, negative, or zero. System (1) includes mixed time-delays, which is more complex than that in [33]. The systems concerned in [26–31] are deterministic, so the stochastic system studied in this paper is more complex and realistic.

When $\sigma_1 = \sigma_2 = 0$, system (1) becomes the following determined system:

$$\frac{dx(t)}{dt} = -Cx(t) + Af(x(t)) + Bf(x(t - \tau(t))) \\ + D \int_{-\infty}^t K(t-s)g(x(s))ds + J. \tag{29}$$

Definition 8. System (29) is said to be uniformly bounded, if, for each $H > 0$, there exists a constant $M = M(H) > 0$ such that $[t_0 \in R, \phi \in C(-\infty, 0], \|\phi\| \leq H, t > t_0]$ imply $\|x(t, t_0, \phi)\| \leq M$, where $\|\phi\| = \sup_{t \leq 0} \|\phi(t)\|$.

Theorem 9. System (29) is uniformly bounded provided that $\tau(t)$ satisfies $0 \leq \tau(t) \leq \tau, \dot{\tau}(t) \leq \mu \leq 1$, and there exist some matrices $P > 0, Q_i \geq 0$, and $U_i = \text{diag}(u_{i1}, \dots, u_{in}) \geq 0 (i = 1, 2, 3, 4)$ such that the following linear matrix inequality holds:

$$\Sigma = \begin{pmatrix} \Sigma_{11} & 0 & PA + L_2 U_1 & PB & M_2 U_4 & PD \\ * & \Sigma_{22} & 0 & L_2 U_2 & 0 & 0 \\ * & * & \Sigma_{33} & 0 & 0 & 0 \\ * & * & * & \Sigma_{44} & 0 & 0 \\ * & * & * & * & \Sigma_{55} & 0 \\ * & * & * & * & * & -U_3 \end{pmatrix} \tag{30}$$

$< 0,$

where $*$ denotes the corresponding symmetric terms, $\Sigma_{ii} (i = 1, 2, 3, 4, 5), L_1, L_2, M_1, M_2$ are the same as in Theorem 3.

Proof. From $\Sigma < 0$, there exists a sufficiently small $\lambda \in (0, \nu)$ such that

$$\Gamma = \begin{pmatrix} \Gamma_{11} & 0 & PA + L_2U_1 & PB & M_2U_4 & PD \\ * & \Gamma_{22} & 0 & L_2U_2 & 0 & 0 \\ * & * & \Gamma_{33} & 0 & 0 & 0 \\ * & * & * & \Gamma_{44} & 0 & 0 \\ * & * & * & * & \Gamma_{55} & 0 \\ * & * & * & * & * & -U_3 \end{pmatrix} \quad (31)$$

< 0 ,

where I is identity matrix, $\Gamma_{11} = e^{\lambda\tau}Q_1 + \tau Q_2 - PC - CP - 2L_1U_1 - 2M_1U_4 + 2\lambda P + 2\lambda I$, $\Gamma_{22} = \Sigma_{22} + \lambda I$, $\Gamma_{33} = \lambda I + e^{\lambda\tau}Q_3 + \tau Q_4 - 2U_1$, $\Gamma_{44} = \Sigma_{44} + \lambda I$, and $\Gamma_{55} = \Sigma_{55} + \lambda I$.

We still consider the Lyapunov-Krasovskii functional $V(x(t))$ in (13). From (16)–(21), one may obtain

$$\begin{aligned} \|x(t)\|^2 &\leq \lambda_{\min}^{-1}(P) \left(e^{-\lambda t} V(x(0), 0) + \lambda^{-1} C_1 \right) \\ &\leq \lambda_{\min}^{-1}(P) \left(V(x(0), 0) + \lambda^{-1} C_1 \right) \\ &\leq \lambda_{\min}^{-1}(P) \left[\lambda_{\max}(P) \|\xi\|^2 \right. \\ &\quad + \sum_{j=1}^n u_{3j} \int_0^\infty k_j(\theta) \\ &\quad \times \int_{-\theta}^0 e^{\lambda(s+\theta)} g_j^2(x_j(s)) ds d\theta \\ &\quad + \int_{-\tau}^0 e^{\lambda(s+\tau)} \\ &\quad \times \left[x^T(s) Q_1 x(s) \right. \\ &\quad \left. + f^T(x(s)) Q_3 f(x(s)) \right] ds \\ &\quad + \int_{-\tau}^0 \int_s^0 e^{\lambda\theta} \\ &\quad \times \left[x^T(\theta) Q_2 x(\theta) \right. \\ &\quad \left. + f^T(x(\theta)) Q_4 f(x(\theta)) \right] d\theta ds \\ &\quad \left. + \lambda^{-1} C_1 \right], \end{aligned} \quad (32)$$

where $\|\xi\|^2 = \sup_{t \leq 0} \|x(t)\|^2$, C_1 is the same as in (24). Note that

$$\begin{aligned} &\sum_{j=1}^n u_{3j} \int_0^\infty k_j(\theta) \int_{-\theta}^0 e^{\lambda(s+\theta)} g_j^2(x_j(s)) ds d\theta \\ &\leq \sum_{j=1}^n 2u_{3j} \int_0^\infty k_j(\theta) \int_{-\theta}^0 e^{\lambda(s+\theta)} \left[\max_{1 \leq j \leq n} \left\{ (m_j^-)^2, (m_j^+)^2 \right\} x_j^2(s) \right. \\ &\quad \left. + g_j^2(0) \right] ds d\theta \end{aligned}$$

$$\begin{aligned} &\leq \sum_{j=1}^n 2u_{3j} \int_0^\infty k_j(\theta) \int_{-\theta}^0 e^{\lambda(s+\theta)} \left[\max_{1 \leq j \leq n} \left\{ (m_j^-)^2, (m_j^+)^2 \right\} \right. \\ &\quad \left. \times \|x(s)\|^2 + \|g(0)\|^2 \right] ds d\theta \\ &\leq \left(\sum_{j=1}^n 2u_{3j} \lambda^{-1} \bar{k}_j(\nu) \right) \\ &\quad \times \left[\max_{1 \leq j \leq n} \left\{ (m_j^-)^2, (m_j^+)^2 \right\} \|\xi\|^2 + \|g(0)\|^2 \right], \\ &\int_{-\tau}^0 e^{\lambda(s+\tau)} \left[x^T(s) Q_1 x(s) + f^T(x(s)) Q_3 f(x(s)) \right] ds \\ &\leq \int_{-\tau}^0 e^{\lambda(s+\tau)} \left[\lambda_{\max}(Q_1) \|x(s)\|^2 \right. \\ &\quad \left. + \lambda_{\max}(Q_3) \|f(x(s))\|^2 \right] ds \\ &\leq \int_{-\tau}^0 e^{\lambda(s+\tau)} \left[\lambda_{\max}(Q_1) \|\xi\|^2 + 2\lambda_{\max}(Q_3) \right. \\ &\quad \left. \times \left(\|f(0)\|^2 \right. \right. \\ &\quad \left. \left. + \max_{1 \leq i \leq n} \left\{ (l_i^-)^2, (l_i^+)^2 \right\} \|\xi\|^2 \right) \right] ds \\ &\leq \lambda^{-1} \left[\lambda_{\max}(Q_1) \|\xi\|^2 + 2\lambda_{\max}(Q_3) \right. \\ &\quad \left. \times \left(\|f(0)\|^2 + \max_{1 \leq i \leq n} \left\{ (l_i^-)^2, (l_i^+)^2 \right\} \|\xi\|^2 \right) \right] ds, \\ &\int_{-\tau}^0 \int_s^0 e^{\lambda\theta} \left[x^T(\theta) Q_2 x(\theta) + f^T(x(\theta)) Q_4 f(x(\theta)) \right] d\theta ds \\ &\leq \int_{-\tau}^0 \int_s^0 e^{\lambda\theta} \left[\lambda_{\max}(Q_2) \|\xi\|^2 + 2\lambda_{\max}(Q_4) \right. \\ &\quad \left. \times \left(\|f(0)\|^2 \right. \right. \\ &\quad \left. \left. + \max_{1 \leq i \leq n} \left\{ (l_i^-)^2, (l_i^+)^2 \right\} \|\xi\|^2 \right) \right] d\theta ds \\ &\leq \tau \lambda^{-1} \left[\lambda_{\max}(Q_2) \|\xi\|^2 + 2\lambda_{\max}(Q_4) \right. \\ &\quad \left. \times \left(\|f(0)\|^2 + \max_{1 \leq i \leq n} \left\{ (l_i^-)^2, (l_i^+)^2 \right\} \|\xi\|^2 \right) \right]. \end{aligned} \quad (33)$$

So, system (29) is uniformly bounded. \square




FIGURE 1: Time trajectories (a) as well as the phase portrait (b) for the system in Example 1.

4. One Example

Example 1. Consider system (1) with $J = (0, 1)^T$, $K(t) = \text{diag}(e^{-t}, 2e^{-2t})$, $\nu = 0.5$, and

$$\begin{aligned} A &= \begin{pmatrix} -0.1 & 0.4 \\ 0.2 & -0.5 \end{pmatrix}, & B &= \begin{pmatrix} 0.1 & -1 \\ -1.4 & 0.4 \end{pmatrix}, \\ C &= \begin{pmatrix} 1.4 & 0 \\ 0 & 1.65 \end{pmatrix}, \\ D &= \begin{pmatrix} -0.6 & 0.7 \\ 1 & 1.15 \end{pmatrix}, & \sigma_1 &= \begin{pmatrix} 0.23 & 0.1 \\ 0.3 & 0.2 \end{pmatrix}, \\ & & \sigma_2 &= \begin{pmatrix} 0.1 & -0.2 \\ 0.2 & 0.3 \end{pmatrix}. \end{aligned} \quad (34)$$

The activation functions $f_i(x_i) = x_i + \sin(x_i)$, $g_i(x_i) = \tanh(x_i)$ ($i = 1, 2$) satisfy, $l_i^- = m_i^- = 0$, $l_i^+ = m_i^+ = 1$. Then, one computes $L_1 = M_1 = 0$, $L_2 = M_2 = \text{diag}(1, 1)$, and $\bar{K}(0.5) = \text{diag}(2, 4/3)$. By using MATLAB's LMI Control Toolbox [34], for $\mu = 0.0035$ and $\tau = 1$, based on Theorem 3, such system is stochastically ultimately bounded when P , U_i , and Q_i ($i = 1, 2, 3, 4$) satisfy

$$\begin{aligned} P &= \begin{pmatrix} 1.7748 & 0.2342 \\ 0.2342 & 1.9398 \end{pmatrix}, & U_1 &= \begin{pmatrix} 1.3091 & 0 \\ 0 & 2.1602 \end{pmatrix}, \\ U_2 &= \begin{pmatrix} 218.3215 & 0 \\ 0 & 274.1607 \end{pmatrix}, & U_3 &= \begin{pmatrix} 1.0825 & 0 \\ 0 & 1.4021 \end{pmatrix}, \\ U_4 &= \begin{pmatrix} 2.1620 & 0 \\ 0 & 1.8493 \end{pmatrix}, & Q_1 &= \begin{pmatrix} 275.3435 & 104.0781 \\ 104.0781 & 397.1452 \end{pmatrix}, \\ Q_2 &= \begin{pmatrix} 31.3129 & 31.4377 \\ 31.4377 & 74.4756 \end{pmatrix}, & Q_3 &= \begin{pmatrix} 1.0857 & -1.2164 \\ -1.2164 & 2.9090 \end{pmatrix}, \\ & & Q_4 &= \begin{pmatrix} 189.3661 & 16.3951 \\ 16.3951 & 92.8987 \end{pmatrix}. \end{aligned} \quad (35)$$

From Figure 1, it is easy to see that $x(t)$ is stochastically ultimately bounded.

5. Conclusions

A proper Lyapunov functional and linear matrix inequalities are employed to investigate the ultimate boundedness, stability, and weak attractor of stochastic Hopfield neural networks with both time-varying and continuously distributed delays. New results and sufficient criteria are derived after extensive deductions. From the proposed sufficient conditions, one can easily prove that zero solution of such network is mean square exponentially stable and almost surely exponentially stable by applying the semimartingale convergence theorem.

Acknowledgments

The authors thank the editor and the reviewers for their insightful comments and valuable suggestions. This work was supported by the National Natural Science Foundation of China (nos. 10801109, 11271295, and 11047114), Science and Technology Research Projects of Hubei Provincial Department of Education (D20131602), and Young Talent Cultivation Projects of Guangdong (LYM09134).

References

- [1] C. Bai, "Stability analysis of Cohen-Grossberg BAM neural networks with delays and impulses," *Chaos, Solitons & Fractals*, vol. 35, no. 2, pp. 263–267, 2008.
- [2] T. Lv and P. Yan, "Dynamical behaviors of reaction-diffusion fuzzy neural networks with mixed delays and general boundary conditions," *Communications in Nonlinear Science and Numerical Simulation*, vol. 16, no. 2, pp. 993–1001, 2011.
- [3] J. Liu, X. Liu, and W.-C. Xie, "Global convergence of neural networks with mixed time-varying delays and discontinuous neuron activations," *Information Sciences*, vol. 183, pp. 92–105, 2012.

- [4] H. Huang and G. Feng, "Delay-dependent stability for uncertain stochastic neural networks with time-varying delay," *Physica A*, vol. 381, no. 1-2, pp. 93–103, 2007.
- [5] W.-H. Chen and X. M. Lu, "Mean square exponential stability of uncertain stochastic delayed neural networks," *Physics Letters A*, vol. 372, no. 7, pp. 1061–1069, 2008.
- [6] M. Hua, X. Liu, F. Deng, and J. Fei, "New results on robust exponential stability of uncertain stochastic neural networks with mixed time-varying delays," *Neural Processing Letters*, vol. 32, no. 3, pp. 219–233, 2010.
- [7] P. Balasubramaniam, S. Lakshmanan, and R. Rakkiyappan, "LMI optimization problem of delay-dependent robust stability criteria for stochastic systems with polytopic and linear fractional uncertainties," *International Journal of Applied Mathematics and Computer Science*, vol. 22, no. 2, pp. 339–351, 2012.
- [8] C. Huang and J. Cao, "Almost sure exponential stability of stochastic cellular neural networks with unbounded distributed delays," *Neurocomputing*, vol. 72, no. 13–15, pp. 3352–3356, 2009.
- [9] H. Zhao, N. Ding, and L. Chen, "Almost sure exponential stability of stochastic fuzzy cellular neural networks with delays," *Chaos, Solitons & Fractals*, vol. 40, no. 4, pp. 1653–1659, 2009.
- [10] C. Huang, P. Chen, Y. He, L. Huang, and W. Tan, "Almost sure exponential stability of delayed Hopfield neural networks," *Applied Mathematics Letters*, vol. 21, no. 7, pp. 701–705, 2008.
- [11] C. Huang and J. Cao, "On p th moment exponential stability of stochastic Cohen-Grossberg neural networks with time-varying delays," *Neurocomputing*, vol. 73, no. 4–6, pp. 986–990, 2010.
- [12] C. X. Huang, Y. G. He, L. H. Huang, and W. J. Zhu, " p th moment stability analysis of stochastic recurrent neural networks with time-varying delays," *Information Sciences*, vol. 178, no. 9, pp. 2194–2203, 2008.
- [13] C. Huang, Y. He, and H. Wang, "Mean square exponential stability of stochastic recurrent neural networks with time-varying delays," *Computers and Mathematics with Applications*, vol. 56, no. 7, pp. 1773–1778, 2008.
- [14] R. Rakkiyappan and P. Balasubramaniam, "Delay-dependent asymptotic stability for stochastic delayed recurrent neural networks with time varying delays," *Applied Mathematics and Computation*, vol. 198, no. 2, pp. 526–533, 2008.
- [15] S. Lakshmanan and P. Balasubramaniam, "Linear matrix inequality approach for robust stability analysis for stochastic neural networks with time-varying delay," *Chinese Physics B*, vol. 20, no. 4, Article ID 040204, 2011.
- [16] S. Lakshmanan, A. Manivannan, and P. Balasubramaniam, "Delay-distribution-dependent stability criteria for neural networks with time-varying delays," *Dynamics of Continuous, Discrete & Impulsive Systems A*, vol. 19, no. 1, pp. 1–14, 2012.
- [17] Q. Song and Z. Wang, "Stability analysis of impulsive stochastic Cohen-Grossberg neural networks with mixed time delays," *Physica A*, vol. 387, no. 13, pp. 3314–3326, 2008.
- [18] X. Li, "Existence and global exponential stability of periodic solution for delayed neural networks with impulsive and stochastic effects," *Neurocomputing*, vol. 73, no. 4–6, pp. 749–758, 2010.
- [19] C. H. Wang, Y. G. Kao, and G. W. Yang, "Exponential stability of impulsive stochastic fuzzy reaction-diffusion Cohen-Grossberg neural networks with mixed delays," *Neurocomputing*, vol. 89, pp. 55–63, 2012.
- [20] Q. Zhu, C. Huang, and X. Yang, "Exponential stability for stochastic jumping BAM neural networks with time-varying and distributed delays," *Nonlinear Analysis*, vol. 5, no. 1, pp. 52–77, 2011.
- [21] P. Balasubramaniam and M. Syed Ali, "Stochastic stability of uncertain fuzzy recurrent neural networks with Markovian jumping parameters," *International Journal of Computer Mathematics*, vol. 88, no. 5, pp. 892–904, 2011.
- [22] Y. F. Wang, P. Lin, and L. S. Wang, "Exponential stability of reaction-diffusion high-order Markovian jump Hopfield neural networks with time-varying delays," *Nonlinear Analysis*, vol. 13, no. 3, pp. 1353–1361, 2012.
- [23] P. Balasubramaniam and G. Nagamani, "Global robust passivity analysis for stochastic interval neural networks with interval time-varying delays and Markovian jumping parameters," *Journal of Optimization Theory and Applications*, vol. 149, no. 1, pp. 197–215, 2011.
- [24] P. Balasubramaniam and G. Nagamani, "Global robust passivity analysis for stochastic fuzzy interval neural networks with time-varying delays," *Expert Systems With Applications*, vol. 39, pp. 732–742, 2012.
- [25] P. Wang, D. Li, and Q. Hu, "Bounds of the hyper-chaotic Lorenz-Stenflo system," *Communications in Nonlinear Science and Numerical Simulation*, vol. 15, no. 9, pp. 2514–2520, 2010.
- [26] X. Liao and J. Wang, "Global dissipativity of continuous-time recurrent neural networks with time delay," *Physical Review E*, vol. 68, no. 1, Article ID 016118, 7 pages, 2003.
- [27] S. Arik, "On the global dissipativity of dynamical neural networks with time delays," *Physics Letters A*, vol. 326, no. 1-2, pp. 126–132, 2004.
- [28] X. Y. Lou and B. T. Cui, "Global robust dissipativity for integro-differential systems modeling neural networks with delays," *Chaos, Solitons & Fractals*, vol. 36, no. 2, pp. 469–478, 2008.
- [29] Q. Song and Z. Zhao, "Global dissipativity of neural networks with both variable and unbounded delays," *Chaos, Solitons & Fractals*, vol. 25, no. 2, pp. 393–401, 2005.
- [30] H. Jiang and Z. Teng, "Global exponential stability of cellular neural networks with time-varying coefficients and delays," *Neural Networks*, vol. 17, no. 10, pp. 1415–1425, 2004.
- [31] H. Jiang and Z. Teng, "Boundedness, periodic solutions and global stability for cellular neural networks with variable coefficients and infinite delays," *Neurocomputing*, vol. 72, no. 10–12, pp. 2455–2463, 2009.
- [32] L. Wan and Q. H. Zhou, "Attractor and ultimate boundedness for stochastic cellular neural networks with delays," *Nonlinear Analysis*, vol. 12, no. 5, pp. 2561–2566, 2011.
- [33] L. Wan, Q. Zhou, P. Wang, and J. Li, "Ultimate boundedness and an attractor for stochastic Hopfield neural networks with time-varying delays," *Nonlinear Analysis*, vol. 13, no. 2, pp. 953–958, 2012.
- [34] S. Boyd, L. El Ghaoui, E. Feron, and V. Balakrishnan, *Linear Matrix Inequalities in System and Control Theory*, SIAM, Philadelphia, Pa, USA.
- [35] X. Mao, *Stochastic Differential Equations and Applications*, Horwood Publishing, 1997.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

