

Corrigendum

Corrigendum to “Stevioside from *Stevia rebaudiana* Bertoni Increases Insulin Sensitivity in 3T3-L1 Adipocytes”

Nabilatul Hani Mohd-Radzman,¹ Wan Iryani Wan Ismail,¹ Siti Safura Jaapar,¹ Zainah Adam,² and Aishah Adam¹

¹Faculty of Pharmacy, Universiti Teknologi MARA, Puncak Alam Campus, 42300 Bandar Puncak Alam, Selangor, Malaysia

²Medical Technology Division, Malaysian Nuclear Agency, Bangi, 43000 Kajang, Selangor, Malaysia

Correspondence should be addressed to Wan Iryani Wan Ismail; waniryani@gmail.com

Received 28 December 2015; Accepted 21 June 2016

Copyright © 2016 Nabilatul Hani Mohd-Radzman et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The authors would like to indicate that there has been an inadvertent error in “Stevioside from *Stevia rebaudiana* Bertoni Increases Insulin Sensitivity in 3T3-L1 Adipocytes” [1].

The Western blot image of Figure 6 that was uploaded for the β -actin protein bands is incorrect. The corrected image of the β -actin protein for Figure 6 is hereby attached. This error does not alter or affect the discussions and conclusions that have been reached in this paper.

Also in Figure 6, the pIRS1 bands were rearranged to match the pY20 group, because the blots were run in a different order.

References

- [1] N. H. Mohd-Radzman, W. I. W. Ismail, S. S. Jaapar, Z. Adam, and A. Adam, “Stevioside from *Stevia rebaudiana* Bertoni increases insulin sensitivity in 3T3-L1 adipocytes,” *Evidence-Based Complementary and Alternative Medicine*, vol. 2013, Article ID 938081, 8 pages, 2013.




FIGURE 6: Band intensities observed via Western blotting, showing the different expression levels of phosphorylated insulin receptor substrate 1 (p-IRS1) and phosphorylated tyrosine (pY20), in groups treated with stevioside (Svs) and rosiglitazone maleate (RM) in comparison to the normal and insulin-resistant (IR) groups. β -actin was used as a loading control. The experiment was repeated thrice.



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

