

## ORAL PRESENTATION

## Open Access

# Cardiovascular impairment in young adults horizontally infected with HIV-1 in childhood

Florentina Dumitrescu<sup>1,2\*</sup>, Dina Maria Cupşa<sup>2</sup>, Andreea Cristina Stoian<sup>1,2</sup>, Augustin Cupşa<sup>1,2</sup>, Irina Niculescu<sup>1,2</sup>, Lucian Giubelan<sup>1,2</sup>, Daniela Culman<sup>2</sup>

From The 9th Edition of the Scientific Days of the National Institute for Infectious Diseases Prof Dr Matei Bals Bucharest, Romania. 23-25 October 2013

## Background

We aimed to evaluate the prevalence and spectrum of cardiovascular impairment (CVI) in young HIV/AIDS patients in evidence at the HIV/AIDS Craiova Regional Center.

## Methods

We performed a retrospective study between 01 January 1996 – 31 December 2005 and a prospective study between 01 June 2005 – 30 June 2010; it targeted epidemiological issues, the clinical spectrum of CVI in patient with horizontal HIV infection, acquired during 1987-1990 (lot P) compared to patients with other transmission routes of infection (lot S).

## Results

CVI was encountered in 61 (10.23%) of 596 patients parenterally-infected with HIV-1 in childhood and 32 (16.16%) of 198 patients infected through sexual route, with statistically significant differences in the prevalence of CVI depending on the transmission route of HIV-1 ( $p\text{Chi}^2=0.02$ ), probably being underestimated in pre-HAART era. The average age at the moment of CVI was  $9.02\pm 4.76$  years in lot P vs.  $41.47\pm 12.41$  years in lot S ( $p < 0.0001$ ). The spectrum of CVI was dominated by arrhythmias and conduction blocks 42 patients (45.16%), dilated cardiomyopathy (DCM) 18 patients (19.35%), pulmonary arterial hypertension 17 patients (18.27%). The most common entity identified in parenterally-infected patients in early childhood was DCM 16 patients lot P vs. 2 patients lot S ( $p$  Fischer test=0.026), found in moderate and severe immunosuppressed patients, with virological failure, often associated with respiratory

disease. In sexually-infected patients diseases due to atherosclerosis prevailed 17 patients ( $p$  Fisher test $<0.0001$ ). The presence of high blood pressure in 2 young people with HIV/AIDS raises the issue of cardiovascular risk and occurrence of coronary heart disease in young patients. Although the survival rate of patients with CVI ( $15.21 \pm 0.62$  years) was similar to that of the patients without CVI ( $13.78 \pm 0.25$  years), for patients with CVI, death due to direct cardiac damage was significant: 9 patients (52.94%).

## Conclusion

The prevalence of CVI was low, probably overlooked and underdiagnosed, at least in early ages and young adults, dominated by arrhythmias and DCM; monitoring and management in individual care programs constitutes a way for improving the quality of life and prognosis for people with HIV/AIDS.

## Authors' details

<sup>1</sup>University of Medicine and Pharmacy Craiova, Romania. <sup>2</sup>"Victor Babes" Clinical Hospital of Infectious Diseases and Pneumology, Craiova, Romania.

Published: 16 December 2013

doi:10.1186/1471-2334-13-S1-O28

Cite this article as: Dumitrescu et al.: Cardiovascular impairment in young adults horizontally infected with HIV-1 in childhood. *BMC Infectious Diseases* 2013 **13**(Suppl 1):O28.

\* Correspondence: [dumitrescu\\_florentina@yahoo.com](mailto:dumitrescu_florentina@yahoo.com)

<sup>1</sup>University of Medicine and Pharmacy Craiova, Romania

Full list of author information is available at the end of the article