

Corrigendum

Corrigendum to “Intraspecific and Intracolonial Variation in the Profile of Venom Alkaloids and Cuticular Hydrocarbons of the Fire Ant *Solenopsis saevissima* Smith (Hymenoptera: Formicidae)”

Eduardo Gonçalves Paterson Fox,¹ Adriana Pianaro,² Daniel Russ Solis,³
Jacques Hubert Charles Delabie,⁴ Bruno Cunha Vairo,⁵
Ednildo de Alcântara Machado,¹ and Odair Correa Bueno³

¹Laboratório de Entomologia Médica e Molecular, Instituto de Biofísica Carlos Chagas Filho,
Universidade Federal do Rio de Janeiro (IBCCF/UFRJ), Rio de Janeiro, RJ, Brazil

²Instituto de Química, Universidade Estadual de Campinas (IQ/UNICAMP), 13083-862 Campinas, SP, Brazil

³Instituto de Biociências, Universidade Estadual Paulista (UNESP),
Campus de Rio Claro and Centro de Estudos de Insetos Sociais (CEIS), 13506-900 Rio Claro, SP, Brazil

⁴Laboratório de Mirmecologia, Centro de Pesquisas do Cacau (CEPLAC), Itabuna, BA, Brazil

⁵Laboratório de Tecido Conjuntivo, Instituto de Bioquímica Médica, UFRJ, RJ, Brazil

Correspondence should be addressed to Eduardo Gonçalves Paterson Fox; ofoxofox@gmail.com





Received 14 July 2016; Accepted 17 October 2016

Copyright © 2016 Eduardo Gonçalves Paterson Fox et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the article titled “Intraspecific and Intracolonial Variation in the Profile of Venom Alkaloids and Cuticular Hydrocarbons of the Fire Ant *Solenopsis saevissima* Smith (Hymenoptera: Formicidae)” [1], the method description was incomplete and inaccurate. Thus, the last part of the “2.4. GC-MS Analyses” section “The injector temperature was 250°C. Oven temperature was programmed to increase at 12°C/min from 50°C to 330°C, with a final hold time of 1 min” should be corrected to “The injector temperature was 250°C. Oven temperature was programmed to increase at 12°C/min from 50°C to 290°C with a hold pause for 6 minutes, followed by an increase of 30°C/min from 290°C to 320°C followed by a final hold time of 1 min.”

References

- [1] E. G. P. Fox, A. Pianaro, D. R. Solis et al., “Intraspecific and intracolonial variation in the profile of venom alkaloids and cuticular hydrocarbons of the fire ant *Solenopsis saevissima* smith (Hymenoptera: Formicidae),” *Psyche*, vol. 2012, Article ID 398061, 10 pages, 2012.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

