



POSTER PRESENTATION

Open Access

Inverse dose-response to gp140 YU2 foldon trimer formulated with aluminum phosphate and ISCOMATRIX[®] adjuvants

A Wilson*, R Powell, S Hoffenberg, A Carpov, H Arendt, J DeStefano, MJ Caulfield

From AIDS Vaccine 2012

Boston, MA, USA. 9-12 September 2012

Background

Conventional vaccine approaches based on delivery of HIV-1 envelope (Env) proteins or peptides derived from Env sequences have failed to generate broadly neutralizing antibodies (bNAbs) to the virus. Even with large doses (200 ug) of adjuvanted gp120 proteins administered multiple times to human volunteers, the subsequent antibody response boosts only moderately with each succeeding vaccination, and titers drop precipitously thereafter. We hypothesized that the usual practice of administering a moderate to high antigen doses may be counter productive to the goal of eliciting durable, high-affinity antibody responses.

Methods

We conducted a rabbit immunogenicity study in which we compared the anti-gp120 antibody response of rabbits immunized with low (1 ug), medium (10 ug) and high (100 ug) quantities of gp140 YU2 foldon trimer (FT) formulated with aluminum phosphate (alum) alone or in combination with ISCOMATRIX[®] adjuvant. In addition, we used a more protracted vaccination regimen by administering the vaccine at 0, 8, and 24 week time points.

Results

Antibody responses elicited by the different YU2 FT vaccination regimens were quantified by ELISA against JRCSF gp120 protein after vaccination showing weak responses after the first two vaccine doses. However, after 3 doses, the responses to vaccines co-formulated with both ISCOMATRIX[®] and alum were markedly higher than the corresponding responses to the antigen

formulated with alum only. Interestingly, at 4 weeks post-dose 3, there was a reverse dose response effect, with the 1 ug dose group having higher titers than the 10 and 100 ug groups. At 12 weeks post-dose 3, the antibody GMTs were 28,078 (1 ug), 9,681 (10 ug) and 7,253 (100 ug).

Conclusion

The results showing that the low dose group maintained a response ~4 fold higher than the high dose group suggests that durability of the antibody response to HIV Env may be a function of antigen dose.

Published: 13 September 2012

doi:10.1186/1742-4690-9-S2-P14

Cite this article as: Wilson *et al.*: Inverse dose-response to gp140 YU2 foldon trimer formulated with aluminum phosphate and ISCOMATRIX[®] adjuvants. *Retrovirology* 2012 **9**(Suppl 2):P14.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

International AIDS Vaccine Initiative, Brooklyn, NY, USA



© 2012 Wilson *et al.*; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.