

RESEARCH

Open Access

Tuberculosis examination using whole blood interferon-gamma release assay among health care workers in a Japanese hospital without tuberculosis-specific wards

Eisuke Adachi^{1*}, Mika Kogayu², Takeshi Fujii¹, Hiroshi Mae³, Shoichi Shimizu¹, Yuichi Iwai³, Hiroko Shibata⁴, Masato Suzuki⁴, Kohzoh Imai⁵ and Tomohiko Koibuchi¹

Abstract

Occupational latent tuberculosis infection (LTBI) among health care workers (HCWs) is an important public health issue. The objective of this study is to assess prevalence and risk factors of LTBI among Japanese HCWs by Quantiferon-TB Gold in Tube (QFT-GIT) and the structured questionnaire. This is a cross-sectional study involving HCWs from a hospital without tuberculosis-specific wards, receiving QFT-GIT for LTBI screening. We reviewed medical records of HCWs and questioned HCWs about exposure to *M. tuberculosis* and employment length in health care industries. 165 HCWs, approximately 80% of the total hospital staff, were enrolled in this study. 18 out of 165 subjects had positive results, suggesting LTBI prevalence rate of 11%. Multiple regression analysis revealed a significant association between the positive or intermediate QFT-GIT results and history of contact investigation for tuberculosis. QFT-GIT positivity rate among HCWs is higher than among general population in Japan.

Keywords: Latent tuberculosis infection; Quantiferon-TB gold in tube; Latent tuberculosis infection baseline screening; Health care workers

Background

Occupational latent tuberculosis infection (LTBI) among health care workers (HCWs) is an important public health issue. World Health Organization (WHO) estimate of tuberculosis incidence of Japan in 2011 was 20 per 100,000 population per year. The Tuberculosis Surveillance Center of The research institute of Tuberculosis, Japan Anti-tuberculosis Association reported the incidence of tuberculosis of Japan was 18.2 in 2010 and 17.7 in 2011 (Rates per 100,000) and had continued to decline. However, approximately 80% of nurses with tuberculosis were speculated to be infected by nosocomial infection in Japan (Ohmori et al. 2007). The screening for LTBI by means of the tuberculin skin test (TST) or whole blood interferon-gamma release assays (IGRAs) and the

administration of chemoprophylaxis have been shown to be effective for prevention to develop active tuberculosis (Ringshausen et al. 2012; Pai et al. 2004). In Japan, TST use is limited for screening of LTBI, because most Japanese residents have received bacille Calmette-Guérin (BCG) vaccination on multiple occasions. Accordingly, The Japanese Society of Tuberculosis recommend the Quantiferon®-TB Gold (QFT-G) test for the screening of *Mycobacterium tuberculosis* (*M. tuberculosis*) infection for HCWs or health care students (Hotta et al. 2007).

Most of the previous studies described the prevalence of LTBI among Japanese HCWs were performed in tuberculosis referral hospitals. The objective of this study is to assess prevalence and risk factors of LTBI among Japanese HCWs by Quantiferon-TB Gold in Tube (QFT-GIT), a new version of QFT-G, and the structured questionnaire in a hospital without tuberculosis-specific wards.

* Correspondence: e-adachi@ims.u-tokyo.ac.jp

¹Department of Infectious Diseases and Applied Immunology, IMSUT Hospital of The Institute of Medical Science, The University of Tokyo, 4-6-1 Shirokanedai, Minato-ku, Tokyo 108-8639, Japan

Full list of author information is available at the end of the article

Methods

Setting and design

This is a cross-sectional study involving HCWs at the hospital affiliated with the Institute of Medical Science, the University of Tokyo (IMSUT hospital), receiving QFT-GIT for LTBI screening between November 2011 and July 2012. IMSUT Hospital has 135 beds and four negative pressure isolation rooms for patients diagnosed with active tuberculosis accidentally. The hospital is not a tuberculosis referral hospital and has no tuberculosis-specific wards. There were three active tuberculosis patients admitted in the IMSUT hospital in 2007. This is the highest number of annual admission of active tuberculosis cases in the hospital. We reviewed medical records of HCWs and questioned HCWs about exposure to *M. tuberculosis* and employment length in health care industries. We utilized the questionnaire consisting of the following questions: age, job category, history of contact investigation for tuberculosis, history of living with families who developed active tuberculosis, past history of active/latent tuberculosis infection, length of working in health care industries (general hospitals, tuberculosis referral hospitals, clinics, nursing home), history of working in tuberculosis ward/out patient department (OPD). Eligible subjects had been working as HCWs for more than 6 months and never had received IGRAs for LTBI screening. We excluded the HCWs with history of active tuberculosis. Written informed consent was obtained from subjects before completing the study questionnaire. This study was approved by the Research Ethics Committee, Institutional Review Board of IMSUT (accession number: 24-6, 24-32).

Quantiferon®-TB Gold in Tube

Blood samples for QFT-GIT tests (Cellestis Limited, Australia) were collected at our hospital and QFT-GIT tests were performed at laboratories in the clinical laboratory testing industry (SRL Inc. Japan). Interferon-gamma (IFN- γ) responses to antigens that are at least 0.35 IU/mL greater than the nil control value are considered *M. tuberculosis* infection ("positive"). We also evaluated a prevalence using the value of 0.10 IU/mL as an alternative cut-off ("intermediate") as the guideline of committees for the preventions of The Japanese Society of Tuberculosis. If the response is less than a cut-off value and their response to the mitogen positive control is greater than 0.5 IU/mL, the test result is considered negative.

Statistical analysis

Comparisons of proportions were analyzed by the chi-square test or the Fisher exact test and a linear trend in proportions of ordered multiple groups were analyzed by Mantel-extension test. Multiple logistic regression

analysis was performed (using JMP 9 software; SAS Institute Inc.) with the following independent variables: male/female, age groups, job category, history of contact investigation for tuberculosis, history of living with families who developed active tuberculosis and history of working in tuberculosis ward or OPD for more than 2 months.

Results

165 HCWs, approximately 80% of the total hospital staff, were enrolled in this study. All subjects were Japanese. Table 1 shows the QFT-GIT positivity rates and characteristics of HCWs. 18 out of 165 subjects (10.9%) had positive results (0.35 IU/mL, cut-off). None of them manifested signs of active tuberculosis by their chest radiograph. 38 subjects (23%) had the value of QFT-GIT showed greater than 0.10 IU/mL. Notably, among HCWs aged 30-39 years, 14% of subjects (10/70) had positive QFT-GIT result. Table 2 shows the results of univariate analysis of tuberculosis infection risk factors. Univariate analysis revealed a significant association between the positive or intermediate QFT-GIT results and history of contact investigation for tuberculosis (odds ratio (OR) 3.4, 95% confidence interval (CI) 1.1 to 10.0). Neither age nor length of working in health care industries was associated with results of QFT-GIT among all HCWs. Among nurses and doctors aged 20-49 years, the percentage of HCWs with positive QFT-GIT results increased for each decade of age (odds ratios relative to persons aged 20-29 years, 2.5

Table 1 Characteristics of HCWs and Quantiferon TB-Gold in tube positivity rate

Total (number)		165
Gender	male	60 (36.4%)
	female	105 (63.4%)
Age (years, (IQR))		38 ^a (32 - 46)
Job category	nurse	72 (43.6%)
	doctor	53 (32.1%)
	laboratory*	12 (7.27%)
	pharmacist	12 (7.27%)
	other**	16 (9.70%)
Length of working in health care industries (years, (IQR))		13 ^a (7 - 18)
History of contact investigation for TB		25 (15.2%)
History of living with families who developed active TB		7 (4.24%)
History of working in TB ward/OPD		14 (8.48%)
QFT-GIT positivity (%)	Cut off: 0.35 IU/mL	18 (10.9%)
	Cut off: 0.10 IU/mL	38 (23.0%)

Note. *IQR*, interquartile range; *TB*, tuberculosis; *OPD*, outpatient department; *QFT-GIT*, Quantiferon TB-Gold in tube; * Laboratory technician ** radiography technicians, nutritionist, clerical, nursing assistants and social workers; ^amedian.

Table 2 Association between results of Quantiferon TB-Gold in tube and risk factors for tuberculosis infection

	No of subjects				No of subjects		
	Cut off: 0.35 IU/mL				Cut off: 0.1 IU/mL		
	≥ 0.35	≤ 0.35	OR(95% CI)		≥ 0.1	≤ 0.1	OR(95% CI)
Male	10	50	2.4 (0.90 - 6.5, P = 0.12 ^b)	15	45	1.2 (0.56 - 2.5, P = 0.7 ^b)	
Female	8	97		23	82		
Job category							
Nurse	6	66	1.0 ^a	14	58	1.0 ^a	P = 0.69 ^b
Doctor	5	48	1.1	12	41	1.2	
Laboratory*	3	9	3.7	4	8	2.1	
Pharmacist	3	9	3.7	3	9	1.4	
other**	1	15	0.73	5	11	1.9	
age							
20-29	2	24	1.0 ^a	6	20	1.0 ^a	P = 0.63 ^b
30-39	10	60	2.0	17	53	1.1	
40-49	4	38	1.3	7	35	0.67	
50-	2	25	0.96	8	19	1.4	
Length of working in health care industries							
0-4	3	25	1.0 ^a	6	22	1.0 ^a	P = 0.87 ^b
5-9	1	30	0.28	5	26	0.71	
10-14	6	37	1.4	11	32	1.3	
15-19	6	23	2.3	7	22	1.2	
20-	2	32	0.52	9	25	1.3	
History of contact investigation for TB							
Yes	6	19	3.4 (1.1 - 10.0, P = 0.034 ^b)	12	13	4.0 (1.66 - 9.9, P = 0.003 ^b)	
No	12	128		26	114		
History of living with families who developed active TB							
Yes	2	5	3.6 (0.64 - 20, P = 0.17 ^b)	3	4	2.6 (0.56 - 12.0, P = 0.20 ^b)	
No	16	142		35	123		
History of working in TB ward/OPD							
Yes	1	13	0.61 (0.075 - 4.9, P = 1.0 ^b)	3	11	0.90 (0.24 - 3.4, P = 1.0 ^b)	
No	17	134		35	116		

Note. OR, odds ratio; CI, confidence interval; TB, tuberculosis; OPD, outpatient department; * Laboratory technician ** radiography technicians, nutritionist, clerical, nursing assistants and social workers; ^areference; ^bFisher Exact test.

(30-39 years), 2.9 (40-49 years)), but it did not reach statistical significance. Multiple logistic regression analysis to risk factors revealed that positive QFT-GIT result was significantly associated with male/female, the history of contact investigation for tuberculosis (Table 3).

Discussion

The utility of IGRAs in countries where BCG vaccine is widely used is supported by many studies (Pai et al. 2004). Harada et al. reported that a prevalence of LTBI among HCWs in a Japanese tuberculosis referral hospital was 9.9% by means of QFT-G (Harada et al. 2006). However, QFT-GIT, a new version of QFT-G, has higher sensitivity than QFT-G with high specificity for *M.*

tuberculosis infection and is the currently recommended method for screening of LTBI in HCWs (Harada et al. 2008). Mori et al. reported that the QFT-G positivity rate of general population was 7.1% (40-69 years); it increased with age from 3.1% for the 40-49 years age group to 9.8% for those aged 60-69 (Mori et al. 2007).

We conducted here a cross-sectional study for HCWs receiving QFT-GIT. It showed the prevalence of LTBI to be 11%, which is remarkably higher than that among the general population (Mori et al. 2007). The QFT-GIT positivity rate in our study is consistent with the previous report from a tuberculosis referral hospital in Japan (Harada et al. 2006). We found that the prevalence of LTBI among HCWs in general hospitals without tubercu-

Table 3 Multivariate analysis of association between results of Quantiferon TB-Gold in tube and risk factors for tuberculosis infection

	Cut off: 0.35 IU/mL		Cut off: 0.1 IU/mL	
	OR	P	OR	P
Male/ female	8.0	0.011	1.3	0.41
Job category				
Nurse	1.0 ^a		1.0 ^a	
Doctor	0.34	0.30	1.4	0.66
Laboratory technician	6.5	0.092	2.6	0.26
Pharmacist	4.9	0.096	1.7	0.51
Other*	1.5	0.76	2.9	0.13
Age				
20-29	1.0 ^a		1.0 ^a	
30-39	2.0	0.42	0.90	0.87
40-49	0.92	0.93	0.33	0.12
50-	0.23	0.26	0.83	0.78
History of contact investigation for TB	6.4	0.0077	4.7	0.0017
History of living with families who developed active TB	6.0	0.11	3.3	0.24
History of working in TB ward/OPD	0.62	0.70	0.67	0.62

Note. OR, odds ratio; TB, tuberculosis; OPD, outpatient department * radiography technicians, nutritionist, clerical, nursing assistants and social workers; ^areference.

losis wards and among HCWs in tuberculosis referral hospitals is similar. The report from the tuberculosis referral hospital also described that the positive QFT-G results were closely associated with age and with a history of working in a tuberculosis ward or an OPD of a tuberculosis clinic (Harada et al. 2006). In our study, positive QFT-GIT results were associated with the history of contact investigation for tuberculosis, that is suggestive of the history of exposure to *M. tuberculosis*. Of note, a high proportion of HCWs with positive/intermediate QFT-GIT results was laboratory technicians, which was consistent with the previous report (Harada et al. 2006). These suggest laboratory technicians are likely to be exposed to infectious agents in handling patients' samples, such as sputum. There were no laboratory technicians with the history of contact investigation for tuberculosis in our study. We would like to offer that laboratory technicians should be included as subjects in performing contact investigation.

Our study has several limitations, including the small sample size and a single-center study. Diagnosis of LTBI by means of single QFT-GIT test remains controversial in accuracy and reliability, because the specific IFN- γ response may wane considerably with time after infection and a substantial technical variability exists (Ringshausen et al. 2012; Mori et al. 2007; Pai and O'Brien 2007). Mori

et al. demonstrated that the probability of infection increases almost exponentially after 30 years of age (Mori 2000). The fact that age was not associated with results of QFT-GIT in our study may be due to the sample-size bias.

Our study is first to offer QFT-GIT positivity rate among Japanese HCWs in a hospital without tuberculosis-specific wards. To establish the evaluation of the LTBI by means of QFT-GIT test, several large, multi-center studies might be warranted.

Conclusions

QFT-GIT positivity rate among HCWs is higher than among general population in Japan. This study showed the prevalence of LTBI among Japanese HCWs was 11%. Occupational LTBI among HCWs is an important public health issue.

Competing interests

The authors declare that they have no competing interests

Authors' contributions

EA and MK designed the project and carried out research. EA drafted manuscript. TF, KI and TK contributed to project design and revised the drafted manuscript. HM, SS, YI, HS and MS contributed to data collections. All authors read and approved the final manuscript.

Acknowledgment

The authors thank Kei Ouchi, MD, for his comments on drafts of the manuscript.

Author details

¹Department of Infectious Diseases and Applied Immunology, IMSUT Hospital of The Institute of Medical Science, The University of Tokyo, 4-6-1 Shirokanedai, Minato-ku, Tokyo 108-8639, Japan. ²Department of Nursing, IMSUT Hospital of The Institute of Medical Science, The University of Tokyo, 4-6-1 Shirokanedai, Minato-ku, Tokyo 108-8639, Japan. ³Department of Pharmacy, IMSUT Hospital of The Institute of Medical Science, The University of Tokyo, 4-6-1 Shirokanedai, Minato-ku, Tokyo 108-8639, Japan. ⁴Department of Laboratory Medicine, IMSUT Hospital of The Institute of Medical Science, The University of Tokyo, 4-6-1 Shirokanedai, Minato-ku, Tokyo 108-8639, Japan. ⁵Director, IMSUT Hospital of The Institute of Medical Science, The University of Tokyo, 4-6-1 Shirokanedai, Minato-ku, Tokyo 108-8639, Japan.

Received: 13 June 2013 Accepted: 2 September 2013

Published: 5 September 2013

References

- Harada N, Nakajima Y, Higuchi K, Sekiya Y, Rothel J, Mori T (2006) Screening for tuberculosis infection using whole-blood interferon-gamma and Mantoux testing among Japanese healthcare workers. *Infect Control Hosp Epidemiol* 27:442-448
- Harada N, Higuchi K, Yoshiyama T, Kawabe Y, Fujita A, Sasaki Y, Horiba M, Mitarai S, Yonemaru M, Ogata H, Ariga H, Kurashima A, Wada A, Takamori M, Yamagishi F, Suzuki K, Mori T, Ishikawa N (2008) Comparison of the sensitivity and specificity of two whole blood interferon-gamma assays for *M. tuberculosis* infection. *J Infect* 56:348-353
- Hotta K, Ogura T, Nishii K, Kodani T, Onishi M, Shimizu Y, Kanehiro A, Kiura K, Tanimoto M, Tobe K (2007) Whole blood interferon-gamma assay for baseline tuberculosis screening among Japanese healthcare students. *PLoS One* 2:e803
- Mori T (2000) Recent trends in tuberculosis, Japan. *Emerg Infect Dis* 6(6):566-568
- Mori T, Harada N, Higuchi K, Sekiya Y, Uchimura K, Shimao T (2007) Waning of the specific interferon-gamma response after years of tuberculosis infection. *Int J Tuberc Lung Dis* 11:1021-1025

- Ohmori M, Hoshino H, Yamauchi Y, Uchimura K (2007) Current epidemiological situation of tuberculosis in the workplace: considering the risk of tuberculosis among nurses. *Kekkaku* 82:85–93
- Pai M, O'Brien R (2007) Serial testing for tuberculosis: can we make sense of T cell assay conversions and reversions? *PLoS Med* 4:e208
- Pai M, Riley LW, Colford JM Jr (2004) Interferon-gamma assays in the immunodiagnosis of tuberculosis: a systematic review. *Lancet Infect Dis* 4:761–776
- Ringshausen FC, Schablon A, Nienhaus A (2012) Interferon-gamma release assays for the tuberculosis serial testing of health care workers: a systematic review. *J Occup Med Toxicol* 7:6

doi:10.1186/2193-1801-2-440

Cite this article as: Adachi et al.: Tuberculosis examination using whole blood interferon-gamma release assay among health care workers in a Japanese hospital without tuberculosis-specific wards. *SpringerPlus* 2013 **2**:440.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
