

Poster presentation

Open Access

Dendritic cell-mediated infection of primary B cells with KSHV

R Bagni*¹, E Barsov², B Ortiz-Conde³, D Dittmer⁴, V Kewalramani⁵, D Ott², C Sadowski⁶, P Tuma⁷, F Ruscetti⁶ and D Whitby¹

Address: ¹Viral Oncology Section, AIDS and Cancer Virus Program, Basic Research Program, SAIC, National Cancer Institute, Frederick, Maryland, USA, ²Retroviral Assembly Section, AIDS and Cancer Virus Program, Basic Research Program, SAIC, National Cancer Institute, Frederick, Maryland, USA, ³Virus Technology Laboratory, Advanced Technology Program, SAIC, National Cancer Institute, Frederick, Maryland, USA, ⁴Department of Microbiology and Immunology and Lineberger Comprehensive Cancer, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA, ⁵Model Development Section, HIV Drug Resistance Program, National Cancer Institute, Frederick, Maryland, USA, ⁶Leukocyte Biology Section, Laboratory of Experimental Immunology, National Cancer Institute, Frederick, Maryland, USA and ⁷Department of Biology, Catholic University of America, Washington, DC, USA

* Corresponding author

from 11th International Conference on Malignancies in AIDS and Other Acquired Immunodeficiencies (ICMAOI): Basic, Epidemiologic, and Clinical Research
Bethesda, MD, USA. 6–7 October 2008

Published: 17 June 2009

Infectious Agents and Cancer 2009, **4**(Suppl 2):P9 doi:10.1186/1750-9378-4-S2-P9

This abstract is available from: <http://www.infectagentscancer.com/content/4/S2/P9>

© 2009 Bagni et al; licensee BioMed Central Ltd.

Circulating B lymphocytes are the major reservoir of KSHV infection in infected subjects. However, B cell lines and primary B cells are resistant to direct KSHV infection *in vitro*. In addition, primary B cells are difficult to propagate for more than a few days. In this study, we combined a novel primary B cell propagation method with efficient infection mediated by dendritic cells to study KSHV *de novo* infection of primary B cells *in vitro*.

Primary monocyte-derived dendritic cells (MDDCs) or plasmacytoid dendritic cells (pDCs) were pulsed with KSHV for 4 hours at which point KSHV DNA was readily detectable. Uptake of KSHV was significantly reduced by pre-incubating cells with antibodies to integrins $\alpha 3$, $\beta 1$ and DC-SIGN. Autologous B cells were grown on a feeder layer of irradiated NIH3T3 cells transduced with a human CD40L retroviral vector. KSHV+ DCs were co-cultivated with primary B cells for 4–8 hours and then separated by CD19+ immunomagnetic isolation. B cell cultures were maintained on feeder cells for >30 days and monitored for KSHV infection.

Efficient KSHV infection of primary B cells was mediated by both MDDCs and pDCs. KSHV LANA protein (ORF73) was detected by IFA in 2–15 percent of B cells through day 14. Viral gene expression analysis using a KSHV whole

genome virus array showed establishment of latent KSHV infection followed by spontaneous reactivation of lytic viral replication in the primary B cell cultures.

These studies suggest that dendritic cells play an important role in the transmission and pathogenesis of KSHV in infected subjects as well as demonstrating a powerful *in vitro* model for studying KSHV infection of B cells.

Acknowledgements

Funded by NCI Contract N01-CO-12400.