

RESEARCH

Open Access

Acetylcholinesterase of the sand fly, *Phlebotomus papatasi* (Scopoli): cDNA sequence, baculovirus expression, and biochemical properties

Kevin B Temeyer^{1*}, Danett K Brake^{1,2}, Alexander P Tuckow¹, Andrew Y Li¹ and Adalberto A Pérez de León¹**Abstract**

Background: Millions of people and domestic animals around the world are affected by leishmaniasis, a disease caused by various species of flagellated protozoans in the genus *Leishmania* that are transmitted by several sand fly species. Insecticides are widely used for sand fly population control to try to reduce or interrupt *Leishmania* transmission. Zoonotic cutaneous leishmaniasis caused by *L. major* is vectored mainly by *Phlebotomus papatasi* (Scopoli) in Asia and Africa. Organophosphates comprise a class of insecticides used for sand fly control, which act through the inhibition of acetylcholinesterase (AChE) in the central nervous system. Point mutations producing an altered, insensitive AChE are a major mechanism of organophosphate resistance in insects and preliminary evidence for organophosphate-insensitive AChE has been reported in sand flies. This report describes the identification of complementary DNA for an AChE in *P. papatasi* and the biochemical characterization of recombinant *P. papatasi* AChE.

Methods: A *P. papatasi* Israeli strain laboratory colony was utilized to prepare total RNA utilized as template for RT-PCR amplification and sequencing of cDNA encoding acetylcholinesterase 1 using gene specific primers and 3'-5'-RACE. The cDNA was cloned into pBlueBac4.5/V5-His TOPO, and expressed by baculovirus in Sf21 insect cells in serum-free medium. Recombinant *P. papatasi* acetylcholinesterase was biochemically characterized using a modified Ellman's assay in microplates.

Results: A 2309 nucleotide sequence of *PpAChE1* cDNA [GenBank: JQ922267] of *P. papatasi* from a laboratory colony susceptible to insecticides is reported with 73-83% nucleotide identity to acetylcholinesterase mRNA sequences of *Culex tritaeniorhynchus* and *Lutzomyia longipalpis*, respectively. The *P. papatasi* cDNA ORF encoded a 710-amino acid protein [GenBank: AFP20868] exhibiting 85% amino acid identity with acetylcholinesterases of *Cx. pipiens*, *Aedes aegypti*, and 92% amino acid identity for *L. longipalpis*. Recombinant *P. papatasi* AChE1 was expressed in the baculovirus system and characterized as an insect acetylcholinesterase with substrate preference for acetylthiocholine and inhibition at high substrate concentration. Enzyme activity was strongly inhibited by eserine, BW284c51, malaoxon, and paraoxon, and was insensitive to the butyrylcholinesterase inhibitors ethopropazine and iso-OMPA.

Conclusions: Results presented here enable the screening and identification of *PpAChE* mutations resulting in the genotype for insensitive *PpAChE*. Use of the recombinant *P. papatasi* AChE1 will facilitate rapid in vitro screening to identify novel *PpAChE* inhibitors, and comparative studies on biochemical kinetics of inhibition.

Keywords: Sand fly, Acetylcholinesterase, *P. papatasi*, cDNA, AChE

GenBank: JQ922267, AFP20868

* Correspondence: Kevin.Temeyer@ars.usda.gov

¹Knipling-Bushland U.S. Livestock Insects Research Laboratory, Agricultural Research Service, U. S. Department of Agriculture, 2700 Fredericksburg Road, Kerville, TX 78028-9184, USA

Full list of author information is available at the end of the article

Background

Over 350 million people are at risk of suffering from leishmaniasis and over 2 million new cases are reported each year, which makes it one of the most neglected diseases globally [1,2]. The sand-fly *Phlebotomus papatasi* (Scopoli) is the vector of the flagellate protozoan *Leishmania major* that causes zoonotic cutaneous leishmaniasis [3-5] in Asia and Africa [1,6,7]. Various species of burrowing rodents are reservoir hosts of *L. major* and sand flies are closely associated with the rodent burrows making vector control by direct insecticidal application difficult to achieve [6,8-10]. Prevention of *Leishmania* spp. transmission to humans is largely reliant on vector control to avoid insect bites by use of screens, insecticide impregnated bed nets or curtains, establishment of barrier zones by insecticide applications, or environmental modification [2,9,11,12]. Use of synthetic sand fly pheromone as an attractant was reported to improve effectiveness of pesticide application for control of the new world sand fly *Lutzomyia longipalpis* in chicken sheds [13]. Attempts to control *Phlebotomus papatasi* by various types of insecticide application has had questionable effectiveness, particularly in the harsh environments reported during combat operations by the U.S. military in Iraq [14,15], although there was no evidence of significant insecticide resistance.

Insecticide resistance is an increasing problem in control of insect vectors of disease and there have been reports of possible resistance in sand fly populations to various insecticides [2,11,16]. Pesticide effectiveness for knockdown of Phlebotomine sand flies in Morocco was found to be variable for different fly populations, reflective of past history of high pesticide use in some areas for malaria control, although all of the sand flies tested as susceptible to the pesticides by bioassay, suggesting possible genetic changes that might lead to development of resistance [17]. Preliminary evidence for an altered acetylcholinesterase (E.C. 3.1.1.7, AChE) was reported in the sand fly *Phlebotomus argentipes* from Sri Lanka [18], and from *Phlebotomus papatasi* in Khartoum State, Sudan [19], each possibly the result of widespread use of insecticides in the Anti-Malarial Campaign. Coutinho-Abreu et al. [20] reported that the AChE gene of *Lutzomyia longipalpis* (Lutz & Neiva) appears to be structurally similar to other insect species in which point mutations result in an altered enzyme insensitive to organophosphate or carbamate inhibitors. Organophosphate pesticides function as quasi-irreversible inhibitors of AChE, resulting in failure of the central nervous system and death of the insect. Point mutations within the sequence encoding AChE resulting in production of an altered, insensitive enzyme have been reported to be a major mechanism of organophosphate resistance in insects [21-26]. Determination of the nucleotide sequence encoding AChE of *P. papatasi* should enable

rapid identification of mutations potentially associated with resistance, and the development of rapid molecular tests to screen *P. papatasi* populations for the presence of specific mutations. Confirmation of AChE mutations resulting in production of an insensitive AChE is expected to facilitate the development of rapid molecular tests for pesticide resistance in sand flies. This report describes the identification of complementary DNA for AChE1 in *P. papatasi* and the biochemical characterization of recombinant *P. papatasi* AChE1 produced using a baculovirus expression system.

Methods

Sand flies

Sand flies used in this study were from a *P. papatasi* colony maintained at the USDA-ARS, Knipling-Bushland U.S. Livestock Insects Research Laboratory in Kerrville, Texas. The colony was established using pupae from an Israeli strain of *P. papatasi* maintained for 30 years and never exposed to pesticides at the Division of Entomology, Walter Reed Army Institute of Research (Silver Spring, Maryland). The colony is therefore considered to be generally susceptible to insecticides and baseline data on pesticide susceptibility by bioassay of colony sand flies is being collected for publication (A. Li, personal communication). Larvae were fed with a sand fly larval diet, a mixture of fermented rabbit feces and rabbit food [27]. Both larvae and adult flies were maintained at $26\pm 2^\circ\text{C}$ and a relative humidity of $85\pm 2\%$. Males were fed with 30% sucrose water and females were fed with defibrinated cattle blood using an in vitro membrane feeding system. Larvae and newly emerged unfed adult male flies were collected and quick-frozen by immersion in liquid nitrogen, and stored at -70°C until use.

RNA preparation

Total RNA was prepared from frozen whole larvae or unfed adults by grinding in a glass tissue homogenizer containing Tri-Reagent (Sigma Chemical, St. Louis, MO) or by use of a MagMAXTM-96 RNA Isolation Kit (Life Technologies, Carlsbad, CA) according to the manufacturers' instructions.

Oligonucleotide synthesis

Synthetic oligodeoxynucleotides were purchased from Sigma-Genosys (The Woodlands, TX).

cDNA synthesis

Complementary DNA (cDNA) was synthesized by reverse transcription from template RNA using the SuperScriptTM Choice System for cDNA Synthesis (Life Technologies) according to the manufacturer's instructions. Oligodeoxynucleotide primers were oligo(dT)₁₈V or gene-specific primers based on known sequence.

Polymerase chain reaction (PCR)

We utilized automatic hot-start PCR using AmpliTaq Gold[®] DNA polymerase (Applied Biosystems, Foster City, CA) or Advantage[®] 2 HF DNA Polymerase (BD Biosciences Clontech, Mountain View, CA) according to the manufacturers' instructions.

Agarose gel electrophoresis

DNA preparations generated from PCR, RT-PCR, or bacterial plasmids were analyzed by agarose gel electrophoresis in tris-borate-EDTA buffer by standard techniques [28]. DNA bands were visualized by staining with GelStar[®] Nucleic Acid Gel Stain (Cambrex, Rockland, MD) and photographed on a Kodak GelLogic 440 Imaging system (Eastman Kodak Co., Rochester, NY) with uv illumination.

Rapid Amplification of cDNA Ends (RACE)

5'- and 3'-RACE was performed using the SMARTer[™] RACE cDNA Amplification Kit (BD Biosciences Clontech, Palo Alto, CA).

DNA cloning

Double-stranded cDNA was inserted into bacterial plasmid DNA using the TOPO TA Cloning[®] Kit for Sequencing (Invitrogen, Carlsbad, CA), transformed into One Shot[®] TOP10 chemically competent *Escherichia coli* (Invitrogen) for amplification, screened for appropriate inserts by PCR, and sequenced by normal sequencing procedures (see below).

DNA sequencing

For sequencing, PCR amplification reactions were scaled up to 50 µl and purified using ExoSAP-IT[®] (Affymetrix, Santa Clara, CA). The sequencing reaction using BigDye[®] terminator (Applied Biosystems) and precipitation of the products with ethanol were performed according to the manufacturer's instructions. Sequencing products were analyzed on a PRISM 3130xl Genetic Analyzer (Applied Biosystems).

Analysis of amino acid or nucleotide sequences

Sequence analysis to design oligonucleotide primers used for PCR or sequencing utilized OLIGO[®] Primer Analysis Software, Version 5.0 (1996, National Biosciences Inc, Plymouth, MN). Sequence chromatograms and sequence assembly utilized ChromasPro ver. 1.32 (Technelysium Pty Ltd., available at http://technelysium.com.au/?page_id=27). Clustal W (<http://www.ebi.ac.uk/Tools/msa/clustalw2/>) was used to perform multiple sequence alignments [29]. BLAST search of GenBank (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>), [30] and on-line protein sequence analyses were done using ExPASy Proteomics tools.

PpAChE expression construct

Double stranded cDNA containing the complete PpAChE1 coding sequence [GenBank: JQ922267] was obtained by PCR amplification of cDNA using PpAce18F-273U23 and PpAce18F-2561 L20 (Table 1) using an annealing temperature of 62.5°C and extension time of 3 min. at 72°C. The cDNA was cloned into pBlueBac4.5/V5-His TOPO[®] (Applied Biosystems/Life Technologies) and sequenced to verify proper construction.

Baculovirus expression of recombinant PpAChE

PpAChE1 clones were constructed, sequenced, and expressed as baculovirus expression clones in Sf21 insect cells in sf900 III serum-free medium (Life Technologies/Gibco) as previously described [31,32]. Serum-free cell culture baculovirus lysates were collected and centrifuged to remove cell debris and supernatants were used for biochemical characterization of recombinant PpAChE1.

Biochemical characterization of rPpAChE1

AChE activity of baculovirus culture supernatant was measured by a modified Ellman's method as described previously [33] except that the reaction was monitored every minute for 30 min to establish initial velocity. For inhibition studies, rPpAChE was preincubated with an appropriate concentration of the inhibitor at 23°C for 10 min in 100 µl in the microtiter plate without substrate or DTNB [(5',5'-dithiobis-(2-nitrobenzoic acid))] and AChE activity was monitored for 10 min after initiation of the reaction by addition of 100 µl substrate and DTNB. Cholinesterase inhibitors included the AChE-specific inhibitors eserine and BW284c51, butyrylcholinesterase-specific inhibitors ethopropazine and iso-OMPA, and oxidized organophosphates, paraoxon and malaoxon. All chemicals were of reagent grade obtained from Sigma Chemical Co. Data were analyzed and plotted using GraphPad Prism ver. 5.0 (GraphPad Prism, Inc., La Jolla, CA).

Results

Beginning with a presumptive acetylcholinesterase partial sequence (111 nucleotides) of *Phlebotomus perniciosus* [GenBank: AJ865843], we constructed oligodeoxynucleotide primers for 5'-, and 3'-RACE to obtain the complete *PpAChE* cDNA sequence [GenBank: JQ922267] (Additional file 1). Both strands were sequenced for the entire presumptive coding sequence of the cDNA at least three times. Oligodeoxynucleotide primers used for sequencing or 3'-/5'-RACE are listed in Table 1. The sequence obtained for the presumptive *PpAChE1* cDNA of *P. papatasi* was comprised of 2309 nucleotides organized as a 71-nucleotide 5'-untranslated region, an open reading frame (ORF) of 2130 nucleotides, and a 105-nucleotide 3'-untranslated region. BLAST homology search

Table 1 Oligodeoxynucleotide primers for AChE of *P. papatasi*

Oligonucleotide name	Position ^a	Nucleotide Sequence ^b
PpAce18F-273U23	1-23 (U)	CAATAACGTGGTATCTCGCATAA
PpAChE458U17	186-202 (U)	TTGGCGGAGGGTCGTCA
PpAChE552U22	280-302 (U)	TCTTAGGCGAATCAACATTAGA
PpAChE550L23	300-278 (L)	CTAATGTTGATTCCGCTAAGACT
PpAChE5R-693 L20	497-378 (L)	CGCTCCAGTGTGCCCAATTC
PpAChE20-828U23	827-849 (U)	CTTCGGTGGTGGATTCTACTCAG
PpAChEp13-435 L21	975-955 (L)	CAGGGGCATCAGGAGTACCAA
PpAChEp13-719U18	1221-1239 (U)	CTAGCCGAAGCCGTGGAG
PpAChE7Jan-590 L22	1226-1205 (L)	GGCTAGGCGAAGGGTCTATTG
PpAChE7J-887U23	1502-1524 (U)	AGAGGAGGGCATAACTGTAAACAC
PpAChE7J-947 L22	1582-1562 (L)	CGCACGGCACCATTGACATAG
PpAChE7J-1100U22	1715-1736 (U)	TGAGGAGGGCAACAATGTCTAC
PpAce2L27	1749-1723 (L)	TGTAGAGATACATGTAGACATTGTTGC
PpAce22L18	1760-1743 (L)	GGTGCATGGGTGTAGAG
PpAce18F-2561 L20	2309-2289 (L)	GAGTAAATCGCGTACTTCA

^aPosition (based on PpAChE sequence numbering, Figure 1).

^bSequence is 5'→3', upper (U) or lower (L) strand.

[30] of GenBank (11,299,630 sequences) found significant nucleotide identity (73-83%) to the AChE mRNA sequences of a mosquito, *Culex tritaeniorhynchus* [GenBank: AB122152] and for the new world sand fly *L. longipalpis* (partial codes, 1188 nucleotides, [GenBank: DQ898276]) respectively, as well as to other arthropod AChEs.

The *P. papatasi* AChE cDNA ORF encodes a 710-amino acid protein [GenBank: AFP20868]; 79451 MW, 5.77 pI, http://web.expasy.org/compute_pi/) exhibiting 85% amino acid identity with AChEs of mosquitoes, *Culex pipiens* [GenBank: Q86GC8], *Aedes aegypti* [GenBank: XP_001656977], and 92% amino acid identity with the new world sand fly, *L. longipalpis* AChE (partial codes, 396 amino acids, [GenBank: ABI74669]). Comparison of the amino acid sequence of PpAChE1 with the AChE protein of *L. longipalpis* (partial codes) and the AChEs of *Cx. pipiens* and *Ae. aegypti* by Clustal W multiple sequence alignment is shown in Figure 1. As seen in Figure 1, there was very high amino acid identity throughout the partial sequence of *L. longipalpis* AChE as well as substantial identity with the two mosquito AChEs. Presumptive disulfide bonds, members of the catalytic triad, and residues lining the catalytic gorge [34,35] are indicated in the Clustal W multiple sequence alignment with AChEs of *Torpedo californica* [GenBank: 1EA5_A] and *Drosophila melanogaster* [GenBank: 1QO9_A] (Additional file 2).

Recombinant PpAChE1 was expressed in the baculovirus system and used for biochemical characterization. Biochemical properties (substrate K_m and relative sensitivities to inhibitors) obtained for rPpAChE1 are presented in Table 2. The enzyme was strongly inhibited by

eserine, paraoxon, malaoxon, and the acetylcholinesterase inhibitor Bw284c51 and exhibited much lower sensitivity to the butyrylcholinesterase inhibitors iso-OMPA and ethopropazine. In addition, rPpAChE1 hydrolyzed 0.1 mM acetylthiocholine (AcSCh) 17-fold faster than 0.1 mM butyrylthiocholine (BuSCh), demonstrating substrate preference for AcSCh over BuSCh and was inhibited at substrate concentrations above 5 mM.

Discussion

Biochemical properties of rPpAChE1 presented in this report are consistent with classification as an insect acetylcholinesterase as defined by Toutant [36]. There is substrate preference for acetylthiocholine over butyrylthiocholine, significant inhibition by the AChE-specific inhibitors eserine and BW284c51 with relative insensitivity to inhibitors (ethopropazine and iso-OMPA) classified as specific for butyrylcholinesterase. The enzyme is strongly inhibited by paraoxon and malaoxon (oxidized forms of parathion and malathion) and exhibits a K_m for acetylthiocholine (37.9 μ M) similar to values reported for recombinant acetylcholinesterases of *Ae. aegypti* (13.79 μ M) [37], *Stomoxys calcitrans* (63.9 μ M) [33], *Haematobia irritans* (31.3 μ M) [38], or *Musca domestica* (150 μ M) [39].

P. papatasi is a significant public health problem in the Middle East, Asia and Africa where it transmits *L. major* that causes zoonotic cutaneous leishmaniasis in humans [40]. Additionally, zoonotic cutaneous leishmaniasis can seriously affect the operational readiness of United States military personnel deployed in Iraq and Afghanistan [41]. Attempts by the U.S. military to control sand flies through the use of chemical pesticides were not highly successful,

<i>P. papatasi</i>	MEIRGIVVTTMRPFTGIHSGVDQMFVICLLSLLGVMCQLAEGRRHDLSENTQSFKSGPKHI	60
<i>L. longipalpis</i>	-----	
<i>C. pipiens</i>	MEIRGLITRLLGPCHLRH-----LILCSLGLYSILVQSVHCRHHDIGSSVAHQLGSKYS	54
<i>A. aegypti</i>	MEIRGLITRLLGPCHIRH-----LILCSLGLYSILVQSVHCRHHDIGSSSTAHLQLGSKYS	54
<i>P. papatasi</i>	ASVEAAAVSVLGESTLEVSSSEDDT--IFTPLYGHGDAVRVDAELGLTLEREGVSAGSDG	118
<i>L. longipalpis</i>	-----	
<i>C. pipiens</i>	QSSSLSSSSQSSSLAEEATLNKDSDAFFTPYIGHGDSVRIVDAELGLTLEREHIHS----	110
<i>A. aegypti</i>	QSSSLSSSSQSSSLVEDPVLNKDSDAFFTPYIGHGDSVRIVDAELGLTLEREHVHS----	110
<i>P. papatasi</i>	TSQPRRRNISRRSNPDAEENDPLIVTTDKGKVRGVTLTSPGKKVDAWMGIPYAQPPLG	178
<i>L. longipalpis</i>	-----	
<i>C. pipiens</i>	-TTTRRRGLTRRESSSDATDSPLVITTDKGIKRGTTLEAPSGKKVDAWMGIPYAQPPLG	169
<i>A. aegypti</i>	-TTTRRRGLTRRESSSDGTDNDPLIITTDKGIKRGTTLEAPSGKKVDAWLGIPYAQPPLG	169
<i>P. papatasi</i>	ALRFRHRPRAERWSGILNATTPPNTCVQIVDTLFGDFPGATMWNPNNTLTEDCLYINVAV	238
<i>L. longipalpis</i>	-----ATMWNPNNTLTEDCLYINVAV	21
<i>C. pipiens</i>	PLRFRHRPRAERWTGVLNATKPPNSCVQIVDTVFGDFPGATMWNPNNTPLSEDCLYINVVV	229
<i>A. aegypti</i>	PLRFRHRPRAERWTGVLNATTPPNSCVQIVDTVFGDFPGATMWNPNNTPLSEDCLYINVVV	229
	1	1
	***** * :***** *	
<i>P. papatasi</i>	PHPRPKNSPVMWLWIFGGGFYSGTSTLDVYDHRTLVAEENIILVSMQYRVASLGLFYLGLTP	298
<i>L. longipalpis</i>	PHPRPKNSPVMWLWIFGGGFYSGTSTLDVYDHRTLVAEENIILVSMQYRVASLGLFYLGLTP	81
<i>C. pipiens</i>	PRPRPKNAAVMLWIFGGGFYSGTATLDVYDHRTLASEENIIVVSLQYRVASLGLFYLGLTP	289
<i>A. aegypti</i>	PHPRPKNSAVMLWIFGGGFYSGTATLDVYDHRTLASEENIIVVSLQYRVASLGLFYLGLTP	289
	†	
<i>P. papatasi</i>	DAPGNAGLFDQHLALRWVRNNIHRFGGDPTRVTLFGESAGAVSVSMHLLSSLSHDLFQRA	358
<i>L. longipalpis</i>	DAPGNAGLFDQNLALRWVRDNIHRFGGDPTRVTLFGESAGAVSVSMHLLSALSLSHDLFQRA	141
<i>C. pipiens</i>	EAPGNAGLFDQNLALRWVRDNIHRFGGDPTRVTLFGESAGAVSVSLHLLSALSLSHDLFQRA	349
<i>A. aegypti</i>	EAPGNAGLFDQNLALRWVRDNIHKFGGDPTRVTLFGESAGAVSVSLHLLSALSLSHDLFQRA	349
	2	2
	***** :*:* * :***** :*:* * :***** :*:* * :*****	
<i>P. papatasi</i>	ILQSGSPTAPWALITRDEAINRTLRLAEAVECPHNRDELSEVLECLRSRDAKQLVNNENW	418
<i>L. longipalpis</i>	ILQSGSPTAPWALMTREEAINRALRLADAVECPHNRDNLPEVVECLRGRDAKQLVNNENW	201
<i>C. pipiens</i>	ILQSGSPTAPWALVSRREATLRLRLAEAVNCPHDATKLSDAVECLRTKDPNVLVDNEWG	409
<i>A. aegypti</i>	ILQSGSPTAPWALVSRREATLRLRLAEAVNCPHDATKLTDTVECLRTKDPNVLVDNEWG	409
	†	
<i>P. papatasi</i>	NLIGICEFPFVVPVVDGSLDESQRAMATGRFEKTDILTGSNTEEGYYPFIYYLTELRLKE	478
<i>L. longipalpis</i>	NLIGICEFPFVVPVVDGAFLDDETPQRSLSASGRFVKTDILTGSNTEEGYYPFIYYLTELRLKE	261
<i>C. pipiens</i>	TLGICEFPFVVPVVDGAFLDDETPQRSLSASGRFVKTDILTGSNTEEGYYPFIYYLTELRLKE	469
<i>A. aegypti</i>	TLGICEFPFVVPVVDGAFLDDETPQRSLSASGRFVKTDILTGSNTEEGYYPFIYYLTELRLKE	469
	3	
	***** :*:* * :***** :*:* * :***** :*:* * :*****	
<i>P. papatasi</i>	EGITVTREEFKAVRELNPYVNGAARQAI VFEYTDWTDPDNAHSNRDALDKMVG DYHFTC	538
<i>L. longipalpis</i>	EGITVTREEFKAVRELDPYVNGAARQAI VFEYTDWTDPDNAHSNRDALDKMVG DYHFTC	321
<i>C. pipiens</i>	EGVTVTREEFKAVRELNPYVNGAARQAI VFEYTDWTDPEPNPNSNRDALDKMVG DYHFTC	529
<i>A. aegypti</i>	EGVTVSREEFKAVRELNPYVNGAARQAI VFEYTDWTEPENPNSNRDALDKMVG DYHFTC	529
	†	
<i>P. papatasi</i>	NVNEFAHRYAEEGNNVYMYLYTHRSTKANPWRPRTGVMHGDEIN YVFGEP LNSLYTDEE	598
<i>L. longipalpis</i>	NVNEFAHRYAEEGNNVYMYLYTHRSKGNPWRPRTGVMHGDEIN YVFGEP LNSALSYTEEE	381
<i>C. pipiens</i>	NVNEFAQR YAEEGNNVYMYLYTHRSKGNPWRPRTGVMHGDEIN YVFGEP LNSALGYQDDE	589
<i>A. aegypti</i>	NVNEFAQR YAEEGNNVYMYLYTHRSKGNPWRPRTGVMHGDEIN YVFGEP LNSDLGYMEDE	589
	*	* :*
<i>P. papatasi</i>	KEFSRRIMRYWVNFAKTGNPNPGFVS-NLPDWPKHTAHGRQYME LGLN TTYLGRGPRLRQ	657
<i>L. longipalpis</i>	KDFSRRIMRYWSKFA-----	396
<i>C. pipiens</i>	KDFSRRIMRYWSNFAKTGNPNPSTPSVDLPEWPKHTAHGRHYLELGLN TTYVGRGPRLRQ	649
<i>A. aegypti</i>	KDFSRRIMRYWSNFAKTGNPNPSPNSDFPEWPKHTAHGRHYLELGLN TTYVGRGPRLRQ	649
	3	
	***** :*:* * :***** :*:* * :***** :*:* * :*****	
<i>P. papatasi</i>	CAFWKYLPQLMAATIENSSTKNCTNVGNQFVRNPNFSIPTLLVILGILSVN----	710
<i>L. longipalpis</i>	-----	
<i>C. pipiens</i>	CAFWKYLPQLVAAT----SNLQVTPAPSVPCSSSTSYRSTLLLI VTL LLLVTRFKI	702
<i>A. aegypti</i>	CAFWKYLPQLVAAT----SNLQATPAPSEPCSSSATPYRPLLFIVSLVIVTFCFKI	702

Figure 1 Clustal W 2.1 multiple sequence alignment of AChE protein sequences for *Phlebotomus papatasi* [GenBank: JQ922267], *Lutzomyia longipalpis* [GenBank: ABI74669], *Culex pipiens* AChE [GenBank: Q86GC8], and *Aedes aegypti* [GenBank: XP_001656977]. The consensus line below the aligned sequences indicates positions of conserved amino acid identity (*) or similarity (: or .). Positions of the 3 disulfide bond linkages are indicated by numbers above participating cysteine pairs. The members of the catalytic triad (S, E, H) which make up the catalytic site are indicated by (†) above the participating amino acid.

Table 2 Biochemical properties of rPpAChE

Property ^a	PpAChE ^b
K_m AcSCh (μ M)	37.9 \pm 1.6
IC ₅₀ Paraoxon (10^{-7} M)	2.3 (1.9-2.7)
IC ₅₀ Malaoxon (10^{-8} M)	2.7 (1.7-4.5)
IC ₅₀ Eserine (10^{-10} M)	7.2 (5.0-10.5)
IC ₅₀ BW284c51 (10^{-8} M)	7.1 (5.9-8.4)
IC ₅₀ Ethopropazine (10^{-6} M)	4.4 (4.0-4.9)
IC ₅₀ Iso-OMPA	NC ^c

^aAcSCh = acetylthiocholine; IC₅₀ = concentration of inhibitor producing 50% inhibition of enzyme activity during 10 min preincubation.

^bData are presented as mean \pm SE or (95% confidence interval).

^cNC = not calculated (no inhibition detected at 10^{-3} M).

possibly due to extreme environmental conditions [14,15], and pesticide bioassays did not reveal evidence of insecticide resistance. Chemical pesticides have been used in several ways for insect control, including area-wide applications for malaria control in the same areas inhabited by *P. papatasi*. Recent reports of pesticide resistance in sand flies, possibly resulting from insensitive AChE [18,19], highlight the urgent need to characterize the mechanisms responsible for insecticide resistance and the relevance of efforts to develop rapid tests to identify resistant pest populations.

Point mutations in pest AChEs have been shown to generate altered enzymes with decreased inhibition by organophosphate [16,21,42-45]. The present report demonstrating very high amino acid sequence identity of the *P. papatasi* AChE to those of *Cx. pipiens* and *Ae. aegypti* strongly suggests that this PpAChE1 is the target site for organophosphate and will facilitate identification of PpAChE1 mutations that produce insensitivity to organophosphate or carbamate pesticides.

In mosquito vectors, very high resistance to OPs and carbamates results from a single amino acid substitution of Serine for Glycine (codon GGC, PpAChE1 pos 837, Gly→AGC, Ser) [46-50]. *Ae. aegypti* is reported to have the GGA codon at this position (as does our PpAChE1), and when converted to AGC, the *A. aegypti* enzyme has the identical high level resistance as *Anopheles* [49]. It has been suggested that GGA at this position is extremely unlikely to evolve to AGC, and that in nature, only Gly or Ser are found at this position, which indicates that no other amino acids are allowed in AChE1 at that position [49]. Our results suggest significant OP-insensitivity in AChE of *P. papatasi* would be unlikely to involve PpAChE1, unless nucleotide pos 839 exists in a polymorphic state (i.e., GGA/C) in nature. Alternatively, it is hypothesized that, if present, other AChEs in *P. papatasi* could provide the mechanism to develop target insensitivity as documented in *Ae. aegypti* where a high level of AChE insensitivity was due to substitutions at positions F455W (*Tc* pos Phe331) and Ile697 of AChE2 [51,52].

Conclusions

Use of the recombinant *P. papatasi* AChE1 will facilitate rapid in vitro screening to identify novel PpAChE inhibitors, and comparative studies on biochemical kinetics of inhibition. Information on the identification and characterization of PpAChE1 presented here will facilitate the development of a rapid molecular assay for the GGC codon (pos. 839) in *P. papatasi* populations, and other molecular tools to screen for mutations giving rise to an organophosphate-insensitive PpAChE1. Molecular data on PpAChE1 could also be used in modeling studies to predict in vivo insecticidal activity for novel inhibitors as described by Naik et al. [53]. Availability of the recombinant PpAChE1 will enable the creation of mechanism-based screens to discover more effective inhibitors that may be developed to innovate safer vector control technologies. Novel synthetic carbamates have already been identified by screening using the recombinant PpAChE that present essentially equivalent inhibition of the target AChE with substantially improved target specificity resulting in significantly enhanced safety [Swale DR, Tong F, Temeyer KB, Li AY, Totrov MM, Carlier PR, unpublished].

Endnotes

¹This article reports the results of research only. Mention of a proprietary product does not constitute an endorsement by the USDA for its use.

²In conducting research described in this report, investigators adhered to the "Guide for the Care and Use of Laboratory Animals," as promulgated by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animal Resources, National Research Council. A protocol describing routing procedures for animal care and use is on file with the Animal Care Committee at the research location.

³USDA is an equal opportunity provider and employer.

⁴Copyright statement: Copyright protection is not available for any work of the United States Government.

⁵Disclaimer: "The views expressed in this article are those of the authors and do not necessarily reflect the official policy or position of the Department of the Navy, Department of Defense, nor the U.S Government."

Additional files

Additional file 1: Figure S1. Nucleotide and amino acid sequences are listed for *Phlebotomus papatasi* AChE for rPpAChE-16 expressed in the baculovirus system.

Additional file 2: Figure S2. Clustal W 2.1 multiple sequence alignment of AChE protein sequences for *Torpedo californica* [GenBank: 1EA5_A], *Phlebotomus papatasi* (rPpAChE-16), and *Drosophila melanogaster* AChE [GenBank: 1QO9_A]. The consensus line below the aligned sequences indicates positions of conserved amino acid identity (*) or similarity (: or). Positions of the 3 disulfide bond linkages are indicated by numbers above

participating cysteine pairs. The members of the catalytic triad (S, E, H) which make up the catalytic site are indicated by (#) above the participating amino acid. Positions lining the catalytic gorge are indicated by (▼) above the participating amino acid [32,33].

Abbreviations

(AChE): Acetylcholinesterase; (DTNB): Dithiobisnitrobenzoic acid; (AcSch): Acetylthiocholine; (BuSch): Butyrylthiocholine; (rPpAChE1): Recombinant *P. papatasi* AChE1.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

KBT, AYL, and AAPdL conceived experiments to clone the PpAChE and biochemically characterize its recombinant form. KBT designed experiments and supervised laboratory tests to identify, clone, sequence, and express rPpAChE in the baculovirus system. DKB and APT conducted biochemical assays and data analysis of rPpAChE1. AYL and AAPdL obtained DWFP funding, maintained the sand fly colony, and provided sand flies for experimental use. All authors contributed to writing and revision of the manuscript and approved the final version.

Acknowledgements

The authors thank Kristie Schlechte, DeEsta Hyatt and Darci Burchers for excellent technical assistance. The authors also thank Dr. Fan Tong and Dr. Donald Thomas for critical review of the manuscript. This work was funded by U.S. Department of Agriculture [USDA] appropriated funds (USDA-ARS CRIS number 6205-32000-033000D, Molecular biology and physiology of biting flies affecting livestock) and USDA-ARS CRADA number 60-0208-9-601 (Project number 6201-32000-033017R, Deployed Warfighter Protection Research Program [DWFP] of the U.S. Department of Defense through the Armed Forces Pest Management Board [AFPMB]).

Author details

¹Knipling-Bushland U.S. Livestock Insects Research Laboratory, Agricultural Research Service, U. S. Department of Agriculture, 2700 Fredericksburg Road, Kerrville, TX 78028-9184, USA. ²Present address: Medical Service Corps, U.S. Navy, Wound Infections Dept., Naval Medical Research Center, 503 Robert Grant Ave, Silver Spring, MD 20910-7500, USA.

Received: 27 September 2012 Accepted: 10 January 2013

Published: 4 February 2013

References

- World Health Organization: *Cutaneous leishmaniasis: Why are you neglecting me?*. 2007 [http://whqlibdoc.who.int/hq/2007/WHO_CDS_NTD_IDM_2007.3_eng.pdf]
- World Health Organization: *Control of the leishmaniases*. WHO Technical Report Series; no. 949: Report of a meeting of the WHO Expert Committee on the Control of Leishmaniases, Geneva, 22–26 March 2010; 2010 [http://whqlibdoc.who.int/trs/WHO_TRS_949_eng.pdf]
- Jaffe CL, Baneth G, Abdeen ZA, Schlein Y, Warburg A: **Leishmaniasis in Israel and the Palestinian Authority**. *Trends Parasitol* 2004, **20**:328–332.
- Kravchenko V, Wasserberg G, Warburg A: **Bionomics of phlebotomine sandflies in the Galilee focus of cutaneous leishmaniasis in northern Israel**. *Med Vet Entomol* 2004, **18**:418–428.
- Killick-Kendrick R, Leaney AJ, Peters W, Rioux J-A, Bray RS: **Zoonotic cutaneous leishmaniasis in Saudi Arabia: the incrimination of *Phlebotomus papatasi* as the vector in the Al-Hassa oasis**. *Trans R Soc Trop Med Hyg* 1985, **79**:252–255.
- Mascari TM, Mitchell MA, Rowton ED, Foil LD: **Evaluation of juvenile hormone analogues as rodent feed-through insecticides for control of immature phlebotomine sandflies**. *Med Vet Entomol* 2011, **25**:227–231.
- Mirzaei A, Rouhani S, Taherkhani H, Farahmand M, Kazemi B, Hedayati M, Baghaei A, Davari B, Parvizi P: **Isolation and detection of *Leishmania* species among naturally infected *Rhombomis opimus*, a reservoir host of zoonotic cutaneous leishmaniasis in Turkeman Sahara, North East of Iran**. *Exp Parasitol* 2011, **129**:375–380.
- Rassi Y, Oshaghi MA, Azani SM, Abaie MR, Rafizadeh S, Mohebbi M, Mohtarami F, Zeinali M: **Molecular detection of *Leishmania* infection due to *Leishmania major* and *Leishmania turanica* in the vectors and reservoir host in Iran**. *Vector Borne Zoonotic Dis* 2011, **11**:145–150.
- Schlein Y, Müller GC: **Experimental control of *Phlebotomus papatasi* by spraying attractive toxic sugar bait (ATSB) on vegetation**. *Trans R Soc Trop Med Hyg* 2010, **104**:766–771.
- Wasserberg G, Poche R, Miller D, Chenault M, Zollner G, Rowton ED: **Imidacloprid as a potential agent for the systemic control of sand flies**. *J Vector Ecol* 2011, **36**(Suppl 1):S148–S156.
- Alexander B, Maroli M: **Control of phlebotomine sandflies**. *Med Vet Entomol* 2003, **17**:1–18.
- Kassi M, Kasi PM, Marri SM, Tareen I, Khawar T: **Vector control in cutaneous leishmaniasis of the old world: a review of literature**. *Dermatol Online J* 2008, **14**:1.
- Bray DP, Alves GB, Dorval ME, Brazil RP, Hamilton JGC: **Synthetic sex pheromone attracts the leishmaniasis vector *Lutzomyia longipalpis* to experimental chicken sheds treated with insecticide**. *Parasit Vectors* 2010, **3**:16.
- Colacicco-Mayhugh MG, Grieco JP, Putnam JL, Burkett DA, Coleman RE: **Impact of phlebotomine sand flies on United States Military Operations at Tallil Air Base, Iraq: 5. Impact of weather on sand fly activity**. *J Med Entomol* 2011, **48**:538–545.
- Coleman RE, Burkett DA, Sherwood V, Caci J, Dennett JA, Jennings BT, Cushing R, Cushing RJ, Ploch J, Hopkins G, Putnam JL: **Impact of phlebotomine sand flies on United State military operations at Tallil Air Base, Iraq: 6. Evaluation of insecticides for the control of sand flies**. *J Med Entomol* 2011, **48**:584–599.
- Hemingway J, Ranson H: **Insecticide resistance in insect vectors of human disease**. *Annu Rev Entomol* 2000, **45**:371–391.
- Faraj C, Ouahabi S, Adlaoui EB, Elkohli ME, Lakraa L, Rhazi ME, Ameur B: **Insecticide susceptibility status of *Phlebotomus (Paraphlebotomus) sergenti* and *Phlebotomus (Phlebotomus) papatasi* in endemic foci of cutaneous leishmaniasis in Morocco**. *Parasit Vectors* 2012, **5**:51.
- Surendran SN, Karunaratne SHPP, Adams J, Hemingway J, Hawkes NJ: **Molecular and biochemical characterization of a sand fly population from Sri Lanka: evidence for insecticide resistance due to altered esterases and insensitive acetylcholinesterase**. *Bull Entomol Res* 2005, **95**:371–380.
- Hassan MM, Widaa SO, Osman OM, Numiary MSM, Ibrahim MA, Abushama HM: **Insecticide resistance in the sand fly, *Phlebotomus papatasi* from Khartoum State, Sudan**. *Parasit Vectors* 2012, **5**:46.
- Coutinho-Abreu IV, Balbino VQ, Valenzuela JG, Sonoda IV, Ramalho-Ortigão JM: **Structural characterization of acetylcholinesterase 1 from the sand fly *Lutzomyia longipalpis* (Diptera: Psychodidae)**. *J Med Entomol* 2007, **44**:639–650.
- Fournier D, Muter A: **Modification of acetylcholinesterase as a mechanism of resistance to insecticides**. *Comp Biochem Physiol C Toxicol Pharmacol* 1994, **108**:19–31.
- Muter A, Pralavorio M, Bride JM, Fournier D: **Resistance-associated point mutations in insecticide insensitive acetylcholinesterase**. *Proc Natl Acad Sci USA* 1994, **91**:5922–5926.
- Zhu KY, Clark JM: **Comparisons of kinetic properties of acetylcholinesterase from azinphosmethyl-susceptible and resistant strains of Colorado potato beetle**. *Pest Biochem Physiol* 1995, **51**:57–67.
- Zhu KY, Lee SH, Clark JM: **A point mutation of acetylcholinesterase associated with azinphosmethyl resistance and reduced fitness in Colorado potato beetle**. *Pest Biochem Physiol* 1996, **55**:100–108.
- Villatte F, Ziliani P, Marcel V, Nenozi P, Fournier D: **A high number of mutations may provide insecticide-resistance in insect acetylcholinesterase**. *Pest Biochem Physiol* 2000, **67**:95–102.
- Menozi P, Shi MA, Lougarre A, Tang AH, Fournier D: **Mutations of acetylcholinesterase which confer insecticide resistance in *Drosophila melanogaster* populations**. *BMC Evol Biol* 2004, **4**:4.
- Young DG, Perkins PV, Endris RG: **A larval diet for rearing Phlebotomine sand flies (Diptera, Psychodidae)**. *J Med Entomol* 1981, **18**:446.
- Ausubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA, Struhl K: *Current Protocols in Molecular Biology*. Hoboken: John Wiley & Sons: Ausubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA, Struhl K; 2003.
- Thompson JD, Higgins DG, Gibson TJ: **CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through**

- sequence weighting, position-specific sequence gap penalties and weight matrix choice. *Nucleic Acids Res* 1994, **22**:4673–4680.
30. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ: Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 1997, **25**:3389–3402.
 31. Temeyer KB, Pruett JH Jr, Untalan PM, Chen AC: Baculovirus expression of *BmAChE3*, a cDNA encoding an acetylcholinesterase of *Boophilus microplus* (Acari: Ixodidae). *J Med Entomol* 2006, **43**:707–712.
 32. Temeyer KB, Pruett JH Jr, Olafson PU, Chen AC: *R86Q*, a mutation in *BmAChE3* yielding a *Rhipicephalus microplus* organophosphate-insensitive acetylcholinesterase. *J Med Entomol* 2007, **44**:1013–1018.
 33. Temeyer KB, Chen AC: Acetylcholinesterase of *Stomoxys calcitrans* (L.) (Diptera: Muscidae): cDNA sequence, baculovirus expression, and biochemical properties. *Vet Parasitol* 2012, **184**:92–95.
 34. Harel M, Kryger C, Tosenberry TL, Mallender WD, Lewis T, Fletcher RJ, Guss JM, Silman I, Sussman JL: Three-dimensional structures of *Drosophila melanogaster* acetylcholinesterase and of its complexes with two potent inhibitors. *Protein Sci* 2000, **9**:1063–1072.
 35. Sussman JL, Harel M, Frolov F, Oefner C, Goldman A, Toker L, Silman I: Atomic structure of acetylcholinesterase from *Torpedo californica*: a prototypic acetylcholine-binding protein. *Science* 1991, **253**:872–879.
 36. Toutant J-P: Insect acetylcholinesterase: catalytic properties, tissue distribution and molecular forms. *Prog Neurobiol* 1989, **32**:423–446.
 37. Anthony N, Rocheleau T, Mocelin G, Lee H-J, French-Constant R: Cloning, sequencing and functional expression of an acetylcholinesterase gene from the yellow fever mosquito *Aedes aegypti*. *FEBS Lett* 1995, **368**:461–465.
 38. Temeyer KB, Brake DK, Schlechte KG: Acetylcholinesterase of *Haematobia irritans* (Diptera: Muscidae): baculovirus expression, biochemical properties, and organophosphate insensitivity of the G262A mutant. *J Med Entomol* 2012, **49**:589–594.
 39. Walsh SB, Dolden TA, Moores GD, Kristensen M, Lewis T, Devonshire AL, Williamson MS: Identification and characterization of mutations in housefly (*Musca domestica*) acetylcholinesterase involved in insecticide resistance. *Biochem J* 2001, **359**:175–181.
 40. Jacobson RL: Leishmaniasis in an era of conflict in the Middle East. *Vector Borne Zoonotic Dis* 2011, **11**:247–258.
 41. Lay JC: Leishmaniasis among U.S. armed forces, January 2003–November 2004. Medical surveillance monthly report. *U S Army Center Health Promotion Preventive Med* 2004, **10**:2–5.
 42. Fournier D, Muterio A, Pralavorio M, Bride J-M: *Drosophila* acetylcholinesterase: mechanisms of resistance to organophosphates. *Chem Biol Interact* 1993, **87**:233–238.
 43. Feyereisen R: Molecular biology of insecticide resistance. *Toxicol Lett* 1995, **82/83**:83–90.
 44. Kozaki T, Tomita T, Taniae K, Yamakawa M, Kono Y: Expression of two acetylcholinesterase genes from organophosphate sensitive- and insensitive-houseflies, *Musca domestica* L. (Diptera: Muscidae), using a baculovirus insect cell system. *Appl Entomol Zool* 2002, **37**:213–218.
 45. Vontas JG, Hejazi MJ, Hawkes NJ, Cosmidis N, Loukas M, Hemingway J: Resistance-associated point mutations of organophosphate insensitive acetylcholinesterase in the olive fruit fly *Bactrocera oleae*. *Insect Mol Biol* 2002, **11**:329–336.
 46. Weill W, Fort P, Berthomieu A, Dubois MP, Pasteur N, Raymond M: A novel acetylcholinesterase gene in mosquitoes codes for the insecticide target and is non-homologous to the *ace* gene in *Drosophila*. *Proc R Soc Lond B* 2002, **269**:2007–2016.
 47. Weill W, Lutfalla G, Mogensen K, Chandre F, Berthomieu A, Berticat C, Pasteur N, Phillips A, Fort P, Raymond M: Corrigendum - Insecticide resistance in mosquito vectors. *Nature* 2003, **425**:366.
 48. Weill W, Lutfalla G, Mogensen K, Chandre F, Berthomieu A, Berticat C, Pasteur N, Phillips A, Fort P, Raymond M: Insecticide resistance in mosquito vectors. *Nature* 2003, **423**:136–137.
 49. Weill W, Berthomieu A, Berticat C, Lutfalla G, Negre V, Pasteur N, Phillips A, Leonetti J-P, Fort P, Raymond M: Insecticide resistance: a silent base prediction. *Curr Biol* 2004, **14**:R552.
 50. Weill W, Malcolm C, Chandre F, Mogensen K, Berthomieu A, Marquie M, Raymond M: The unique mutation in *ace-1* giving high insecticide resistance is easily detectable in mosquito vectors. *Insect Mol Biol* 2004, **13**:1–7.
 51. Nabeshima T, Kozaki T, Tomita T, Kono Y: An amino acid substitution on the second acetylcholinesterase in the pirimicarb-resistant strains of the peach potato aphid, *Myzus persicae*. *Biochem Biophys Res Commun* 2003, **307**:15–22.
 52. Nabeshima T, Mori A, Kozaki T, Iwata Y, Hidoh O, Harada S, Kasai S, Severson DW, Kono Y, Tomita T: An amino acid substitution attributable to insecticide-insensitivity of acetylcholinesterase in a Japanese encephalitis vector mosquito, *Culex tritaeniorhynchus*. *Biochem Biophys Res Commun* 2004, **313**:794–801.
 53. Naik PK, Sindhura TS, Singh H: Quantitative structure-activity relationship (QSAR) for insecticides: development of predictive in vivo insecticide activity models. *SAR QSAR Environ Res* 2009, **20**:551–566.

doi:10.1186/1756-3305-6-31

Cite this article as: Temeyer et al.: Acetylcholinesterase of the sand fly, *Phlebotomus papatasi* (Scopoli): cDNA sequence, baculovirus expression, and biochemical properties. *Parasites & Vectors* 2013 **6**:31.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

