

RESEARCH ARTICLE

Open Access



Perceived neighborhood safety related to physical activity but not recreational screen-based sedentary behavior in adolescents

Clare M. Lenhart¹, Andrew Wiemken², Alexandra Hanlon², Mackenzie Perket³ and Freda Patterson^{3*}

Abstract

Background: A growing proportion of adolescents have poor cardiovascular health behaviors, including low levels of physical activity and high levels of sedentary behavior, thus increasing the likelihood of poor heart health in later years. This study tested the hypothesis that low perceived neighborhood safety would be associated with low levels of physical activity and high levels of recreational sedentary behavior in high-school students.

Methods: Using cross-sectional, weighted data from the 2015 Pennsylvania (USA) State and Philadelphia city Youth Risk Behavior Survey, multivariable logistic regression modeling was used to examine the association between perceived neighborhood safety, and physical activity levels and recreational screen-based sedentary behavior time respectively, while controlling for potential confounders.

Results: After adjustment for other significant correlates of physical activity, students with low perceived neighborhood safety had a 21% reduced odds of being physically active on 5 or more days of the last week as compared to those who felt safe ($p = 0.044$). Perceived safety was not related to sedentary behavior; but sports team participation emerged as a strong correlate of low screen-based sedentary behavior ($OR = 0.73, p = .002$).

Conclusion: These data add to a growing body of work demonstrating the importance of perceived safety with physical activity levels in youth. Sports team participation may be a viable target to reduce screen-based sedentary time.

Background

Cardiovascular disease is the leading cause of death among adults globally, and in the United States [1, 2]. Stemming adult cardiovascular disease through the promotion of heart health behaviors, such as increased physical activity, and decreased recreational sedentary behavior in youth, is a viable public health approach [3]. Underpinning the importance of this work are data showing that cardiovascular disease and the associated risk factors (e.g., obesity) is on the rise among youth [4–7]. For example, between 2001 and 2009 a 30.5% increase in type 2 diabetes in youth was reported [5].

Approximately three in five 2–19 year olds are overweight or obese [8], while one in five youth aged 8 to 17 years have an adverse lipid concentration and slightly more than 1 in 10 have either borderline high, or high, blood pressure [9]. Combined, less than 1% of adolescents achieve ideal heart health as defined by the American Heart Association, with fewer Black/African American youth meeting the recommendations for physical activity, blood pressure and cholesterol than Non-Hispanic Whites [10, 11].

Ecological models of health behavior have helped elucidate the role of built environment and neighborhood factors on health [12]. In the context of physical activity and sedentary behavior, access and proximity to green space and recreation facilities, safe neighborhoods with low crime as well as streetscape features such as walkable sidewalks have all been significantly associated with increased physical activity and lower sedentary behavior

* Correspondence: fredap@udel.edu

³Department of Behavioral Health and Nutrition, College of Health Sciences, University of Delaware, 019 Carpenter Sports Building, Newark, DE, USA
Full list of author information is available at the end of the article

(e.g., screen-based sedentary behavior) in both adults and youth [13–15]. Lower-income, urban neighborhoods tend to be characterized by features not conducive to higher physical activity and lower recreational sedentary behavior such as limited green space, unappealing aesthetics, damaged sidewalks and poor quality recreation facilities [16–18]. Residents of these neighborhoods, regardless of race, have a lower likelihood of engaging in habitual heart health behaviors such as physical activity. For example, Wilson and colleagues reported that in low-income, urban environments, racial differences in physical inactivity were ameliorated with Whites and Blacks having similar odds of low activity levels [19]. Racially diverse, urban populations represent a particularly high-risk group for cardiovascular disease and poor heart health behaviors.

Neighborhood safety, and in particular, perceived neighborhood safety, are distinct environmental constructs that are relevant to urban residents. In adult populations, neighborhood safety has been associated with physical activity in some [20] but not all studies [21]. Perceptions of neighborhood safety may account for this difference [22, 23]. For example, perception of neighborhood safety is a determinant of physical activity, even after adjustment for objective markers of safety such as neighborhood crime [22–24]. Higher levels of perceived safety and access to recreation facilities predicted higher physical activity levels in adults of low but not high socioeconomic status [19]. Given that perceived safety is a salient and highly associated construct of physical activity and to a lesser extent, sedentary behavior, in adults, the same may be true for youth.

To add to this body of research, we used a population sample of adolescents aged 14–18 years ($n = 4020$) to examine the extent to which lower levels of perceived safety was associated with lower levels of physical activity and higher recreational screen-based sedentary behavior. We adjusted for other known determinants of physical activity and sedentary behavior including gender, race/ethnicity [25, 26], poor mental health [27], age [28], overweight [29], sleep [30], fruit and vegetable consumption [31], sports team and physical education class participation [32].

Methods

Study sample

Data from the 2015 Centers for Disease Control (CDC) Youth Risk Behaviors Survey (YRBS) administered in the state of Pennsylvania and the School District of Philadelphia were used for this analysis. A two-stage, cluster sample design was used to produce a representative sample of high school students across Pennsylvania. In the first stage, proportional sampling of schools was based on school enrollment size; in the second stage,

classroom-level sampling was conducted in which specific classrooms were randomly sampled from a comprehensive list of all classes available in a given required period; all students in sampled classes were eligible to participate [33]. Data for the current analysis include classes from 64 high schools across the state of Pennsylvania and 25 additional high schools from the urban center of Philadelphia. Student response rate from the Pennsylvania state high schools was 64% ($n = 2899$) and from the Philadelphia schools it was 68% ($n = 1896$). Parental consent and student assent were obtained prior to surveying and all study procedures were approved by the Institutional Review Board and by each participating school district (IRB# 21251).

For the present study, students providing complete data for study demographics (age, sex, race), the independent variable of perceived safety, and dependent variables of physical activity and sedentary behavior were retained. The resulting sample, $n = 4020$ ($N = 474,768$), represents 83.8% of all respondents in the original Philadelphia and Pennsylvania samples.

Instrument

The YRBS is a school-based biennial, self-administered paper-pencil survey of US high school students, administered at the national level by CDC and by states, territories, tribal areas or large urban school districts. Conducted since 1991, the surveys include representative samples of students in grades 9–12 and monitor six categories of priority health-risk behaviors: behaviors relating to injuries and violence, sexual risk behaviors, tobacco use, alcohol and other drug use, unhealthy diet, and physical inactivity [33, 34]. Reliability assessments of the YRBS instrument have demonstrated moderate reliability (mean kappa 61%) [35].

Measures

The study outcome of physical activity was measured using the single item “During the past 7 days, on how many days were you physically active for a total of at least 60 minutes per day? (Add up all the time you spent in any kind of physical activity that increased your heart rate and made you breathe hard some of the time?)” Responses were dichotomized: participants reporting 5 or more days of moderate physical activity per week were considered physically active [36].

The study outcome of recreational screen-based sedentary behavior was estimated using two questions that queried students to self-report television watching and recreational computer use. The survey items were: “On an average school day, how many hours do you watch TV?” and “On an average school day, how many hours do you play video or computer games or use a computer

for something that is not school work?" Seven response options were presented for each question ranging from, "I do not [watch TV/play video or computer games] on an average school day" to "5 or more hours per day." Responses to both questions were added together and based on a median split of the resulting combined data, those students reporting four or less hours of combined TV and video/computer game usage were compared to those reporting more than 4 h in recreational screen-based sedentary behavior. The main exposure variable of neighborhood safety was assessed using the item, "How often do you feel safe and secure in your neighborhood?" Those responding "always" or "most of the time" were compared to those responding "sometimes," "rarely," or "never" [37].

Several health behaviors were assessed using single survey items that were coded dichotomously (yes/no). These behaviors included: current smoking (one or more cigarettes in the last 30 days); consumption of three or more servings of vegetables per day, consumption of two or more servings of fruit per day; adequate sleep (≥ 8 h) on a school night; any sports team participation in the last year; and physical education class participation on at least 1 day per week [33].

Participants' socio-demographic characteristics including age, sex and ethnicity/race were self-reported. In the current analyses the ethnicity/race items were collapsed to generate the following categories: White, Black or African American, Asian, Hispanic and Other. The "Other" category was comprised of students who self-identified as more than one race, American Indian or Alaskan Native, Native Hawaiian or Other Pacific Islander.

Statistical analysis

Descriptive statistics were generated for all study variables. Tests of association between each independent variable with the dichotomous outcomes of physical activity and recreational screen-based sedentary behavior were examined using chi-square test of independence. The independent variable of perceived neighborhood safety along with those variables modestly associated with the outcome measures ($p < 0.2$) were entered into separate multivariable logistic regression models to identify variables uniquely associated with physical activity and screen-based sedentary behavior after controlling for other related variables. The multivariable model of physical activity included perceived neighborhood safety, gender, race/ethnicity, academic achievement, weight status, area of residence, sports team participation, sleep, fruit consumption, vegetable consumption, and screen-based sedentary behavior. The multivariable model of screen-based sedentary behavior included perceived neighborhood safety, gender, race/ethnicity, academic achievement, weight status, physical

education participation, sports team participation, vegetable consumption, and physical activity. Analyses were run using SPSS version 23. Having achieved necessary overall response rate within Philadelphia and Pennsylvania, defined as $\geq 60\%$ after multiplying the school response rate and the student response rate, data were weighted to account for student non-response and demographic distributions in each district such that the weight reflects the number of students the respondent represents [33, 34]. Sampling errors were estimated from the primary sampling units and strata and using Taylor series linearization for SPSS version 23 (IBM, New York).

Results

Participant characteristics

Approximately half the sample (50.5%) was female, 59.5% were 16 years or older, 70.9% were non-Hispanic white, 15.2% were non-Hispanic black, 8.4% self-identified as Hispanic and 2.4% were Asian. Eighty-three (82.8%) percent earned mostly A and B grades in school, while 29.5% were overweight or obese. Only 5.0% of the sample were part of the city of Philadelphia YRBS data collection while 7.6% reported rarely or never feeling safe in their neighborhood. Approximately one half of the sample (44.9%) reported at least 60-min of moderate physical activity on 5 or more days of the week. High sedentary behavior (≥ 4 h/day recreational screen time) was reported by 51.7% of the sample. (see Table 1 for a full listing of variables). These data are consistent with those reported by other data sources.

Variables associated with five or more days per week of physical activity

At the bivariate level, perceived safety was associated with physical activity such that adolescents who reported rarely or never feeling safe had a significantly higher proportion of low physical activity (8.9% reported low physical activity versus 5.9% who reported high physical activity; $\chi^2 = 19.35$, $p \leq .001$). High levels of physical activity was also significantly associated with being male, being younger than 16 years, self-identifying as non-Hispanic white, earning mostly A and B grades, not being overweight or obese, and not participating in the Philadelphia city data collection. In terms of other health-related variables, engaging in physical education class at least 1 day per week, participating in one or more sports teams, accruing eight or more hours of sleep, eating two or more servings of fruit and three or more servings of vegetables per day as well as not accruing four or more hours of recreational screen time per day were all significantly associated with high levels of physical activity (Table 1).

Table 1 Bivariate associations with Physical Activity (PA)

	Total sample N = 474,768 (n = 4020) CI = 405,872 - 543,663	High PA N = 213,443 (n = 1623) CI = 177,470 - 249,415	Low PA N = 261,324 (n = 2397) CI = 222,873 - 299,776	χ^2
Demographic variables				
Sex				
Female	50.5% (48.0% – 52.9%)	40.7% (38.0% – 43.6%)	58.4% (55.0% – 61.8%)	80.44***
Male	49.5% (47.1%–52.0%)	59.3% (56.4%– 62.0%)	41.6% (38.2% – 45.0%)	
Age				
< 16 years	40.5% (33.6% – 47.8%)	43.3% (35.5% – 51.4%)	38.2% (31.6% – 45.4%)	4.88*
≥ 16 years	59.5% (52.2% – 66.4%)	56.7% (48.6% – 64.5%)	61.8% (54.6% – 68.4%)	
Race				
Non-Hispanic white	70.9% (63.5% – 77.3%)	75.7% (69.1% – 81.2%)	67.0% (58.5% – 74.5%)	17.77***
Non-Hispanic black	15.2% (10.6% – 21.4%)	12.3% (8.2% – 17.9%)	17.6% (12.2% – 24.7%)	9.61**
Hispanic	8.4% (6.2% – 11.3%)	6.9% (4.9% – 9.6%)	9.6% (7.1% – 13.0%)	11.68***
Asian	2.4% (1.8% – 3.2%)	1.6% (1.0% – 2.6%)	3.0% (2.2% – 4.1%)	7.25**
Other	3.1% (2.6% – 3.8%)	3.5% (2.6% – 4.7%)	2.8% (2.2% – 3.5%)	1.54
Academic achievement				
Earned mostly A's & B's	82.8% (79.3% – 85.7%)	85.2% (81.2%– 88.5%)	80.8% (77.0% – 84.1%)	6.34*
Overweight/obese	29.5% (27.2% – 32.0%)	25.3% (22.6% – 28.2%)	33.1% (30.3% – 36.0%)	23.62***
Environmental level variables				
Dataset (PA State vs. Philadelphia City)				
Philadelphia City	5.0% (3.9% – 6.3%)	3.6% (2.8% – 4.6%)	6.1% (4.7% – 7.8%)	34.26***
Perceived safety				
Rarely/never feel safe	7.6% (6.1% – 9.4%)	5.9% (4.6% – 7.7%)	8.9% (7.1% – 11.1%)	19.35***
Health related variables				
≥ 4 h screen time/day	51.7% (48.7% – 54.6%)	45.1% (40.8% – 49.3%)	57.1% (53.3% – 60.8%)	19.34***
≥ 1 day PE/week	58.6% (50.7% – 66.1%)	65.3% (56.9% – 72.7%)	53.2% (45.0% – 61.2%)	23.20***
≥ 1 sports teams	60.9% (57.9% – 63.8%)	79.4% (76.1% – 82.4%)	45.8% (42.6% – 49.0%)	258.78***
≥ 8 h sleep/night	25.6% (23.4% – 27.9%)	28.8% (25.7% – 32.0%)	23.1% (20.6% – 25.7%)	9.78**
Current alcohol use	30.6% (27.7% – 33.7%)	31.0% (27.4% – 34.9%)	30.3% (26.7% – 34.1%)	0.12
Current cigarette use	11.9% (9.7% – 14.6%)	11.8% (9.1% – 15.2%)	12.6% (9.4% – 15.3%)	0.02
≥ 2 servings fruit/day	28.9% (26.5% – 31.3%)	36.3% (33.0% – 39.7%)	22.8% (20.6% – 25.2%)	84.88***
≥ 3 servings veg/day	10.5% (8.8% – 12.4%)	15.2% (12.2% – 18.7%)	6.6% (5.2% – 8.4%)	31.43***

The association between each dichotomous independent variable and the dichotomous outcomes of physical activity (column 3 & 4) was examined using chi-square test of independence

* $p \leq .05$ ** $p \leq .01$ *** $p \leq .001$

Variables associated with recreational screen-based sedentary behavior

High levels (≥4 h/day) of recreational screen-based sedentary behavior was not significantly associated with perceptions of neighborhood safety ($\chi^2 = 0.51, p > .05$). Being male, aged less than 16 years, self-identifying as non-Hispanic Black, being obese or overweight and residing in the city of Philadelphia were significantly associated with high levels (≥4 h/day) of recreational screen-based sedentary behavior (Table 2). High levels of physical activity (≥5 days of physical activity), playing on one or more sports

teams and eating three or more servings of vegetables per day was associated with low levels of recreational screen-based sedentary behavior (Table 1).

Independent association between perceived safety and high physical activity

In a multivariable logistic regression model of high physical activity (≥ 5 days of activity per week) with perceived neighborhood safety as the independent variable of interest, adolescents who reported low levels of perceived safety had a 21% lower odds of being highly physical activity (OR = 0.79, CI = 0.56-0.89, $p = .044$) even

Table 2 Bivariate associations with recreational screen based sedentary behavior

	≥4 Hours Recreational Screen time per Day N = 245,305; n = 2152 CI = 206,106 – 284,503	<4 Hours Recreational Screen-time per Day N = 229,463; n = 1868 CI = 194,206 – 264,719	χ^2
Demographic variables			
Sex			
Female	48.1% (44.8% – 51.3%)	53.0% (49.4% – 56.6%)	4.17*
Male	51.9% (48.7% – 55.2%)	47.0% (43.4 – 50.6%)	4.17*
Age			
< 16 years	42.9% (35.6% – 50.6%)	37.9% (30.8% – 45.7%)	
≥ 16 years	57.1% (49.4% – 64.4%)	62.1% (54.3% – 69.2%)	3.88*
Race			
Non-Hispanic white	67.8% (59.3% – 75.2%)	74.2% (67.6% – 79.8%)	12.67***
Non-Hispanic black	18.1% (12.5% – 25.5%)	12.1% (8.1% – 17.7%)	9.65**
Hispanic	9.2% (6.9% – 12.2%)	7.6% (5.2% – 10.8%)	3.57
Asian	1.7% (1.2% – 2.4%)	3.1% (2.2% – 4.3%)	10.28***
Other	3.2% (2.5% – 4.0%)	3.1% (2.3% – 4.1%)	0.02
Academic achievement			
Earned mostly A's & B's	78.4% (74.4% – 81.9%)	87.5% (83.6% – 90.5%)	21.28***
Overweight or obese	34.2% (31.1% – 37.5%)	24.6% (21.2% – 28.3%)	14.28***
Environmental level variables			
Philadelphia City	5.6% (4.3% – 7.1%)	4.4% (3.4% – 5.7%)	8.07**
Perceived safety			
Rarely/never feel safe	7.9% (6.3% – 9.9%)	7.2% (5.3% – 9.6%)	0.51
Health related variables			
≥ 5 days physical activity	39.5% (35.7% – 43.5%)	51.1% (47.2% – 55.0%)	19.34***
≥ 1 day PE/week	59.3% (51.3% – 66.9%)	57.8% (49.0% – 66.1%)	0.29
≥ 1 sports teams	55.0% (51.5% – 58.4%)	67.2% (63.1% – 71.0%)	27.09***
≥ 8 h sleep/night	25.0% (22.5% – 27.7%)	26.3% (23.3% – 29.6%)	0.55
Current alcohol use	29.5% (26.0% – 33.3%)	31.8% (28.0% – 35.8%)	0.97
Current cigarette use	12.4% (9.7% – 15.6%)	11.4% (9.0% – 14.4%)	0.47
≥ 2 servings fruit/day	27.2% (24.5% – 30.0%)	30.7% (27.5% – 34.0%)	3.38
≥ 3 servings veg/ day	8.6% (6.8% – 10.9%)	12.4% (10.5% – 14.7%)	12.30***

The association between each dichotomous independent variable and the dichotomous outcome of high vs. low recreational screen based sedentary behavior (column 2 & 3) was examined using chi-square test of independence

* $p \leq .05$ ** $p \leq .01$ *** $p \leq .001$

after adjustment for other known determinants of physical activity (Table 3).

Other variables significantly associated with higher levels of physical activity included being male (OR = 2.11, CI = 1.78-2.51, $p < 0.001$), self-reporting as non-Hispanic white relative to all other racial and ethnic groups (OR = 1.42, CI = 1.12-1.80, $p = 0.004$), sports team participation (OR = 4.40, CI = 3.56-5.43, $p < 0.001$), adequate fruit consumption (OR = 1.62, CI = 1.36-1.92, $p < 0.001$), adequate vegetable consumption (OR = 2.17, CI = 1.48-3.20, $p < 0.001$). In contrast, high levels (≥ 4 h/day) of screen-based sedentary behavior was significantly

associated with reduced odds of being active on five or more days per week (OR = 0.68, CI = 0.54-0.88, $p < 0.001$) (Table 3).

Independent association between perceived safety and recreational screen-based sedentary behavior

In a multivariable model of high levels of recreational screen-based sedentary behavior, perceived safety did not emerge as a significant correlate, after adjustment for other significant correlates of recreational screen-based sedentary behavior. Instead, being male (OR = 1.25, CI = 1.01-1.53, $p = 0.039$) and

Table 3 Multivariable associations with physical activity on five or more days of the last week

	OR	95% Confidence Interval	P
Gender (1 = Male)	2.11	1.78 - 2.51	<0.001
Race (1 = Non-Hispanic White)	1.42	1.12 - 1.80	0.004
Academic achievement	0.94	0.73 - 1.21	0.640
Overweight or obese	0.87	0.72-1.04	0.129
City of Philadelphia only	1.11	0.82-1.48	0.499
Sports team participation	4.40	3.56-5.43	<0.001
≥ 8 or more hours of sleep	1.17	0.10-1.41	0.117
Adequate fruit consumption	1.62	1.36-1.92	<0.001
Adequate vegetable consumption	2.17	1.48-3.20	<0.001
High screen based sedentary behavior	0.69	0.54-0.88	<0.001
Low perceived safety	0.79	0.56-0.89	0.044

being overweight or obese (OR = 1.39, CI = 1.10-1.75, $p = 0.006$) were associated with high levels of screen-based sedentary behavior. Additionally, high academic achievement (OR = 0.60, CI = 0.44-0.80, $p < 0.001$), sports team participation (OR = 0.73, CI = 0.52-0.86, $p = 0.002$), adequate vegetable consumption (OR = 0.67, CI = 0.52-0.86, $p = 0.002$) and high physical activity (OR = 0.67, CI = 0.54-0.85, $p = 0.001$) was independently associated with low recreational screen-based sedentary behavior (Table 4).

Discussion

Increasing physical activity and reducing recreational screen-based sedentary behavior (i.e., television viewing and leisure computer use), are particularly important to improving cardiovascular health indices in youth and to forestall the incidence and progression of poor heart health in later years [38, 39]. Low neighborhood safety and perceptions of neighborhood safety are associated with lower levels of physical activity in adult populations

Table 4 Multivariable associations with high recreational screen based sedentary behavior (≥4 h/day)

	OR	95% Confidence interval	P
Gender (1 = Male)	1.25	1.01-1.53	0.039
Race (1 = Non-Hispanic White)	0.83	0.68-1.01	0.066
Academic achievement	0.60	0.44-0.80	<0.001
Overweight or obese	1.39	1.10-1.75	0.006
Physical Education class participation	1.18	0.94-1.49	0.152
Sports team participation	0.73	0.61-0.89	0.002
Adequate vegetable consumption	0.67	0.52-0.86	0.002
High physical activity	0.67	0.54-0.85	0.001
Low perceived safety	0.84	0.60-1.18	0.311

[40]. In expanding on this literature to examine the association between recreational screen-based sedentary behavior with perceived safety in a population sample of adolescents, these data showed that while perceived safety was not associated with recreational screen-based sedentary behavior, it was significantly associated with physical activity. Specifically, youth who perceived their neighborhoods as unsafe had a 21% reduced odds of achieving five or more days of physical activity in the last week after adjustment for other known determinants of physical activity including sex, obesity status, and dietary intake. Other notable findings include data showing that participation in sports teams were strong correlates of sufficient physical activity and reduced recreational screen-based sedentary time. Together these population data suggest a role of perceived safety in physical activity promotion and underscore the importance of organized, formal physical activity opportunities for youth in achieving recommended levels of leisure-time physical activity and lower screen-based sedentary behavior.

In adults, high levels of neighborhood crime and low perceived safety have consistently been shown to relate to lower levels of physical activity [41]. Similarly, low levels of perceived neighborhood safety in parents is associated with lower levels of outdoor play and physical activity in their young children [42, 43], while high levels of perceived neighborhood safety correlated positively with child activity levels [44, 45]. Data have also shown children’s perceptions of neighborhood features (i.e., traffic density) to correspond with their parent’s perceptions [46]. Studies mostly conducted outside the US have reported adolescent perceptions of safety to be associated with physical activity levels [47, 48]. Increased fear of going outside to exercise [49], and lower feelings of community trust or social cohesion [50], are possible mechanisms that may help explain why low levels of perceived safety may be significantly associated with low levels of physical activity. Alternatively, higher levels of physical activity may lead to improved neighborhood perceptions. Evaluation of these, and other possible mechanisms for this association, is necessary in adolescents.

That perceived safety was significantly related to physical activity but not recreational screen-based sedentary behavior is consistent with data showing that physical activity and sedentary behavior in youth are not always related [51, 52]. In terms of other studies that have related perceived safety to sedentary behavior, data have shown that children whose parents perceived their neighborhood as unsafe had significantly higher levels of sedentary behavior than children whose parents perceived their neighborhood as safe [44, 53]. Our finding that adolescent perceived safety did not relate to recreational screen-based sedentary time diverges

from this parental literature, and suggests that targeting parental perceived safety as opposed to youth perceived safety may be more important to reducing sedentary time in youth.

High levels of physical activity, sports team participation, and vegetable consumption emerged as being significantly related to reduced recreation screen-based sedentary behavior in this sample of youth. These data add to previous work showing increased physical activity [54], no participation on sports teams [55], and low vegetable consumption [56], to be associated with increased sedentary behavior. Together, this work aligns with a growing consideration of the clustering of health behaviors, and the recognition that negative health behaviors, such as poor diet and physical inactivity, tend to co-occur [57]. Addressing multiple risk factors in adolescents has demonstrated feasibility [58]; the current data provide candidate behaviors (i.e., physical activity, sports team participation and vegetable consumption) to be addressed in the context of reducing sedentary behavior.

The clinical and policy implications of these data are three-fold. First, these data reiterate the concept that sedentary behavior is not simply the other side of the physical activity coin, and instead is a separate construct that can and should be addressed independently [51]. Second, improving perceived neighborhood safety has been identified by these data as a potentially important intervention target to increase physical activity in youth. Changes to the physical and built environment may impact how spaces are used and by whom. For example, “greening” approaches (i.e., cleaning vacant lots, planting grass and trees, building a wooden fence around the perimeter) have been shown to hold multiple benefits for a community including increased feelings of perceived safety [59] and activity levels in urban adult residents [60]. Whether such an approach could increase levels of perceived safety and physical activity in adolescents has yet to be examined but such exploration holds relevance in the broader context of increasing community-level safety and physical activity. Third, these data highlight how critical organized physical activity and team sports are to youth activity levels. Concerning national trends include the drastic reductions in the required physical education class (i.e., 25% in 8th grade and 5% of 12th grade) [61] and declining quality of physical education instruction with only a third of adolescents being physically active in PE class for more than 20 min 3 to 5 days a week [62]. Efforts to improve perceived neighborhood safety and increase the requirement and quality of physical education class as well as promotion of sport team participation could incite far-reaching benefits for cardiovascular health in the short and longer term among youth [63].

While the current study is one of the largest population level examinations of the association between

perceived safety in physical activity and recreational screen-based sedentary behavior in adolescents, these data should be interpreted with consideration to the fact that these data are based on self-report estimations and that several study constructs, including physical activity, was measured using single items. Importantly, observed neighborhood safety was not measured or considered in the multivariable models thus multilevel modeling based on neighborhood-level factors was not feasible. Physical activity and dietary habits were queried with the brief, standard YRBS questions rather than with longer construct-specific inventories. Finally, lack of sample homogeneity limits the generalizability of these results. Despite these limitations, these data point to the importance of considering sedentary behavior and physical activity as separate entities and present sports team participation and perceived safety, respectively, as being possible intervention targets to achieve improvements in these cardiovascular risk behaviors in adolescents.

Conclusions

Findings from this study of a large and diverse sample of adolescents add to a growing body of work demonstrating the importance of perceived safety with physical activity levels in youth. Perceived safety was associated with physical activity but not screen-based sedentary behavior, underscoring the notion that physical activity and sedentary behavior are distinct constructs. Sports team participation may be a viable target to reduce screen-based sedentary time among adolescents and should be assessed with future interventional research.

Acknowledgements

The authors are indebted to YRBS coordinators at the School District of Philadelphia and the Pennsylvania Department of Health as well as to the students and classroom teachers who were impacted by data collection.

Funding

Funding for this work was provided by the School District of Philadelphia (FP). The study sponsor had no role in the analysis of these data or the preparation of this manuscript.

Availability of data and materials

The datasets analyzed for the current study are not publicly available due contractual obligations between the study team, the School District of Philadelphia, and the Pennsylvania Department of Health but may be available from the corresponding author on reasonable request.

Authors' contributions

CML contributed to study conceptualization, lead data analysis, and contributed to the final manuscript. AW contributed to data analysis and development of the manuscript. AH contributed to data analysis, the final manuscript, and study conceptualization. MP contributed to data preparation and to manuscript development. FP oversaw data collection, analysis, and manuscript development and was instrumental on study conceptualization. All authors read and approved the final manuscript.

Ethics approval and consent to participate

The present study received IRB approval from East Stroudsburg University. Parental consent and student assent were obtained prior to collection of the original study data, a process then overseen by the IRB at Temple University.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹East Stroudsburg University, East Stroudsburg, PA, USA. ²University of Pennsylvania, Philadelphia, PA, USA. ³Department of Behavioral Health and Nutrition, College of Health Sciences, University of Delaware, 019 Carpenter Sports Building, Newark, DE, USA.

Received: 29 September 2016 Accepted: 13 September 2017

Published online: 18 September 2017

References

- Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, Cushman M, de Ferranti S, Despres JP, Fullerton HJ, Howard VJ, et al. Heart disease and stroke statistics—2015 update: a report from the American Heart Association. *Circulation*. 2015;131(4):e29–322.
- Mendis S, Puska P, Norrving B. *Global Atlas on Cardiovascular Disease Prevention and Control*. Geneva: World Health Organization; 2011.
- Kavey RE, Daniels SR, Lauer RM, Atkins DL, Hayman LL, Taubert K, American Heart A. American Heart Association guidelines for primary prevention of atherosclerotic cardiovascular disease beginning in childhood. *Circulation*. 2003;107(11):1562–6.
- Ogden CL, Lamb MM, Carroll MD, Flegal KM. Obesity and socioeconomic status in children and adolescents: United States. *NCHS Data Brief*. 2005–2008;2010(51):1–8.
- Dabelea D, Mayer-Davis EJ, Saydah S, Imperatore G, Linder B, Divers J, Bell R, Badaru A, Talton JW, Crume T, et al. Prevalence of type 1 and type 2 diabetes among children and adolescents from 2001 to 2009. *JAMA*. 2014; 311(17):1778–86.
- Hannon TS, Arslanian SA. The changing face of diabetes in youth: lessons learned from studies of type 2 diabetes. *Ann N Y Acad Sci*. 2015;1353:113–37.
- Urbina EM, Gao Z, Khoury PR, Martin LJ, Dolan LM. Insulin resistance and arterial stiffness in healthy adolescents and young adults. *Diabetologia*. 2012;55(3):625–31.
- Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011–2012. *JAMA*. 2014;311(8):806–14.
- Kit BK, Kuklina E, Carroll MD, Ostchega Y, Freedman DS, Ogden CL. Prevalence of and trends in dyslipidemia and blood pressure among US children and adolescents, 1999–2012. *JAMA Pediatr*. 2015;169(3):272–9.
- Lenhart CM, Manjeliuevskaia J, Echeverri A, Patterson F. Cardiovascular health among Philadelphia adolescents: analysis of youth risk behaviour data, 2011. *Cardiol Young*. 2014;24(4):748–51.
- Shay CM, Ning H, Daniels SR, Rooks CR, Gidding SS, Lloyd-Jones DM. Status of cardiovascular health in US adolescents: prevalence estimates from the National Health and Nutrition Examination Surveys (NHANES) 2005–2010. *Circulation*. 2013;127(13):1369–76.
- Sallis JF, Cervero RB, Ascher W, Henderson KA, Kraft MK, Kerr J. An ecological approach to creating active living communities. *Annu Rev Public Health*. 2006;27:297–322.
- Barnett A, Cerin E, Ching CS, Johnston JM, Lee RS. Neighbourhood environment, sitting time and motorised transport in older adults: a cross-sectional study in Hong Kong. *BMJ Open*. 2015;5(4):e007557.
- Cain KL, Millstein RA, Sallis JF, Conway TL, Gavand KA, Frank LD, Saelens BE, Geremia CM, Chapman J, Adams MA, et al. Contribution of streetscape audits to explanation of physical activity in four age groups based on the Microscale Audit of Pedestrian Streetscapes (MAPS). *Soc Sci Med*. 2014;116: 82–92.
- Sanders T, Feng X, Fahey PP, Lonsdale C, Astell-Burt T. The influence of neighbourhood green space on children's physical activity and screen time: findings from the longitudinal study of Australian children. *Int J Behav Nutr Phys Act*. 2015;12:126.
- Gordon-Larsen P, Nelson MC, Page P, Popkin BM. Inequality in the built environment underlies key health disparities in physical activity and obesity. *Pediatrics*. 2006;117(2):417–24.
- Hoehner CM, Brennan Ramirez LK, Elliott MB, Handy SL, Brownson RC. Perceived and objective environmental measures and physical activity among urban adults. *Am J Prev Med*. 2005;28(2 Suppl 2):105–16.
- McCormack G, Giles-Corti B, Lange A, Smith T, Martin K, Pikora TJ. An update of recent evidence of the relationship between objective and self-report measures of the physical environment and physical activity behaviours. *J Sci Med Sport*. 2004;7(1 Suppl):81–92.
- Wilson-Frederick SM, Thorpe RJ Jr, Bell CN, Bleich SN, Ford JG, LaVeist TA. Examination of race disparities in physical inactivity among adults of similar social context. *Ethn Dis*. 2014;24(3):363–9.
- Gomez JE, Johnson BA, Selva M, Sallis JF. Violent crime and outdoor physical activity among inner-city youth. *Prev Med*. 2004;39(5):876–81.
- Ruijsbroek A, Droomers M, Groenewegen PP, Hardyns W, Stronks K. Social safety, self-rated general health and physical activity: changes in area crime, area safety feelings and the role of social cohesion. *Health Place*. 2015;31:39–45.
- Bailey EJ, Malecki KC, Engelman CD, Walsh MC, Bersch AJ, Martinez-Donate AP, Peppard PE, Nieto FJ. Predictors of discordance between perceived and objective neighborhood data. *Ann Epidemiol*. 2014;24(3):214–21.
- Giles-Corti B, Donovan RJ. The relative influence of individual, social and physical environment determinants of physical activity. *Soc Sci Med*. 2002; 54(12):1793–812.
- Janssen I. Crime and perceptions of safety in the home neighborhood are independently associated with physical activity among 11–15 year olds. *Prev Med*. 2014;66:113–7.
- Belcher BR, Berrigan D, Dodd KW, Ermen BA, Chou CP, Spruijt-Metz D. Physical activity in US youth: effect of race/ethnicity, age, gender, and weight status. *Med Sci Sports Exerc*. 2010;42(12):2211–21.
- Smith NR, Lewis DJ, Fahy A, Eldridge S, Taylor SJ, Moore DG, Clark C, Stansfeld SA, Cummins S. Individual socio-demographic factors and perceptions of the environment as determinants of inequalities in adolescent physical and psychological health: the Olympic Regeneration in East London (ORIEL) study. *BMC Public Health*. 2015;15:150.
- Kremer P, Elshaug C, Leslie E, Toumbourou JW, Patton GC, Williams J. Physical activity, leisure-time screen use and depression among children and young adolescents. *J Sci Med Sport*. 2014;17(2):183–7.
- Lenhart C, Patterson F, Brown M, O'Brien M, Nelson D. Disparity in physical activity among urban youth: an ecologically guided assessment. *Am J Health Educ*. 2014;45
- Hughes AR, Henderson A, Ortiz-Rodriguez V, Artinou ML, Reilly JJ. Habitual physical activity and sedentary behaviour in a clinical sample of obese children. *Int J Obes*. 2006;30(10):1494–500.
- Foti KE, Eaton DK, Lowry R, McKnight-Ely LR. Sufficient sleep, physical activity, and sedentary behaviors. *Am J Prev Med*. 2011;41(6):596–602.
- Lowry R, Michael S, Demissie Z, Kann L, Galuska DA. Associations of physical activity and sedentary behaviors with dietary behaviors among US high school students. *J Obes*. 2015;2015:876524.
- Lenhart CM, Hanlon A, Daly B, Brown MD, Patterson F. Gender disparity in structured physical activity and overall activity level in adolescence: evaluation of youth risk behavior surveillance data. *ISRN Public Health*. 2012;
- Centers for Disease C, Prevention, Brener ND, Kann L, Shanklin S, Kinchen S, Eaton DK, Hawkins J, Flint KH, Centers for Disease C, et al. Methodology of the Youth Risk Behavior Surveillance System—2013. *MMWR Recomm Rep*. 2013;62(RR-1):1–20.
- Brener ND, Kann L, Kinchen SA, Grunbaum JA, Whalen L, Eaton D, Hawkins J, Ross JG. Methodology of the youth risk behavior surveillance system. *MMWR Recomm Rep*. 2004;53(RR-12):1–13.
- Brener ND, Kann L, McManus T, et al. Reliability of the 1999 youth risk behavior survey questionnaire. *J Adolesc Health*. 2002;31(4):336–42.
- Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, et al. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Circulation*. 2007;116(9):1081–93.

37. Sallis JF, Johnson MF, Calfas KJ, Caparosa S, Nichols JF. Assessing perceived physical environmental variables that may influence physical activity. *Res Q Exerc Sport*. 1997;68(4):345–51.
38. Janz KF, Dawson JD, Mahoney LT. Tracking physical fitness and physical activity from childhood to adolescence: the muscatine study. *Med Sci Sports Exerc*. 2000;32(7):1250–7.
39. Nicklas TA, von Duvillard SP, Berenson GS. Tracking of serum lipids and lipoproteins from childhood to dyslipidemia in adults: the Bogalusa Heart Study. *Int J Sports Med*. 2002;23(Suppl 1):S39–43.
40. Bennett GG, McNeill LH, Wolin KY, Duncan DT, Puleo E, Emmons KM. Safe to walk? Neighborhood safety and physical activity among public housing residents. *PLoS Med*. 2007;4(10):1599–606. discussion 1607
41. Wilson DK, Kirtland KA, Ainsworth BE, Addy CL. Socioeconomic status and perceptions of access and safety for physical activity. *Ann Behav Med*. 2004;28(1):20–8.
42. Carver A, Timperio A, Hesketh K, Crawford D. Are children and adolescents less active if parents restrict their physical activity and active transport due to perceived risk? *Soc Sci Med*. 2010;70(11):1799–805.
43. Weir LA, Etelson D, Brand DA. Parents' perceptions of neighborhood safety and children's physical activity. *Prev Med*. 2006;43(3):212–7.
44. Santos MP, Pizarro AN, Mota J, Marques EA. Parental physical activity, safety perceptions and children's independent mobility. *BMC Public Health*. 2013;13:584.
45. Tappe KA, Glanz K, Sallis JF, Zhou C, Saelens BE. Children's physical activity and parents' perception of the neighborhood environment: neighborhood impact on kids study. *Int J Behav Nutr Phys Act*. 2013;10:39.
46. Dunton GF, Intille SS, Wolch J, Pentz MA. Children's perceptions of physical activity environments captured through ecological momentary assessment: a validation study. *Prev Med*. 2012;55(2):119–21.
47. Nichol M, Janssen I, Pickett W. Associations between neighborhood safety, availability of recreational facilities, and adolescent physical activity among Canadian youth. *J Phys Act Health*. 2010;7(4):442–50.
48. Carver A, Timperio A, Crawford D. Perceptions of neighborhood safety and physical activity among youth: the CLAN study. *J Phys Act Health*. 2008;5(3):430–44.
49. Harrison RA, Gemmell I, Heller RF. The population effect of crime and neighbourhood on physical activity: an analysis of 15,461 adults. *J Epidemiol Community Health*. 2007;61(1):34–9.
50. Craddock AL, Kawachi I, Colditz GA, Gortmaker SL, Buka SL. Neighborhood social cohesion and youth participation in physical activity in Chicago. *Soc Sci Med*. 2009;68(3):427–35.
51. Owen N, Leslie E, Salmon J, Fotheringham MJ. Environmental determinants of physical activity and sedentary behavior. *Exerc Sport Sci Rev*. 2000;28(4):153–8.
52. Biddle SJ, Gorely T, Marshall SJ, Murdey I, Cameron N. Physical activity and sedentary behaviours in youth: issues and controversies. *J R Soc Promot Heal*. 2004;124(1):29–33.
53. Datar A, Nicosia N, Shier V. Parent perceptions of neighborhood safety and children's physical activity, sedentary behavior, and obesity: evidence from a national longitudinal study. *Am J Epidemiol*. 2013;177(10):1065–73.
54. Lowry R, Lee SM, Fulton JE, Demissie Z, Kann L. Obesity and other correlates of physical activity and sedentary behaviors among US high school students. *J Obes*. 2013;2013:276318.
55. Silva DA, Dos Santos Silva RJ. Association between sports participation and sedentary behavior during school recess among Brazilian adolescents. *J Hum Kinet*. 2015;45:225–32.
56. Borraccino A, Lemma P, Berchiolla P, Cappello N, Inchley J, Dalmaso P, Charrier L, Cavallo F, Italian HG. Unhealthy food consumption in adolescence: role of sedentary behaviours and modifiers in 11-, 13- and 15-year-old Italians. *Eur J Pub Health*. 2016;26(4):650–6.
57. Prochaska JJ, Prochaska JO. A review of multiple health behavior change interventions for primary prevention. *Am J Lifestyle Med*. 2011;5(3):208–21.
58. Mauriello LM, Ciavatta MM, Paiva AL, Sherman KJ, Castle PH, Johnson JL, Prochaska JM. Results of a multi-media multiple behavior obesity prevention program for adolescents. *Prev Med*. 2010;51(6):451–6.
59. Garvin EC, Cannuscio CC, Branas CC. Greening vacant lots to reduce violent crime: a randomised controlled trial. *Inj Prev*. 2013;19(3):198–203.
60. Branas CC, Cheney RA, MacDonald JM, Tam VW, Jackson TD, Ten Have TR. A difference-in-differences analysis of health, safety, and greening vacant urban space. *Am J Epidemiol*. 2011;174(11):1296–306.
61. Grunbaum JA, Kann L, Kinchen S, Ross J, Hawkins J, Lowry R, Harris WA, McManus T, Chyen D, Collins J. Youth risk behavior surveillance—United States, 2003. *MMWR Surveill Summ*. 2004;53(2):1–96.
62. Story M, Kaphingst KM, French S. The role of schools in obesity prevention. *Futur Child*. 2006;16(1):109–42.
63. Kriemler S, Meyer U, Martin E, van Sluijs E, Andersen L, Martin B. Effect of school-based interventions on physical activity and fitness in children and adolescents: a review of reviews and systematic update. *Br J Sports Med*. 2011;45(11). <https://doi.org/10.1136/bjsports-2011-090186>.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

