

## RESEARCH

## Open Access



# A new half-discrete Mulholland-type inequality with multi-parameters

Qiliang Huang and Bicheng Yang\*

\*Correspondence:  
bcyang@gdei.edu.cn  
Department of Mathematics,  
Guangdong University of  
Education, Guangzhou, Guangdong  
510303, P.R. China

**Abstract**

By means of weight functions and Hermite-Hadamard's inequality, a new half-discrete Mulholland-type inequality with a best constant factor is given. A best extension with multi-parameters, some equivalent forms, the operator expressions as well as some particular cases are considered.

**MSC:** 26D15; 47A07

**Keywords:** Mulholland-type inequality; weight function; equivalent form

**1 Introduction**

Assuming that  $f, g \in L^2(\mathbf{R}_+)$ ,  $\|f\| = \{\int_0^\infty f^2(x) dx\}^{\frac{1}{2}} > 0$ ,  $\|g\| > 0$ , we have the following Hilbert integral inequality (cf. [1]):

$$\int_0^\infty \int_0^\infty \frac{f(x)g(y)}{x+y} dx dy < \pi \|f\| \|g\|, \quad (1.1)$$

where the constant factor  $\pi$  is best possible. If  $a = \{a_m\}_{m=1}^\infty, b = \{b_n\}_{n=1}^\infty \in l^2$ ,  $\|a\| = \{\sum_{m=1}^\infty a_m^2\}^{\frac{1}{2}} > 0$ ,  $\|b\| > 0$ , then we have the following discrete Hilbert inequality:

$$\sum_{m=1}^\infty \sum_{n=1}^\infty \frac{a_m b_n}{m+n} < \pi \|a\| \|b\|, \quad (1.2)$$

with the same best constant factor  $\pi$ . Inequalities (1.1) and (1.2) are important in analysis and its applications (cf. [2, 3]). On the other hand, we have the following Mulholland inequality with the same best constant factor  $\pi$  (cf. [1, 4]):

$$\sum_{m=2}^\infty \sum_{n=2}^\infty \frac{a_m b_n}{\ln mn} < \pi \left\{ \sum_{m=2}^\infty m a_m^2 \sum_{n=2}^\infty n b_n^2 \right\}^{\frac{1}{2}}. \quad (1.3)$$

In 1998, by introducing an independent parameter  $\lambda \in (0, 1]$ , Yang [5] gave an extension of (1.1). Generalizing the results from [5], Yang [3] gave some extensions of (1.1) and (1.2) as follows: If  $p > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $\lambda_1 + \lambda_2 = \lambda \in \mathbf{R}$ ,  $k_\lambda(x, y)$  is a non-negative homogeneous function of degree  $-\lambda$  satisfying

$$k(\lambda_1) = \int_0^\infty k_\lambda(t, 1) t^{\lambda_1-1} dt \in \mathbf{R}_+,$$

$$\phi(x) = x^{p(1-\lambda_1)-1}, \psi(x) = x^{q(1-\lambda_2)-1}, f(x), g(y) \geq 0,$$

$$f \in L_{p,\phi}(\mathbf{R}_+) = \left\{ f \mid \|f\|_{p,\phi} := \left\{ \int_0^\infty \phi(x) |f(x)|^p dx \right\}^{\frac{1}{p}} < \infty \right\},$$

$g \in L_{q,\psi}(\mathbf{R}_+)$ , and  $\|f\|_{p,\phi}, \|g\|_{q,\psi} > 0$ , then

$$\int_0^\infty \int_0^\infty k_\lambda(x, y) f(x) g(y) dx dy < k(\lambda_1) \|f\|_{p,\phi} \|g\|_{q,\psi}, \tag{1.4}$$

where the constant factor  $k(\lambda_1)$  is best possible. Moreover, if  $k_\lambda(x, y)$  is finite and  $k_\lambda(x, y)x^{\lambda_1-1} (k_\lambda(x, y)y^{\lambda_2-1})$  is decreasing for  $x > 0 (y > 0)$ , then for  $a_m, b_n \geq 0$ ,

$$a = \{a_m\}_{m=1}^\infty \in l_{p,\phi} = \left\{ a \mid \|a\|_{p,\phi} := \left\{ \sum_{m=1}^\infty \phi(m) |a_m|^p \right\}^{\frac{1}{p}} < \infty \right\},$$

and  $b = \{b_n\}_{n=1}^\infty \in l_{q,\psi}, \|a\|_{p,\phi}, \|b\|_{q,\psi} > 0$ , we have

$$\sum_{m=1}^\infty \sum_{n=1}^\infty k_\lambda(m, n) a_m b_n < k(\lambda_1) \|a\|_{p,\phi} \|b\|_{q,\psi}, \tag{1.5}$$

where the constant factor  $k(\lambda_1)$  is still the best possible. Clearly, for  $p = q = 2, \lambda = 1$ ,  $k_1(x, y) = \frac{1}{x+y}$  and  $\lambda_1 = \lambda_2 = \frac{1}{2}$ , (1.4) reduces to (1.1), while (1.5) reduces to (1.2).

Some other results about Hilbert-type inequalities can be found in [6–13]. On half-discrete Hilbert-type inequalities with the general non-homogeneous kernels, Hardy *et al.* provided a few results in Theorem 351 of [1]. But they did not prove that the constant factors are best possible. In 2005, Yang [14] gave a result with the kernel  $\frac{1}{(1+nx)^\lambda}$  by introducing a variable and proved that the constant factor is best possible. Recently, Wang and Yang [15] gave a more accurate reverse half-discrete Hilbert-type inequality, and Yang [16] provided the following half-discrete Hilbert inequality with best constant factor:

$$\int_0^\infty f(x) \sum_{n=1}^\infty \frac{a_n}{x+n} dx < \pi \|f\| \|a\|. \tag{1.6}$$

In this paper, by means of weight functions and Hermite-Hadamard’s inequality, a new half-discrete Mulholland-type inequality similar to (1.3) and (1.6) with a best possible constant factor is given as follows:

$$\int_1^\infty f(x) \sum_{n=2}^\infty \frac{a_n}{1 + \ln x \ln n} dx < \pi \left\{ \int_1^\infty x f^2(x) dx \sum_{n=2}^\infty n a_n^2 \right\}^{\frac{1}{2}}. \tag{1.7}$$

Moreover, a best extension of (1.7) with multi-parameters, some equivalent forms, the operator expressions as well as some particular cases are considered.

### 2 Some lemmas

**Lemma 2.1** *If  $0 < \sigma < \lambda$  ( $\sigma \leq 1$ ),  $\alpha > 0$ ,  $\beta \geq \frac{2}{3}$ ,  $\delta \in \{-1, 1\}$ , the weight functions  $\omega(n)$  and  $\varpi(x)$  are defined by*

$$\omega(n) := (\ln \beta n)^\sigma \int_{\frac{1}{\alpha}}^\infty \frac{(\ln \alpha x)^{\delta\sigma-1}}{x(1 + \ln^\delta \alpha x \ln \beta n)^\lambda} dx, \quad n \in \mathbf{N} \setminus \{1\}, \tag{2.1}$$

$$\varpi(x) := (\ln \alpha x)^{\delta\sigma} \sum_{n=2}^\infty \frac{(\ln \beta n)^{\sigma-1}}{n(1 + \ln^\delta \alpha x \ln \beta n)^\lambda}, \quad x \in \left(\frac{1}{\alpha}, \infty\right), \tag{2.2}$$

then we have

$$\varpi(x) < \omega(n) = B(\sigma, \lambda - \sigma). \tag{2.3}$$

*Proof* Substituting  $t = \ln^\delta \alpha x \ln \beta n$  in (2.1), and by a simple calculation, for  $\delta \in \{-1, 1\}$ , we have

$$\omega(n) = \int_0^\infty \frac{1}{(1+t)^\lambda} t^{\sigma-1} dt = B(\sigma, \lambda - \sigma).$$

For fixed  $x > \frac{1}{\alpha}$ , in view of the conditions, it is easy to find that

$$h(x, y) := \frac{(\ln \beta y)^{\sigma-1}}{y(1 + \ln^\delta \alpha x \ln \beta y)^\lambda} = \frac{1}{y(1 + \ln^\delta \alpha x \ln \beta y)^\lambda (\ln \beta y)^{1-\sigma}}$$

is decreasing and strictly convex with  $h'_y(x, y) < 0$  and  $h''_{y^2}(x, y) > 0$ , for  $y \in (\frac{3}{2}, \infty)$ . Hence by the Hermite-Hadamard inequality (cf. [17]), we find

$$\begin{aligned} \varpi(x) &< (\ln \alpha x)^{\delta\sigma} \int_{\frac{3}{2}}^\infty \frac{1}{y(1 + \ln^\delta \alpha x \ln \beta y)^\lambda (\ln \beta y)^{1-\sigma}} dy \\ &\stackrel{t = \ln^\delta \alpha x \ln \beta y}{=} \int_{\ln^\delta \alpha x \ln(\frac{3}{2}\beta)}^\infty \frac{t^{\sigma-1}}{(1+t)^\lambda} dt \leq B(\sigma, \lambda - \sigma), \end{aligned}$$

and then (2.3) follows. □

**Lemma 2.2** *Let the assumptions of Lemma 2.1 be fulfilled and, additionally, let  $p > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $a_n \geq 0$ ,  $n \in \mathbf{N} \setminus \{1\}$ ,  $f(x)$  is a non-negative measurable function in  $(\frac{1}{\alpha}, \infty)$ . Then we have the following inequalities:*

$$\begin{aligned} J &:= \left\{ \sum_{n=2}^\infty \frac{1}{n} (\ln \beta n)^{p\sigma-1} \left[ \int_{\frac{1}{\alpha}}^\infty \frac{f(x)}{(1 + \ln^\delta \alpha x \ln \beta n)^\lambda} dx \right]^p \right\}^{\frac{1}{p}} \\ &\leq [B(\sigma, \lambda - \sigma)]^{\frac{1}{q}} \left\{ \int_{\frac{1}{\alpha}}^\infty \varpi(x) x^{p-1} (\ln \alpha x)^{p(1-\delta\sigma)-1} f^p(x) dx \right\}^{\frac{1}{p}}, \end{aligned} \tag{2.4}$$

$$\begin{aligned} L_1 &:= \left\{ \int_{\frac{1}{\alpha}}^\infty \frac{(\ln \alpha x)^{q\delta\alpha-1}}{x[\varpi(x)]^{q-1}} \left[ \sum_{n=2}^\infty \frac{a_n}{(1 + \ln^\delta \alpha x \ln \beta n)^\lambda} \right]^q dx \right\}^{\frac{1}{q}} \\ &\leq \left\{ B(\sigma, \lambda - \sigma) \sum_{n=2}^\infty n^{q-1} (\ln \beta n)^{q(1-\sigma)-1} a_n^q \right\}^{\frac{1}{q}}. \end{aligned} \tag{2.5}$$

*Proof* By Hölder’s inequality (cf. [17]) and (2.3), it follows that

$$\begin{aligned}
 & \left[ \int_{\frac{1}{\alpha}}^{\infty} \frac{f(x) dx}{(1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda}} \right]^p \\
 &= \left\{ \int_{\frac{1}{\alpha}}^{\infty} \frac{1}{(1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda}} \left[ \frac{(\ln \alpha x)^{(1-\delta\sigma)/q} x^{\frac{1}{q}} f(x)}{(\ln \beta n)^{(1-\sigma)/p} n^{\frac{1}{p}}} \right] \right. \\
 & \quad \times \left. \left[ \frac{(\ln \beta n)^{(1-\sigma)/p} n^{\frac{1}{p}}}{(\ln \alpha x)^{(1-\delta\sigma)/q} x^{\frac{1}{q}}} \right] dx \right\}^p \leq \int_{\frac{1}{\alpha}}^{\infty} \frac{x^{p-1} (\ln \alpha x)^{(1-\delta\sigma)(p-1)} f^p(x) dx}{(1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda} n (\ln \beta n)^{1-\sigma}} \\
 & \quad \times \left\{ \int_{\frac{1}{\alpha}}^{\infty} \frac{n^{q-1}}{(1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda}} \frac{(\ln \beta n)^{(1-\sigma)(q-1)}}{x (\ln \alpha x)^{1-\delta\sigma}} dx \right\}^{p-1} \\
 &= \left\{ \omega(n) n^{q-1} (\ln \beta n)^{q(1-\sigma)-1} \right\}^{p-1} \int_{\frac{1}{\alpha}}^{\infty} \frac{x^{p-1} (\ln \alpha x)^{(1-\delta\sigma)(p-1)} f^p(x) dx}{(1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda} n (\ln \beta n)^{1-\sigma}} \\
 &= [B(\sigma, \lambda - \sigma)]^{p-1} n (\ln \beta n)^{1-p\sigma} \int_{\frac{1}{\alpha}}^{\infty} \frac{x^{p-1} (\ln \alpha x)^{(1-\delta\sigma)(p-1)} f^p(x) dx}{(1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda} n (\ln \beta n)^{1-\sigma}}.
 \end{aligned}$$

Then by Lebesgue term-by-term integration theorem (cf. [18]), we have

$$\begin{aligned}
 J &\leq [B(\sigma, \lambda - \sigma)]^{\frac{1}{q}} \left\{ \sum_{n=2}^{\infty} \int_{\frac{1}{\alpha}}^{\infty} \frac{x^{p-1} (\ln \alpha x)^{(1-\delta\sigma)(p-1)} f^p(x) dx}{(1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda} n (\ln \beta n)^{1-\sigma}} \right\}^{\frac{1}{p}} \\
 &= [B(\sigma, \lambda - \sigma)]^{\frac{1}{q}} \left\{ \int_{\frac{1}{\alpha}}^{\infty} \sum_{n=2}^{\infty} \frac{x^{p-1} (\ln \alpha x)^{(1-\delta\sigma)(p-1)} f^p(x) dx}{(1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda} n (\ln \beta n)^{1-\sigma}} \right\}^{\frac{1}{p}} \\
 &= [B(\sigma, \lambda - \sigma)]^{\frac{1}{q}} \left\{ \int_{\frac{1}{\alpha}}^{\infty} \varpi(x) x^{p-1} (\ln \alpha x)^{p(1-\delta\sigma)-1} f^p(x) dx \right\}^{\frac{1}{p}},
 \end{aligned}$$

hence, (2.4) follows.

By Hölder’s inequality again, we have

$$\begin{aligned}
 & \left[ \sum_{n=2}^{\infty} \frac{a_n}{(1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda}} \right]^q \\
 &= \left\{ \sum_{n=2}^{\infty} \frac{1}{(1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda}} \left[ \frac{(\ln \alpha x)^{(1-\delta\sigma)/q} x^{\frac{1}{q}}}{(\ln \beta n)^{(1-\sigma)/p} n^{\frac{1}{p}}} \right] \right. \\
 & \quad \times \left. \left[ \frac{(\ln \beta n)^{(1-\sigma)/p} n^{\frac{1}{p}} a_n}{(\ln \alpha x)^{(1-\delta\sigma)/q} x^{\frac{1}{q}}} \right] \right\}^q \leq \left\{ \sum_{n=2}^{\infty} \frac{x^{p-1} (\ln \alpha x)^{(1-\delta\sigma)(p-1)}}{n (1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda} (\ln \beta n)^{1-\sigma}} \right\}^{q-1} \\
 & \quad \times \sum_{n=2}^{\infty} \frac{n^{q-1}}{(1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda}} \frac{(\ln \beta n)^{(1-\sigma)(q-1)}}{x (\ln \alpha x)^{1-\delta\sigma}} a_n^q \\
 &= \frac{x [\varpi(x)]^{q-1}}{(\ln \alpha x)^{q\delta\sigma-1}} \sum_{n=2}^{\infty} \frac{n^{q-1}}{x (1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda}} (\ln \alpha x)^{\delta\sigma-1} (\ln \beta n)^{(1-\sigma)(q-1)} a_n^q.
 \end{aligned}$$

By the Lebesgue term-by-term integration theorem, we have

$$\begin{aligned} L_1 &\leq \left\{ \int_{\frac{1}{\alpha}}^{\infty} \sum_{n=2}^{\infty} \frac{n^{q-1}}{x(1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda}} (\ln \alpha x)^{\delta\sigma-1} (\ln \beta n)^{(1-\sigma)(q-1)} a_n^q dx \right\}^{\frac{1}{q}} \\ &= \left\{ \sum_{n=2}^{\infty} \left[ (\ln \beta n)^{\sigma} \int_{\frac{1}{\alpha}}^{\infty} \frac{(\ln \alpha x)^{\delta\sigma-1} dx}{x(1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda}} \right] n^{q-1} (\ln \beta n)^{q(1-\sigma)-1} a_n^q \right\}^{\frac{1}{q}} \\ &= \left\{ \sum_{n=2}^{\infty} \omega(n) n^{q-1} (\ln \beta n)^{q(1-\sigma)-1} a_n^q \right\}^{\frac{1}{q}}, \end{aligned}$$

and in view of (2.3), inequality (2.5) follows. □

### 3 Main results

We introduce the functions

$$\begin{aligned} \Phi_{\delta}(x) &:= x^{p-1} (\ln \alpha x)^{p(1-\delta\sigma)-1} \quad \left( x > \frac{1}{\alpha} \right), \\ \Psi(n) &:= n^{q-1} (\ln \beta n)^{q(1-\sigma)-1} \quad (n \in \mathbf{N} \setminus \{1\}), \end{aligned}$$

wherefrom  $[\Phi_{\delta}(x)]^{1-q} = \frac{1}{x} (\ln \alpha x)^{q\delta\sigma-1}$ , and  $[\Psi(n)]^{1-p} = \frac{1}{n} (\ln \beta n)^{p\sigma-1}$ .

**Theorem 3.1** *If  $0 < \sigma < \lambda$  ( $\sigma \leq 1$ ),  $\alpha > 0$ ,  $\beta \geq \frac{2}{3}$ ,  $\delta \in \{-1, 1\}$ ,  $p > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $f(x)$ ,  $a_n \geq 0$ ,  $f \in L_{p,\Phi}(\frac{1}{\alpha}, \infty)$ ,  $a = \{a_n\}_{n=2}^{\infty} \in l_{q,\Psi}$ ,  $\|f\|_{p,\Phi_{\delta}} > 0$ , and  $\|a\|_{q,\Psi} > 0$ , then we have the following equivalent inequalities:*

$$\begin{aligned} I &:= \sum_{n=2}^{\infty} a_n \int_{\frac{1}{\alpha}}^{\infty} \frac{f(x) dx}{(1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda}} \\ &= \int_{\frac{1}{\alpha}}^{\infty} f(x) \sum_{n=2}^{\infty} \frac{a_n dx}{(1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda}} < B(\sigma, \lambda - \sigma) \|f\|_{p,\Phi_{\delta}} \|a\|_{q,\Psi}, \end{aligned} \tag{3.1}$$

$$J = \left\{ \sum_{n=2}^{\infty} [\Psi(n)]^{1-p} \left[ \int_{\frac{1}{\alpha}}^{\infty} \frac{f(x) dx}{(1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda}} \right]^p \right\}^{\frac{1}{p}} < B(\sigma, \lambda - \sigma) \|f\|_{p,\Phi_{\delta}}, \tag{3.2}$$

$$L := \left\{ \int_{\frac{1}{\alpha}}^{\infty} [\Phi_{\delta}(x)]^{1-q} \left[ \sum_{n=2}^{\infty} \frac{a_n}{(1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda}} \right]^q dx \right\}^{\frac{1}{q}} < B(\sigma, \lambda - \sigma) \|a\|_{q,\Psi}, \tag{3.3}$$

where the constant  $B(\sigma, \lambda - \sigma)$  is the best possible in the above inequalities.

*Proof* The two expressions for  $I$  in (3.1) follow from Lebesgue’s term-by-term integration theorem. By (2.4) and (2.3), we have (3.2). By Hölder’s inequality, we have

$$I = \sum_{n=2}^{\infty} \left[ \Psi^{\frac{-1}{q}}(n) \int_{\frac{1}{\alpha}}^{\infty} \frac{f(x) dx}{(1 + \ln^{\delta} \alpha x \ln \beta n)^{\lambda}} \right] \left[ \Psi^{\frac{1}{p}}(n) a_n \right] \leq J \|a\|_{q,\Psi}. \tag{3.4}$$

Then by (3.2), we have (3.1).

On the other hand, assuming that (3.1) is valid, we set

$$a_n := [\Psi(n)]^{1-p} \left[ \int_{\frac{1}{\alpha}}^{\infty} \frac{f(x) dx}{(1 + \ln^\delta \alpha x \ln \beta n)^\lambda} \right]^{p-1}, \quad n \in \mathbf{N} \setminus \{1\}.$$

It follows that  $J^{p-1} = \|a\|_{q,\Psi}$ . By (2.4), we find  $J < \infty$ . If  $J = 0$ , then (3.2) is trivially valid; if  $J > 0$ , then by (3.1), we have

$$\|a\|_{q,\Psi}^q = J^{q(p-1)} = J^p = I < B(\sigma, \lambda - \sigma) \|f\|_{p,\Phi_\delta} \|a\|_{q,\Psi},$$

namely,  $\|a\|_{q,\Psi}^{q-1} = J < B(\sigma, \lambda - \sigma) \|f\|_{p,\Phi_\delta}$ . That is, (3.2) is equivalent to (3.1).

By (2.3) we have  $[\varpi(x)]^{1-q} > [B(\sigma, \lambda - \sigma)]^{1-q}$ . Then in view of (2.5), we have (3.3). By Hölder's inequality, we find

$$I = \int_{\frac{1}{\alpha}}^{\infty} [\Phi_\delta^{\frac{1}{p}}(x) f(x)] \left[ \Phi_\delta^{\frac{-1}{p}}(x) \sum_{n=2}^{\infty} \frac{a_n}{(1 + \ln^\delta \alpha x \ln \beta n)^\lambda} \right] dx \leq \|f\|_{p,\Phi_\delta} L. \tag{3.5}$$

Then by (3.3), we have (3.1).

On the other hand, assume that (3.1) is valid. Setting

$$f(x) := [\Phi_\delta(x)]^{1-q} \left[ \sum_{n=2}^{\infty} \frac{a_n}{(1 + \ln^\delta \alpha x \ln \beta n)^\lambda} \right]^{q-1}, \quad x \in \left(\frac{1}{\alpha}, \infty\right),$$

then  $L^{q-1} = \|f\|_{p,\Phi_\delta}$ . By (2.5), we find  $L < \infty$ . If  $L = 0$ , then (3.3) is trivially valid; if  $L > 0$ , then by (3.1), we have

$$\|f\|_{p,\Phi_\delta}^p = L^{p(q-1)} = L^q = I < B(\sigma, \lambda - \sigma) \|f\|_{p,\Phi_\delta} \|a\|_{q,\Psi},$$

therefore  $\|f\|_{p,\Phi_\delta}^{p-1} = L < B(\sigma, \lambda - \sigma) \|a\|_{q,\Psi}$ , that is, (3.3) is equivalent to (3.1). Hence, inequalities (3.1), (3.2), and (3.3) are equivalent.

For  $0 < \varepsilon < p(\lambda - \sigma)$ , setting  $E_\delta := \{x; x > \frac{1}{\alpha}, \ln^\delta \alpha x \in (0, 1)\}$ ,

$$\tilde{f}(x) = \frac{1}{x} (\ln \alpha x)^{\delta(\sigma + \frac{\varepsilon}{p}) - 1}, \quad x \in E_\delta; \quad \tilde{f}(x) = 0, \quad \left\{x; x > \frac{1}{\alpha}\right\} \setminus E_\delta,$$

and  $\tilde{a}_n = \frac{1}{n} (\ln \beta n)^{\sigma - \frac{\varepsilon}{q} - 1}$ ,  $n \in \mathbf{N} \setminus \{1\}$ , if there exists a positive number  $k (\leq B(\sigma, \lambda - \sigma))$ , such that (3.1) is valid when replacing  $B(\sigma, \lambda - \sigma)$  with  $k$ , then in particular, for  $\delta = \pm 1$ , setting  $u = \ln^\delta \alpha x$ , it follows that

$$\begin{aligned} \int_{E_\delta} \frac{dx}{x(\ln \alpha x)^{-\delta\varepsilon+1}} &= \int_0^1 \frac{|\delta| u^{\delta-1}}{u^{-\varepsilon+\delta}} du = \frac{1}{\varepsilon}, \\ \tilde{I} &:= \sum_{n=2}^{\infty} \int_{\frac{1}{\alpha}}^{\infty} \frac{1}{(1 + \ln^\delta \alpha x \ln \beta n)^\lambda} \tilde{a}_n \tilde{f}(x) dx < k \|\tilde{f}\|_{p,\Phi_\delta} \|\tilde{a}\|_{q,\Psi} \\ &= k \left\{ \int_{E_\delta} \frac{dx}{x(\ln \alpha x)^{-\delta\varepsilon+1}} \right\}^{\frac{1}{p}} \left\{ \frac{1}{2(\ln 2\beta)^{\varepsilon+1}} + \sum_{n=3}^{\infty} \frac{1}{n(\ln \beta n)^{\varepsilon+1}} \right\}^{\frac{1}{q}} \end{aligned}$$

$$\begin{aligned}
 &< k \left( \frac{1}{\varepsilon} \right)^{\frac{1}{p}} \left\{ \frac{1}{2(\ln 2\beta)^{\varepsilon+1}} + \int_2^\infty \frac{1}{x(\ln \beta x)^{\varepsilon+1}} dx \right\}^{\frac{1}{q}} \\
 &= \frac{k}{\varepsilon} \left\{ \frac{\varepsilon}{2(\ln 2\beta)^{\varepsilon+1}} + \frac{1}{(\ln 2\beta)^\varepsilon} \right\}^{\frac{1}{q}}, \tag{3.6} \\
 \tilde{I} &= \sum_{n=2}^\infty \frac{1}{n} (\ln \beta n)^{\sigma - \frac{\varepsilon}{q} - 1} \int_{E_\delta} \frac{(\ln \alpha x)^{\delta(\sigma + \frac{\varepsilon}{p}) - 1}}{x(1 + \ln^\delta \alpha x \ln \beta n)^\lambda} dx \\
 &\stackrel{t = \ln^\delta \alpha x \ln \beta n}{=} \sum_{n=2}^\infty \frac{1}{n(\ln \beta n)^{\varepsilon+1}} \int_0^{\ln \beta n} \frac{1}{(1+t)^\lambda} t^{\sigma + \frac{\varepsilon}{p} - 1} dt \\
 &= B\left(\sigma + \frac{\varepsilon}{p}, \lambda - \sigma - \frac{\varepsilon}{p}\right) \sum_{n=2}^\infty \frac{1}{n(\ln \beta n)^{\varepsilon+1}} - A(\varepsilon) \\
 &> B\left(\sigma + \frac{\varepsilon}{p}, \lambda - \sigma - \frac{\varepsilon}{p}\right) \int_2^\infty \frac{1}{y(\ln \beta y)^{\varepsilon+1}} dy - A(\varepsilon) \\
 &= \frac{1}{\varepsilon(\ln 2\beta)^\varepsilon} B\left(\sigma + \frac{\varepsilon}{p}, \lambda - \sigma - \frac{\varepsilon}{p}\right) - A(\varepsilon), \\
 A(\varepsilon) &:= \sum_{n=2}^\infty \frac{1}{n(\ln \beta n)^{\varepsilon+1}} \int_{\ln \beta n}^\infty \frac{1}{(t+1)^\lambda} t^{\sigma + \frac{\varepsilon}{p} - 1} dt. \tag{3.7}
 \end{aligned}$$

We find

$$\begin{aligned}
 0 < A(\varepsilon) &\leq \sum_{n=2}^\infty \frac{1}{n(\ln \beta n)^{\varepsilon+1}} \int_{\ln \beta n}^\infty \frac{1}{t^\lambda} t^{\sigma + \frac{\varepsilon}{p} - 1} dt \\
 &= \frac{1}{\frac{\lambda}{2} - \frac{\varepsilon}{p}} \sum_{n=2}^\infty \frac{1}{n(\ln \beta n)^{\sigma + \frac{\varepsilon}{q} + 1}} < \infty,
 \end{aligned}$$

and so  $A(\varepsilon) = O(1)(\varepsilon \rightarrow 0^+)$ . Hence by (3.6) and (3.7), it follows that

$$\frac{1}{(\ln 2\beta)^\varepsilon} B\left(\sigma + \frac{\varepsilon}{p}, \lambda - \sigma - \frac{\varepsilon}{p}\right) - \varepsilon O(1) < k \left\{ \frac{\varepsilon}{2(\ln 2\beta)^{\varepsilon+1}} + \frac{1}{(\ln 2\beta)^\varepsilon} \right\}^{\frac{1}{q}},$$

and  $B(\sigma, \lambda - \sigma) \leq k(\varepsilon \rightarrow 0^+)$ . Hence  $k = B(\sigma, \lambda - \sigma)$  is the best value of (3.1).

By the equivalence of the inequalities, the constant factor  $B(\sigma, \lambda - \sigma)$  in (3.2) ((3.3)) is the best possible. Otherwise, we would reach the contradiction by (3.4) ((3.5)) that the constant factor in (3.1) is not the best possible.  $\square$

**Remark 3.2** (i) Define the first type half-discrete Hilbert-type operator  $T_1 : L_{p, \Phi_\delta}(\frac{1}{\alpha}, \infty) \rightarrow l_{p, \Psi^{1-p}}$  as follows: For  $f \in L_{p, \Phi_\delta}(\frac{1}{\alpha}, \infty)$ , we define  $T_1 f \in l_{p, \Psi^{1-p}}$  by

$$T_1 f(n) = \int_{\frac{1}{\alpha}}^\infty \frac{1}{(1 + \ln^\delta \alpha x \ln \beta n)^\lambda} f(x) dx, \quad n \in \mathbf{N} \setminus \{1\}.$$

Then by (3.2),  $\|T_1 f\|_{p, \Psi^{1-p}} \leq B(\sigma, \lambda - \sigma) \|f\|_{p, \Phi_\delta}$  and so  $T_1$  is a bounded operator with  $\|T_1\| \leq B(\sigma, \lambda - \sigma)$ . Since by Theorem 3.1, the constant factor in (3.2) is best possible, we have  $\|T_1\| = B(\sigma, \lambda - \sigma)$ .

(ii) Define the second type half-discrete Hilbert-type operator  $T_2 : l_{q,\Psi} \rightarrow L_{q,\Phi_\delta^{1-q}}(\frac{1}{\alpha}, \infty)$  as follows: For  $a \in l_{q,\Psi}$ , we define  $T_2 a \in L_{q,\Phi_\delta^{1-q}}(\frac{1}{\alpha}, \infty)$  by

$$T_2 a(x) = \sum_{n=2}^{\infty} \frac{1}{(1 + \ln^\delta \alpha x \ln \beta n)^\lambda} a_n, \quad x \in \left(\frac{1}{\alpha}, \infty\right).$$

Then by (3.3),  $\|T_2 a\|_{q,\Phi_\delta^{1-q}} \leq B(\sigma, \lambda - \sigma) \|a\|_{q,\Psi}$  and so  $T_2$  is a bounded operator with  $\|T_2\| \leq B(\sigma, \lambda - \sigma)$ . Since by Theorem 3.1, the constant factor in (3.3) is best possible, we have  $\|T_2\| = B(\sigma, \lambda - \sigma)$ .

**Remark 3.3** For  $p = q = 2, \lambda = 1, \sigma = \frac{1}{2}, \delta = 1$  in (3.1), (3.2), and (3.3), (i) if  $\alpha = \beta = 1$ , then we have (1.7) and the following equivalent inequalities:

$$\sum_{n=2}^{\infty} \frac{1}{n} \left( \int_1^{\infty} \frac{f(x)}{1 + \ln x \ln n} dx \right)^2 < \pi^2 \int_1^{\infty} x f^2(x) dx, \tag{3.8}$$

$$\int_1^{\infty} \frac{1}{x} \left( \sum_{n=2}^{\infty} \frac{a_n}{1 + \ln x \ln n} \right)^2 dx < \pi^2 \sum_{n=2}^{\infty} n a_n^2; \tag{3.9}$$

(ii) if  $\alpha = \beta = \frac{2}{3}$ , then we have the following equivalent inequalities:

$$\int_{\frac{3}{2}}^{\infty} \sum_{n=2}^{\infty} \frac{a_n f(x) dx}{1 + \ln \frac{2}{3} x \ln \frac{2}{3} n} < \pi \left\{ \int_{\frac{3}{2}}^{\infty} x f^2(x) dx \sum_{n=2}^{\infty} n a_n^2 \right\}^{\frac{1}{2}}, \tag{3.10}$$

$$\sum_{n=2}^{\infty} \frac{1}{n} \left( \int_{\frac{3}{2}}^{\infty} \frac{f(x)}{1 + \ln \frac{2}{3} x \ln \frac{2}{3} n} dx \right)^2 < \pi^2 \int_{\frac{3}{2}}^{\infty} x f^2(x) dx, \tag{3.11}$$

$$\int_{\frac{3}{2}}^{\infty} \frac{1}{x} \left( \sum_{n=2}^{\infty} \frac{a_n}{1 + \ln \frac{2}{3} x \ln \frac{2}{3} n} \right)^2 dx < \pi^2 \sum_{n=2}^{\infty} n a_n^2. \tag{3.12}$$

**Remark 3.4** For  $\delta = -1$  in (3.1), (3.2), and (3.3), setting  $F(x) = \ln^\lambda(\alpha x)f(x), \mu = \lambda - \sigma (> 0)$ , and  $\Phi(x) := x^{p-1}(\ln \alpha x)^{p(1-\mu)-1}$ , we have the following new equivalent inequalities with the same best possible constant factor  $B(\sigma, \mu)$ :

$$\begin{aligned} \sum_{n=2}^{\infty} a_n \int_{\frac{1}{\alpha}}^{\infty} \frac{F(x) dx}{\ln^\lambda(\alpha \beta n x)} &= \int_{\frac{1}{\alpha}}^{\infty} F(x) \sum_{n=2}^{\infty} \frac{a_n}{\ln^\lambda(\alpha \beta n x)} dx \\ &< B(\sigma, \mu) \|F\|_{p,\Phi} \|a\|_{q,\Psi}, \end{aligned} \tag{3.13}$$

$$\left\{ \sum_{n=2}^{\infty} [\Psi(n)]^{1-p} \left[ \int_{\frac{1}{\alpha}}^{\infty} \frac{F(x) dx}{\ln^\lambda(\alpha \beta n x)} \right]^p \right\}^{\frac{1}{p}} < B(\sigma, \mu) \|F\|_{p,\Phi}, \tag{3.14}$$

$$\left\{ \int_{\frac{1}{\alpha}}^{\infty} [\Phi(x)]^{1-q} \left[ \sum_{n=2}^{\infty} \frac{a_n}{\ln^\lambda(\alpha \beta n x)} \right]^q dx \right\}^{\frac{1}{q}} < B(\sigma, \mu) \|a\|_{q,\Psi}. \tag{3.15}$$

**Competing interests**

The authors declare that they have no competing interests.



**Authors' contributions**

BY carried out the mathematical studies, participated in the sequence alignment and drafted the manuscript. QH participated in the design of the study and performed the numerical analysis. All authors read and approved the final manuscript.

**Acknowledgements**

The authors wish to express their thanks to the referees for their careful reading of the manuscript and for their valuable suggestions. This work is supported by the National Natural Science Foundation (No. 61370186), and 2013 Knowledge Construction Special Foundation Item of Guangdong Institution of Higher Learning College and University (No. 2013KJCX0140).

Received: 18 March 2015 Accepted: 10 July 2015 Published online: 30 July 2015

**References**

- Hardy, GH, Littlewood, JE, Pólya, G: *Inequalities*. Cambridge University Press, Cambridge (1934)
- Mitrinović, DS, Pečarić, JE, Fink, AM: *Inequalities Involving Functions and Their Integrals and Derivatives*. Kluwer Academic, Boston (1991)
- Yang, BC: *The Norm of Operator and Hilbert-Type Inequalities*. Science Press, Beijing (2009)
- Yang, BC: An extension of Mulholland's inequality. *Jordan J. Math. Stat.* **3**(3), 151-157 (2010)
- Yang, BC: On Hilbert's integral inequality. *J. Math. Anal. Appl.* **220**, 778-785 (1998)
- Yang, BC, Brnetić, I, Krnić, M, Pečarić, J: Generalization of Hilbert and Hardy-Hilbert integral inequalities. *Math. Inequal. Appl.* **8**(2), 259-272 (2005)
- Krnić, M, Pečarić, J: Hilbert's inequalities and their reverses. *Publ. Math. (Debr.)* **67**(3-4), 315-331 (2005)
- Jin, JJ, Debnath, L: On a Hilbert-type linear series operator and its applications. *J. Math. Anal. Appl.* **371**, 691-704 (2010)
- Azar, L: On some extensions of Hardy-Hilbert's inequality and applications. *J. Inequal. Appl.* **2009**, 546829 (2009)
- Arpad, B, Choonghong, O: Best constant for certain multi-linear integral operator. *J. Inequal. Appl.* **2006**, 28582 (2006)
- Kuang, JC, Debnath, L: On Hilbert's type inequalities on the weighted Orlicz spaces. *Pac. J. Appl. Math.* **1**(1), 95-103 (2007)
- Zhong, WY: The Hilbert-type integral inequality with a homogeneous kernel of Lambda-degree. *J. Inequal. Appl.* **2008**, 917392 (2008)
- Li, YJ, He, B: On inequalities of Hilbert's type. *Bull. Aust. Math. Soc.* **76**(1), 1-13 (2007)
- Yang, BC: A mixed Hilbert-type inequality with a best constant factor. *Int. J. Pure Appl. Math.* **20**(3), 319-328 (2005)
- Wang, AZ, Yang, BC: A more accurate reverse half-discrete Hilbert-type inequality. *J. Inequal. Appl.* **2015**, 85 (2015)
- Yang, BC: A half-discrete Hilbert's inequality. *J. Guangdong Univ. Educ.* **31**(3), 1-7 (2011)
- Kuang, JC: *Applied Inequalities*. Shangdong Science Technic Press, Jinan (2004)
- Kuang, JC: *Introduction to Real Analysis*. Hunan Education Press, Chansha (1996)

Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

---

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)

---