

This is the author's final, peer-reviewed manuscript as accepted for publication. The publisher-formatted version may be available through the publisher's web site or your institution's library.

D-lactic acid biosynthesis from biomass-derived sugars via *Lactobacillus delbrueckii* fermentation

Yixing Zhang, Praveen V. Vadlani

How to cite this manuscript

If you make reference to this version of the manuscript, use the following information:

Zhang, Y., & Vadlani, P. V. (2013). D-lactic acid biosynthesis from biomass-derived sugars via *Lactobacillus delbrueckii* fermentation. Retrieved from <http://krex.ksu.edu>

Published Version Information

Citation: Zhang, Y., & Vadlani, P. V. (2013). D-lactic acid biosynthesis from biomass-derived sugars via *Lactobacillus delbrueckii* fermentation. *Bioprocess and Biosystems Engineering*, 36(12), 1897-1904.

Copyright: © Springer-Verlag Berlin Heidelberg 2013

Digital Object Identifier (DOI): doi:10.1007/s00449-013-0965-8

Publisher's Link: <http://link.springer.com/article/10.1007/s00449-013-0965-8>

This item was retrieved from the K-State Research Exchange (K-REx), the institutional repository of Kansas State University. K-REx is available at <http://krex.ksu.edu>

1 **D-lactic acid biosynthesis from biomass-derived sugars**
2 ***via Lactobacillus delbrueckii* fermentation**

3 Yixing Zhang · Praveen V. Vadlani *

4 Bioprocessing and Renewable Energy Laboratory, Department of Grain Science and
5 Industry, Kansas State University, Manhattan, Kansas. USA

6 *Corresponding Author, Tel: +1-785-532-5012; Fax: +1-785-532-7193; e-mail:

7 vadlani@ksu.edu

8

1 **Abstract** Poly-lactic acid (PLA) derived from renewable resources is considered to be a
2 good substitute for petroleum-based plastics. The number of poly L-lactic acid
3 applications is increased by the introduction of a stereocomplex PLA, which consists of
4 both poly-L and D-lactic acid and has a higher melting temperature. To date, several
5 studies have explored the production of L-lactic acid, but information on biosynthesis of
6 D-lactic acid is limited. Pulp and corn stover are abundant, renewable lignocellulosic
7 materials that can be hydrolyzed to sugars and used in biosynthesis of D-lactic acid. In
8 our study, saccharification of pulp and corn stover was done by cellulase CTec2 and
9 sugars generated from hydrolysis were converted to D-lactic acid by a homofermentative
10 strain, *L. delbrueckii*, through a sequential hydrolysis and fermentation process (SHF)
11 and a simultaneous saccharification and fermentation process (SSF). 36.3 g L⁻¹ of D-
12 lactic acid with 99.8% optical purity was obtained in the batch fermentation of pulp and
13 attained highest yield and productivity of 0.83 g g⁻¹ and 1.01 g L⁻¹ h⁻¹, respectively.
14 Luedeking-Piret model described the mixed growth-associated production of D-lactic
15 acid with a maximum specific growth rate 0.2 h⁻¹ and product formation rate 0.026 h⁻¹
16 ¹, obtained for this strain. The efficient synthesis of D-lactic acid having high optical
17 purity and melting point will lead to unique stereo-complex PLA with innovative
18 applications in polymer industry.

19 **Keywords** D-lactic acid, fermentation, corn stover, pulp, biosynthesis

20 **List of symbols**

21 μ_{\max} Maximum specific growth rate (h⁻¹)

22 C_0 Initial glucose concentration (g L⁻¹)

23 C_p Product concentration (g L⁻¹)

- 1 Y_{PS} Product yield (g lactic acid g^{-1} glucose)
- 2 Y'_{PS} Product overall yield (g lactic acid g^{-1} biomass)
- 3 Y_{XS} Yield of cell dry mass from substrate (g cell dry mass g^{-1} glucose)
- 4 Y_{PX} Yield of product from cell dry mass (g D-lactic acid g^{-1} cell dry mass)
- 5 q_{PS} Product formation rate (h^{-1}) calculated based on the equation $q_{PS} = \frac{1}{S} \times \frac{dP}{dt}$
- 6 Q_p Productivity (g $L^{-1} h^{-1}$)
- 7

1 **Introduction**

2 Lignocellulosic biomass is gaining importance as a potential source of renewable energy
3 and chemicals as the fossil fuel reserves are eventually getting depleted. Demand
4 continues to increase for production of high-value chemicals and materials from
5 renewable resources to attain domestic self-sufficiency and enhanced national security.
6 Lactic acid is an important and multifunctional organic acid that has wide applications in
7 the food, pharmaceutical, and chemical industries [1, 2]. It exists in two optical isomeric
8 forms, L (+) and D (-) poly-lactic acid (PLA), which are being developed as a substitute
9 for petroleum-derived plastics. The high chemical resistance of poly lactic acid is
10 advantageous in the manufacture of fibers, nonwoven fabrics, and films [3]; however, the
11 application of poly L-lactic acid (PLLA) is limited by its melting point [4]. This problem
12 can be obviated by blending it with poly D-lactic acid (PDLA). The melting point of the
13 resulting stereocomplex polymer is approximately 50 °C higher than that of the
14 respective single polymers [5]. The optical purity of lactic acid accentuates the physical
15 properties of poly D-lactic acid-based polymers [6]. The chemical process of making
16 lactic acids usually yields a mixture of these two enantiomers, which is an undesirable
17 feature; therefore, the biological process of making pure lactic acid is preferred [7].
18 To date, intense studies have been conducted on the production of L-lactic acid from
19 different biomass through microbial fermentation [8-11], but information on biosynthesis
20 of D-lactic acid from biomass is limited. A few wild-type strains such as *Lactobacillus*
21 *delbrueckii* subsp. *delbrueckii*, *Sporolactobacillus inulinus* [12], *Lactobacillus*
22 *coryniformis* subsp. *torquens* [13], and *Lactobacillus delbrueckii* subsp. *lactis* QU41 [14]

1 have been identified as D-lactic acid producers. Traditional production of lactic acids
2 typically uses starch derived from food crops as the fermentation substrate [12, 15], but
3 this process may affect the global food supply. Lignocellulosic materials are favorably
4 structured to produce lactic acids, which require the breakdown of cellulose to sugars
5 [16]. This step usually can be done by acid hydrolysis and enzymatic hydrolysis. The
6 enzymatic hydrolysis method is preferred, because it can be done under mild reaction
7 conditions avoiding the use of toxic and corrosive chemicals [17]. The hydrolysis and
8 fermentation steps can be done sequentially (SHF) or simultaneously (SSF). The SSF
9 process offers better yields because it avoids product inhibition and results in higher
10 productivity [18, 19].

11 Production of D-lactic acid from cardboard [20, 21], cellulose [13], peanut meal [22], and
12 rice bran [3] has been studied. Other sources include pulp and corn stover, which have
13 the potential to become cheap and abundant sources for production of ethanol, organic
14 acids, and other chemicals [7, 21]. Pulp is prepared by chemically or mechanically
15 separating cellulose fibers from wood, fiber crops, or waste paper [23]. Corn stover,
16 which includes the leaves, stalks, and cobs of corn plant, is the most abundant
17 agricultural residue in the U.S. [24]; to the best of our knowledge, no research has been
18 reported on D-lactic acid fermentation via pulp and corn stover as substrates.

19 The purpose of this study was to produce D-lactic acid with high yield and optical purity
20 from pulp and corn stover by *Lactobacillus delbrueckii* ATCC 9649. *L. delbrueckii* is a
21 homofermentative lactic acid bacterium that can provide a continuous bioprocess with
22 high volumetric productivity and optically high purity of D-lactic acid under anaerobic

1 conditions [25]. In addition, kinetic analyses of enzyme hydrolysis and fermentation of
2 glucose to D- lactic acid also have been studied in this work.

3 **Materials and methods**

4 Raw materials and chemical treatment

5 Regular pulp and mechanically modified pulp were obtained from the MeadWestvaco's
6 Crompton mill. Corn stover was obtained from fields in Manhattan and Tribune, Kansas.
7 Alkali treatment was performed on corn stover before hydrolysis. Corn stover was
8 suspended in 20 g L⁻¹ NaOH and heated at 121 °C for 30 min in an autoclave (Tomy SS-
9 325E, Tomy SEKO CO., LTD, Tokyo, Japan), then washed under running distilled water
10 and filtered through muslin cloth until no color was visible in the wash water. The alkali-
11 treated corn stover was dried at 80 °C for 24 h and ground to fine particle size in a
12 laboratory mill (3303, Perten Instruments, Springfield, IL) for further enzymatic
13 hydrolysis.

14 Enzyme hydrolysis

15 CTec2 (cellulase) obtained from Novozymes Inc. (Franklinton, NC) was used in this
16 experiment. Enzyme hydrolysis assays were carried out at 45 °C in 250 mL screw capped
17 plastic conical flasks with orbital agitation (150 rpm). The substrate concentration was 2%
18 (w/v). pH was kept at 4.8 using 0.05 mol L⁻¹ citric acid-sodium citrate buffer. The
19 cellulase activity of CTec2 was measured by the filter paper assay [26], and the activity
20 was expressed in terms of filter paper units (FPU). CTec2 was added on a dosage of 2, 4,

1 and 8 FPU g⁻¹ of dry biomass, respectively. Product yield is based on the amount of
2 glucose released divided by the amount of biomass consumed.

3 Microorganism and culture conditions

4 *Lactobacillus delbrueckii* ATCC 9649 obtained from the American Type Culture
5 Collection (Manassas, VA) was used in this work. *L. delbrueckii* inoculum was prepared
6 by growing cells in a 100 mL Wheaton serum bottle containing 50 mL of liquid MRS
7 medium (MRS broth, Difco Laboratories, Detroit, MI) and incubated at 37 °C in a
8 temperature-controlled shaker (Innova 4300, New Brunswick scientific, NJ) at 120 rpm
9 for 15 h. CO₂ (3 vvm) was sparged into the bottle to create anaerobic growing conditions.

10 Sequential hydrolysis and fermentation (SHF)

11 Shake flask fermentation was modified according to the procedure described by
12 Mukhopadhyay [27]. Fermentation was performed in 100 mL Wheaton serum bottles
13 containing 50 mL of synthetic medium, pulp, modified pulp, or corn stover hydrolyzate,
14 and lasted for 30 h. The synthetic medium consisted of 10 g L⁻¹ of glucose, 10 g L⁻¹ of
15 peptone, 5 g L⁻¹ of yeast extract, 2 g L⁻¹ of ammonium citrate, 2 g L⁻¹ of sodium acetate, 2
16 g L⁻¹ of ammonium citrate, 2 g L⁻¹ of K₂HPO₄, 0.1 g L⁻¹ of MgSO₄·7H₂O, 0.05 g L⁻¹ of
17 MnSO₄·4H₂O, and 1 g L⁻¹ of Tween 80. Pulp, modified pulp, and corn stover hydrolyzate
18 were supplemented with all the components (except glucose) of the synthetic medium.
19 pH of the media was adjusted to 6.5 by 10 mol L⁻¹ NaOH, and 3% (w/v) of calcium
20 carbonate was added to control the pH. Temperature was maintained at 37 °C, and
21 agitation was 120 rpm. Batch and fed-batch fermentation were performed in a 7 L
22 fermenter with a working volume of 5 L (Bioflo 110, New Brunswick Scientific Inc.

1 Enfield, CT). In the batch fermentation experiment, paper pulp was added in quantity
2 (270 g) that would possibly achieve a glucose concentration of 40 g L⁻¹ in the medium.
3 After hydrolysis, the pulp hydrolyzate was supplemented with all the components (except
4 glucose) of the synthetic medium. The synthetic medium was used in fed-batch
5 fermentation as a control. After 36 h, 1 L of fermentation medium was taken out and 1 L
6 of feeding medium, which consisted of 40 g L⁻¹ of glucose, 2 g L⁻¹ of ammonium citrate,
7 2 g L⁻¹ of sodium acetate, 2 g L⁻¹ of ammonium citrate, 2 g L⁻¹ of K₂HPO₄, 0.1 g L⁻¹ of
8 MgSO₄·7H₂O, and 0.05 g L⁻¹ of MnSO₄·4H₂O, was added. During the fermentation, the
9 temperature was maintained at 37 °C; agitation speed at 100 rpm; and pH at 6.5. CO₂ was
10 sparged at 3 vvm through the vessel to maintain anaerobic conditions.

11 Simultaneous saccharification and fermentation (SSF)

12 SSF process was modified according to the procedure described by Mukhopadhyay
13 [27]The optimal temperature and pH for the enzymatic hydrolysis and the bacterial
14 fermentation are different; In SSF, temperature was at 40 °C and pH was at 5.5, which
15 were conducive for both enzymatic hydrolysis and bacterial activity. 2 g of dried pulp
16 and corn stover was suspended in 50 ml 0.05 mol L⁻¹ sodium citrate buffer (pH 5.5) with
17 all the components (except glucose) of the synthetic medium. 3% (w/v)calcium carbonate
18 was added to control the pH. CTec2 was added at 8 FPU g⁻¹ of biomass, and
19 *L.delbrueckii* was inoculated at 5% (v/v), and agitation rate was 150 rpm.

1 Analyses

2 Fermentation samples were centrifuged at 15,000×g for 10 min in an Eppendorf
3 centrifuge (5415R, Eppendorf, Hauppauge, NY). The supernatant was collected in
4 sample vials and stored at -4 °C for product and residue glucose analyses.

5 Sugars were quantified by a binary HPLC system (Shimadzu Scientific Instruments,
6 Columbia, MD) equipped with a refractive Index detector (RID-10A) and phenomenex
7 RPM monosaccharide column (300×7.8 mm, Phenomenex, Torrance, CA). Deionised
8 water was used as the mobile phase at a flow rate of 0.6 mL min⁻¹. The oven (Prominence
9 CTD-20A) temperature was maintained at 80 °C.

10 Lactic acids were quantified by a Chirex Chiral column (150×4.6 mm, Phenomenex,
11 Torrance, CA) with isocratic 1 mmol L⁻¹ copper (II) sulfate mobile phase at 1 mL min⁻¹.

12 Peaks were monitored using a UV detector at 254 nm (Shimadzu, PDA).

13 **Results and discussion**

14 Enzymatic hydrolysis

15 Experiments with different loads of cellulase were performed to determine a suitable
16 enzyme loading for enzymatic hydrolysis of pulp, modified pulp, and alkali-treated corn
17 stover. The maximum reaction rate (v_{\max}) was calculated from the Michaelis-Menten

18 equation ($v = \frac{v_{\max}[S]}{K_m + [S]}$). v_{\max} increased almost linearly with the increase of enzyme

19 concentration in all three biomass cases (Fig. 1). The hydrolysis rate of corn stover and
20 modified pulp was about to reach a plateau when the enzyme loading increased, perhaps
21 due to substrate saturation [28]. Increased enzyme loading from 2 to 8 FPU g⁻¹ of

1 substrate increased glucose yield by 24% after 48 h of pulp saccharification (Fig.2a);
2 however, increasing the enzyme dosage did not significantly change the final glucose
3 yield in the saccharification of mechanically modified pulp (12%) (Fig. 2b) and alkali
4 treated corn stover (11%) (Fig. 2c). The highest glucose yield was observed at 24 h for
5 mechanically modified pulp as well as corn stover. The initial saccharification rate of
6 mechanically modified pulp and corn stover was higher than that of pulp. Mechanically
7 modified pulp had finer fiber size, which made it much easier for the enzymes to break
8 down. Alkali treatment caused the cellulose in corn stover to swell, which led to an
9 increase in the internal surface area and a decrease in the degree of crystallinity of
10 cellulose [29], therefore making cellulose in alkali-treated corn stover much easier for the
11 enzyme to access.

12 Production of D-lactic acid by SHF

13 The purpose of this portion of the study was to produce D-lactic acid by *L. delbrueckii*
14 using sugars derived from biomass as a cheap carbon source. We also tested another
15 strain *Sporolactobacillus inulinus* ATCC 15538. Unlike in the results obtained by
16 Fukushima *et al.* [12], *S. inulinus* produced L-lactic acid instead of D-lactic acid in our
17 experiments. This result may be due to the difference in strain or the possible alternation
18 of bacterial character after receiving it.

19 In shake flask fermentation, the amount of pulp (1 g), mechanically modified pulp (1.3 g),
20 and corn stover (1.2 g) was set up to obtain 10 g L⁻¹ glucose after enzymatic hydrolysis.
21 No residual glucose was observed after 30 h fermentation, and the final pH of the
22 medium was between 5 to 5.5. The optical purity of D-lactic acid was 99.9%. These

1 results were in close agreement with Demirci and Pometto [30]. The highest yield of D-
2 lactic acid was observed in corn stover hydrolyzate (Table 1). Besides glucose, 5.6 g L⁻¹
3 xylose and 1.7 g L⁻¹ arabinose were also present in the corn stover hydrolyzate; however,
4 xylose remained unused, and arabinose was below detectable levels at the end of
5 fermentation. *L.delbrueckii* cannot use xylose due to the lack of xylose isomerase and
6 xylulokinase, two key enzymes in xylose assimilation [31].

7 In fed-batch fermentation, almost all glucose was consumed within the first 36 h (first
8 stage). In the second stage, feeding medium was added, and fermentation was completed
9 within 80 h. The Luedeking-Piret equation ($\frac{1}{X} \frac{dP}{dt} = \alpha \frac{1}{X} \frac{dX}{dt} + \beta$) was used to describe
10 the D-lactic acid production from synthetic medium in the first stage. Growth-associated
11 constant (α) and non-growth associated constant (β) can be calculated from the graph of
12 the specific production rate (q_p) versus the specific growth rate (μ); the correlation
13 coefficient (R^2) was 0.88 (Fig. 3). Compared with other strains listed in Table 2, in our
14 study *L. delbrueckii* had lower μ_{\max} and higher α values. Lower μ_{\max} suggests lower
15 growth efficiency, and a high α value indicates a higher contribution of the cell growth to
16 D-lactic acid production [32]. The value of α multiplied by μ_{\max} was 1.56, which was
17 larger than the β value, indicating that the specific growth rate played an important role in
18 specific D-lactic acid production.

19 Figures 4 and 5 show the fermentation profile of the synthetic medium and pulp
20 hydrolyzate, respectively. Table 3 summarizes the results of the first stage of fed-batch
21 fermentation and batch fermentation. 37.4 g L⁻¹ of D-lactic acid was obtained by the end
22 of first-stage fermentation, and the product yield and productivity obtained were 0.93 g g⁻¹

1 1 and $1.04 \text{ g L}^{-1} \text{ h}^{-1}$, respectively. These results were in agreement with other studies in
2 literature [8, 33]. Fed-batch fermentation was completed within 80 h; at the end of
3 fermentation, about 5.5 g L^{-1} glucose was left and up to 57.3 g L^{-1} D-lactic acid with
4 optical purity of 99.8% was accumulated, which led to a productivity of $0.72 \text{ g L}^{-1} \text{ h}^{-1}$.
5 After pulp hydrolysis, the glucose concentration was 50 g L^{-1} and was used in the batch
6 fermentation. After 30 h, glucose was hardly consumed, and even if we extended the
7 fermentation time to 36 h, 6.2 g L^{-1} residual glucose remained. At the end of fermentation,
8 36.3 g L^{-1} lactic acid was produced, the yield of D-lactic acid was calculated by the
9 amount of D-lactic acid produced divided by the amount of glucose consumed, which
10 was 0.83 g g^{-1} , and productivity was $1.01 \text{ g L}^{-1} \text{ h}^{-1}$. In a similar study undertaken in our
11 laboratory, L-Lactic acid was synthesized from cheese whey and a yield (0.98 g g^{-1}) and
12 productivity ($1.14 \text{ g L}^{-1} \text{ h}^{-1}$) was obtained [8]. The product formation rate of batch
13 fermentation of pulp hydrolyzate was quite close to the product formation rate of first-
14 stage fed-batch fermentation using the synthetic medium. The yield of D-lactic acid (0.83
15 g g^{-1}) from pulp hydrolyzate was lower than the first-stage yield (0.93 g g^{-1}) from
16 synthetic medium. The reason might be due to substrate inhibition; therefore, the SSF
17 process was preferred in subsequent experiments.

18 Production of D-lactic acid by SSF

19 After demonstrating the feasibility of producing D-lactic acid from biomass hydrolyzate
20 in the batch process, SSF was carried out using pulp and corn stover in a shake flask. In
21 SSF, samples were collected after 4 h of incubation; the profiles obtained for corn stover
22 and pulp SSF experiments are shown in Figure 6.. In SSF, cellulose hydrolysis and

1 glucose assimilation were combined into a single fermentation process [34]. During the
2 first 8 h, bacteria were in low activity and glucose accumulated to around 8 g L⁻¹ and 14 g
3 L⁻¹ in the case of pulp and corn stover, respectively. After the first 8 h cultivation,
4 glucose concentration was kept low, which indicated that bacterial cells were
5 metabolically active during the entire course of the fermentation and also meant that
6 enzymatic hydrolysis of cellulose was the rate limiting step for D-lactic acid production
7 as already observed by other groups [35, 36]. Xylose accumulated and remained nearly
8 constant throughout the process. It was impossible to know the exact amount of glucose
9 consumed in the SSF process; therefore, in order to compare SSF and SHF, results were
10 expressed as an overall yield (the amount of D-lactic acid produced divided by the
11 amount of biomass used)(Table 1). The highest D-lactic acid overall yield was 0.48 and
12 0.38 g g⁻¹ of pulp in SSF and SHF, respectively. For corn stover, the maximum D-lactic
13 acid overall yield was 0.58 and 0.41 g g⁻¹ in SSF and SHF, respectively, demonstrating
14 that the SSF process was more efficient than the SHF process. The reason for the higher
15 overall yield in SSF may be that glucose released during the hydrolysis step was rapidly
16 consumed as substrate during the fermentation step, therefore reducing the end-product
17 inhibition of hydrolysis [37].

18 **Conclusions**

19 In this study, we demonstrated efficient D-lactic acid production with high optical purity
20 from pulp, modified pulp, and corn stover by *L. delbrueckii* ATCC 9649. Enzymatic
21 hydrolysis of biomass was achieved effectively by CTec2 enzyme system. D-lactic acid
22 productivity was not only high, but also cost-effective because pulp and modified pulp

1 need no pretreatment. The SSF process demonstrated the advantages of avoiding
2 substrate inhibition and increasing the productivity and yield of D-lactic acid. The yield
3 obtained in the present study would have been even higher if xylose from corn stover
4 hydrolyzate could be completely used by the microorganism. Future study should be
5 directed toward complete use of the available carbohydrate for efficient D-lactic acid
6 production.

7 **Acknowledgements**

8 This work was funded by the Consortium for Plant Biotechnology Research and
9 supported by the Department of Grain Science and Industry at Kansas State University.
10 The authors are grateful to Novozymes Inc. for the donation of enzymes. This is
11 contribution number 13-180-J from the Kansas Agricultural Experiment Station.

12 13 **References**

- 14 1. Shen X, Xia L (2006) Lactic acid production from cellulosic material by synergetic
15 hydrolysis and fermentation. *Appl Biochem Biotechnol* 133:251-262
- 16 2. Datta R, Tsai S, Bonsignore P, Moon S, Frank J (1995) Technological and Economic-
17 Potential of Poly (lactic Acid) and Lactic-Acid Derivatives. *FEMS Microbiol Rev*
18 16:221-231
- 19 3. Tanaka T, Hoshina M, Tanabe S, Sakai K, Ohtsubo S, Taniguchi M (2006) Production
20 of D-lactic acid from defatted rice bran by simultaneous saccharification and
21 fermentation. *Bioresour Technol* 97:211-217

- 1 4. Brizzolara D, Cantow H, Diederichs K, Keller E, Domb A (1996) Mechanism of the
2 stereocomplex formation between enantiomeric poly(lactide)s. *Macromolecules* 29:191-
3 197
- 4 5. Ikada Y, Jamshidi K, Tsuji H, Hyon S (1987) Stereocomplex formation between
5 enantiomeric poly (lactides). *Macromolecules* 20:904-906
- 6 6. Tsuji F (2002) Autocatalytic hydrolysis of amorphous-made polylactides: effects of L-
7 lactide content, tacticity, and enantiomeric polymer blending. *Polymer* 43:1789-1796
- 8 7. Yadav AK, Chaudhari AB, Kothari RM (2011) Bioconversion of renewable resources
9 into lactic acid: an industrial view. *Crit Rev Biotechnol* 31:1-19
- 10 8. Vadlani PV, Mathews AP, Karr GS (2008) Low-cost propionate salt as road deicer:
11 evaluation of cheese whey and other media components. *World J Microbiol Biotechnol*
12 24:825-832
- 13 9. Vadlani PV, Matthews AP, Karr GS (2008) A two-stage fermentation process:
14 production of propionate and acetate salt as road deicer from cheese whey. *Biological*
15 *Engineering* 1:95-104
- 16 10. Moon SK, Lee J, Song H, Cho JH, Choi GW, Seung D (2012) Characterization of
17 ethanol fermentation waste and its application to lactic acid production by *Lactobacillus*
18 *paracasei*. *Bioprocess Biosyst Eng*. Doi: 10.1007/s00449-012-0810-5
- 19 11. Phruksawan P, Kulpreecha S, Sooksai S, Thongchul N (2012) Direct fermentation of
20 L (+) -lactic acid from cassava pulp by solid state culture of *Rhizopus oryzae*. *Bioprocess*
21 *and Biosyst Eng*. 35:1429-1436
- 22 12. Fukushima K, Sogo K, Miura S, Kimura Y (2004) Production of D-lactic acid by
23 bacterial fermentation of rice starch. *Macromolecular Bioscience* 4:1021-1027

- 1 13. Yanez R, Moldes AB, Alonso JL, Parajo JC (2003) Production of D-lactic acid from
2 cellulose by simultaneous saccharification and fermentation using *Lactobacillus*
3 *coryniformis* subsp *torquens*. Biotechnol Lett 25:1161-1164
- 4 14. Tashiro Y, Kaneko W, Sun Y, Shibata K, Inokuma K, Zendo T, Sonomoto K (2011)
5 Continuous D-lactic acid production by a novel thermotolerant *Lactobacillus delbrueckii*
6 subsp *lactis* QU 41. Appl Microbiol Biotechnol 89:1741-1750
- 7 15. Shinkawa S, Okano K, Tanaka T, Ogino C, Kondo A (2009) Efficient D-lactic acid
8 production from raw starch. J Biosci Bioeng 108:S47-S48
- 9 16. Schmidt S, Padukone N (1997) Production of lactic acid from wastepaper as a
10 cellulosic feedstock. J Ind Microbiol Biotechnol 18:10-14
- 11 17. Xu Z, Wang Q, Jiang Z, Yang X, Ji Y (2007) Enzymatic hydrolysis of pretreated
12 soybean straw. Biomass & Bioenergy 31:162-167
- 13 18. Marques S, Santos JAL, Girio FM, Roseiro JC (2008) Lactic acid production from
14 recycled paper sludge by simultaneous saccharification and fermentation. Biochem Eng J
15 41:210-216
- 16 19. Kim K, Kim W, Seo D, Yoo I, Kim E, Yoon H (2003) Production of lactic acid from
17 food wastes. Appl Biochem Biotechnol 105:637-647
- 18 20. Yanez R, Alonso JL, Parajo JC (2005) Study on the suitability of untreated
19 corrugated cardboard for D-lactic acid production by SSF using *Lactobacillus*
20 *coryniformis* subsp *torquens*. Afinidad 62:295-301
- 21 21. Yanez R, Alonso JL, Parajo JC (2005) D-Lactic acid production from waste
22 cardboard. J Chem Technol Biotechnol 80:76-84

- 1 22. Wang L, Zhao B, Li F, Xu K, Ma C, Tao F, Li Q, Xu P (2011) Highly efficient
2 production of D-lactate by *Sporolactobacillus* sp. CASD with simultaneous enzymatic
3 hydrolysis of peanut meal. *Appl Microbiol Biotechnol* 89:1009-1017.
- 4 23. Biermann CJ (1996) *Handbook of pulping and papermaking*. Elsevier Science, San
5 Diego.
- 6 24. Li Y, Ruan R, Chen PL, Liu Z, Pan X, Lin X, Liu Y, Mok CK, Yang T (2004)
7 Enzymatic hydrolysis of corn stover pretreated by combined dilute alkaline treatment and
8 homogenization. *Trans ASAE* 47:821-825
- 9 25. Calabia BP, Tokiwa Y (2007) Production of D-lactic acid from sugarcane molasses,
10 sugarcane juice and sugar beet juice by *Lactobacillus delbrueckii*. *Biotechnol Lett*
11 29:1329-1332
- 12 26. Ghose T (1987) Measurement of cellulase activities. *Pure Appl Chem* 59:257-268
- 13 27. Mukhopadhyay A, Bioconversion of paper mill lignocellulosic materials to lactic acid
14 using cellulase enzyme complex and microbial cultures. [MS Thesis]. Manhattan,
15 KS: Kansas State University; 2009. Available from: K-State Research Exchange
- 16 28. Lee Y, Fan L (1982) Kinetic-studies of enzymatic-hydrolysis of insoluble cellulose -
17 analysis of the initial rates. *Biotechnol Bioeng* 24:2383-240624
- 18 29. Chandra RP, Au-Yeung K, Chanis C, Roos AA, Mabee W, Chung PA, Ghatora S,
19 Saddler JN (2011) The influence of pretreatment and enzyme loading on the effectiveness
20 of batch and fed-batch hydrolysis of corn stover. *Biotechnol Prog* 27:77-85
- 21 30. Demirci A, Pometto AL (1992) Enhanced production of D-lactic acid by mutants of
22 *Lactobacillus delbrueckii* ATCC 9649. *J Ind Microbiol* 11:23-28

- 1 31. Okano K, Yoshida S, Yamada R, Tanaka T, Ogino C, Fukuda H, Kondo A (2009)
2 Improved production of homo-D-Lactic acid via xylose fermentation by introduction of
3 xylose assimilation genes and redirection of the phosphoketolase pathway to the pentose
4 phosphate pathway in L-Lactate dehydrogenase gene-deficient *Lactobacillus plantarum*.
5 Appl Environ Microbiol 75:7858-7861
- 6 32. Zhao B, Wang L, Li F, Hua D, Ma C, Ma Y, Xu P (2010) Kinetics of D-lactic acid
7 production by *Sporolactobacillus* sp strain CASD using repeated batch fermentation.
8 Bioresour Technol 101:6499-6505
- 9 33. Garde A, Jonsson G, Schmidt AS, Ahring BK (2002) Lactic acid production from
10 wheat straw hemicellulose hydrolysate by *Lactobacillus pentosus* and *Lactobacillus*
11 *brevis*. Bioresour Technol 81(3):217-223
- 12 34. Patel M, Ou M, Harbrucker R, Aldrich H, Buszko M, Ingram L, Shanmugam KT
13 (2006) Isolation and characterization of acid-tolerant, thermophilic bacteria for effective
14 fermentation of biomass-derived sugars to lactic acid. Appl Environ Microbiol 72:3228-
15 3235
- 16 35. Parajo J, Alonso J, Moldes A (1997) Production of lactic acid from lignocellulose in a
17 single stage of hydrolysis and fermentation. Food Biotechnol 11:45-58
- 18 36. Nakasaki K, Adachi T (2003) Effects of intermittent addition of cellulase for
19 production of L-lactic acid from waste water sludge by simultaneous saccharification and
20 fermentation. Biotechnol Bioeng 82:263-270
- 21 37. Akerberg C, Zacchi G (2000) An economic evaluation of the fermentative production
22 of lactic acid from wheat flour. Bioresour Technol 75:119-126

- 1 38. Boonmee M, Leksawasdi N, Bridge W, Rogers P (2003) Batch and continuous
2 culture of *Lactococcus lactis* NZ133: experimental data and model development.
3 Biochem Eng J 14:127-135
- 4 39. Nandasana AD, Kumar S (2008) Kinetic modeling of lactic acid production from
5 molasses using *Enterococcus faecalis* RKY1. Biochem Eng J 15:277-284
- 6 40. Amrane A (2005) Analysis of the kinetics of growth and lactic acid production for
7 *Lactobacillus helveticus* growing on supplemented whey permeate. J Chem Technol
8 Biotechnol 80:345-352
- 9

1 **Table 1** D-lactic acid production through SHF and SSF process in shake flask

		C_0^a	C_P^b	Y_{PS}^c	Y'_{PS}^d	Q_P^e
SHF	Synthetic medium	10	7.7±0.05	0.77±0.01	---	0.25±0.01
	Pulp	9.7±0.17	7.5±0.47	0.77±0.66	0.38±0.02	0.25±0.03
	Modified pulp	11.2±0.09*	8.5±0.39	0.76±0.03	0.42±0.02	0.28±0.01
	Corn stover	9.9±0.05	8.3 ±0.04	0.83±0.01	0.41±0.01	0.27±0.01
SSF	Pulp	---	19.2±1.63*	---	0.48±0.04*	0.31±0.04
	Corn stover	---	20.1±0.65*	---	0.58±0.03*	0.32±0.07

2 Each mean is based on three replications ($p < 0.05$; REGWQ; one-way ANOVA)

3 ^a Initial glucose of modified pulp hydrolyzate was significantly different

4 ^b D-lactic acid concentration in SSF process was significantly different in SHF process

5 ^c Product yield was not significantly different in SHF process; product yield was calculated by the amount of D-lactic acid produced divided by the amount of glucose consumed.

7 ^d Product overall yield was significantly different between SSF and SHF; product overall yield was calculated by the amount of D-lactic acid produced divided by the amount of biomass used.

9 ^e Productivity was not significantly different.

10

11 **Table 2** Kinetic parameters of different lactic acid bacteria


Microorganism	Substrate	μ_{max}	α	β
<i>L. delbrueckii</i> (this study)	Glucose	0.2	7.8	0.18
<i>L. lactis</i> [38]	Lactose	1.1	0.392	3.02
<i>E. faecalis</i> RKY1 [39]	Molasses	1.6	0.26	---
<i>Lactobacillus helveticus</i> [40]	Whey permeate	0.48	2.33	0.77

12

13 **Table 3** Kinetic parameters of fed-batch and batch fermentation


	C_p	Y_{PS}	Y_{PX}	Y_{XS}	q_{PS}	Q_P
Fed-batch (stage I)	37.4	0.93	10.9	0.086	0.026	1.04
Pulp hydrolyzate batch	36.3	0.83	---	---	0.023	1.01

14




1
2


Fig. 1 Plot of v_{\max} of different biomass versus enzyme concentration



3
4




1
2



3
4
5
6
7


Fig. 2 Enzymatic hydrolysis of pulp (a), mechanically modified pulp (b), and alkali-treated corn stover (c) at varying cellulase levels



1


2 **Fig. 3** Specific production rate versus specific growth rate for *L. delbrueckii* growing on the synthetic
 3 medium

4




5

6 **Fig. 4** Fed-batch fermentation profile of D-lactic acid from the synthetic medium



1


2 **Fig. 5** Batch fermentation profile of D-lactic acid production from pulp hydrolyzate



3

4

(a)



1

(b)

2

3

Fig. 6 Time course of SSF process with *L. delbrueckii* using pulp (a) and alkali-treated corn stover (b)