

Hindawi
Advances in Mathematical Physics
Volume 2018, Article ID 4523512, 6 pages
<https://doi.org/10.1155/2018/4523512>



Research Article

Spacelike Hypersurfaces in Weighted Generalized Robertson-Walker Space-Times

Ximin Liu  and Ning Zhang 

School of Mathematical Sciences, Dalian University of Technology, Dalian 116024, China

Correspondence should be addressed to Ximin Liu; ximinliu@dlut.edu.cn

Received 28 October 2017; Accepted 6 February 2018; Published 5 March 2018

Academic Editor: Remi Léandre

Copyright © 2018 Ximin Liu and Ning Zhang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Applying generalized maximum principle and weak maximum principle, we obtain several uniqueness results for spacelike hypersurfaces immersed in a weighted generalized Robertson-Walker (GRW) space-time under suitable geometric assumptions. Furthermore, we also study the special case when the ambient space is static and provide some results by using Bochner's formula.

1. Introduction

In recent years, spacelike hypersurfaces in Lorentzian manifolds have been deeply studied not only from their mathematical interest, but also from their importance in general relativity.

Particularly, there are many articles that study spacelike hypersurfaces in weighted warped product space-times. A weighted manifold is a Riemannian manifold with a measure that has a smooth positive density with respect to the Riemannian one. More precisely, the weighted manifold M_f associated with a complete n -dimensional Riemannian manifold (M^n, g) and a smooth function f on M^n is the triple $(M^n, g, d\mu = e^{-f} dM)$, where dM stands for the volume element of M^n . In this setting, we will take into account the so-called Bakry-Émery Ricci tensor (see [1]) which as an extension of the standard Ricci tensor Ric , which is defined by

$$\text{Ric}_f = \text{Ric} + \text{Hess } f. \quad (1)$$

Therefore, it is natural to extend some results of the Ricci curvature to analogous results for the Bakry-Émery Ricci tensor. Before giving more details on our work we present a brief outline of some recent results related to our one.

In [2], Wei and Wylie considered the complete n -dimensional weighted Riemannian manifold and proved mean curvature and volume comparison results on the assumption that

the ∞ -Bakry-Émery Ricci tensor is bounded from below and f or $|\nabla f|$ is bounded. Later, Cavalcante et al. [3] researched the Bernstein-type properties concerning complete two-sided hypersurfaces immersed in a weighted warped product space using the appropriated generalized maximum principles. Moreover, [4] obtained new Calabi-Bernstein's type results related to complete spacelike hypersurfaces in a weighted GRW space-time. More recently, some rigidity results of complete spacelike hypersurfaces immersed into a weighted static GRW space-time are given in [5].

In this paper we study spacelike hypersurfaces in a weighted *generalized Robertson-Walker* (GRW) space-times. Moreover, a GRW space-time is a space-time regarding a warped product of a negative definite interval as a base, a Riemannian manifold as a fiber, and a positive smooth function as a warped function. Furthermore, there exists a distinguished family of spacelike hypersurfaces in a GRW space-time, that is, the so-called slices, which are defined as level hypersurfaces of the time coordinate of the space-time. Notice that any slice is totally umbilical and has constant mean curvature.

We have organized this paper as follows. In Section 2, we introduce some basic notions to be used for spacelike hypersurfaces immersed in weighted GRW space-times. In Section 3, we prove some uniqueness results of spacelike hypersurface in a weighted GRW space-time under appropriate conditions on the weighted mean curvature and the

weighted function by using the generalized Omori-Yau maximum principle or the weak maximum principle. Finally, in Section 4, applying the weak maximum principle, we obtain some rigidity results for the special case when the ambient space is static.

2. Preliminaries

Let M^n be a connected n -dimensional oriented Riemannian manifold and I be an open interval in \mathbb{R} endowed with the metric $-dt^2$. We let $\rho : I \rightarrow \mathbb{R}^+$ be a positive smooth function. Denote $-I \times_\rho M^n$ to be the warped product endowed with the Lorentzian metric

$$\langle \cdot, \cdot \rangle = -\pi_I^*(dt^2) + \rho(\pi_I)^2 \pi_M^*(\langle \cdot, \cdot \rangle_M), \quad (2)$$

where π_I and π_M are the projections onto I and M , respectively. This space-time is a *warped product* in the sense of ([6], Chap. 7), with *fiber* $(M, \langle \cdot, \cdot \rangle)$, *base* $(I, -dt^2)$, and *warping function* ρ . Furthermore, for a fixed point $t_0 \in I$, we say that $M_{t_0}^n = -\{t_0\} \times M^n$ is a *slice* of $-I \times_\rho M^n$. Following the terminology used in [7], we will refer to $-I \times_\rho M^n$ as a *generalized Robertson-Walker (GRW) space-time*. Particularly, if the fiber M^n has constant section curvature, it is called a *Robertson-Walker (RW) space-time*.

Recall that a smooth immersion $\psi : \Sigma^n \rightarrow -I \times_\rho M^n$ of an n -dimensional connected manifold Σ^n is called a *spacelike hypersurface* if the induced metric via ψ is a Riemannian metric on Σ^n , which will be also denoted for $\langle \cdot, \cdot \rangle$.

In the following, we will deal with two particular functions naturally attached to spacelike hypersurface Σ^n , namely, the *angle (or support) function* $\Theta = \langle N, \partial_t \rangle$ and the *height function* $h = (\pi_I)|_\Sigma$, where $\partial_t := \partial/\partial t$ is a (unitary) timelike vector field globally defined on \bar{M} and N is a unitary timelike normal vector field globally defined on Σ .

Let $\bar{\nabla}$ and ∇ stand for gradients with respect to the metrics of $-I \times_\rho M^n$ and Σ^n , respectively. By a simple computation, we have

$$\bar{\nabla}\pi_I = -\langle \bar{\nabla}\pi_I, \partial_t \rangle \partial_t = -\partial_t. \quad (3)$$

Therefore, the gradient of h on Σ^n is

$$\nabla h = (\bar{\nabla}\pi_I)^\top = -\partial_t^\top = -\partial_t - \Theta N. \quad (4)$$

Particularly, we have

$$|\nabla h|^2 = -1 + \Theta^2, \quad (5)$$

where $|\cdot|$ denotes the norm of a vector field on Σ^n .

Now, we consider that a GRW space-time $-I \times_\rho M^n$ is endowed with a weighted function f , which will be called a weighted GRW space-time $-I \times_\rho M_f^n$. In this setting, for a spacelike hypersurface Σ^n immersed into $-I \times_\rho M_f^n$, the *f-divergence operator* on Σ^n is defined by

$$\operatorname{div}_f(X) = e^f \operatorname{div}(e^{-f}X), \quad (6)$$

where X is a tangent vector field on Σ^n .

For a smooth function $u : \Sigma^n \rightarrow \mathbb{R}$, we define its *drifting Laplacian* by

$$\Delta_f u = \operatorname{div}_f(\nabla u) = \Delta u - \langle \nabla u, \nabla f \rangle, \quad (7)$$

and we will also denote such an operator as the *f-Laplacian* of Σ^n .

According to Gromov [8], the weighted mean curvature or *f-mean curvature* H_f of Σ^n is given by

$$nH_f = nH - \langle \bar{\nabla}f, N \rangle, \quad (8)$$

where H is the standard mean curvature of hypersurface Σ^n with respect to the Gauss map N .

It follows from a splitting theorem due to Case (see [9] Theorem 1.2) that if a weighted GRW space-time $-I \times_\rho M_f^n$ is endowed with a bounded weighted function f such that $\bar{\operatorname{Ric}}_f(V, V) \geq 0$ for all timelike vector fields V on $-I \times_\rho M_f^n$, then f must be constant along \mathbb{R} . In the same spirit of this result, in the following we will consider weighted GRW space-times $-I \times_\rho M_f^n$ whose weighted function f does not depend on the parameter $t \in I$; that is, $\langle \bar{\nabla}f, \partial_t \rangle = 0$. Moreover, for simplicity, we will refer to them as $\bar{M}^{n+1} := -I \times_\rho M_f^n$.

In the following, we give some technical lemmas that will be essential for the proofs of our main results in weighted GRW space-times $\bar{M}^{n+1} = -I \times_\rho M_f^n$ (for further details on the proof, see Lemma 1 in [4]).

Lemma 1. *Let Σ^n be a spacelike hypersurface immersed in a weighted GRW spacetime $\bar{M}^{n+1} = -I \times_\rho M_f^n$, with height function h . Then,*

$$\Delta_f h = -(\log \rho)'(h)(n + |\nabla h|^2) - nH_f \Theta, \quad (9)$$

$$\begin{aligned} \Delta_f \rho(h) &= -n \frac{\rho'(h)^2}{\rho(h)} + |\nabla h|^2 \rho(h) (\log \rho)''(h) \\ &\quad - n \rho'(h) H_f \Theta. \end{aligned} \quad (10)$$

If we denote \mathcal{L}_f^1 as the space of the integrable functions on Σ^n with respect to the weighted volume element $d\mu = e^{-f} d\Sigma$, using the relation of $\operatorname{div}_f(X) = e^f \operatorname{div}(e^{-f}X)$ and Proposition 2.1 in [10], we can obtain the following extension of a result in [11].

Lemma 2. *Let u be a smooth function on a complete weighted Riemannian manifold Σ^n with weighted function f such that $\Delta_f u$ does not change sign on Σ^n . If $|\nabla u| \in \mathcal{L}_f^1$, then $\Delta_f u$ vanishes identically on Σ^n .*

In the following, we will introduce the weak maximum principle for the drifted Laplacian. By the fact in [12], that is, the Riemannian manifold M satisfies the weak maximum principle if and only if M is stochastically complete, we can have the next lemma which extended a result of [13].

Lemma 3. Let $(M^n, \langle \cdot, \cdot \rangle, e^{-f} dM)$ be an n -dimensional stochastically complete weighted Riemannian manifold and $u : M \rightarrow \mathbb{R}$ be a smooth function which is bounded from below on M^n . Then there is a sequence of points $p_k \in M^n$ such that

$$\begin{aligned} \lim_k u(p_k) &= \inf u, \\ \lim_k \Delta_f u(p_k) &\geq 0. \end{aligned} \quad (11)$$

Equivalently, for any smooth function $u : \Sigma^n \rightarrow \mathbb{R}$ which is bounded from above on M^n , there is a sequence of points $q_k \in M^n$ such that

$$\begin{aligned} \lim_k u(q_k) &= \sup u, \\ \lim_k \Delta_f u(q_k) &\leq 0. \end{aligned} \quad (12)$$

3. Uniqueness Results in Weighted GRW Space-Times

In this section, we will state and prove our main results in weighted GRW space-times $\overline{M}^{n+1} = -I \times_\rho M_f^n$. We point out that, to prove the following results, we do not require that the f -mean curvature H_f of the spacelike hypersurface Σ^n is constant.

Recall that a slab of a weighted GRW spacetime $-I \times_\rho M_f^n$ is a region of the type

$$[t_1, t_2] \times M_f^n = \{(t, p) \in -I \times_\rho M_f^n : t_1 \leq t \leq t_2\}. \quad (13)$$

Theorem 4. Let $\overline{M}^{n+1} = -I \times_\rho M_f^n$ be a weighted GRW spacetime which obeys $(\log \rho)''(h) \leq 0$. Let $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ be a complete spacelike hypersurface that lies in a slab of \overline{M}^{n+1} . If the f -mean curvature H_f satisfies $H_f^2 \leq (1/\Theta^2)(\rho'(h)^2/\rho(h)^2)$ and $|\nabla h| \in \mathcal{L}_f^1(\Sigma)$, then Σ^n is a slice of \overline{M}^{n+1} .

Proof. From (10), we have

$$\begin{aligned} \frac{1}{\rho(h)} \Delta_f \rho(h) &= -n \frac{\rho'(h)^2}{\rho(h)^2} + (\log \rho)''(h) |\nabla h|^2 \\ &\quad - n \frac{\rho'(h)}{\rho(h)} H_f \Theta \\ &\leq -n \frac{\rho'(h)^2}{\rho(h)^2} + (\log \rho)''(h) |\nabla h|^2 \\ &\quad + \frac{n}{2} \left(\frac{\rho'(h)^2}{\rho(h)^2} + H_f^2 \Theta^2 \right) \\ &\leq \frac{n}{2} \left(H_f^2 \Theta^2 - \frac{\rho'(h)^2}{\rho(h)^2} \right) \\ &\quad + (\log \rho)''(h) |\nabla h|^2. \end{aligned} \quad (14)$$

By the hypotheses, we have $\Delta_f \rho(h) \leq 0$. Moreover, since Σ^n lies in a slab, there is a positive constant C such that

$$|\nabla \rho(h)| \leq C \rho'(h) |\nabla h| \in \mathcal{L}_f^1(\Sigma). \quad (15)$$

Therefore, we can apply Lemma 2 to get $\Delta_f \rho(h) = 0$; that is, $\rho(h)$ is constant. Therefore Σ^n is a slice. \square

Theorem 5. Let $\overline{M}^{n+1} = -I \times_\rho M_f^n$ be a weighted GRW spacetime which obeys $\rho''(h) \leq 0$. Let $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ be a complete spacelike hypersurface that lies in a slab of \overline{M}^{n+1} . If the f -mean curvature H_f satisfies $H_f^2 \leq \rho'(h)^2/\rho(h)^2$ and $|\nabla h| \in \mathcal{L}_f^1(\Sigma)$, then Σ^n is a slice of \overline{M}^{n+1} .

Proof. By a similar reasoning as in the proof of Theorem 4, we have

$$\begin{aligned} \frac{1}{\rho(h)} \Delta_f \rho(h) &\leq -n \frac{\rho'(h)^2}{\rho(h)^2} + (\log \rho)''(h) |\nabla h|^2 \\ &\quad + \frac{n}{2} \left(\frac{\rho'(h)^2}{\rho(h)^2} + H_f^2 \Theta^2 \right) \\ &\leq \frac{n}{2} H_f^2 \Theta^2 - \frac{n}{2} \frac{\rho'(h)^2}{\rho(h)^2} + \frac{\rho''(h)}{\rho(h)} |\nabla h|^2 \\ &\leq \frac{n}{2} \left(H_f^2 - \frac{\rho'(h)^2}{\rho(h)^2} \right) \Theta^2 \\ &\quad + \frac{\rho''(h)}{\rho(h)} |\nabla h|^2, \end{aligned} \quad (16)$$

where the last inequality is due to $\Theta^2 \geq 1$.

Taking into account the assumptions, we have $\Delta_f \rho(h) \leq 0$. Now in the same argument as in Theorem 4, we have that Σ^n is a slice. \square

Next, we will use the weak maximum principle to study the rigidity of the spacelike hypersurfaces in weighted GRW space-times.

Theorem 6. Let $\overline{M}^{n+1} = -I \times_\rho M_f^n$ be a weighted GRW spacetime which satisfies $(\log \rho)'' \leq 0$ and there is a point $h_0 \in I$ such that $\rho'(h_0) = 0$. Let $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ be a stochastically complete constant f -mean curvature H_f spacelike hypersurface such that $\sup h \geq h_0$, $\inf h \leq h_0$ which is contained in a slab; then Σ^n is f -maximal. In addition, if Σ^n is complete and $|\nabla h| \in \mathcal{L}_f^1$, then Σ^n is a slice.

Proof. We take the Gauss map N of the hypersurface Σ^n such that $\Theta > 0$; from (7) we have $\Theta \geq 1$.

By Lemma 3, the weak maximum principle for the drifted Laplacian holds on Σ^n ; then there exist two sequences $\{p_j\}, \{q_j\} \subset \Sigma^n$ such that

$$\lim_j h(p_j) = \inf h,$$

$$\begin{aligned}
\lim_j \Delta_f h(p_j) &\geq 0, \\
\lim_j h(q_j) &= \sup h, \\
\lim_j \Delta_f h(q_j) &\leq 0.
\end{aligned} \tag{17}$$

On the other hand, from (9), we have

$$H_f = \frac{(-\rho'(h)/\rho(h))\{n + |\nabla h|^2\} - \Delta_f h}{n\Theta}. \tag{18}$$

Since Σ^n lies in a slab, if h is bounded from below, then

$$H_f \leq \frac{(-\rho'(\inf h)/\rho(\inf h))\{n + |\nabla h(p_j)|^2\}}{n\Theta(p_j)}. \tag{19}$$

Moreover, if h is bounded from above, we get

$$H_f \geq \frac{(-\rho'(\sup h)/\rho(\sup h))\{n + |\nabla h(q_j)|^2\}}{n\Theta(q_j)}. \tag{20}$$

Considering that the function $-\rho'/\rho$ is increasing, then

$$\begin{aligned}
H_f &\leq \frac{(-\rho'(\inf h)/\rho(\inf h))\{n + |\nabla h(p_j)|^2\}}{n\Theta(p_j)} \\
&\leq \frac{(-\rho'(h_0)/\rho(h_0))\{n + |\nabla h(p_j)|^2\}}{n\Theta(p_j)} \leq 0, \\
H_f &\geq \frac{(-\rho'(\sup h)/\rho(\sup h))\{n + |\nabla h(q_j)|^2\}}{n\Theta(q_j)} \\
&\geq \frac{(-\rho'(h_0)/\rho(h_0))\{n + |\nabla h(q_j)|^2\}}{n\Theta(q_j)} \geq 0.
\end{aligned} \tag{21}$$

Hence, $H_f = 0$; that is, Σ^n is a f -maximal spacelike hypersurface. Using (10), we have

$$\Delta_f \rho(h) = -n \frac{\rho'(h)^2}{\rho(h)} + |\nabla h|^2 \rho(h) (\log \rho)''(h) \leq 0. \tag{22}$$

In the following, by the same argument as in Theorem 4, we have that Σ^n is a slice. \square

4. Weighted Static GRW Space-Times

In this section, we obtain some rigidity results of stochastically complete hypersurfaces in weighted static GRW spacetimes $-I \times M_f^n$ by the weak maximal principle. Firstly, we give the following technical result which extended the corresponding conclusion in [12].

Lemma 7. *Let M be a stochastically complete Riemannian manifold and $u : M \rightarrow \mathbb{R}$ be a nonnegative smooth function on M . If there exists a positive constant λ such that $\Delta_f u \geq \lambda u$, then $u = 0$.*

Theorem 8. *Let $\psi : \Sigma^n \rightarrow -I \times M_f^n$ be a stochastically complete hypersurface with constant f -mean curvature H_f in a weighted static GRW spacetime $-I \times M_f^n$. Assume that $K_M \geq -k$ for some positive constant k and the weighted function f is convex. If $|\nabla h|^2 \leq \alpha |A|^2 / (n-1)k$ for some constant $0 < \alpha < 1$, then Σ^n is a slice.*

Proof. Let E_1, \dots, E_n be a (local) orthonormal frame in $\mathfrak{X}(\Sigma)$; using the Gauss equation, we have that

$$\begin{aligned}
\text{Ric}(X, X) &= \sum_{i=1}^n \langle \bar{R}(X, E_i)X, E_i \rangle + nH \langle AX, X \rangle \\
&\quad + |AX|^2,
\end{aligned} \tag{23}$$

for $X \in \mathfrak{X}(\Sigma)$. Moreover, we also have

$$\begin{aligned}
\langle \bar{R}(X, E_i)X, E_i \rangle \\
= K_M(X^*, E_i^*) (\langle X^*, X^* \rangle \langle E_i^*, E_i^* \rangle - \langle X^*, E_i^* \rangle^2),
\end{aligned} \tag{24}$$

where K_M is the sectional curvature of the fiber M^n and $X^* = X + \langle X, \partial_t \rangle \partial_t$ and $E_i^* = E_i + \langle E_i, \partial_t \rangle \partial_t$ are the projections of the tangent vector fields X and E_i onto M^n .

By a direct computation and considering the hypothesis $K_M > -k$, we get

$$\begin{aligned}
\sum_{i=1}^n \langle \bar{R}(X, E_i)X, E_i \rangle \\
\geq -k((n-1)|X|^2 + (n-2)\langle X, \nabla h \rangle^2 + |X|^2 |\nabla h|^2).
\end{aligned} \tag{25}$$

Substituting (25) into (23),

$$\begin{aligned}
\text{Ric}(X, X) \\
\geq -k((n-1)|X|^2 + (n-2)\langle X, \nabla h \rangle^2 + |X|^2 |\nabla h|^2) \\
+ nH \langle AX, X \rangle + |AX|^2.
\end{aligned} \tag{26}$$

Furthermore, taking into account that the weighted function f is convex, we have

$$\begin{aligned}
\text{Hess } f(X, X) &= \overline{\text{Hess } f}(X, X) - \langle \bar{\nabla} f, N \rangle \langle AX, X \rangle \\
&\geq -\langle \bar{\nabla} f, N \rangle \langle AX, X \rangle.
\end{aligned} \tag{27}$$

Therefore,

$$\begin{aligned}
\text{Ric}_f(X, X) \\
\geq -k((n-1)|X|^2 + (n-2)\langle X, \nabla h \rangle^2 + |X|^2 |\nabla h|^2) \\
+ nH_f \langle AX, X \rangle + |AX|^2.
\end{aligned} \tag{28}$$

In particular, we have

$$\begin{aligned} \text{Ric}_f(\nabla h, \nabla h) &\geq -k(n-1)|\nabla h|^2(1+|\nabla h|^2) \\ &\quad + nH_f \langle A(\nabla h), \nabla h \rangle + |A(\nabla h)|^2. \end{aligned} \quad (29)$$

Now we recall the Bochner-Lichnerowicz formula (see [2]):

$$\begin{aligned} \frac{1}{2}\Delta_f(|\nabla h|^2) &= |\text{Hess } h|^2 + \text{Ric}_f(\nabla h, \nabla h) \\ &\quad + \langle \nabla \Delta_f h, \nabla h \rangle. \end{aligned} \quad (30)$$

From the fact that H_f is a constant, we have

$$\nabla \Delta_f h = -nH_f A(\nabla h). \quad (31)$$

By [14], we get

$$|\text{Hess } h|^2 = |A|^2(1+|\nabla h|^2). \quad (32)$$

Using (29), (31), and (32) in (30), we have

$$\frac{1}{2}\Delta_f |\nabla h|^2 \geq (|A|^2 - (n-1)k|\nabla h|^2)(1+|\nabla h|^2). \quad (33)$$

Finally, considering the hypothesis $|\nabla h|^2 \leq \alpha|A|^2/(n-1)k$, we obtain

$$\Delta_f |\nabla h|^2 \geq 2(1-\alpha)|A|^2|\nabla h|^2. \quad (34)$$

Thus, there is a positive constant λ such that

$$\Delta_f |\nabla h|^2 \geq \lambda |\nabla h|^2. \quad (35)$$

Therefore, h is constant by Lemma 7. \square

Theorem 9. *Let $\psi : \Sigma^n \rightarrow -I \times M_f^n$ be a stochastically complete hypersurface with constant f -mean curvature H_f in a weighted static GRW space-time $-I \times M_f^n$. Assume that the sectional curvature K_M is nonnegative and the weighted function f is convex. If $|\nabla h|^2$ is bounded from above, then Σ^n is f -maximal.*

Proof. As in the proof of Theorem 8, taking into account that the hypothesis K_M is nonnegative, there is a constant $k \geq 0$ such that

$$\frac{1}{2}\Delta_f |\nabla h|^2 \geq |\text{Hess } h|^2 + (n-1)k|\nabla h|^2(1+|\nabla h|^2). \quad (36)$$

Moreover, considering the relation $n|\text{Hess } h|^2 \geq (\Delta_f h)^2$, we have

$$\Delta_f |\nabla h|^2 \geq \frac{2}{n}(\Delta_f h)^2. \quad (37)$$

Using (9) and (37), we obtain

$$\Delta_f |\nabla h|^2 \geq 2nH_f^2 \Theta^2 \geq 2nH_f^2 \geq 0. \quad (38)$$

By the hypothesis that $|\nabla h|^2$ is bounded from above, applying Lemma 3, the weak maximum principle, we get

$$0 \geq \lim_k \Delta_f |\nabla h|^2(q_k) \geq 2nH_f^2 \geq 0. \quad (39)$$

Therefore Σ^n is f -maximal. \square

As a consequence of the proof of Theorem 8, we can get the following corollary.

Corollary 10. *Let $\psi : \Sigma^n \rightarrow -I \times M_f^n$ be a stochastically complete hypersurface with constant f -mean curvature H_f in a weighted static GRW space-time $-I \times M_f^n$. Assume that $K_M \geq -k$ and $\overline{\text{Hess } f} \geq -\beta$ for some positive constants k and β . If $|\nabla h|^2 \leq \alpha|A|^2/((n-1)k+\beta)$ for some constant $0 < \alpha < 1$, then Σ^n is a slice.*

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Acknowledgments

This work is supported by National Natural Science Foundation of China (no. 11371076).

References

- [1] D. Bakry and M. Mery, "Diffusions hypercontractives," in *Seminaire de probabilités, XIX*, vol. 1123 of *Lecture Notes in Math.*, pp. 177–206, Springer, Berlin, Germany, 1983.
- [2] G. Wei and W. Wylie, "Comparison geometry for the Bakry-Emery Ricci tensor," *Journal of Differential Geometry*, vol. 83, no. 2, pp. 377–405, 2009.
- [3] M. P. Cavalcante, H. F. de Lima, and M. S. Santos, "On Bernstein-type properties of complete hypersurfaces in weighted warped products," *Annali di Matematica Pura ed Applicata. Series IV*, vol. 195, no. 2, pp. 309–322, 2016.
- [4] M. P. Cavalcante, H. F. de Lima, and M. S. Santos, "New Calabi-Bernstein type results in weighted generalized Robertson-Walker spacetimes," *Acta Mathematica Hungarica*, vol. 145, no. 2, pp. 440–454, 2015.
- [5] H. F. de Lima, A. M. Oliveira, and M. S. Santos, "Rigidity of complete spacelike hypersurfaces with constant weighted mean curvature," *Beiträge zur Algebra und Geometrie/Contributions to Algebra and Geometry*, vol. 57, no. 3, pp. 623–635, 2016.
- [6] B. O'Neill, *Semi-Riemannian Geometry with Applications to Relativity*, Academic Press, San Diego, Calif, USA, 1983.
- [7] L. J. Alas, A. Romero, and M. Sánchez, "Uniqueness of complete spacelike hypersurfaces of constant mean curvature in generalized Robertson-Walker spacetimes," *General Relativity and Gravitation*, vol. 27, no. 1, pp. 71–84, 1995.
- [8] M. Gromov, "Isoperimetry of waists and concentration of maps," *Geometric and Functional Analysis*, vol. 13, no. 1, pp. 178–215, 2003.
- [9] J. S. Case, "Singularity theorems and the Lorentzian splitting theorem for the Bakry-Emery-Ricci tensor," *Journal of Geometry and Physics*, vol. 60, no. 3, pp. 477–490, 2010.
- [10] A. Caminha, "The geometry of closed conformal vector fields on Riemannian spaces," *Bulletin of the Brazilian Mathematical Society. New Series. Boletim da Sociedade Brasileira de Matemática*, vol. 42, no. 2, pp. 277–300, 2011.
- [11] S. T. Yau, "Some function-theoretic properties of complete Riemannian manifold and their applications to geometry," *Indiana University Mathematics Journal*, vol. 25, no. 7, pp. 659–670, 1976.
- [12] S. Pigola, M. Rigoli, and A. G. Setti, "A remark on the maximum principle and stochastic completeness," *Proceedings of the*

American Mathematical Society, vol. 131, no. 4, pp. 1283–1288, 2003.

- [13] M. Rimoldi, *Rigidity Results for Lichnerowicz Bakry-Émery Ricci Tensors [Ph.D. thesis]*, Università degli Studi di Milano, Milano, Italy, 2011.
- [14] J. M. Latorre and A. Romero, “Uniqueness of noncompact spacelike hypersurfaces of constant mean curvature in generalized Robertson-Walker spacetimes,” *Geometriae Dedicata*, vol. 93, pp. 1–10, 2002.



Hindawi

Submit your manuscripts at
www.hindawi.com

