

Franco, S., He, Y., Herzog, C. & Walcher, J. (2004). Chaotic cascades for D-branes on singularities. Paper presented at the Cargese Summer School, Cargèse, France, 7th - 19th 2004.



**CITY UNIVERSITY
LONDON**

[City Research Online](http://www.city.ac.uk/researchonline)

Original citation: Franco, S., He, Y., Herzog, C. & Walcher, J. (2004). Chaotic cascades for D-branes on singularities. Paper presented at the Cargese Summer School, Cargèse, France, 7th - 19th 2004.

Permanent City Research Online URL: <http://openaccess.city.ac.uk/854/>

Copyright & reuse

City University London has developed City Research Online so that its users may access the research outputs of City University London's staff. Copyright © and Moral Rights for this paper are retained by the individual author(s) and/ or other copyright holders. All material in City Research Online is checked for eligibility for copyright before being made available in the live archive. URLs from City Research Online may be freely distributed and linked to from other web pages.

Versions of research

The version in City Research Online may differ from the final published version. Users are advised to check the Permanent City Research Online URL above for the status of the paper.

Enquiries

If you have any enquiries about any aspect of City Research Online, or if you wish to make contact with the author(s) of this paper, please email the team at publications@city.ac.uk.

CHAOTIC CASCADES FOR D-BRANES ON SINGULARITIES

Sebastián Franco¹, Yang-Hui He², Christopher Herzog³, Johannes Walcher⁴

1. Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

2. Dept. of Physics and Math/Physics RG, Univ. of Pennsylvania, Philadelphia, PA 19104, USA

3. Kavli Institute for Theoretical Physics, University of California, Santa Barbara, CA 93106, USA

4. Institute for Advanced Study, Princeton, NJ 08540, USA

sfranco@mit.edu, yanghe@physics.upenn.edu, herzog@kitp.ucsb.edu, walcher@ias.edu

Abstract We briefly review our work on the cascading renormalization group flows for gauge theories on D-branes probing Calabi-Yau singularities. Such RG flows are sometimes chaotic and exhibit duality walls. We construct supergravity solutions dual to logarithmic flows for these theories. We make new observations about a surface of conformal theories and more complicated supergravity solutions.

1. INTRODUCTION

Extending the revolutionary AdS/CFT correspondence [1] beyond the original relation between $\mathcal{N} = 4$ SYM on N D3-branes and Type IIB supergravity (sugra) in $AdS_5 \times S^5$ with N units of RR 5-form flux on the S^5 is important to understanding realistic strongly coupled field theories such as QCD.

Two standard extensions have been (1) *reducing the SUSY to $\mathcal{N} = 1$* by placing the D3-branes transverse to a Calabi-Yau singularity (the dual sugra background becomes $AdS_5 \times X^5$, where X^5 is some non-spherical horizon); and (2) *breaking conformal invariance and inducing an RG flow*, by introducing fractional branes, i.e., D5-branes wrapped over collapsing 2-cycles of the singularity (in the sugra dual, 3-form fluxes are turned on). A fascinating type of RG flow is the *duality cascade*: Seiberg duality is used to switch to an alternative description whenever infinite coupling is reached. This idea was introduced in [2] for the gauge theory on D-branes probing the conifold.




Figure 1.1

2. CASCADES IN COUPLING SPACE

There is an interesting way to look at cascading RG flows. In a gauge theory described by a quiver with k gauge groups, the inverse squared couplings $x_i \equiv 1/g_i^2$ are positive and define a k -dimensional cone $(\mathbb{R}_+)^k$. Inside this cone, the RG flow generates a trajectory dictated by the beta functions and satisfying $\sum_i r^i/g_i^2 = \text{constant}$. Each step between dualizations then corresponds to a straight line in the simplex defined by the intersection between this hyperplane and the $(\mathbb{R}_+)^k$ cone. We show such a trajectory in Figure 1.1(A). Now, each wall of the cone corresponds to one of the gauge couplings going to infinity. Therefore, whenever one of them is reached, we switch to a Seiberg dual theory at weak coupling. There will be then a different simplex associated to the dual theory. The entire cascade corresponds to a flow in the space of glued simplices. From this perspective, which resembles a billiard

bouncing in coupling space, one foresees that cascading RG flows will exhibit chaotic behavior.

3. DUALITY WALLS AND FRACTALS

After introducing the notion of a duality cascade, it is natural to wonder whether some supersymmetric extension of the Standard Model, such as the MSSM, can sit at the IR endpoint of a cascade. This question was posed by Matthew Strassler [3]. Generically, while trying to reconstruct such a RG flow, one encounters a UV accumulation point beyond which Seiberg duality cannot proceed. This phenomenon is dubbed a *duality wall* and has been constructed for gauge theories engineered with D-branes on singularities [4]. Figure 1.1(B) shows the behavior of couplings for a cascade with a duality wall for the theory on D-branes over a complex cone over the Zeroth-Hirzebruch surface F_0 ¹.

Postponing the question of a possible UV completion of duality walls, we can study the dependence of its position on initial couplings. Illustrating with F_0 , the result is remarkable and is presented in Figure 1.1(C). The curve is a fractal, with concave and convex cusps. Whenever we zoom in on a convex cusp, an infinite, self-similar structure of more cusps emerges.

One subtle point which was not emphasized in [5] involves the existence in coupling space of a codimension two surface of conformal theories for F_0 and the other del Pezzo quiver gauge theories. If the number of gauge couplings is $n + 2$, then a naive counting of the linearly independent β -functions constrains only two combinations of gauge couplings when the theory is conformal, leaving an n -dimensional surface of conformal theories. This n -dimensional surface is parametrized on the gravity side by the dilaton and the integral of the NSNS B_2 form through $n - 1$ independent 2-cycles.

The existence of this codimension two surface may well affect the existence and behavior of the duality wall for F_0 . In [4], it was assumed that a generic choice of initial couplings would lie on the conformal surface. However, if the initial conditions do not lie on the conformal surface, one expects large coupling constant corrections to the anomalous dimensions, which will in turn affect the strengths of the β -functions.

4. SUPERGRAVITY DUALS

The main support for the idea of a cascading RG flow in the original case of the conifold comes from a supergravity dual construction. This

¹We refer the reader to [4, 5] for a detailed description of the associated quiver theory.

dual reproduces the logarithmic decrease in the effective number of colors towards the IR and also matches the beta functions for the gauge couplings.

In [5], analog supergravity solutions were constructed describing logarithmic cascades for the gauge theories on D-branes probing complex cones over del Pezzo surfaces. The fact that this was possible is remarkable, since they were obtained without knowing the explicit metric. These supergravity solutions are of the general type studied by Graña and Polchinski [6].

The general form of the metric is a warped product of flat four-dimensional Minkowski space and a Calabi-Yau \mathbf{X}

$$ds^2 = Z^{-1/2} \eta_{\mu\nu} dx^\mu dx^\nu + Z^{1/2} ds_{\mathbf{X}}^2, \quad (1.1)$$

The solution also carries 3-form flux $G_3 = F_3 - \frac{i}{g_s} H_3$. In order to preserve $\mathcal{N} = 1$ supersymmetry, G_3 must be supported only on X , imaginary self-dual, a $(2, 1)$ form and harmonic. Indeed, it is possible to construct a G_3 satisfying all these conditions. It has the form

$$G_3 = \sum_{I=1}^n a^I \left(\eta + i \frac{dr}{r} \right) \wedge \phi_I \quad (1.2)$$

where the ϕ_I , $I = 1 \dots n$, are a basis of $(1, 1)$ forms orthogonal to the Kähler class of the del Pezzo and $\eta = \left(\frac{1}{3} d\psi + \sigma \right)$. The one-form σ satisfies $d\sigma = 2\omega$, with ω the Kähler form on dP_n , and $0 \leq \psi < 2\pi$ is the angular coordinate on the circle bundle over dP_n .

The intersection product between the ϕ_I is $\int_{dP_n} \phi_I \wedge \phi_J = -A_{IJ}$, where A_{IJ} is the Cartan matrix for the exceptional Lie algebra \mathcal{E}_n . There is a different type of fractional brane associated to each ϕ_I , given by D5-branes wrapping the 2-cycle in the del Pezzo Poincaré dual to ϕ_I .

Let us now study the number of D5-branes and D3-branes associated to these solutions

D5-Branes: The number of D5-branes is given by the Dirac quantization of the RR 3-form F_3 : $a^J = 6\pi\alpha' M^J$. Hence, this family of solutions are dual to cascades in which the number of fractional branes of each type remains constant.

D3-Branes: Similarly, the effective number of D3-branes is computed from $F_5 = \mathcal{F}_5 + *\mathcal{F}_5$ where $\mathcal{F}_5 = d^4x \wedge d(Z^{-1})$ and Z is the warp factor in (1.1). The factor Z satisfies the equation

$$\nabla_{\mathbf{X}}^2 Z = -\frac{1}{6} |H_3|^2. \quad (1.3)$$

In [5], $|F_3|^2$ was assumed to be a function only of the radius, in which case

$$Z(r) = \frac{2 \cdot 3^4}{9-n} \alpha'^2 g_s^2 \left(\frac{\ln(r/r_0)}{r^4} + \frac{1}{4r^4} \right) \sum_{i,j} M^I A_{IJ} M^J \quad (1.4)$$

and from Dirac quantization, the number of D3-branes will grow logarithmically: $N = \frac{3}{2\pi} g_s \ln(r/r_0) \sum_{I,J} M^I A_{IJ} M^J$. However, generically, Z may depend on other coordinates on the Calabi-Yau cone \mathbf{X} .² The function Z averaged over the other coordinates may still be logarithmic in r [7].

5. RECENT DEVELOPMENTS

Recently, there has been further progress in the study of quiver theories and their sugra duals. In [8], a-maximization [9] was used to compute the volume of the 5d horizon of the dual of the dP_1 gauge theory, yielding an irrational value. This result corrected previous computations in the literature and was obtained by carefully taking into account the global symmetries that are actually preserved by the superpotential. In [5], the duality cascade for dP_1 was analyzed using naive R-charges that did not take into account these global symmetries. A stable elliptical region in coupling space was found with a self-similar logarithmic cascade. Re-doing the analysis with the new R-charges, we find the same elliptical region albeit with a slightly different shape and center.

The 5d horizon for the complex cone over dP_1 is called $Y^{2,1}$ and is a member of an infinite family of Sasaki-Einstein geometries denoted $Y^{p,q}$. They have $S^2 \times S^3$ topology. Their metrics were first found, locally, in [10] and then the global properties were analysed in [11]. In [12], their toric description was worked out. Furthermore, the gauge theory duals to the entire $Y^{p,q}$ family have been constructed [13]. These developments change profoundly the status of the AdS/CFT, providing an infinite number of field theories with explicit sugra duals.

Acknowledgments

We would like to thank Qudisia Jabeen Ejaz, Ami Hanany, Pavlos Kazakopoulos, Igor Klebanov and Joe Polchinski for useful discussions. This research is funded in part by the CTP and the LNS of MIT and by the department of Physics at UPenn. The research of S. F. was supported in part by U.S. DOE Grant #DE-FC02-94ER40818. The research of Y.-H. H. was supported in part by U.S. DOE Grant #DE-FG02-95ER40893 as well as an NSF Focused Research Grant DMS0139799 for “The Geometry of Superstrings”. C. H. was supported in part by the NSF under

²We would like to thank Q. J. Ejaz for telling us about this possibility.

Grant No. PHY99-07949. The research of J. W. was supported by U.S. DOE Grant #DE-FG02-90ER40542. S. F. would like to thank the organizers of Cargese Summer School, where this material was presented.

References

- [1] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity,” *Adv. Theor. Math. Phys.* **2**, 231 (1998) [*Int. J. Theor. Phys.* **38**, 1113 (1999)] [arXiv:hep-th/9711200].
- [2] I. R. Klebanov and M. J. Strassler, “Supergravity and a confining gauge theory: Duality cascades and chiSB-resolution of naked singularities,” *JHEP* **0008**, 052 (2000) [arXiv:hep-th/0007191].
- [3] M. J. Strassler, “Duality in Supersymmetric Field Theory and an Application to Real Particle Physics,” Talk given at International Workshop on Perspectives of Strong Coupling Gauge Theories (SCGT 96), Nagoya, Japan. Available at <http://www.eken.phys.nagoya-u.ac.jp/Scgt/proc/>
- [4] S. Franco, A. Hanany, Y. H. He and P. Kazakopoulos, “Duality walls, duality trees and fractional branes,” arXiv:hep-th/0306092.
- [5] S. Franco, Y. H. He, C. Herzog and J. Walcher, *Phys. Rev. D* **70**, 046006 (2004) [arXiv:hep-th/0402120].
- [6] M. Grana and J. Polchinski, “Supersymmetric three-form flux perturbations on AdS(5),” *Phys. Rev. D* **63**, 026001 (2001) [arXiv:hep-th/0009211].
- [7] C. Herzog, Q. J. Ejaz, and I. Klebanov, “Cascading RG Flows for New Sasaki-Einstein Manifolds,” arXiv:hep-th/0412193.
- [8] M. Bertolini, F. Bigazzi and A. L. Cotrone, “New checks and subtleties for AdS/CFT and a-maximization,” arXiv:hep-th/0411249.
- [9] K. Intriligator and B. Wecht, “The exact superconformal R-symmetry maximizes a,” *Nucl. Phys. B* **667**, 183 (2003) [arXiv:hep-th/0304128].
- [10] J. P. Gauntlett, D. Martelli, J. Sparks and D. Waldram, “Supersymmetric AdS(5) solutions of M-theory,” *Class. Quant. Grav.* **21**, 4335 (2004) [arXiv:hep-th/0402153].
- [11] J. P. Gauntlett, D. Martelli, J. Sparks and D. Waldram, “Sasaki-Einstein metrics on $S(2) \times S(3)$,” arXiv:hep-th/0403002.
- [12] D. Martelli and J. Sparks, “Toric geometry, Sasaki-Einstein manifolds and a new infinite class of AdS/CFT duals,” arXiv:hep-th/0411238, and references therein.

- [13] S. Benvenuti, S. Franco, A. Hanany, D. Martelli and J. Sparks, “An infinite family of superconformal quiver gauge theories with Sasaki-Einstein duals,” arXiv:hep-th/0411264.