

## MECHANISMS OF INSECTICIDE RESISTANCE IN FIELD POPULATIONS OF *Aedes aegypti* (L.) FROM QUINTANA ROO, SOUTHERN MEXICO

ADRIANA E. FLORES,<sup>1</sup> JAIME SALOMON GRAJALES,<sup>1</sup> ILDEFONSO FERNANDEZ SALAS,<sup>1</sup>  
GUSTAVO PONCE GARCIA,<sup>1</sup> MA. HAYDEE LOAIZA BECERRA,<sup>1</sup> SAUL LOZANO,<sup>1</sup>  
WILLIAM G. BROGDON,<sup>2</sup> WILLIAM C. BLACK IV,<sup>3</sup> AND BARRY BEATY<sup>4</sup>

**ABSTRACT.** Potential insecticide-resistance mechanisms were studied with the use of biochemical assays in *Aedes aegypti* (L.) collected from 5 municipalities representing the north part of Quintana Roo: Benito Juárez, Cozumel, Isla Mujeres, Lazaro Cardenas, and Solidaridad. The activities of  $\alpha$  and  $\beta$  esterases, mixed-function oxidases (MFO), glutathione-S-transferase (GST), acetylcholinesterase (AChE), and insensitive acetylcholinesterase (iAChE) were assayed in microplates. Three replicates were performed for each enzyme and 60 males and 60 females were analyzed in each population. The New Orleans (NO) susceptible strain of *Ae. aegypti* was used as a susceptible reference and the threshold criteria for each enzyme were the highest NO absorbance values. In none of the 6 tests were absorbance values correlated in males and females.  $\alpha$  esterases were elevated in Benito Juárez, Cozumel females and in Lazaro Cardenas males and females.  $\beta$  esterases were elevated in Benito Juárez, Cozumel females and in Cozumel and Lazaro Cardenas males. Elevated esterases suggest potential insecticide-resistance mechanisms against organophosphate, carbamate, and some pyrethroid insecticides. Slightly elevated levels of MFOs appeared in Lazaro Cardenas females and in Cozumel, Isla Mujeres, and Solidaridad males. Mechanisms involving iAChE or GST were not apparent.

**KEY WORDS.** *Aedes aegypti*, insecticide resistance, insecticide surveillance, esterases, permethrin.

### INTRODUCTION

Dengue fever (DF) and Dengue hemorrhagic fever (DHF), vector-borne diseases transmitted by the mosquito *Aedes aegypti*, are increasing in prevalence in Quintana Roo state, in southern Mexico (Boletín de Epidemiología SSA, 2002). In 2002, 65 positive locations (including 12 municipalities) were reported in Quintana Roo. There were 172 cases of DF and 10 of DHF, in 5 municipalities in northern areas of the state and 479 DF cases and 80 DHF cases total.

Insecticides have played an important role in the control of *Ae. aegypti* in Mexico. However, the resistance of vectors to insecticides is one of the most important factors contributing to the ineffectiveness of control programs. Therefore, a critique of the potential for increases in resistance is essential (Georghiou and Mellom 1983). Since 1960 the vector program of the Ministry of Health in Mexico has used a series of insecticides for the control of dengue and malaria (Official Regulations of Mexico, NOM-032-

SSA). Dichloro-diphenyl-trichloroethane (DDT) is currently restricted, but had seen prolonged use for indoor house spraying. Malathion was used for ultra-low volume (ULV) space spraying of wide areas for both *Anopheles* and *Aedes* mosquitoes from 1981 to 1999. Ministry of Health vector-control programs have switched to pyrethroids (permethrin) in the last 6 years for adult control. Temephos or 1% Abate<sup>®</sup> granules have been applied to bodies of water and domestic containers for control of larvae. The insecticides utilized in public health and those directed at agricultural pests have provided prolonged and intense selection pressure for resistance evolution in *Ae. aegypti* and other vectors. Herein we report the first investigations of the principal biochemical mechanisms responsible for resistance in *Ae. aegypti* in Quintana Roo. This information will assist in the management and control of *Ae. aegypti* in Mexico.

### MATERIALS AND METHODS

**Study sites.** Quintana Roo adjoins the states of Yucatan and Campeche, and shares international borders with Guatemala and Belize. The geographic location of Quintana Roo is between latitudes 21°37'N and 17°49'N, and between longitudes 86°44'W and 89°24'52"W. The total area of the state is 50,844 km<sup>2</sup>. The history of *Ae. aegypti* larvae and eggs were sampled from 5 municipalities and transported to the Medical Entomology Laboratory of UANL, Mexico. The eggs were placed in plastic containers with water and, after hatching, larvae were provided with dog food, yeast, and brain-heart infusion agar in

<sup>1</sup> Laboratorio de Entomología Médica, Facultad de Ciencias Biológicas, Universidad Autónoma de Nuevo León, Ap. Postal 109-F, San Nicolás de los Garza, NL 66450, Mexico.

<sup>2</sup> Secretaría Estatal de Salud de Quintana Roo, Avenida Miguel Hidalgo 75, Cancun QRoo, CP 77510, Mexico.

<sup>3</sup> Centers for Disease Control and Prevention, Entomology Branch, F22, DPD, NCID, 4770 Buford Highway, Atlanta, GA 30341.

<sup>4</sup> Arthropod-Borne and Infectious Diseases Laboratory, Microbiology, Immunology and Pathology Department, Colorado State University, Fort Collins, CO 80523.

a finely ground mixture. Emerged postteneral adults were provided with water and cotton soaked in 10% honey. Females were provided with rat blood for egg production. Once the colonies were established, 60  $F_1$  unfed 1-day-old females and 60 males were placed individually in numbered vials, and stored at  $-70^\circ\text{C}$ .

**Biochemical assay:** Each mosquito was homogenized in 100  $\mu\text{l}$  0.01 M potassium phosphate buffer, pH 7.2, and then diluted to 2 ml with the same buffer. Aliquots of 100  $\mu\text{l}$  were transferred to individual microtiter plate wells. Sixty female and 60 male adults per location were analyzed in triplicate per plate. Six different resistance enzymes were evaluated for each mosquito (Brogdon and McAllister 1988a, 1988b, 1997):  $\alpha$  and  $\beta$  esterases, mixed-function oxidases (MFO), glutathione-S-transferase (GST), acetylcholinesterase (AChE), and insensitive acetylcholinesterase (iAChE). A minimum of 2 positive and negative controls, depending upon the test, were used per plate. Absorbance was measured with a Benchmark spectrophotometer (Biorad Laboratories, Philadelphia, USA). Results were expressed as a frequency distribution of the absorbance values. The New Orleans strain (NO) was used as the susceptible reference standard. The maximum absorbance value for NO for females and males was used as the susceptibility threshold.

Protein concentration was determined to correct for size variation among the specimens (Brogdon 1984). Statistical analysis was performed with ANOVA ( $\alpha = 0.05$ ) to test for differences in enzymes among populations and between sexes.

## RESULTS AND DISCUSSION

Maximum absorbance values for the NO strain appear on Table 1. Table 2 lists, by location and sex, the percentage of mosquitoes that exceeded the NO thresholds. Benito Juarez females exceeded the thresholds for  $\alpha$  and  $\beta$  esterases; however, in males, none of the enzyme activities was greater than the susceptibility threshold. All Cozumel females had increased levels of  $\alpha$  and  $\beta$  esterases, but only a few individuals had

Table 1. Susceptibility threshold based on maximum absorbance of different detoxification mechanisms in New Orleans strain.

Biochemical test	Females	Males
$\alpha$ -esterases	0.65	0.60
$\beta$ -esterases	0.95	1.02
MFO	0.16	0.12
GST	0.23	0.16
AChE	0.28	0.27
iAChE	0.02	0.02

increased iAChE and MFO. Most Cozumel males had an increase in  $\beta$  esterase and just 35% had elevated MFO activity, whereas increases in iAChE and  $\alpha$  esterase were negligible. Neither Isla Mujeres females nor males exceeded the threshold. Lazaro Cardenas females did not exhibit increased esterase activity, but MFO activity was elevated. In contrast, Lazaro Cardenas males had increased  $\alpha$  and  $\beta$  esterase activities and a few had increased MFO activity. Solidaridad females did not exhibit higher activity in any enzyme system, but one quarter of males had increased MFO activity.

Correlation coefficients were calculated between population means for males and females for each enzyme system. In none of the 6 systems was absorbance value correlated in males and females (Fig. 1). Benito Juarez and Cozumel females had significantly greater  $\alpha$  esterase activities than NO (Fig. 2). Only Lazaro Cardenas males had significantly greater  $\alpha$  esterase activities (Fig. 3). Benito Juarez and Cozumel females also had the highest  $\beta$  esterase activities, and Lazaro Cardenas and Cozumel males had significantly higher  $\beta$  esterase activities (Figs. 2 and 3). The rest of the populations also had significantly greater mean activity levels, but differences were only slight (Figs. 2 and 3, Table 3).

Broad-spectrum organophosphate resistance is conferred by the elevated esterases of *Culex* sp. Work is in progress on esterase-resistance mechanisms in a range of *Anopheles* and *Aedes* species. All of these esterases act by rapidly binding and then slowly metabolizing the insecticide (Kadous et al. 1983). Esterases have diverse functions in insects, including proteolysis, nervous system

Table 2. Percentage of female and male *Ae. aegypti* that exceeded the susceptibility threshold established in New Orleans.

Biochemical test	B. Juarez		Cozumel		I. Mujeres		L. Cardenas		Solidaridad	
	Female	Male	Female	Male	Female	Male	Female	Male	Female	Male
$\alpha$ esterases	100.00	0	100.00	5.00	0	0	1.67	100.00	0	1.67
$\beta$ esterases	98.33	0	100.00	65.00	1.67	0	0	65.00	3.33	0
MFO	10.00	3.33	18.34	66.67	11.66	16.67	98.33	20.00	0	58.34
GST	0	0	0	6.67	0	0	0	0	0	0
AChE	0	0	0	0	0	0	0	0	0	0
iAChE	5.00	0	1.67	1.67	0	0	5.00	1.67	0	1.67






Fig. 1. Correlations between males and females according to absorbance values.

function, hormone metabolism, and xenobiotic metabolism/sequestration (Aldridge 1993). Studies of insecticide resistance have indicated the specific relevance of esterases with regard to xenobiotic metabolism in several insect species, including *Ae. aegypti*. The resistance of vectors to insecticides has indicated that it is one of the most important factors contributing to the ineffectiveness of mosquito control programs. Nonspecific esterases have been reported to be involved in pyrethroid metabolism in several insects (Soderlund et al. 1983, Ruigt 1985) and could play a role in the metabolism of permethrin in *Ae. aegypti*. Even in some cases, there is evidence that esterases are involved in conferring cross resistance to fenitrothion and deltamethrin in the larvae and adults of *Anopheles albimanus* in Guatemala (Brogdon and Barber 1990). Other reports indicate that populations of *Cx p. quinquefasciatus* that were highly resistant to

lambda-cyhalothrin and deltamethrin were also highly cross resistant to malathion (Bisset et al. 1997, Bisset et al. 1998). All the populations studied showed the consistent presence of  $\alpha$ -esterases, with elevated levels in permethrin-selected populations.

Only Lazaro Cardenas females had a mean MFO absorbance value greater than that of the NO strain (Fig. 2). Cozumel, Solidaridad, and Isla Mujeres males displayed greater MFO activity than the NO strain (Fig. 3). Mixed-function oxidases (MFOs) and nonspecific esterases (NSEs) are commonly involved in the detoxification of permethrin (Miller 1988, Zerba 1988, Ishaaya 1993). Biochemical assays have been developed to measure levels of these enzymes in mosquitoes, and elevated levels of such enzymes are known to enhance insecticide tolerance (Brogdon and McAllister 1988a). Vulule et al. (1999) found evidence that elevated




Fig. 2. Mean absorbance values of adult females of *Aedes aegypti* from 5 populations in study and NO strain obtained in biochemical assays.

MFO levels were associated with permethrin tolerance in *An. gambiae*. First, in Kenyan villages (Vulule et al. 1994); after introduction of permethrin-impregnated nets, found that *An. gambiae* males had higher MFO levels than in males from nearby villages where *An. gambiae* remained susceptible to permethrin and impreg-

nated nets were not used. They assumed that high MFO levels in males can be inherited from maternal females selected for permethrin by exposure to permethrin-impregnated nets when foraging for a blood meal. Brogdon et al. (1999) found that only adult females of *An. albimanus* expressed an oxidase mechanism, although they




Fig. 3. Mean absorbance values of adult males of *Aedes aegypti* from 5 populations in study and NO strain obtained in biochemical assays.

Table 3. Means and standard deviations of the mean absorbance in each biochemical test of 6 populations of female and male *Ae. aegypti*.

	$\alpha$ esterases			$\beta$ esterases			MFO	GST	AChE	iAChE		
<b>Females</b>												
New Orleans	0.4979D**	0.0775	0.7913C*	0.0829	0.1236B*	0.0238	0.1207A*	0.0402	0.2005A*	0.0305	0.0091A*	0.0043
Benito Juarez	0.9632B*	0.0565	1.1990A*	0.1082	0.1027D*	0.0298	0.0739C*	0.0220	0.1106D*	0.0640	0.0138B*	0.0440
Cozumel	0.8479A*	0.0413	1.1444B*	0.0565	0.1170BC*	0.0254	0.0872B*	0.0273	0.1515C*	0.0278	0.0087A*	0.0037
Isla Mujeres	0.3714E*	0.0284	0.6752D*	0.0853	0.1109CD*	0.0220	0.0035E*	0.0043	0.1098D*	0.0265	0.0046A*	0.0043
Lazaro Cardenas	0.5787C*	0.0316	0.6019E*	0.0378	0.1621A*	0.0120	0.1087A*	0.0346	0.1727B*	0.0253	0.0011A*	0.0097
Solidaridad	0.4979D*	0.0490	0.6588D*	0.1208	0.0809E*	0.0238	0.0455D*	0.0174	0.1388C*	0.0271	0.0028A*	0.0033
<b>Males</b>												
New Orleans	0.5408B*	0.0225	0.9645C*	0.0302	0.0872CD*	0.0087	0.1010AB*	0.0276	0.2202A*	0.0193	0.0075A*	0.0060
Benito Juarez	0.5150C*	0.0234	0.8088D*	0.0477	0.0750D*	0.0106	0.0932B*	0.0233	0.1238B*	0.0248	0.0061AB*	0.0031
Cozumel	0.5475B*	0.0338	1.0396B*	0.1008	0.1200A*	0.0533	0.1099A*	0.0319	0.1582B*	0.0353	0.0078A*	0.0030
Isla Mujeres	0.4317E*	0.0342	0.7539E*	0.0703	0.0994BC*	0.0285	0.0027D*	0.0038	0.0909D*	0.0211	0.0026C*	0.0019
Lazaro Cardenas	0.7548A*	0.0947	1.1987A*	0.2476	0.0797D*	0.0522	0.0069D*	0.0021	0.1153C*	0.0189	0.0069AB*	0.0021
Solidaridad	0.4743D*	0.0559	0.9260C*	0.0328	0.1064AB*	0.0207	0.0486C*	0.0196	0.0672E*	0.0217	0.0054B*	0.0070

\* Different letters in columns mean significant difference.

\*\* P &lt; 0.01.

said that further investigation is needed to provide the genetic explanation for the absence of expression in males. Many studies have shown that resistant insects have elevated levels of glutathione S-transferase activity in crude homogenates, which suggests a role for GSTs in resistance (Grant 1991, Grant et al. 1991). Multiple forms of these enzymes have been reported for mosquitoes, house fly, *Drosophila*, sheep blow fly, and grass grub (Clark et al. 1984, 1985; Toung et al. 1990). In *Ae. aegypti* at least 2 GSTs are elevated in DDT-resistant insects (Grant and Hammock 1992, Grant and Matsuura 1989), and in *An. gambiae* a large number of different GSTs are elevated, some of which are class I GSTs (Prapanthadara et al. 1993, 1995). The *Ae. aegypti* and *An. gambiae* GSTs in resistant insects are constitutively overexpressed. The GST-2 of *Ae. aegypti* is overexpressed in all tissues except the ovaries of resistant insects (Grant and Hammock 1992). Only Cozumel males had elevated GST over that of NO.

In examining AChE and iAChE (Figs. 2 and 3) activity, all populations showed lower absorbance than the NO strain.

Both  $\alpha$  and  $\beta$  esterase assays showed elevated enzyme levels characteristic of resistance mechanisms. These were detected in females of *Ae. aegypti* and occurred at a much higher frequency in areas of intensive use of insecticide, such as in Benito Juarez and Cozumel (tourist areas). Levels of MFO in females were elevated in Lazaro Cardenas, suggesting that they have been selected as a detoxifying mechanism. These elevated levels were focal, but with a high degree of intersite variability. These data are consistent with the view that these resistance mechanisms have been under selection by the insecticides used in Quintana Roo.

#### ACKNOWLEDGMENT

This study was supported by CONACyT, México through Grant No. 37408-N within the project "Assessing the significance of insecticide resistance in *Aedes aegypti* (L.) in Mexico."

#### REFERENCES CITED

- Aldridge WN. 1993. The esterases: perspectives and problems. *Chem Biol Interact* 87:5-13.  
 Bisset JA, Rodríguez MM, Díaz C, Soca A. 1998. Estudio de la resistencia en una cepa de *Culex quinquefasciatus*, procedente de Medellín, Colombia. *Rev Cubana Med Trop* 50(2):133-137.  
 Bisset JA, Rodríguez MM, Soca A, Pasteur N, Raymond M. 1997. Cross-resistance to pyrethroids and organophosphorus insecticides in the southern house mosquito (Diptera: Culicidae) from Cuba. *J Med Entomol* 34:244-246.



- Boletín de Epidemiología. 2002. Sistema Nacional de Vigilancia Epidemiológica. Secretaría de Salud: Mexico. Vol. 52, No. 16.
- Brogdon WG. 1984. Mosquito protein microassay. I. Protein determinations from small portions of single-mosquito homogenates. *Comp Biochem Physiol* 79:457-459.
- Brogdon WG, Barber AM. 1990. Microplate assay of glutathione S-transferase activity for resistance detection in single-mosquito triturates. *Comp Biochem Physiol* 96:339-342.
- Brogdon WG, McAllister JC. 1997. Heme peroxidase activity measured in single mosquitoes identifies individuals expressing an elevated oxidase for insecticide resistance. *J Am Mosq Control Assoc* 13:233-237.
- Brogdon WG, McAllister JC. 1998a. Insecticide resistance and vector control. Centers for Disease Control: Atlanta, GA, Vol. 4, No. 4, 12 p.
- Brogdon WG, McAllister JC. 1998b. Simplification of adult mosquito bioassays through use of time-mortality determinations in bottles. *J Am Mosq Control Assoc* 14(2):159-164.
- Brogdon WG, McAllister JC, Corwin AM, Cordon RC. 1999. Oxidase-based DDT-pyrethroid cross-resistance in Guatemalan *Anopheles albimanus*. *Pestic Biochem Physiol* 64:101-111.
- Clark AG, Dick GL, Martindale SM, Smith JN. 1985. Glutathione S transferases from the New Zealand grass grub, *Costelytra zealandica*. *Insect Biochem* 15:35-44.
- Clark AG, Shamaan NA, Dauterman WC, Hayaoka T. 1984. Characterization of multiple glutathione transferases from the housefly, *Musca domestica* (L.). *Pestic Biochem Physiol* 22:51-59.
- Georghiou GP, Mellom RB. 1983. Pesticide resistance in time and space. In: Georghiou GP, Saito T, eds. *Pest resistance to pesticides*. New York: Plenum Press, p 1-46.
- Grant DF. 1991. Evolution of glutathione S-transferase subunits in Culicidae and in the picrotoxinin receptor between the cyclodiene-resistant and susceptible strains of the German-cockroach. *Pest Biochem Physiol* 19:157-166.
- Grant DF, Dietze EC, Hammock BD. 1991. Glutathione S-transferase isozymes in *Aedes aegypti*: purification, characterization, and isozyme specific regulation. *Insect Biochem* 4:421-33.
- Grant DF, Hammock BD. 1992. Genetic and molecular evidence for a trans-acting regulatory locus controlling glutathione-s transferase-2 expression in *Aedes aegypti*. *Mol Gen Genet* 234:169-176.
- Grant DF, Matsumura F. 1989. Glutathione S-transferase 1 and 2 in susceptible and insecticide resistant *Aedes aegypti*. *Pestic Biochem Physiol* 33:132-143.
- Ishaaya I. 1993. Insect detoxifying enzymes: their importance in pesticide synergism and resistance. *Arch Insect Biochem Physiol* 22:263-276.
- Kadous AA, Ghiasuddin SM, Matsumura F, Scott JG, Tanaka K. 1983. Difference in the picrotoxinin receptor between the cyclodiene resistant and susceptible strains of the German-cockroach. *Pest Biochem Physiol* 19:157-166.
- Miller TM. 1988. Mechanisms of resistance to pyrethroid insecticides. *Parasitol Today* 4:S8-12.
- Prapanthadara L, Hemingway J, Ketterman AJ. 1993. Partial purification and characterization of glutathione S-transferase involved in DDT resistance from the mosquito *Anopheles gambiae*. *Pest Biochem Physiol* 47:119-133.
- Prapanthadara L, Hemingway J, Ketterman AJ. 1995. DDT-resistance in *Anopheles gambiae* Giles from Zanzibar Tanzania, based on increased DDT-dehydrochlorinase activity of glutathione S-transferases. *Bull Entomol Res* 85:267-274.
- Ruigt DF. 1985. Pyrethroids. In: Kerkut GA, Gilbert LI, eds. *Comprehensive insect physiology, biochemistry and pharmacology*. Volume 12. Oxford, United Kingdom: Pergamon Press, p 183-262.
- Soderlund DM, Sanborn JR, Lee PW. 1983. Metabolism of pyrethrins and pyrethroids in insects. In: Hutson D, Roberts TR, eds. *Progress in pesticide biochemistry and toxicology*. Vol. 3. New York: Wiley, p 401-435.
- Toung YS, Hsieh T, Tu CD. 1990. *Drosophila* glutathione S-transferase 1-1 shares a region of sequence homology with maize glutathione S-transferase III. *Proc Natl Acad Sci USA* 87:31-35.
- Vulule JM, Beach RF, Atieli FK, McAllister JC, Brogdon WG, Roberts JM, Mwangi RW, Hawley WA. 1999. Elevated oxidase and esterase levels associated with permethrin tolerance in *Anopheles gambiae* from Kenyan villages using permethrin-impregnated nets. *Med Vet Entomol* 13:239-244.
- Vulule JM, Beach RF, Atieli FK, Roberts JM, Mount DL, Mwangi RW. 1994. Reduced susceptibility of *Anopheles gambiae* to permethrin associated with the use of permethrin impregnated bednets and curtains in Kenya. *Med Vet Entomol* 8:71-75.
- Zerba E. 1988. Insecticidal activity of pyrethroids on insects of medical importance. *Parasitol Today* 4:S3-S7.