

Hindawi Publishing Corporation
Abstract and Applied Analysis
Volume 2009, Article ID 957475, 15 pages
doi:10.1155/2009/957475

Research Article

Existence and Exponential Stability of Periodic Solution for a Class of Generalized Neural Networks with Arbitrary Delays

Yimin Zhang,¹ Yongkun Li,² and Kuohui Ye²

¹ Department of Mathematics, Zhaotong Teacher's College, Zhaotong, Yunnan 657000, China

² Department of Mathematics, Yunnan University, Kunming, Yunnan 650091, China

Correspondence should be addressed to Yongkun Li, yklic@ynu.edu.cn

Received 4 August 2009; Accepted 1 December 2009

Recommended by Jean Mawhin

By the continuation theorem of coincidence degree and M -matrix theory, we obtain some sufficient conditions for the existence and exponential stability of periodic solutions for a class of generalized neural networks with arbitrary delays, which are milder and less restrictive than those of previous known criteria. Moreover our results generalize and improve many existing ones.

Copyright © 2009 Yimin Zhang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Consider the following generalized neural networks with arbitrary delays:

$$x'(t) = A(t, x(t))[B(t, x(t)) + F(t, x_t)], \quad (1.1)$$

where $A(t, x(t)) = \text{diag}(a_1(t, x_1(t)), a_2(t, x_2(t)), \dots, a_n(t, x_n(t)))$, $B(t, x(t)) = (b_1(t, x_1(t)), b_2(t, x_2(t)), \dots, b_n(t, x_n(t)))^T$, $F(t, x_t) = (f_1(t, x_t), f_2(t, x_t), \dots, f_n(t, x_t))^T$, $f_i(t, x_t) = f_i(t, x_{1t}, x_{2t}, \dots, x_{nt})$, $x_t = (x_{1t}, x_{2t}, \dots, x_{nt})^T$ is defined by $x_i(\theta) = x_i(t + \theta) = (x_1(t + \theta), x_2(t + \theta), \dots, x_n(t + \theta))^T$, $\theta \in E$, and E is a subset of $R^- = (-\infty, 0]$.

System (1.1) contains many neural networks, for examples, the higher-order Cohen-Grossberg type neural networks with delays (see [1])

$$\begin{aligned}
 x'_i(t) = -a_i(x_i(t)) & \left[b_i(x_i(t)) - \sum_{j=1}^n a_{ij}(t)g_j(x_j(t)) - \sum_{j=1}^n b_{ij}(t)g_j(x_j(t - \tau_j(t))) \right. \\
 & \left. - \sum_{j=1}^n \sum_{l=1}^n b_{ijl}(t)g_j(x_j(t - \tau_j(t)))g_l(x_l(t - \tau_l(t))) + I_i(t) \right], \quad (1.2) \\
 & i = 1, 2, \dots, n,
 \end{aligned}$$

the Cohen-Grossberg neural networks with bounded and unbounded delays (see [2])

$$\begin{aligned}
 x'_i(t) = -a_i(t, x_i(t)) & \left[b_i(t, x_i(t)) - \sum_{j=1}^n c_{ij}(t)f_j(x_j(t - \tau_{ij}(t))) \right. \\
 & \left. - \sum_{j=1}^n d_{ij}(t)g_j\left(\int_0^\infty K_{ij}(u)x_j(t - u)du\right) + I_i(t) \right], \quad i = 1, 2, \dots, n, \quad (1.3)
 \end{aligned}$$

the Cohen-Grossberg neural networks with time-varying delays (see [3])

$$\begin{aligned}
 x'_i(t) = -a_i(t, x_i(t)) & \left[b_i(t, x_i(t)) - \sum_{j=1}^n c_{ij}(t)f_j(x_j(t)) \right. \\
 & \left. - \sum_{j=1}^n d_{ij}(t)f_j(x_j(t - \tau_{ij}(t))) + I_i(t) \right], \quad i = 1, 2, \dots, n, \quad (1.4)
 \end{aligned}$$

the cellular neural networks (see [4, Page 193]):

$$\begin{aligned}
 x'_i(t) = -r_i(t)x_i(t) + \sum_{j=1}^n a_{ij}(t)g_j(x_j(t)) \\
 + \sum_{j=1}^n b_{ij}(t)g_j(x_j(t - \tau_{ij}(t))) + I_i(t), \quad i = 1, 2, \dots, n, \quad (1.5)
 \end{aligned}$$

and so on.

Since the model of Cohen-Grossberg neural networks was first introduced by Cohen and Grossberg in [5], the dynamical characteristics (including stable, unstable, and periodic oscillatory) of Cohen-Grossberg neural networks have been widely investigated for the sake of theoretical interest as well as application considerations. Many good results have already been obtained by some authors in [6–15] and the references cited therein. Moreover, the existing results are based on the assumption that demanding either the activation functions, the behaved functions, or delays is bounded in the above-mentioned literature. However, to the best of our knowledge, few authors have discussed the existence and exponential stability of periodic solutions of (1.1). In this paper, by using the continuation theorem of coincidence

degree and M -matrix theory, we study model (1.1), and get some sufficient conditions for the existence and exponential stability of the periodic solution of system (1.1); our results generalize and improve many existing ones.

Let $A = (a_{ij})_{n \times n}$, $B = (b_{ij})_{n \times n} \in R^{n \times n}$ be two matrices, $u = (u_1, u_2, \dots, u_n)^T \in R^n$, $v = (v_1, v_2, \dots, v_n)^T \in R^n$ be two vectors. For convenience, we introduce the following notations.

- (i) $A \geq 0$ ($A > 0$) means that each element a_{ij} is nonnegative (positive) respectively,
- (ii) $A \geq B$ ($> B$) means $A - B \geq 0$ (> 0),
- (iii) $u \geq 0$ ($u > 0$) means each element $u_i \geq 0$ ($u_i > 0$),
- (iv) $u \leq v$ ($u < v$) means $v - u \geq 0$ ($v - u > 0$),
- (v) $|u| = (|u_1|, |u_2|, \dots, |u_n|)^T$.

For continuous ω -periodic function $g : R \rightarrow R$, we denote $\overline{|g|} = \max_{0 \leq t \leq \omega} |g(t)|$, $C_E = C[E, R^n]$ is the family of continuous functions $\phi = (\phi_1, \phi_2, \dots, \phi_n)^T$ from $E \subset (-\infty, 0]$ to R^n . Clearly, it is a Banach space with the norm $\|\phi\| = \max_{0 \leq i \leq n} |\phi_i|$, where $|\phi_i| = \sup_{\theta \in E} |\phi_i(\theta)|$. The initial conditions of system (1.1) are of the form

$$x_0 = \phi, \quad \text{that is, } x_i(\theta) = \phi_i(\theta), \quad \theta \in E, \quad i = 1, 2, \dots, n, \tag{1.6}$$

where $\phi = (\phi_1, \phi_2, \dots, \phi_n)^T \in C_E$. For $V(t) \in C((a, +\infty), R)$, let

$$D^-V(t) = \limsup_{h \rightarrow 0^+} \frac{D(t+h) - D(t)}{h}, \quad D^-V(t) = \liminf_{h \rightarrow 0^+} \frac{D(t+h) - D(t)}{h}, \quad t \in (a, +\infty). \tag{1.7}$$

Throughout this paper, we assume the following:

- (H₁) For $i = 1, 2, \dots, n$, $a_i, b_i \in C[R^2, R]$, $f_i \in C[R \times C_E, R]$ and are ω -periodic for their first arguments, respectively, that is, $a_i(t+\omega, u) = a_i(t, u)$, $b_i(t+\omega, u) = b_i(t, u)$, $f_i(t+\omega, \phi) = f_i(t, \phi)$ so $A(t+\omega, u) = A(t, u)$, $B(t+\omega, u) = B(t, u)$, $F(t+\omega, \phi) = F(t, \phi)$, for all $t \in R$, $u \in R^n$, $\phi \in C_E$.
- (H₂) There exists a positive diagonal matrix $A = \text{diag}(a_1, a_2, \dots, a_n)$ such that $A(t, u) \geq A$, for all $(t, u) \in R^{n+1}$.
- (H₃) There is a positive diagonal matrix $B = \text{diag}(b_1, b_2, \dots, b_n)$ such that $|B(t, u)| \geq B|u|$, and $b_i(t, u_i)u_i > 0$ or $b_i(t, u_i)u_i < 0$ for all $(t, u) \in R^{n+1}$, $i = 1, 2, \dots, n$.
- (H₄) There exist a nonnegative matrix $C = (c_{ij})_{n \times n} \in R^{n \times n}$ and a nonnegative vector $D = (D_1, D_2, \dots, D_n)^T$ such that $|F(t, \phi)| \leq C|\phi| + D$, for all $(t, \phi) \in R \times C_E$, where $\phi = (\phi_1, \phi_2, \dots, \phi_n)^T \in C_E$, $|\phi| = (|\phi_1|, |\phi_2|, \dots, |\phi_n|)^T$.

2. Preliminaries

In this section, we first introduce some definitions and lemmas which play an important role in the proof of our main results in this paper.

Definition 2.1. Let $\tilde{x}(t) = (\tilde{x}_1(t), \tilde{x}_2(t), \dots, \tilde{x}_n(t))^T$ be an ω -periodic solution of system (1.1) with initial value $\tilde{\phi} \in C_E$, if there exist two constants $\alpha > 0$ and $M > 1$ such that for every solution $x(t) = (x_1(t), x_2(t), \dots, x_n(t))^T$ of system (1.1) with initial value (1.6),

$$|x_i(t) - \tilde{x}_i(t)| \leq M \|\phi - \tilde{\phi}\| e^{-\alpha t}, \quad \forall t > 0, \quad i = 1, 2, \dots, n. \quad (2.1)$$

Then $\tilde{x}(t)$ is said to be globally exponential stable.

Definition 2.2. A real matrix $W = (w_{ij})_{n \times n} \in R^{n \times n}$ is said to be an M -matrix if $w_{ij} \leq 0$, $i, j = 1, 2, \dots, n$, $i \neq j$, and $W^{-1} \geq 0$.

Lemma 2.3 (see [15, 16]). *Assume that A is an M -matrix and $Au \leq d$, $u, d \in R^n$, then $u \leq A^{-1}d$.*

Lemma 2.4 (see [15, 16]). *Let $W = (w_{ij})_{n \times n}$ with $w_{ij} \leq 0$, $i, j = 1, 2, \dots, n$, $i \neq j$, then the following statements are equivalent.*

- (i) W is an M -matrix.
- (ii) There exists a positive vector $\eta = (\eta_1, \eta_2, \dots, \eta_n) > 0$ such that $\eta W > 0$.
- (iii) There exists a positive vector $\xi = (\xi_1, \xi_2, \dots, \xi_n)^T > 0$ such that $W\xi > 0$.

Lemma 2.5 (see [15, 16]). *Let $A \geq 0$ be an $n \times n$ matrix and $\rho(A) < 1$, then $(E_n - A)^{-1} \geq 0$, where E_n denotes the identity matrix of size n , so $E_n - A$ is an M -matrix.*

Now we introduce Mawhin's continuation theorem which will be fundamental in this paper.

Lemma 2.6 (see [17]). *Let X and Y be two Banach spaces and L a Fredholm mapping of index zero. Assume that $\Omega \subset X$ is an open bounded set and $N : X \rightarrow Z$ is a continuous operator which is L -compact on $\overline{\Omega}$. Then $Lx = Nx$ has at least one solution in $\text{Dom } L \cap \overline{\Omega}$, if the following conditions are satisfied:*

- (1) $Lx \neq \lambda Nx$, for all $(x, \lambda) \in (\text{Dom } L \cap \partial\Omega) \times (0, 1)$,
- (2) $QNx \neq 0$, for all $x \in \partial\Omega \cap \text{Ker } L$,
- (3) $\text{deg}\{JQN|_{\text{Ker } L \cap \overline{\Omega}}, \Omega \cap \text{Ker } L, 0\} \neq 0$.

Let

$$X = Y = \left\{ x = (x_1, x_2, \dots, x_n)^T \in C(R, R^n) : x(t + \omega) = x(t), t \in R \right\} \quad (2.2)$$

with the norm defined by $\|x\| = \max_{1 \leq i \leq n} \overline{|x_i|}$, where $\overline{|x_i|} = \max_{t \in [0, \omega]} |x_i(t)|$. Clearly, X and Y are two Banach spaces. Let $x = (x_1, x_2, \dots, x_n)^T \in X = Y$, we define the linear operator $L : \text{Dom } L \subset X \rightarrow Y$ as

$$(Lx)(t) = x'(t) = (x'_1(t), x'_2(t), \dots, x'_n(t))^T, \quad \text{Dom } L = \{x \in X : x' \in Y\}, \quad (2.3)$$

and the operators $N : X \rightarrow X$, $P : X \rightarrow X$, $Q : Y \rightarrow Y$ as

$$\begin{aligned} (Nx)(t) &= A(t, x(t))[B(t, x(t)) + F(t, x_t)] := \Delta(t, x_t), \\ Px = Qx &= \frac{1}{\omega} \int_0^\omega x(t) dt \\ &= \left(\frac{1}{\omega} \int_0^\omega x_1(t) dt, \frac{1}{\omega} \int_0^\omega x_2(t) dt, \dots, \frac{1}{\omega} \int_0^\omega x_n(t) dt \right)^T. \end{aligned} \quad (2.4)$$

It is not difficult to show that P and Q are continuous projectors and the following conditions are satisfied:

$$\begin{aligned} \text{Ker } L &= R^n = \text{Im } P = \text{Im } Q, \\ \text{Im } L &= \left\{ y \in Y = X : \int_0^\omega y(t) dt = 0 \right\} = \text{Ker } Q = \text{Im } (I - Q), \\ \text{Im } L &\text{ is closed in } Y, \quad \dim \text{Ker } L = n = \text{codim Im } L. \end{aligned} \quad (2.5)$$

Thus, the mapping L is a Fredholm mapping of index zero and the isomorphism $J : \text{Im } Q \rightarrow \text{Ker } L$ is the identity operator; the generalized inverse (of $L|_{\text{Dom } L \cap \text{Ker } P}$) $K_P : \text{Im } L \rightarrow \text{Ker } P \cap \text{Dom } L$ exists, which has the form

$$(K_P x)(t) = \int_0^t x(s) ds - \frac{1}{\omega} \int_0^\omega \int_0^t x(s) ds dt, \quad \forall x \in \text{Im } L. \quad (2.6)$$

Therefore

$$\begin{aligned} (QNx)(t) &= \frac{1}{\omega} \int_0^\omega \Delta(t, x_t) dt, \\ (K_P(I - Q)Nx)(t) &= \int_0^t \Delta(s, x_s) ds - \frac{1}{\omega} \int_0^\omega \int_0^t \Delta(s, x_s) ds dt + \left(\frac{1}{2} - \frac{t}{\omega} \right) \int_0^\omega \Delta(s, x_s) ds. \end{aligned} \quad (2.7)$$

3. Existence of Periodic Solutions

In this section, we shall use Lemma 2.6 to study the existence of at least one periodic solution of system (1.1).

Theorem 3.1. *Let (H_1) – (H_4) hold. Moreover, suppose that*

$$(H_5) \quad E - K \text{ is a } M\text{-matrix, where the matrix } K = (k_{ij})_{n \times n} = B^{-1}C.$$

then

- (i) system (1.1) has at least one ω -periodic solution;
- (ii) there exists a nonnegative constant δ such that for all ω -periodic solution $x(t) = (x_1(t), x_2(t), \dots, x_n(t))^T$ of system (1.1), $|x_i(t)| \leq \delta$, $i = 1, 2, \dots, n$.

Proof. Clearly, QN and $K_P(I - Q)N$ are continuous functions and for every bounded subset $\Omega \subset X$, $QN(\overline{\Omega})$, $K_P(I - Q)N(\overline{\Omega})$, and $(K_P(I - Q)Nx)'$, $x \in \overline{\Omega}$ are bounded. By using the Arzela-Ascoli theorem, $QN(\overline{\Omega})$ and $K_P(I - Q)N(\overline{\Omega})$ are compact, therefore N is L -compact on $\overline{\Omega}$. Consider the following operator equation:

$$Lx = \lambda Nx, \quad \lambda \in (0, 1). \quad (3.1)$$

That is,

$$x'(t) = \lambda A(t, x(t))[B(t, x(t)) + F(t, x_t)], \quad (3.2)$$

or

$$x'_i(t) = \lambda a_i(t, x_i(t))[b_i(t, x_i(t)) + f_i(t, x_t)], \quad i = 1, 2, \dots, n. \quad (3.3)$$

Assume that $x(t) = (x_1(t), x_2(t), \dots, x_n(t))^T \in X$ is a solution of (3.3) for some $\lambda \in (0, 1)$. Then, for any $i = 1, 2, \dots, n$, $x_i(t)$ are all continuous ω -periodic functions, and there exist $t_i \in [0, \omega]$, such that

$$|x_i(t_i)| = \max_{t \in [0, \omega]} |x_i(t)| = \overline{|x_i|}, \quad x'_i(t_i) = 0, \quad i = 1, 2, \dots, n, \quad (3.4)$$

from (H_2) , we have

$$b_i(t_i, x_i(t_i)) + f_i(t_i, x_{t_i}) = 0, \quad i = 1, 2, \dots, n. \quad (3.5)$$

It follows from (H_3) that

$$\begin{aligned} b_i \overline{|x_i|} &\leq |b_i(t_i, x_i(t_i))| = |f_i(t_i, x_{t_i})| \\ &\leq \sum_{j=1}^n c_{ij} |x_{jt_i}| + D_i \leq \sum_{j=1}^n c_{ij} \overline{|x_j|} + D_i, \quad i = 1, 2, \dots, n. \end{aligned} \quad (3.6)$$

Thus

$$\overline{|x_i|} \leq b_i^{-1} \sum_{j=1}^n c_{ij} \overline{|x_j|} + b_i^{-1} D_i, \quad i = 1, 2, \dots, n, \quad (3.7)$$

we denote the vector $d = (d_1, d_2, \dots, d_n)^T$, $\overline{|x|} = (\overline{|x_1|}, \overline{|x_2|}, \dots, \overline{|x_n|})^T$, where $d_i = b_i^{-1}(D_i + 1) > 0$, $i = 1, 2, \dots, n$. It follows from (3.7) that

$$(E - K)\overline{|x|} < d. \quad (3.8)$$

Since (H_5) , and application of Lemma 2.3 yields

$$\overline{|x|} < (E - K)^{-1}d = (h_1, h_2, \dots, h_n)^T = h, \quad (3.9)$$

where h satisfies the equation $h = Kh + d$, that is, $h_i = \sum_{j=1}^n k_{ij}h_j + d_i > 0$.

Take

$$\Omega = \left\{ x \in X : \overline{|x_i|} < h_i, \quad i = 1, 2, \dots, n \right\}. \quad (3.10)$$

It is easy to see that Ω satisfies condition (1) in Lemma 2.6.

For all $x = (x_1, x_2, \dots, x_n)^T \in \partial\Omega \cap \text{Ker } L$, x is a constant vector in R^n and there exists some $i \in \{1, 2, \dots, n\}$ such that $|x_i| = \overline{|x_i|} = h_i$, we claim that

$$|(QNx)_i| > 0, \quad \text{so that } QNx \neq 0. \quad (3.11)$$

We firstly claim that

- (1) if $b_i(t, u_i)u_i > 0$, then $x_i(QNx)_i > 0$,
- (2) if $b_i(t, u_i)u_i < 0$, then $x_i(QNx)_i < 0$.

We only prove (1), since the proof of (2) is similar. If $b_i(t, u_i)u_i > 0$, we have

$$\begin{aligned} x_i [b_i(t, x_i(t)) + f_i(t, x_t)] &\geq b_i x_i^2 - |x_i| \left[\sum_{j=1}^n c_{ij} \overline{|x_j|} + D_i \right] \\ &> b_i h_i \left[h_i - \left(\sum_{j=1}^n b_i^{-1} c_{ij} h_j + d_i \right) \right] \\ &= b_i h_i \left[h_i - \left(\sum_{j=1}^n k_{ij} h_j + d_i \right) \right] = 0. \end{aligned} \quad (3.12)$$

Therefore

$$x_i(QNx)_i = \omega^{-1} x_i \int_0^\omega a_i(t, x_i(t)) [b_i(t, x_i(t)) + f_i(t, x_t)] dt > 0. \quad (3.13)$$

Thus (3.11) is valid.

Next, we define continuous functions $H_i : (\Omega \cap \text{Ker } L) \times [0, 1] \rightarrow \Omega \cap \text{Ker } L$, $i = 1, 2$, by

$$\begin{aligned} H_1(x, t) &= tx + (1 - t)QNx, \quad \forall (x, t) \in (\Omega \cap \text{Ker } L) \times [0, 1], \\ H_2(x, t) &= -tx + (1 - t)QNx, \quad \forall (x, t) \in (\Omega \cap \text{Ker } L) \times [0, 1], \end{aligned} \quad (3.14)$$

respectively. If $b_i(t, u_i)u_i > 0$, from (i) we have

$$H_1(x, t) \neq 0, \quad \forall (x, t) \in \text{Ker } L \cap \partial\Omega \times [0, 1], \quad (3.15)$$

If $b_i(t, u_i)u_i < 0$, from (2) we can get

$$H_2(x, t) \neq 0, \quad \forall (x, t) \in \text{Ker } L \cap \partial\Omega \times [0, 1]. \quad (3.16)$$

Using the homotopy invariance theorem, we obtain if $b_i(t, u_i)u_i > 0$,

$$\begin{aligned} \deg\{JQN|_{\overline{\Omega \cap \text{Ker } L}}, \Omega \cap \text{Ker } L, 0\} &= \deg\{H_1(\cdot, 0), \Omega \cap \text{Ker } L, 0\} \\ &= \deg\{H_1(\cdot, 1), \Omega \cap \text{Ker } L, 0\} \\ &= \deg\{x, \Omega \cap \text{Ker } L, 0\} = 1, \end{aligned} \quad (3.17)$$

or if $b_i(t, u_i)u_i < 0$,

$$\begin{aligned} \deg\{JQN|_{\overline{\Omega \cap \text{Ker } L}}, \Omega \cap \text{Ker } L, 0\} &= \deg\{H_2(\cdot, 0), \Omega \cap \text{Ker } L, 0\} \\ &= \deg\{H_2(\cdot, 1), \Omega \cap \text{Ker } L, 0\} \\ &= \deg\{-x, \Omega \cap \text{Ker } L, 0\} = (-1)^n. \end{aligned} \quad (3.18)$$

To summarize, Ω satisfies all the conditions of Lemma 2.6. This completes the proof of (i).

For all ω -periodic solution $x(t) = (x_1(t), x_2(t), \dots, x_n(t))^T$ of system (1.1), from (3.3)–(3.7) we have

$$\begin{aligned} \overline{|x_i|} &\leq b_i^{-1} \sum_{j=1}^n c_{ij} \overline{|x_j|} + b_i^{-1} D_i, \\ \overline{|x|} &\leq (E - K)^{-1} v = v = (v_1, v_2, \dots, v_n)^T, \end{aligned} \quad (3.19)$$

where $v = (v_1, v_2, \dots, v_n)^T$, $\overline{|x|} = (\overline{|x_1|}, \overline{|x_2|}, \dots, \overline{|x_n|})^T$, $v_i = b_i^{-1} D_i \geq 0$, $i = 1, 2, \dots, n$. Notes $\delta = \max_{1 \leq i \leq n} v_i \geq 0$, thus $|x_i(t)| \leq \delta$, for all $i = 1, 2, \dots, n$. This completes the proof of (ii). \square

From the proof of Theorem 3.1, we can easily obtain the following corollary.

Corollary 3.2. *Suppose that (H_1) – (H_5) hold, and $D = 0$ in (H_4) , then system (1.1) has only one ω -periodic solution $x(t) = 0$.*

Some special cases of Theorem 3.1 are in what follows.

Corollary 3.3. *Equation (1.3) has at least one ω -periodic solution, if the following conditions are satisfied.*

(A₁) For $i, j = 1, 2, \dots, n$, $a_i, b_i, a_{ij}, b_{ij}, \tau_j, I_i : \mathbb{R} \rightarrow \mathbb{R}$ are continuous ω -periodic ($\omega > 0$) functions.

(A₂) For $i = 1, 2, \dots, n$, $a_i(x)$ are positive, and there exist $a_i > 0$ such that $a_i(x) \geq a_i > 0$.

(A₃) For $i = 1, 2, \dots, n$, there exist $b_i > 0$ such that

$$|b_i(x)| \geq b_i|x|, \quad b_i(x)x > 0, \quad \text{or} \quad b_i(x)x < 0, \quad \forall x \in \mathbb{R}. \quad (3.20)$$

(A₄) For $i = 1, 2, \dots, n$, there exist $G_i, p_i, q_i \geq 0$ such that

$$|g_i(x)| \leq G_i, \quad |g_i(x)| \leq p_i|x| + q_i. \quad (3.21)$$

(A₅) $\rho(K) < 1$, $K = (k_{ij})_{n \times n}$, and $k_{ij} = b_i^{-1}(\overline{|a_{ij}|} + \overline{|b_{ij}|} + \sum_{l=1}^n \overline{|b_{ijl}|} G_l) p_j$, $i, j = 1, 2, \dots, n$.

Proof. It is clear that

$$\begin{aligned} A(t, x) &= \text{diag}(a_1(x_1), a_2(x_2), \dots, a_n(x_n)) \\ &\geq \text{diag}(a_1, a_2, \dots, a_n) = A, \\ |B(t, x)| &= (|b_1(x_1)|, |b_2(x_2)|, \dots, |b_n(x_n)|)^T \\ &\geq (b_1|x_1|, b_2|x_2|, \dots, b_n|x_n|)^T \\ &= \text{diag}(b_1, b_2, \dots, b_n)(|x_1|, |x_2|, \dots, |x_n|)^T = B|x|, \\ |f_i(t, \phi)| &= \left| \sum_{j=1}^n a_{ij}(t) g_j(\phi_j(0)) + \sum_{j=1}^n b_{ij}(t) g_j(\phi_j(-\tau_j(t))) \right. \\ &\quad \left. + \sum_{j=1}^n \sum_{l=1}^n b_{ijl}(t) g_j(\phi_j(-\tau_j(t))) g_l(\phi_l(-\tau_l(t))) - I_i(t) \right| \\ &\leq \sum_{j=1}^n \overline{|a_{ij}|} [p_j|\phi_j| + q_j] + \sum_{j=1}^n \overline{|b_{ij}|} [p_j|\phi_j| + q_j] \\ &\quad + \sum_{j=1}^n \sum_{l=1}^n G_l \overline{|b_{ijl}|} (p_j|\phi_j| + q_j) + \overline{|I_i|} \\ &= \sum_{j=1}^n \left[\overline{|a_{ij}|} + \overline{|b_{ij}|} + \sum_{l=1}^n \overline{|b_{ijl}|} G_l \right] p_j |\phi_j| \\ &\quad + \sum_{j=1}^n \left[\overline{|a_{ij}|} + \overline{|b_{ij}|} + \sum_{l=1}^n \overline{|b_{ijl}|} G_l \right] q_j + \overline{|I_i|} = \sum_{j=1}^n c_{ij} |\phi_j| + D_i, \quad i = 1, 2, \dots, n. \end{aligned} \quad (3.22)$$

Thus

$$|f(t, \phi)| \leq C|\phi| + D, \quad (3.23)$$

where $C = (c_{ij})_{n \times n} \in R^{n \times n}$, $D = (D_1, D_2, \dots, D_n)^T$, $c_{ij} = [|\overline{a_{ij}}| + |\overline{b_{ij}}| + \sum_{l=1}^n |\overline{b_{ijl}}| G_l] p_j \geq 0$, $D_i = \sum_{j=1}^n [|\overline{a_{ij}}| + |\overline{b_{ij}}| + \sum_{l=1}^n |\overline{b_{ijl}}| G_l] q_j + |\overline{I_i}| \geq 0$, $i, j = 1, 2, \dots, n$. Therefore, by using Lemma 2.5 and Theorem 3.1, we know that (1.3) has an ω -periodic solution. The proof is complete. \square

Remark 3.4. For [1, Equation (1.2)], $\tau_j(t), j = 1, 2, \dots, n$ are continuous differentiable ω -periodic solutions and $0 \leq \tau'_j(t) \leq 1$, this implies that $\tau_j(t), j = 1, 2, \dots, n$ are constant functions, thus $\xi_j = 1, j = 1, 2, \dots, n$. It is not difficult to verify that all of conditions of Corollary 3.3 are satisfied under the conditions of [1, Theorem 1] moreover the other requirements of [1, Theorem 1] are more restrictive than ours. Therefore, Corollary 3.3 improves the corresponding result obtained in [1].

Corollary 3.5. *If the following conditions are satisfied:*

- (B₁) for $i, j = 1, 2, \dots, n$, $c_{ij}, d_{ij}, \tau_{ij}, I_i : R \rightarrow R$ are continuous ω -periodic ($\omega > 0$) functions, a_i, b_i are continuous functions on R^2 , and are ω -periodic for their first arguments, respectively,
- (B₂) for $i = 1, 2, \dots, n$, there exist positive constants a_i such that $a_i(t, u) \geq a_i$, for all $t, u \in R$,
- (B₃) for $i = 1, 2, \dots, n$, there exist positive constants b_i such that $|b_i(t, u)| \geq b_i|u|$, $b_i(u)u > 0$ or $b_i(u)u < 0$, for all $t, u \in R$,
- (B₄) there exist nonnegative constants $p_j^f, q_j^f, p_j^g, q_j^g$ such that

$$|f_j(u)| \leq p_j^f |u| + q_j^f, \quad |g_j(u)| \leq p_j^g |u| + q_j^g, \quad \forall u \in R, \quad j = 1, 2, \dots, n, \quad (3.24)$$

- (B₅) the delay kernels $K_{ij} : [0, \infty] \rightarrow R$ satisfy

$$\int_0^\infty |K_{ij}(s)| ds \leq k_{ij}, \quad i, j = 1, 2, \dots, n, \quad (3.25)$$

- (B₆) $\rho(K) < 1$, $K = (k_{ij})_{n \times n} \in R^{n \times n}$, where $k_{ij} = b_i^{-1}(|\overline{c_{ij}}| p_j^f + |\overline{d_{ij}}| k_{ij} p_j^g)$, $i, j = 1, 2, \dots, n$.

then (1.3) has at least one ω -periodic solution.

Remark 3.6. In [2, Theorem 3.1], the activation functions $f_j(u), g_j(u), j = 1, 2, \dots, n$, are required to be Lipschitzian, which implies that condition (B₃) in Corollary 3.5 holds. Therefore, Corollary 3.5 improves Theorem 3. In 2.

Corollary 3.7. *Assume that the following conditions are satisfied:*

- (C₁) $c_{ij}, d_{ij}, \tau_{ij}, I_i : R \rightarrow R$ are continuous ω -periodic ($\omega > 0$) functions, a_i, b_i are continuous functions on R^2 , and are ω -periodic in the first variable,
- (C₂) there exist positive constants a_i such that

$$a_i(t, u) \geq a_i, \quad \forall t, u \in R, \quad i = 1, 2, \dots, n, \quad (3.26)$$

(C₃) there exist positive constants b_i such that

$$|b_i(t, u)| \geq b_i|u|, \quad b_i(u)u > 0 \quad \text{or} \quad b_i(u)u < 0, \quad \forall t, u \in \mathbb{R}, \quad i = 1, 2, \dots, n, \quad (3.27)$$

(C₄) There exist nonnegative constants p_i, q_i such that

$$|f_i(u)| \leq p_i|u| + q_i, \quad \forall u \in \mathbb{R}, \quad i = 1, 2, \dots, n, \quad (3.28)$$

(C₅) $\rho(K) < 1$, where $K = (k_{ij})_{n \times n} \in \mathbb{R}^{n \times n}$ and $k_{ij} = b_i^{-1}(\overline{|c_{ij}|} + \overline{|d_{ij}|})p_j$, $i, j = 1, 2, \dots, n$.

Then (1.4) has at least one ω -periodic solution.

Remark 3.8. In [3, Theorem 3.1], the activation functions $f_j(u)$, $j = 1, 2, \dots, n$, are Lipschitzian (which also implies that condition (C₄) in Corollary 3.7 holds) and the behaved functions $b_i(t, u)$ are required to satisfy that there exist positive constants $\underline{b}_i, \bar{b}_i$ such that $0 \leq ub_i(t, u)$, $\underline{b}_i|u| \leq |b_i(t, u)| \leq \bar{b}_i|u|$ for all $t, u \in \mathbb{R}$, $i = 1, 2, \dots, n$, which are more restrictive than that of Corollary 3.7.

Corollary 3.9. Assume that the following conditions are satisfied

(D₁) For $i, j = 1, 2, \dots, n$, $I_i, a_{ij}, b_{ij}, \tau_{ij} : \mathbb{R} \rightarrow \mathbb{R}$ are continuous ω -periodic solution ($\omega > 0$) functions.

(D₂) For $j = 1, 2, \dots, n$, $g_j : \mathbb{R} \rightarrow \mathbb{R}$ are continuous functions and there exist nonnegative constants p_j, q_j such that

$$|g_j(v)| \leq p_j|v| + q_j, \quad \forall v \in \mathbb{R}, \quad j = 1, 2, \dots, n, \quad (3.29)$$

(D₃) $\rho(K) < 1$, $K = (k_{ij})_{n \times n} \in \mathbb{R}^{n \times n}$ and $k_{ij} = r_i^{-1}(\overline{|a_{ij}|} + \overline{|b_{ij}|})p_j$, $i, j = 1, 2, \dots, n$,

then (1.5) has at least one ω -periodic solution.

The proofs of Corollaries 3.5–3.9 are the same as that of Corollary 3.3.

4. Uniqueness and Exponential Stability of Periodic Solution

In this section, we establish some results for the uniqueness and exponential stability of the ω -periodic solution of (1.1).

Theorem 4.1. Assume that E is a bounded subset of \mathbb{R}^- , and (H₁)–(H₃) and (H₅) hold. Suppose also the following conditions are satisfied.

(H₄)' There exists a nonnegative matrix $C = (c_{ij})_{n \times n} \in \mathbb{R}^{n \times n}$ such that

$$|F(t, \phi) - F(t, \varphi)| \leq C|\phi - \varphi|, \quad \forall (t, \phi), (t, \varphi) \in \mathbb{R} \times C_E, \quad (4.1)$$

where $\phi = (\phi_1, \phi_2, \dots, \phi_n)^T$, $\varphi = (\varphi_1, \varphi_2, \dots, \varphi_n)^T \in C_E$, $|\phi - \varphi| = (|\phi_1 - \varphi_1|, |\phi_2 - \varphi_2|, \dots, |\phi_n - \varphi_n|)^T$.

(H₆) a_i , $i = 1, 2, \dots, n$, are Lipschitzian with Lipschitz constants $L_i^a > 0$, and there exist \bar{a}_i such that

$$a_i(t, u) \leq \bar{a}_i, \quad |a_i(t, u) - a_i(t, v)| \leq L_i^a |u - v|, \quad \forall (t, u), (t, v) \in \mathbb{R}^2, \quad i = 1, 2, \dots, n. \quad (4.2)$$

(H₇) For all $t, u, v \in \mathbb{R}$, $i = 1, 2, \dots, n$, there exist positive constants L_i^{ab} such that

$$\begin{aligned} [a_i(t, u)b_i(t, u) - a_i(t, v)b_i(t, v)](u - v) &\leq 0, \quad i = 1, 2, \dots, n, \\ |a_i(t, u)b_i(t, u) - a_i(t, v)b_i(t, v)| &\geq L_i^{ab} |u - v|, \quad i = 1, 2, \dots, n. \end{aligned} \quad (4.3)$$

(H₈) For $i = 1, 2, \dots, n$, set $\Delta_i = \max_{0 \leq t \leq \omega} |f_i(t, 0)|$, and assume that $E_n - W$ is an M-matrix, where $W = (w_{ij})_{n \times n} \in \mathbb{R}^{n \times n}$, and

$$w_{ij} = \left(L_i^{ab} - L_i^a \Delta_i \right)^{-1} (\bar{a}_i + L_i^a \delta) c_{ij}, \quad L_i^{ab} - L_i^a \Delta_i > 0, \quad i, j = 1, 2, \dots, n. \quad (4.4)$$

Proof. Obviously, (H₄)' implies (H₄), since (H₁)–(H₅) hold, it follows from Theorem 3.1 that system (1.1) has at least one ω -periodic solution

$$\tilde{x}(t) = (\tilde{x}_1(t), \tilde{x}_2(t), \dots, \tilde{x}_n(t))^T \quad (4.5)$$

with the initial value $\tilde{\phi} = (\tilde{\phi}_1, \tilde{\phi}_2, \dots, \tilde{\phi}_n)^T \in C_E$. Let

$$x(t) = (x_1(t), x_2(t), \dots, x_n(t))^T \quad (4.6)$$

be an arbitrary solution of system (1.1) with the initial value (1.6), set $y(t) = x(t) - \tilde{x}(t)$. Then for $i = 1, 2, \dots, n$,

$$\begin{aligned} y_i'(t) &= a_i(t, y_i(t) + \tilde{x}_i(t))b_i(t, y_i(t) + \tilde{x}_i(t)) - a_i(t, \tilde{x}_i(t))b_i(t, \tilde{x}_i(t)) \\ &\quad + a_i(t, y_i(t) + \tilde{x}_i(t)) [f_i(t, y_t + \tilde{x}_t) - f_i(t, \tilde{x}_t)] \\ &\quad + f_i(t, \tilde{x}_t) [a_i(t, x_t) - a_i(t, \tilde{x}_t)]. \end{aligned} \quad (4.7)$$

Thus, for $i = 1, 2, \dots, n$,

$$\begin{aligned} D^- |y_i(t)| &\leq -L_i^{ab} |y_i(t)| + \bar{a}_i \sum_{j=1}^n c_{ij} |y_{jt}| + L_i^a |y_i(t)| \left[\sum_{j=1}^n c_{ij} |\tilde{x}_{jt}| + |f_i(t, 0)| \right] \\ &\leq -\left(L_i^{ab} - L_i^a \Delta_i \right) |y_i(t)| + (\bar{a}_i + L_i^a \delta) \sum_{j=1}^n c_{ij} |y_{jt}|, \end{aligned} \quad (4.8)$$

for (H_8) and Lemma 2.4, there exist a positive constant $\sigma > 0$ and a positive constant vector $\xi = (\xi_1, \xi_2, \dots, \xi_n)^T > 0$ such that $(E_n - W)\xi > (\sigma, \sigma, \dots, \sigma)^T$. Hence

$$\xi_i - \sum_{j=1}^n w_{ij}\xi_j > \sigma, \tag{4.9}$$

where $w_{ij} = (L_i^{ab} - L_i^a \Delta_i)^{-1}(\bar{a}_i + L_i^a \delta)c_{ij}$, $i, j = 1, 2, \dots, n$. Moreover for all $i = 1, 2, \dots, n$,

$$-\left(L_i^{ab} - L_i^a \Delta_i\right)\xi_i + (\bar{a}_i + L_i^a \delta)\sum_{j=1}^n c_{ij}\xi_j < \left(L_i^{ab} - L_i^a \Delta_i\right)\sigma. \tag{4.10}$$

Since, E is a bounded subset of R^- , we can choose a positive constant $\alpha < 1$, such that $\forall \theta \in E$

$$\alpha\xi_i + \left[-\left(L_i^{ab} - L_i^a \Delta_i\right)\xi_i + (\bar{a}_i + L_i^a \delta)\sum_{j=1}^n c_{ij}\xi_j e^{-\alpha\theta}\right] < 0, \quad i = 1, 2, \dots, n, \tag{4.11}$$

and also can choose a positive constant $\beta > 1$ such that

$$\beta\xi_i e^{-\alpha\theta} > 1, \quad \forall \theta \in E, \quad i = 1, 2, \dots, n. \tag{4.12}$$

Set, for all $\varepsilon > 0$, for all $t \in E$,

$$Z_i(t) = \beta\xi_i \left[\sum_{j=1}^n |y_{j0}| + \varepsilon \right] e^{-\alpha t}, \quad i = 1, 2, \dots, n. \tag{4.13}$$

It follows from (4.11) and (4.13) that

$$\begin{aligned} D_- Z_i(t) &= -\alpha\beta\xi_i \left[\sum_{j=1}^n |y_{j0}| + \varepsilon \right] e^{-\alpha t} \\ &> \left[-\left(L_i^{ab} - L_i^a \Delta_i\right)\xi_i + (\bar{a}_i + L_i^a \delta)\sum_{j=1}^n c_{ij}\xi_j e^{-\alpha\theta} \right] \beta \left[\sum_{j=1}^n |y_{j0}| + \varepsilon \right] e^{-\alpha t} \\ &= -\left(L_i^{ab} - L_i^a \Delta_i\right)\xi_i \beta \left[\sum_{j=1}^n |y_{j0}| + \varepsilon \right] e^{-\alpha t} \\ &\quad + (\bar{a}_i + L_i^a \delta)\sum_{j=1}^n c_{ij}\xi_j e^{-\alpha(\theta+t)} \beta \left[\sum_{j=1}^n |y_{j0}| + \varepsilon \right], \quad \forall \theta \in E. \end{aligned} \tag{4.14}$$

Thus

$$D_- Z_i(t) \geq -\left(L_i^{ab} - L_i^a \Delta_i\right) Z_i(t) + (\bar{a}_i + L_i^a \delta) \sum_{j=1}^n c_{ij} |Z_{jt}|, \quad (4.15)$$

where $|Z_{jt}| = \sup_{\theta \in E} Z_j(t + \theta)$, from (4.12) and (4.13), we can get

$$Z_i(t) = \beta \xi_i \left[\sum_{j=1}^n |y_{j0}| + \varepsilon \right] e^{-at} > \sum_{j=1}^n |y_{j0}| + \varepsilon > |y_i(t)|, \quad \forall t \in E. \quad (4.16)$$

We claim that

$$|y_i(t)| < Z_i(t), \quad \forall t > 0, \quad i = 1, 2, \dots, n. \quad (4.17)$$

Suppose that it is not true, then there exists some $i \in \{1, 2, \dots, n\}$ and $t_i > 0$ such that

$$|y_i(t_i)| = Z_i(t_i), \quad |y_j(t)| \leq Z_j(t), \quad \forall t < t_i, \quad j = 1, 2, \dots, n. \quad (4.18)$$

Thus

$$\begin{aligned} 0 &\leq D^- (|y_i(t_i)| - Z_i(t_i)) \\ &= \limsup_{h \rightarrow 0^-} \frac{[|y_i(t_i + h)| - Z_i(t_i + h)] - [|y_i(t_i)| - Z_i(t_i)]}{h} \\ &\leq \limsup_{h \rightarrow 0^-} \frac{|y_i(t_i + h)| - |y_i(t_i)|}{h} - \liminf_{h \rightarrow 0^-} \frac{Z_i(t_i + h) - Z_i(t_i)}{h} \\ &\leq D^- |y_i(t_i)| - D_- Z_i(t_i). \end{aligned} \quad (4.19)$$

It follows from (4.8), (4.15), and (4.18) that

$$\begin{aligned} D^- |y_i(t_i)| &\leq -\left(L_i^{ab} - L_i^a \Delta_i\right) |y_i(t_i)| + (\bar{a}_i + L_i^a \delta) \sum_{j=1}^n c_{ij} |y_{jt_i}| \\ &\leq -\left(L_i^{ab} - L_i^a \Delta_i\right) |Z_i(t_i)| + (\bar{a}_i + L_i^a \delta) \sum_{j=1}^n c_{ij} |Z_{jt_i}| < D_- Z_i(t_i), \end{aligned} \quad (4.20)$$

which contradicts to (4.19), thus (4.17) holds. Set $\varepsilon \rightarrow 0^+$ and $M = n \max_{1 \leq i \leq n} \{\beta \xi_i + 1\} > 1$, from (4.17), we have

$$|x_i(t) - \tilde{x}_i(t)| = |y_i(t)| \leq \beta \xi_i \sum_{j=1}^n |y_{j0}| e^{-at} \leq \beta \xi_i n \|\phi - \tilde{\phi}\| e^{-at} \leq M \|\phi - \tilde{\phi}\| e^{-at}, \quad (4.21)$$

where $i = 1, 2, \dots, n$. This completes the proof of Theorem 4.1. \square

5. Conclusion

In this paper, a class of generalized neural networks with arbitrary delays have been studied. Some sufficient conditions for the existence and exponential stability of the periodic solutions have been established. These obtained results are new and they improve and complement previously known results.

Acknowledgment

This work is supported by the National Natural Sciences Foundation of China under Grant 10971183.

References

- [1] F. Ren and J. Cao, "Periodic solutions for a class of higher-order Cohen-Grossberg type neural networks with delays," *Computers & Mathematics with Applications*, vol. 54, no. 6, pp. 826–839, 2007.
- [2] F. Zhang, B. Liu, and L. Huang, "Existence and exponential stability of periodic solutions for a class of Cohen-Grossberg neural networks with bounded and unbounded delays," *Computers & Mathematics with Applications*, vol. 53, no. 9, pp. 1325–1338, 2007.
- [3] B. Liu and L. Huang, "Existence and exponential stability of periodic solutions for a class of Cohen-Grossberg neural networks with time-varying delays," *Chaos, Solitons & Fractals*, vol. 32, no. 2, pp. 617–627, 2007.
- [4] H. Hong and M. Li, *Dynamics in Celler Neural Network*, Academic Press, Beijing, China, 2007.
- [5] M. A. Cohen and S. Grossberg, "Absolute stability of global pattern formation and parallel memory storage by competitive neural networks," *IEEE Transactions on Systems, Man, and Cybernetics*, vol. 13, no. 5, pp. 815–826, 1983.
- [6] K. Gopalsamy and X. Z. He, "Delay-independent stability in bidirection associative memory networks," *IEEE Transactions on Neural Networks*, vol. 5, pp. 998–1002, 1994.
- [7] T. Chen and L. Rong, "Delay-independent stability analysis of Cohen-Grossberg neural networks," *Physics Letters A*, vol. 317, no. 5-6, pp. 436–449, 2003.
- [8] Y. Li, "Existence and stability of periodic solutions for Cohen-Grossberg neural networks with multiple delays," *Chaos, Solitons & Fractals*, vol. 20, no. 3, pp. 459–466, 2004.
- [9] X. Liao, C. Li, and K.-W. Wong, "Criteria for exponential stability of Cohen-Grossberg neural networks," *Neural Networks*, vol. 17, no. 10, pp. 1401–1414, 2004.
- [10] Z. Yuan, L. Huang, D. Hu, and G. Dong, "Existence and global exponential stability of periodic solution for Cohen-Grossberg neural networks with delays," *Nonlinear Analysis: Real World Applications*, vol. 7, no. 4, pp. 572–590, 2006.
- [11] L. Wang, "Stability of Cohen-Grossberg neural networks with distributed delays," *Applied Mathematics and Computation*, vol. 160, no. 1, pp. 93–110, 2005.
- [12] Y. Li and X. Fan, "Existence and globally exponential stability of almost periodic solution for Cohen-Grossberg BAM neural networks with variable coefficients," *Applied Mathematical Modelling*, vol. 33, no. 4, pp. 2114–2120, 2009.
- [13] C. Li and S. Yang, "Existence and attractivity of periodic solutions to non-autonomous Cohen-Grossberg neural networks with time delays," *Chaos, Solitons & Fractals*, vol. 41, no. 3, pp. 1235–1244, 2009.
- [14] Z. Yuan and L. Yuan, "Existence and global convergence of periodic solution of delayed neural networks," *Mathematical and Computer Modelling*, vol. 48, no. 1-2, pp. 101–113, 2008.
- [15] X. H. Tang and X. Zou, "The existence and global exponential stability of a periodic solution of a class of delay differential equations," *Nonlinearity*, vol. 22, no. 10, pp. 2423–2442, 2009.
- [16] J. P. LaSalle, *The Stability of Dynamical Systems*, SIAM, Philadelphia, Pa, USA, 1976.
- [17] A. Berman and R. J. Plemmons, *Nonnegative Matrices in the Mathematical Sciences*, Computer Science and Applied Mathematic, Academic Press, New York, NY, USA, 1979.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

