



University  
of Glasgow

Veltman, D.M., Lemieux, M.G., Knecht, D.A., and Insall, R.H. (2014) PIP3-dependent macropinocytosis is incompatible with chemotaxis. *Journal of Cell Biology*, 204 (4). pp. 497-505. ISSN 0021-9525

Copyright © 2014 The Authors

<http://eprints.gla.ac.uk/92810>

Deposited on: 28 March 2014

Enlighten – Research publications by members of the University of Glasgow\_  
<http://eprints.gla.ac.uk>

# PIP<sub>3</sub>-dependent macropinocytosis is incompatible with chemotaxis

Douwe M. Veltman,<sup>1</sup> Michael G. Lemieux,<sup>2</sup> David A. Knecht,<sup>2</sup> and Robert H. Insall<sup>1</sup>

<sup>1</sup>Beatson Institute for Cancer Research, Glasgow G61 1BD, Scotland, UK

<sup>2</sup>Department of Molecular and Cell Biology, University of Connecticut, Storrs, CT 06269

In eukaryotic chemotaxis, the mechanisms connecting external signals to the motile apparatus remain unclear. The role of the lipid phosphatidylinositol 3,4,5-trisphosphate (PIP<sub>3</sub>) has been particularly controversial. PIP<sub>3</sub> has many cellular roles, notably in growth control and macropinocytosis as well as cell motility. Here we show that PIP<sub>3</sub> is not only unnecessary for *Dictyostelium discoideum* to migrate toward folate, but actively inhibits chemotaxis. We find that macropinosomes, but not pseudopods, in growing cells are dependent on PIP<sub>3</sub>. PIP<sub>3</sub> patches in these cells show no directional bias, and overall only PIP<sub>3</sub>-free pseudopods orient up-gradient. The pseudopod driver

suppressor of cAR mutations (SCAR)/WASP and verprolin homologue (WAVE) is not recruited to the center of PIP<sub>3</sub> patches, just the edges, where it causes macropinosome formation. Wild-type cells, unlike the widely used axenic mutants, show little macropinocytosis and few large PIP<sub>3</sub> patches, but migrate more efficiently toward folate. Tellingly, folate chemotaxis in axenic cells is rescued by knocking out phosphatidylinositide 3-kinases (PI 3-kinases). Thus PIP<sub>3</sub> promotes macropinocytosis and interferes with pseudopod orientation during chemotaxis of growing cells.

## Introduction

Eukaryotic cells sense chemoattractants using transmembrane receptors. When receptors detect attractants, they activate intracellular second messengers that relay information to the molecules that drive cell movement. One principal group of attractants, including cAMP in *Dictyostelium discoideum* and fMLP in mammalian neutrophils, act through on G proteins. Receptors with bound attractant cause G proteins to split from heterotrimers into active GTP-bound  $\alpha$  subunits and  $\beta\gamma$  subunits, and these free subunits in turn activate further intracellular messengers. In cells such as neutrophils, with a vast range of receptors, this process simplifies the cell's responses by limiting the number of active species that cause intracellular effects. In *D. discoideum*, one of the best organisms for studying chemotaxis (King and Insall, 2009; Cai and Devreotes, 2011), this simplification is even clearer: there are single genes for  $\beta$  and  $\gamma$  subunits (Wu et al., 1995; Zhang et al., 2001), and the  $\beta\gamma$  subunit is thought to mediate most responses. Thus, in theory all attractants (for example, cAMP and folate) should elicit fundamentally similar chemotaxis (Srinivasan et al., 2013). However

folate chemotaxis is often difficult to measure, for reasons explored in this paper.

Cells can be steered using many mechanisms. In shallow gradients, the migration process is fundamentally random, and attractants can bias multiple features of the normal motility cycle, for example the direction of new protrusions after pseudopod splitting (Bosgraaf and Van Haastert, 2009), and the probability of old pseudopods being retracted (Andrew and Insall, 2007). However, the second messengers that are used are still not understood. In one model, phosphatidylinositol 3,4,5-trisphosphate (PIP<sub>3</sub>) is a "compass needle" (Bourne and Weiner, 2002); intracellular signal processing somehow resolves the ideal future direction of the cell, resulting in local generation of PIP<sub>3</sub> by the p110 phosphatidylinositide 3-kinase (PI 3-kinase) family. Discrete patches of PIP<sub>3</sub> appear at the leading edge of some cells during chemotaxis (Postma et al., 2004), which suggests that these represent the outcome of a nonlinear response, perhaps by discretizing spatial information to control protrusion (Cai and Devreotes, 2011). However, newer data have shown that PIP<sub>3</sub> is not essential for chemotaxis in *D. discoideum*

Correspondence to Robert H. Insall: R.Insall@beatson.gla.ac.uk

Abbreviations used in this paper: CRAC, cytoplasmic regulator of adenylyl cyclase; DIC, differential interference contrast; PI 3-kinase, phosphatidylinositide 3-kinase; PIP<sub>3</sub>, phosphatidylinositol 3,4,5-trisphosphate; SCAR, suppressor of cAR mutations; TIRF, total internal reflection fluorescence.

© 2014 Veltman et al. This article is distributed under the terms of an Attribution-Noncommercial-Share Alike-No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms>). After six months it is available under a Creative Commons License (Attribution-Noncommercial-Share Alike 3.0 Unported license, as described at <http://creativecommons.org/licenses/by-nc-sa/3.0/>).

Supplemental Material can be found at:  
<http://jcb.rupress.org/content/suppl/2014/02/14/jcb.201309081.DC1.html>  
Original image data can be found at:  
<http://jcb-dataviewer.rupress.org/jcb/browse/7753>




Figure 1. **Folate chemotaxis is inefficient in axenic cells.** Cells were grown under different conditions, then examined responding to linear attractant gradients in Insall chambers. Numbers are the means  $\pm$  SEM of at least four independent experiments of at least 20 cells each. Tracks of cells from the same experiment have the same color. (A) Axenically cultivated AX2 cells responding to folate (left) and 4 h-starved cells responding to cAMP (right). (B) Bacterially grown NC4 cells (the parent of AX2) migrating toward folate. (C) Bacterially grown AX2 cells migrating toward folate.

(Hoeller and Kay, 2007) or in neutrophils (Ferguson et al., 2007). This, among other data, has led to alternate models such as the pseudopod-centered model (Insall, 2010), in which multiple signaling processes act by biasing a normal, random pseudopod cycle. Despite this, the concept of a chemotactic compass (perhaps operating through an alternative second messenger) and a central role for PIP<sub>3</sub> in chemotaxis are frequently cited in the literature.

More recently, PIP<sub>3</sub> has been strongly associated with macropinocytosis (Posor et al., 2013), in which cells use actin-driven cups to endocytose large volumes of liquid. Macropinosomes are induced by growth factors such as PDGF in mammalian cells, and the small GTPase Ras, which directly activates PI 3-kinases, causes massive macropinocytosis when it is inappropriately activated (Commisso et al., 2013). Thus, macropinocytosis and PIP<sub>3</sub> are associated.

Macropinocytosis also occurs in *D. discoideum*. In the wild, cells feed by phagocytosing bacteria. However, most laboratory work is done in axenic strains (“axenic” in this case means “able to grow in the absence of living prey”). These strains have been selected by growth in medium without bacteria, and are able to grow because of a constitutive high rate of macropinocytosis. *D. discoideum* macropinosomes are large structures (up to 5  $\mu$ m) that efficiently take up liquid and soluble

nutrients (Swanson, 2008). The mutations that allow axenic growth have been mapped to three separate loci, but their identity remains unknown (Clarke and Kayman, 1987). The use of axenic *D. discoideum* strains is so widespread that they are often, incorrectly, referred to as wild-type cells. In this work we show an unexpected conflict between chemotaxis and macropinocytosis that raises fundamental questions about the physiological role of PIP<sub>3</sub>.

## Results and discussion


### Axenic *D. discoideum* cells are defective in chemotaxis to folate

The main reason for studying chemotaxis in model organisms like *D. discoideum* is to find simple but generalizable results. It is therefore desirable to study multiple attractants to separate global from agonist-specific mechanisms. We have therefore studied chemotaxis toward folate. Compared with the well-studied cAMP system, folate uses different receptors and G protein  $\alpha$  subunits (Srinivasan et al., 2013), which are expressed in growing cells that are not responsive to cAMP. However it has been difficult to measure folate chemotaxis with most assays (though under-agar and some micropipette assays succeed). To discover why, we exposed AX2 cells to folate gradients in a standard chemotaxis chamber (Muinonen-Martin et al., 2010). Under these conditions, starved cells chemotax efficiently toward cAMP. However, growing cells consistently failed to migrate up the folate gradient (Fig. 1 A). This was surprising, as folate is thought to be a potent attractant. To analyze the problem, we switched to using wild-type NC4, the parent strain of AX2, which does not have axenic mutations and must thus be grown on bacteria. In contrast to axenic cells, wild-type cells robustly migrated up the folate gradient (Fig. 1 B). AX2 cells grown on bacteria also show a little chemotaxis (Fig. 1 C), but still far less than NC4, which indicates that the loss of folate chemotaxis is caused by genetic differences between the wild type and the axenic AX2 strain.


### Actin, pseudopod orientation, and PIP<sub>3</sub>

This defect in a strain that is widely used to study motility and chemotaxis is worrying, so we determined its cause. Chemoattractants steer cells by altering pseudopod behavior, so we measured pseudopod directions in axenic and nonaxenic cells. We also examined PIP<sub>3</sub> because of its clear association with both macropinocytosis during axenic growth and chemotaxis. Cells were transfected with Lifeact-mRFP (Riedl et al., 2008) and PH-CRAC-GFP (Parent et al., 1998) to show actin pseudopods in red and PIP<sub>3</sub> in green.

We first determined the fraction of F-actin-rich protrusions labeled with PIP<sub>3</sub> during random migration (as observed by others, the PIP<sub>3</sub>-positive and -negative populations were essentially discrete, allowing an all-or-nothing score). In axenically cultivated AX2 cells,  $\sim$ 50% of random protrusions accumulate PIP<sub>3</sub> at the edge (Fig. 2, A and B). In AX2 grown on bacteria, PIP<sub>3</sub>-containing pseudopods are only about half as frequent, though these cells migrate more efficiently.



**Figure 2. PIP<sub>3</sub>-labeled protrusions do not orient up folate gradients.** (A) AX2 and NC4 cells transfected with the F-actin reporter Lifeact-mRFP and the PIP<sub>3</sub> reporter PH-CRAC-GFP were examined by confocal microscopy, and the presence or absence of a PIP<sub>3</sub> patch in each cell was determined during random migration. Bar, 10 μm. (B) Numbers of pseudopods with and without PIP<sub>3</sub> patches during random migration. Percentages are the means and standard deviations from three experiments. (C and D) Mean direction of protrusions during folate chemotaxis. (C) The displacements from the migration vectors of all pseudopods were measured, and the mean is shown in a circular plot. If all protrusions were extended in the same direction, a vector with length 1 would result. PIP<sub>3</sub>-containing pseudopods from NC4 were too rare to give a reliable direction. (D) Rose plots of the directions of all protrusions with and without PIP<sub>3</sub>. The shaded area indicates the 95% confidence interval of the mean angle. (E) Frequency and direction of pseudopods in C and D. (F) Rose plot of the direction of PIP<sub>3</sub> patches that form macropinosomes during folate chemotaxis.



**Figure 3. PIP<sub>3</sub> patches organize macropinosomes.** (A and B) Confocal series showing turnover of PIP<sub>3</sub> patches in NC4 and axenically grown AX2 cells expressing the PIP<sub>3</sub> marker PH-CRAC-GFP. Arrowheads mark budding. Bar, 10  $\mu$ m. (C) Confocal series showing protrusions in an axenically cultivated AX2 cell expressing PH-CRAC-GFP and the F-actin marker Lifectin-mRFP. Arrows indicate negative curvature. Bar, 5  $\mu$ m. This corresponds with Video 1. (D) Kymograph of a line drawn through the protrusion in C.

#### Expression of PIP<sub>3</sub> markers in wild-type cells

It has previously not been practical to express ectopic genes in wild-type *D. discoideum* because standard selectable markers and promoters work poorly in bacterially grown cells. We constructed new vectors to allow expression of markers for PIP<sub>3</sub> and F-actin in wild-type cells, and developed methods for transfection and selection using bacteria as food (see Materials and methods). We successfully cotransfected NC4 cells with Lifectin-mRFP and PH-CRAC-GFP to examine the distribution of F-actin and PIP<sub>3</sub>. To our surprise, however, PIP<sub>3</sub>-containing

pseudopods were extremely rare in randomly moving wild-type cells (Fig. 2 B); 98% of pseudopods had no discernible PIP<sub>3</sub>. Thus, in growing wild-type *D. discoideum*, unlike in axenic cells, PIP<sub>3</sub> appears not to be used at all in random migration, even though the cells migrate far more robustly.

#### PIP<sub>3</sub> and folate chemotaxis

We next examined PIP<sub>3</sub> during folate chemotaxis. Cells were exposed to a relatively steep folate gradient in a chemotaxis chamber, and the angle between each protrusion and the gradient was determined. As before, axenic AX2 cells did not show chemotaxis. The mean cosine of the angle between the protrusions and the chemoattractant gradient was close to zero, which indicates no alignment, regardless of whether PIP<sub>3</sub> was present. Also, protrusions with PIP<sub>3</sub> made a negligible contribution to net migration (Fig. 2, C, D, and E). Macropinosomes were not oriented by the folate gradient (Fig. 2 F).

As before, AX2 cells grown on bacteria chemotaxed weakly, and the proportion of pseudopods with PIP<sub>3</sub> patches decreased substantially (from 63% to 29%; Fig. 2 E). Furthermore, PIP<sub>3</sub>-labeled protrusions did not orient up the gradient (Fig. 2 E). In contrast, the protrusions that showed no discernible PIP<sub>3</sub> oriented up the folate gradient with a mean cosine of 0.40. Thus, unexpectedly, absence of PIP<sub>3</sub> patches correlated with improved chemotaxis.

During chemotaxis, pseudopods in vegetative wild-type cells were still almost never labeled with PIP<sub>3</sub> (Fig. 2 E). In a folate gradient, the protrusions of wild-type cells showed high directionality (mean cosine 0.59) despite the lack of PIP<sub>3</sub> at the front (Fig. 2, C and D). Thus, in all conditions we observed, only PIP<sub>3</sub>-negative pseudopods are directed by folate.

#### PIP<sub>3</sub> and macropinocytosis in wild-type and axenic cells

The negative correlation between the number of PIP<sub>3</sub> patches and the chemotaxis efficiency was unexpected, as PIP<sub>3</sub> signaling is thought to be fundamentally involved in directional sensing. However, PIP<sub>3</sub> is involved in several other physiological processes, in particular macropinocytosis (Dormann et al., 2004; Hoeller et al., 2013). We speculated that this could negatively impact chemotaxis efficiency, and we therefore examined macropinocytosis and its regulation by PIP<sub>3</sub> in more detail.

In wild-type cells, macropinocytosis could readily be visualized with the PIP<sub>3</sub> marker PH-CRAC-GFP (Fig. 3 A), despite the substantially lower rate of fluid uptake. The PIP<sub>3</sub> is strictly confined to the nascent vesicle; levels increase during invagination and reach a maximum as the vesicle closes. The entire process is very fast, with a mean time of 8 s between appearance of PIP<sub>3</sub> and budding of the vesicle ( $n = 27$ ,  $SD = 4$  s). Macropinosomes in wild-type cells are small and do not contribute to cell movement, and are readily distinguishable from the large, convex, and spiky F-actin pseudopods.

Unlike wild-type cells, axenic AX2 cells generate macropinosomes from preassembled PIP<sub>3</sub> patches. The PI 3-kinases responsible for this process have recently been analyzed in depth (Hoeller et al., 2013); the patches are presumably the structures organized by PIKs 1 and 2. The diameter of the patches is a

substantial fraction of the size of the cell (Fig. 3 B). Progression of negative curvature to drive vesicle formation frequently stalls, causing patches to remain planar or slightly inwardly curved for prolonged periods of time. Moreover, PIP<sub>3</sub> patches often disappear without any vesicle formation, which indicates that macropinocytosis was aborted.

The PIP<sub>3</sub> patches that are involved in axenic macropinocytosis are a similar size to pseudopods and recruit large amounts of F-actin. Nascent macropinosomes are therefore difficult to discriminate from pseudopods, especially if they abort before vesicle formation. This is further complicated because pseudopods and macropinosomes are able to interconvert. Fig. 3 C and Video 1 show an example. The protrusion is initially free of PIP<sub>3</sub> and resembles a normal pseudopod. At 20 s, a PIP<sub>3</sub> patch appears at the leading edge, and shortly after, negative curvature is induced at the site of the patch. The image plainly resembles a forming macropinocytic cup. However, it does not fully close and aborts after 80 s. A kymograph of this event (Fig. 3 D) shows that the forward movement of the leading edge reduces after the appearance of PIP<sub>3</sub>.

#### Suppressor of cAR mutations (SCAR) and F-actin organization in macropinosomes

The ambiguous character of these actin structures in axenic cells complicates analysis. To more accurately distinguish the potential role of PIP<sub>3</sub> in pseudopods and macropinosomes, we examined the behavior of the SCAR/WASP and verprolin homologue (WAVE) complex, which is the definitive marker for pseudopods. It remains associated with the leading edge throughout the progression of a pseudopod and disappears from the leading edge when the pseudopod stops advancing (Fig. 4 A; Hahne et al., 2001; Veltman et al., 2012). Interestingly, high concentrations of PIP<sub>3</sub> do not colocalize with the SCAR complex, which is recruited to a small margin surrounding each patch (Fig. 4, B–D; and Video 2). This distribution is not only visible in the PIP<sub>3</sub> patches that develop to form macropinosomes, but also in ones that remain planar and that cannot readily be classified as either a pseudopod or a nascent macropinosome (Video 1 and Fig. 4 B, asterisk). We were unable to find any event where a PIP<sub>3</sub> patch was followed by recruitment of SCAR to the leading edge in the fashion that is typical for pseudopods.

A close-up view demonstrates the complexity of the macropinocytic cup. The region labeled with PIP<sub>3</sub> remains static, with little movement perpendicular to the membrane (Fig. 4 E, green dots). SCAR remains associated with the rim of the cup throughout the event. The SCAR-containing region moves outward and drives closure of the cup (Fig. 4 E, other dots). This outward movement at the edge of the PIP<sub>3</sub> patch can generate cell movement, though it is retarded by the slow movement at the center of the PIP<sub>3</sub> patch. It is also clear that the edge of the patch is a major site of actin filament nucleation (Fig. 4 F). The Lifeact-mRFP signal overlies the PIP<sub>3</sub>, though it is strongest at the extremities (Fig. 4 G), and there is a clear difference between the edges and the center; the SCAR-labeled edges are rough and their thickness is variable, like pseudopods, whereas the central F-actin is thinner and smooth.

Collectively, these data indicate that PIP<sub>3</sub> marks membrane regions that are destined for invagination during axenic growth. In axenic cells, this results in a complex pattern of inward and outward movement of the membrane, which combines with F-actin to mediate both macropinocytosis and a subset of cell movement.

#### Inverse correlation between macropinocytosis and chemotaxis

We assessed the relationship between macropinocytosis and chemotaxis by altering the macropinocytic rate. Axenically grown AX2 cells form FITC-labeled vesicles at a rate of 0.44/min (Fig. 5 A). However, AX2 cells that have been cultivated on bacteria have a diminished rate of macropinocytosis. Macropinosomes in NC4 cells are even rarer, with a mean rate of 0.05/min. These changes correlate with the efficiency of chemotaxis toward folate (Fig. 5 A).

Folate chemotaxis is robust under a sheet of agarose (for example Laevsky and Knecht, 2001; and Blagg et al., 2003). We examined whether the difference between under-agar and chamber assays was connected with changes in macropinocytosis. Surprisingly, macropinocytosis in vegetative axenic cells is almost completely blocked under agar, with the number of macropinosomes dropping below those in nonaxenic cells (Fig. 5 A). It is not clear why physical constraint causes such a large suppression of macropinocytosis, but it is reminiscent of the induction of autophagy by compression (King et al., 2011). However, the effect on chemotaxis is clear: axenic cells become chemotactic when macropinocytosis is blocked under agarose despite being nonchemotactic when unrestricted.


#### Loss of PI 3-kinase blocks macropinocytosis and rescues migration and folate chemotaxis

The inverse correlation between macropinocytosis and chemotaxis raises an interesting conflict. Despite the belief that PIP<sub>3</sub> is fundamental to chemotaxis, we have found that PIP<sub>3</sub>-driven macropinocytosis is not consistent with chemotaxis to folate. We therefore examined a strain lacking all five PI 3-kinases (Hoeller and Kay, 2007). These cells have no PIP<sub>3</sub> patches, and their axenic growth is slow due to a defect in macropinocytosis (Hoeller et al., 2013). In our hands, the rate of macropinocytosis of PI 3-kinase-null cells was only 0.05/min, similar to that of nonaxenic cells. However, chemotaxis toward folate was greatly improved (Fig. 5 A). Bacterially grown AX2 migrated toward folate with a weak chemotaxis index of 0.23 and a speed of 5.6 μm/min; knocking out all the PI 3-kinases restored this to 0.44 and 19.1 μm/min, respectively (Fig. 5 B). Similarly, 10 μM LY294002, enough to inhibit PI 3-kinase without blocking the target of rapamycin (TOR) complex, significantly diminished macropinocytosis and increased cell speed (Fig. S2). Thus, it appears that PIP<sub>3</sub> is not only inessential for folate chemotaxis in *D. discoideum*, but actively opposes it.

#### Axenic mutations, macropinocytosis, and chemotaxis

Our data clearly show that PIP<sub>3</sub> patches are not used at all by cells chemotaxing to one of the two principal attractants used

**Figure 4. SCAR, actin, and PIP<sub>3</sub> in pseudopods and macropinosomes.** (A) TIRF and DIC image of HSPC300-GFP during progression of a normal pseudopod of axenic AX3 cells. Cells were overlaid with 0.4% agarose to ensure contact with the glass surface. (B) Confocal series of an axenically grown AX2 cell expressing PH-CRAC-RFP and the SCAR complex marker HSPC300-GFP. Arrowheads show regions marked by HSPC300-GFP. Asterisk indicates a PIP<sub>3</sub> patch that does not form a macropinosome. The boxed regions are shown in detail (2.2× magnification) in the bottom row. This corresponds with Video 2. (C) TIRF image of an axenically cultivated AX2 cell expressing PH-CRAC-RFP and HSPC300-GFP. (D) Profile plot of the line indicated in C. (E) Confocal image series of an axenic AX2 cell expressing PH-CRAC-RFP and HSPC300-GFP during macropinocytosis. Colored dots indicate the edges (red, purple) and center (green) of the PIP<sub>3</sub> patch. (F) Confocal image of an axenic AX2 cell expressing Lifeact-mRFP and PH-CRAC-GFP with a pseudopod (top left) and a macropinosome (top right). (G) Profile plot of a line drawn along the perimeter of the developing macropinosome in F. All indicated times are in seconds. Bars: (A, B, C, and F) 10 μm; (E) 2 μm.



by *D. discoideum* cells. Thus, at the very least, PIP<sub>3</sub> is not a universal player in chemotaxis. This is not a surprise, as a great deal of data has accumulated over the past six years showing

that PI 3-kinases are not required for chemotaxis. What was a surprise, however, was that PIP<sub>3</sub> actually opposes folate chemotaxis. Folate and cAMP mostly use the same signaling

components, in particular the single G protein  $\beta\gamma$  subunit, which is thought to mediate most of the responses downstream of chemoattractant receptors. The key difference is therefore not likely to be in signaling pathways, but rather the developmental stages of the cells. While cells are growing, axenic mutations lead to massive, PIP<sub>3</sub>-driven macropinocytosis, which is clearly inconsistent with chemotaxis. Starved axenic cells cease macropinocytosis and thus regain chemotaxis at the time they become sensitive to cAMP.

The identity of the axenic mutations is not yet known, but it will be fascinating to unravel their influence on chemotaxis at all stages. This may fundamentally alter our understanding of chemotaxis in *D. discoideum* and metazoan cells.

## Materials and methods

### Cell strains and cultivation

*D. discoideum* AX2 cells were used as an axenic strain and NC4 cells were used as nonaxenic, wild-type cells. The PI3 kinase quintuple knockout strain HM1200 (Hoeller and Kay, 2007) was obtained from the Dicty Stock Center. For axenic growth, cells were cultivated on Petri dishes under HL5 medium (ForMedium). Nonaxenic cultivation was done on Petri dishes under SorMC buffer supplemented with live *Klebsiella aerogenes* at an OD<sub>600</sub> of 2. SorMC buffer consists of 15 mM KH<sub>2</sub>PO<sub>4</sub>, 2 mM Na<sub>2</sub>HPO<sub>4</sub>, 50  $\mu$ M MgCl<sub>2</sub>, and 50  $\mu$ M CaCl<sub>2</sub>, pH 6.0. To reduce autofluorescence of axenically cultivated cells in preparation for fluorescence microscopy, cells were incubated overnight in LoFlo medium (ForMedium). For microscopy, cells were put on a 3.5-cm acid-washed glass-bottom MatTek dish.

### DNA constructs and transfection


In brief, the act6 promoter driving the resistance marker of the modular pDM expression vector set (Veltman et al., 2009) was replaced by the *coaA* promoter (bp -293 to bp -1 relative to the start codon of *coaA*). The PIP<sub>3</sub>-binding PH domain of cytoplasmic regulator of adenyl cyclase (CRAC; residue 1–126 of the protein encoded by *dagA*) was used as a PIP<sub>3</sub> sensor. Full-length HSPC300 was used as a marker for the SCAR complex, and Lifeact (Riedl et al., 2008) was used as a marker for filamentous actin. Gene fragments were cloned into the BglII–SpeI site of the expression vector. For coexpression of two genes the shuttle vector was used as described previously (Veltman et al., 2009). avGFP S65T and mCherry were used for fluorescence labeling.

Transfection of nonaxenic cells was performed as follows:  $5 \times 10^6$  cells were harvested from the plate, washed once in SorMC buffer, and resuspended in 400  $\mu$ l SorMC buffer. Cells were mixed with 5  $\mu$ l miniprep DNA (~0.5–1  $\mu$ g total) and put on ice. Cells were then electroporated at 500 V using the high-voltage setting of an ECM 399 Exponential Decay Wave Electroporation System (Harvard Apparatus) and immediately transferred to a Petri dish with SorMC buffer and live *K. aerogenes* at an OD<sub>600</sub> of 2. Selection marker was added after 5 h (10  $\mu$ g/ml G418 or 100  $\mu$ g/ml hygromycin).

### Microscopy and image analysis

All microscopy was performed at room temperature. Confocal laser scanning microscopy was performed on a confocal microscope (A1R; Nikon) using a 1.4 NA Plan-Apochromat 60x oil immersion objective lens and the manufacturer's NIS Elements acquisition software (Nikon). Total internal reflection fluorescence (TIRF) microscopy was performed on an inverted microscope (Eclipse TE2000-U; Nikon) that was fitted with a custom TIRF condenser and a 1.45 NA 100x Plan-Apochromat TIRF objective lens. Red and green fluorescent signals were separated using a beam splitter and projected onto an EM charge-coupled device camera (Evolve512; Photometrics). MetaMorph software was used to control the camera, shutters, and light sources. Spinning disk confocal microscopy was performed on an Andor system using an Eclipse Ti body (Nikon) fitted with a Plan-Apochromat VC 100x oil immersion objective lens (NA 1.4). Images were captured on a Neo 5.5 sCMOS camera using the manufacturer's iQ software (both from Andor Technology).

Fluorescence images were analyzed using the standard tools available in ImageJ. Images were not processed other than brightness/contrast adjustments and bleach correction for mRFP in Video 2. The Angle tool



**Figure 5. PIP<sub>3</sub>-induced macropinocytosis is incompatible with chemotaxis.** (A) Negative correlation of macropinocytosis and chemotaxis. The rate of macropinosome formation under different conditions was measured using FITC-dextran and fluorescence microscopy. Each data point represents a single cell that was imaged for at least 5 min. Data were taken from at least two independent experiments. Chemotaxis is represented qualitatively to avoid quantitative comparisons made under noncomparable conditions. (B) Deletion of PI 3-kinases from axenically growing cells restores chemotaxis to folate. (B, right) Insall chamber chemotaxis of axenic AX2 cells quintuply deleted for all p110 PI 3-kinases. (B, left) Chemotaxis of axenic wild-type cells (see data from Fig. 1 C for comparison).

was used to determine the angle between a protrusion and the gradient. Graphs of the collected angle data were generated in MATLAB (MathWorks). Statistical analysis was done using the Circular Statistics Toolbox (Berens, 2009).

### Chemotaxis assays

Chemotaxis assays were performed on an inverted microscope (Eclipse TE2000) using a 10x phase contrast objective lens (NA 0.3). The microscope was controlled using Micromanager (Edelstein et al., 2010). Insall chambers (Muinonen-Martin et al., 2010) were used to expose cells to a chemoattractant gradient of 0–100  $\mu$ M folate across a bridge that was 1 mm wide. Typically, a recording of 60 min was captured, starting immediately after the assembly of the chemotaxis chamber. The position of at least 20 randomly selected cells in each recording was determined every minute using the ImageJ plugin mTrackJ (Meijering et al., 2012). The chemotaxis index of the population was calculated for each time



point using a custom-written Excel (Microsoft) spreadsheet that uses the following formula:

$$CI(t) = \frac{\sum_{i=1}^n \cos\theta(t)_i \cdot a(t)_i}{\sum_{i=1}^n a(t)_i}$$

where  $n$  is the number of cells analyzed,  $\cos\theta(t)_i$  is the cosine of the angle between the displacement vector of cell  $i$  and the direction of the gradient at time point  $t$ , and  $a(t)_i$  is the length of the displacement vector of cell  $i$  at time point  $t$ . The chemotaxis index of the population is then plotted over time and a 40-min window is selected where the chemotaxis is optimal. Typically, this would exclude the start of the recording, where the gradient has not yet stabilized, and sometimes the end of the recording, when the chamber dried out and the gradient collapsed. The mean chemotaxis index of the population was then calculated as follows:

$$CI_{population} = \frac{\sum_{t=1}^{40} CI_t \cdot a_t \cdot n_t}{\sum_{t=1}^{40} a_t \cdot n_t}$$

where  $CI_t$  is the mean  $CI$  of the cells at time point  $t$ ,  $a_t$  is the mean displacement of the cells at time point  $t$ , and  $n_t$  is the number of analyzed cells at time point  $t$ . The chemotaxis indices shown are the means of the population means of at least three experiments  $\pm$  the standard error.

#### Quantification of the rate of macropinocytosis

Cells were seeded onto an acid-washed glass-bottom dish. The medium was supplemented with 800  $\mu\text{g}/\text{ml}$  FITC-dextran. For the under-agarose condition, cells were overlaid with a 1-mm thick slab of 0.8% agarose in LoFlo medium + 800  $\mu\text{g}/\text{ml}$  FITC-dextran. Cells were allowed to equilibrate for 20 min. A 15-min recording was made, simultaneously capturing the fluorescent FITC signal and the corresponding differential interference contrast (DIC) image on a line-scanning confocal microscope. Frames were taken every 2.5 s and across three different  $z$  planes spanning a total of 4  $\mu\text{m}$ . Individual uptake events of fluorescent dextran were then identified by eye using both the data from the fluorescence channel and the DIC channel (see Fig. S1).

#### Macropinosome and speed experiment and quantification

Cells expressing Lifeact-mRFP were harvested in growth phase and centrifuged for 5 min at 400  $g$ , then resuspended in LoFlo medium at a titer between  $0.5 \times 10^6$  and  $10^6$  cells/ml, and 0.5 ml was added to a glass-bottom dish (Bioprotech, Inc.). Cells were incubated for 3 h to allow growth medium to be exported and then imaged every 20 s for 40 min on a confocal microscope (A1R; Nikon) as described in the Microscopy and image analysis section. LY294002 was added to a final concentration of 10.8  $\mu\text{M}$  and imaging was continued.

Macropinosome quantification was done by counting the number of internalized macropinosomes (circular and Lifeact-mRFP labeled) formed in 30 min in each cell ( $n = 31$ ). Cells were also tracked for speed, before and after LY294002 addition using the ImageJ and the MTrackJ plugin.

#### Online supplemental material

Fig. S1 shows the detailed process for determination of the rates of macropinocytosis. Fig. S2 shows that moderate levels of LY294002, an inhibitor of PI 3-kinases and PIP<sub>3</sub> accumulation, cause axenically growing cells to migrate faster and macropinocytose more slowly. Video 1 shows a PIP<sub>3</sub> patch inducing a negative (i.e., concave) curvature at the leading edge, unlike a normal pseudopod that shows positive/convex curvature. Video 2 shows that the SCAR complex, the principal driver of pseudopod extension, is only recruited to the edges, not the center, of PIP<sub>3</sub> patches. Online supplemental material is available at <http://www.jcb.org/cgi/content/full/jcb.201309081/DC1>. Additional data are available in the JCB Data-Viewer at <http://dx.doi.org/10.1083/jcb.201309081.dv>.

We are grateful to Dr. G. Kalna for help with statistics and data presentation and Prof. L. Machesky for criticism of the manuscript. We are grateful to all the members of Beatson Advanced Imaging Resource (BAIR) for assistance with microscopy.

We thank Cancer Research UK for core funding to R.H. Insall.

The authors declare no competing financial interests.

Submitted: 16 September 2013

Accepted: 7 January 2014

## References

- Andrew, N., and R.H. Insall. 2007. Chemotaxis in shallow gradients is mediated independently of PtdIns 3-kinase by biased choices between random protrusions. *Nat. Cell Biol.* 9:193–200. <http://dx.doi.org/10.1038/ncb1536>
- Berens, Philipp. 2009. CircStat: A MATLAB Toolbox for Circular Statistics. *Journal of Statistical Significance.* 31:1–21. <http://www.jstatsoft.org/v31/i10> (accessed January 30, 2014).
- Blagg, S.L., M. Stewart, C. Sambles, and R.H. Insall. 2003. PIR121 regulates pseudopod dynamics and SCAR activity in *Dictyostelium*. *Curr. Biol.* 13:1480–1487. [http://dx.doi.org/10.1016/S0960-9822\(03\)00580-3](http://dx.doi.org/10.1016/S0960-9822(03)00580-3)
- Bosgraaf, L., and P.J. Van Haastert. 2009. Navigation of chemotactic cells by parallel signaling to pseudopod persistence and orientation. *PLoS ONE.* 4:e6842. <http://dx.doi.org/10.1371/journal.pone.0006842>
- Bourne, H.R., and O. Weiner. 2002. Cell Polarity: A chemical compass. *Nature.* 419:21. <http://dx.doi.org/10.1038/419021a>
- Cai, H., and P.N. Devreotes. 2011. Moving in the right direction: how eukaryotic cells migrate along chemical gradients. *Semin. Cell Dev. Biol.* 22:834–841. <http://dx.doi.org/10.1016/j.semcdb.2011.07.020>
- Clarke, M., and S.C. Kayman. 1987. The axenic mutations and endocytosis in *Dictyostelium*. *Methods Cell Biol.* 28:157–176. [http://dx.doi.org/10.1016/S0091-679X\(88\)61642-8](http://dx.doi.org/10.1016/S0091-679X(88)61642-8)
- Commisso, C., S.M. Davidson, R.G. Soydaner-Azeloglu, S.J. Parker, J.J. Kamphorst, S. Hackett, E. Grabocka, M. Nofal, J.A. Drebin, C.B. Thompson, et al. 2013. Macropinocytosis of protein is an amino acid supply route in Ras-transformed cells. *Nature.* 497:633–637. <http://dx.doi.org/10.1038/nature12138>
- Dormann, D., G. Weijer, S. Dowler, and C.J. Weijer. 2004. In vivo analysis of 3-phosphoinositide dynamics during *Dictyostelium* phagocytosis and chemotaxis. *J. Cell Sci.* 117:6497–6509. <http://dx.doi.org/10.1242/jcs.01579>
- Edelstein, A., N. Amodaj, K. Hoover, R. Vale, and N. Stuurman. 2010. Computer control of microscopes using  $\mu$ Manager. *Curr. Protoc. Mol. Biol.* Chapter 14:20.
- Ferguson, G.J., L. Milne, S. Kulkarni, T. Sasaki, S. Walker, S. Andrews, T. Crabbe, P. Finan, G. Jones, S. Jackson, et al. 2007. PI(3)K $\gamma$  has an important context-dependent role in neutrophil chemokinesis. *Nat. Cell Biol.* 9:86–91. <http://dx.doi.org/10.1038/ncb1517>
- Hahne, P., A. Sechi, S. Benesch, and J.V. Small. 2001. Scar/WAVE is localised at the tips of protruding lamellipodia in living cells. *FEBS Lett.* 492:215–220. [http://dx.doi.org/10.1016/S0014-5793\(01\)02239-6](http://dx.doi.org/10.1016/S0014-5793(01)02239-6)
- Hoeller, O., and R.R. Kay. 2007. Chemotaxis in the absence of PIP<sub>3</sub> gradients. *Curr. Biol.* 17:813–817. <http://dx.doi.org/10.1016/j.cub.2007.04.004>
- Hoeller, O., P. Bolourani, J. Clark, L.R. Stephens, P.T. Hawkins, O.D. Weiner, G. Weeks, and R.R. Kay. 2013. Two distinct functions for PIP<sub>3</sub>-kinases in macropinocytosis. *J. Cell Sci.* 126:4296–4307. <http://dx.doi.org/10.1242/jcs.134015>
- Insall, R.H. 2010. Understanding eukaryotic chemotaxis: a pseudopod-centred view. *Nat. Rev. Mol. Cell Biol.* 11:453–458. <http://dx.doi.org/10.1038/nrm2905>
- King, J.S., and R.H. Insall. 2009. Chemotaxis: finding the way forward with *Dictyostelium*. *Trends Cell Biol.* 19:523–530. <http://dx.doi.org/10.1016/j.tcb.2009.07.004>
- King, J.S., D.M. Veltman, and R.H. Insall. 2011. The induction of autophagy by mechanical stress. *Autophagy.* 7:1490–1499. <http://dx.doi.org/10.4161/autophagy.7.12.17924>
- Laevsky, G., and D.A. Knecht. 2001. Under-agarose folate chemotaxis of *Dictyostelium discoideum* amoebae in permissive and mechanically inhibited conditions. *Biotechniques.* 31:1140–1142; 1144: 1146–1149.
- Meijering, E., O. Dzyubachyk, and I. Smal. 2012. Methods for cell and particle tracking. *Methods Enzymol.* 504:183–200. <http://dx.doi.org/10.1016/B978-0-12-391857-4.00009-4>
- Muinenen-Martin, A.J., D.M. Veltman, G. Kalna, and R.H. Insall. 2010. An improved chamber for direct visualisation of chemotaxis. *PLoS ONE.* 5:e15309. <http://dx.doi.org/10.1371/journal.pone.0015309>
- Parent, C.A., B.J. Blacklock, W.M. Froehlich, D.B. Murphy, and P.N. Devreotes. 1998. G protein signaling events are activated at the leading edge of chemotactic cells. *Cell.* 95:81–91. [http://dx.doi.org/10.1016/S0092-8674\(00\)81784-5](http://dx.doi.org/10.1016/S0092-8674(00)81784-5)
- Posor, Y., M. Eichhorn-Gruenig, D. Puchkov, J. Schöneberg, A. Ullrich, A. Lampe, R. Müller, S. Zarbakhsh, F. Gulluni, E. Hirsch, et al. 2013. Spatiotemporal control of endocytosis by phosphatidylinositol-3,4-bisphosphate. *Nature.* 499:233–237. <http://dx.doi.org/10.1038/nature12360>
- Postma, M., J. Roelofs, J. Goedhart, H.M. Looovers, A.J. Visser, and P.J. Van Haastert. 2004. Sensitization of *Dictyostelium* chemotaxis by phosphoinositide-3-kinase-mediated self-organizing signalling patches. *J. Cell Sci.* 117:2925–2935. <http://dx.doi.org/10.1242/jcs.01143>

- Riedl, J., A.H. Crevenna, K. Kessenbrock, J.H. Yu, D. Neukirchen, M. Bista, F. Bradke, D. Jenne, T.A. Holak, Z. Werb, et al. 2008. Lifeact: a versatile marker to visualize F-actin. *Nat. Methods*. 5:605–607. <http://dx.doi.org/10.1038/nmeth.1220>
- Srinivasan, K., G.A. Wright, N. Hames, M. Housman, A. Roberts, K.J. Aufderheide, and C. Janetopoulos. 2013. Delineating the core regulatory elements crucial for directed cell migration by examining folic-acid-mediated responses. *J. Cell Sci.* 126:221–233. <http://dx.doi.org/10.1242/jcs.113415>
- Swanson, J.A. 2008. Shaping cups into phagosomes and macropinosomes. *Nat. Rev. Mol. Cell Biol.* 9:639–649. <http://dx.doi.org/10.1038/nrm2447>
- Veltman, D.M., G. Akar, L. Bosgraaf, and P.J. Van Haastert. 2009. A new set of small, extrachromosomal expression vectors for *Dictyostelium discoideum*. *Plasmid*. 61:110–118. <http://dx.doi.org/10.1016/j.plasmid.2008.11.003>
- Veltman, D.M., J.S. King, L.M. Machesky, and R.H. Insall. 2012. SCAR knock-outs in *Dictyostelium*: WASP assumes SCAR's position and upstream regulators in pseudopods. *J. Cell Biol.* 198:501–508. <http://dx.doi.org/10.1083/jcb.201205058>
- Wu, L., R. Valkema, P.J. Van Haastert, and P.N. Devreotes. 1995. The G protein beta subunit is essential for multiple responses to chemoattractants in *Dictyostelium*. *J. Cell Biol.* 129:1667–1675. <http://dx.doi.org/10.1083/jcb.129.6.1667>
- Zhang, N., Y. Long, and P.N. Devreotes. 2001. G $\gamma$  in *Dictyostelium*: Its Role in Localization of G $\beta\gamma$  to the Membrane is Required for Chemotaxis in Shallow Gradients. *Mol. Biol. Cell.* 12:3204–3213. <http://dx.doi.org/10.1091/mbc.12.10.3204>