

## RESEARCH

## Open Access

# Preconditioning methods for solving a general split feasibility problem

Peiyuan Wang<sup>1,2\*</sup>, Haiyun Zhou<sup>2,3</sup> and Yu Zhou<sup>2</sup>

Dedicated to Professor Shih-sen Chang on the occasion of his 80th birthday.

\*Correspondence:

[wangpeiyuan629@163.com](mailto:wangpeiyuan629@163.com)<sup>1</sup>The Second Training Base, Naval Aviation Institution, Huludao, 125001, China<sup>2</sup>Department of Mathematics, Shijiazhuang Mechanical Engineering College, Shijiazhuang, 050003, China

Full list of author information is available at the end of the article

## Abstract

We introduce and study a new general split feasibility problem (GSFP) in a Hilbert space. This problem generalizes the split feasibility problem (SFP). The GSFP extends the SFP with a nonlinear continuous operator. We apply the preconditioning methods to increase the efficiency of the CQ algorithm, two general preconditioning CQ algorithms for solving the GSFP are presented. We also propose a new inexact method to approximate the preconditioner. The convergence theorems are established under the projections with respect to special norms. Some numerical results illustrate the efficiency of the proposed methods.

**MSC:** 46E20; 47J20; 47J25; 47L50**Keywords:** general split feasibility problem; general variational inequality; preconditioning method; projection method

## 1 Introduction

As preconditioning methods can improve the condition number of the ill-posed system matrix, the convergence rate of the iterative algorithm can also be improved [1]. In [2, 3], a preconditioning method is applied to modify the projected Landweber algorithm for solving a linear feasibility problem (LFP). The modified algorithm is

$$x_{n+1} = P_C[x_n - \tau DA^*(Ax_n - b)], \quad n \geq 0,$$

where  $\tau \in (0, 2/\|DA^*A\|)$ ,  $A : X \rightarrow Y$  is a linear and continuous operator,  $\|\cdot\|$  means 2-norm,  $X$  and  $Y$  are Hilbert spaces and  $b \in Y$  is the datum of the problem, corrupted by noise or experimental errors.

While under the nonlinear conditions, Auslender and Dafermos [4, 5] proposed an algorithm to solve variational inequalities (VI),

$$x_{n+1} = P_S[x_n - \tau_n G^{-1}F(x_n)], \quad n \geq 0, \quad (1.1)$$

where  $P_S$  is the projection operator onto  $S$  with respect to the norm  $\|\cdot\|_G$ . Bertsekas and Gafni [6] and Marcotte and Wu [7] improved it with variable symmetric positive defined matrices  $G_n$ , Fukushima [8] modified it by a relaxed projection method with half-space;

then in [9], Yang established the convergence of Auslender’s algorithm under the weak co-coercivity of  $F$ .

Further, the general variational inequality problem (GVIP) has been investigated by many authors (see [10–13]). It is to find  $u^* \in \mathbb{R}^n$  such that  $g(u^*) \in K$  and

$$\langle F(u^*), g(u) - g(u^*) \rangle \geq 0, \quad g(u) \in K,$$

where  $K$  is a nonempty closed convex set in  $\mathbb{R}^n$ ,  $F, g : \mathbb{R}^n \rightarrow \mathbb{R}^n$  are nonlinear operators. In [12], Santos and Scheimberg extended and applied (1.1) to solve the GVI.

However, a general split feasibility problem (GSFP) equals to a GVI, and preconditioning methods for solving the GSFP have not been studied. By introducing a convex minimization problem, the split feasibility problem (SFP) is equivalent to a variational inequality problem (VIP), which involves a Lipschitz continuous and inversely strong monotone (ism) operator, see [14–22]. Similarly, by the same way, in this paper we introduce that a new GSFP equals to a GVI involving a Lipschitz continuous and co-coercive operator [9, 17–19].

Otherwise, Mohammad and Abdul [23] considered a general split feasibility in infinite-dimensional real Hilbert spaces. It is to find  $x^*$  such that

$$x^* \in \bigcap_{i=1}^{\infty} C_i, \quad Ax^* \in \bigcap_{j=1}^{\infty} Q_j,$$

where  $A : H_1 \rightarrow H_2$  is a bounded linear operator,  $\{C_i\}_{i=1}^{\infty}$  and  $\{Q_j\}_{j=1}^{\infty}$  are the families of nonempty closed convex subsets of  $H_1$  and  $H_2$ , respectively.

Let  $C$  and  $Q$  be nonempty closed convex subsets in real Hilbert spaces  $H_1$  and  $H_2$ , respectively. We consider a general split feasibility problem which is different from the one in [23]. Our GSFP is to find

$$x^* \in H_1, g(x^*) \in C \text{ such that } Ag(x^*) \in Q, \tag{1.2}$$

where  $A : H_1 \rightarrow H_2$  is a bounded linear operator and  $g : H_1 \rightarrow C$  is a continuous operator. We see that the SFP in [24] and the GSFP in [23] are particular cases of GSFP (1.2). It has applications in many special fields such as signal decryption, demodulating the digital signal and noise processing, *etc.* In order to solve GSFP (1.2), two preconditioning algorithms are developed in this paper following the iterative scheme

$$g(x_{n+1}) = P_C[g(x_n) - \gamma DA^*(I - P_Q)Ag(x_n)], \quad n \geq 0,$$

where the two general constraints  $C$  and  $Q$  deal with projections with respect to the norms corresponding to some symmetric positive definite matrices. Define the solution set of (1.2)  $\Gamma = \{x^* \in H_1 \mid Ag(x^*) \in Q\}$ , as  $\Gamma$  is nonempty, and by virtue of the related  $G$ -co-coercive operator, we can establish the convergence of the proposed algorithms.

The paper is organized as follows. Section 2 presents two useful propositions. In Section 3, we define the algorithms with fixed preconditioner, variable preconditioner, relaxed projection and preconditioner approximation and analyze the convergence. Numerical results are reported in Section 4. Finally, Section 5 gives some concluding remarks.

## 2 Preliminaries

In what follows, we state some concepts and propositions.

The SFP is to find a point  $x^* \in C$  such that  $Ax^* \in Q$ , where  $A : H_1 \rightarrow H_2$  is a bounded linear operator [24].

Let  $G$  be a symmetric positive definite matrix, and set  $D = G^{-1}$ . Then the norm  $\|\cdot\|_G$  is defined by  $\|x\|_G^2 = \langle x, Gx \rangle$  for  $\forall x \in H$ . We denote by  $P_C$  the projection operator onto  $C$  with respect to the norm  $\|\cdot\|_G$  [6], i.e.,

$$P_C(x) = \arg \min_{y \in C} \{\|x - y\|_G\}.$$

Let  $\lambda_{\min}$  and  $\lambda_{\max}$  be the minimum and the largest eigenvalues of  $G$ , respectively. Then for the 2-norm  $\|\cdot\|$  [18, 19], we have

$$\lambda_{\min}\|x\|^2 \leq \|x\|_G^2 \leq \lambda_{\max}\|x\|^2, \quad \forall x \in H. \tag{2.1}$$

**Proposition 2.1** [7, 10] *Let  $C$  be a nonempty closed convex subset in  $H$ , for  $\forall x, y \in H$  and  $\forall z \in C$ , the  $G$ -projection operator onto  $C$  has the following properties:*

- (i)  $\langle G(x - P_Cx), z - P_C(x) \rangle \leq 0$ ;
- (ii)  $\|x \pm y\|_G^2 = \|x\|_G^2 \pm 2\langle x, Gy \rangle + \|y\|_G^2$ ;
- (iii)  $\|P_C(x) - P_C(y)\|_G^2 \leq \|x - y\|_G^2 - \|(P_C(x) - x) - (P_C(y) - y)\|_G^2$ .

Let  $\tilde{D}$  be a symmetric positive definite matrix, and  $A^T \tilde{D} = DA^T$ . Then the norm  $\|\cdot\|_{\tilde{D}}$  is defined by  $\|y\|_{\tilde{D}}^2 = \langle y, \tilde{D}y \rangle$  for  $\forall y \in H$ . We denote by  $P_Q$  the projection operator onto  $Q$  with respect to the norm  $\|\cdot\|_{\tilde{D}}$ . According to the SFP, when GSFP (1.2) has no solution (refer to [2, 14, 15]), we can define

$$f_g(x) = \frac{1}{2} \|(I - P_Q)Ag(x)\|^2$$

and

$$f_D^g(x) = \frac{1}{2} \|Ag(x) - P_QAg(x)\|_D^2 = \langle \tilde{D}(Ag(x) - P_QAg(x)), Ag(x) - P_QAg(x) \rangle,$$

$f_D^g(x)$  is also convex and continuously differentiable in  $H$ . Its gradient operator is

$$\nabla f_D^g(x) = DA^T(I - P_Q)Ag(x).$$

As  $D = I$ , we define

$$\nabla f_g(x) = A^T(I - P_Q)Ag(x),$$

$\nabla f_g(x)$  is also Lipschitz continuous.

**Proposition 2.2** *If we consider the constrained minimization problem*

$$\min\{f_D^g(x) \mid x \in H_1 \text{ s.t. } g(x) \in C\},$$

its stationary point  $x^* \in H_1$  satisfies

$$\begin{cases} x^* \in H, g(x^*) \in C \text{ such that} \\ \langle \nabla f_D^g(x^*), g(x) - g(x^*) \rangle \geq 0, \quad \forall g(x) \in C, \end{cases}$$

which is a general variational inequality involving a Lipschitz continuous and  $G$ -co-coercive operator.

*Proof* For  $\forall x, y \in H$ , from (2.1) and Lemma 8.1 in [15], we have

$$\begin{aligned} \|\nabla f_D^g(x) - \nabla f_D^g(y)\|_G^2 &\leq \lambda_{\max}(G) \|\nabla f_D^g(x) - \nabla f_D^g(y)\|^2 \\ &\leq \frac{\lambda_{\max}^2(D)}{\lambda_{\min}(D)} \|\nabla f_g(x) - \nabla f_g(y)\|^2 \\ &\leq \frac{\lambda_{\max}^2(D)}{\lambda_{\min}(D)} L^2 \|g(x) - g(y)\|^2 \\ &\leq \frac{\lambda_{\max}^3(D)}{\lambda_{\min}(D)} L^2 \|g(x) - g(y)\|_G^2, \end{aligned}$$

where  $L$  is the largest eigenvalue of  $A^T A$ ; therefore, the operator  $\nabla f_D^g$  is Lipschitz continuous,

$$\begin{aligned} &\langle \nabla f_D^g(x) - \nabla f_D^g(y), g(x) - g(y) \rangle \\ &\geq \lambda_{\min}(D) \langle \nabla f_g(x) - \nabla f_g(y), g(x) - g(y) \rangle \\ &\geq \frac{\lambda_{\min}(D)}{L} \|\nabla f_g(x) - \nabla f_g(y)\|^2 \\ &\geq \frac{\lambda_{\min}(D)}{L \cdot \lambda_{\max}(D)} \|\nabla f_g(x) - \nabla f_g(y)\|_D^2 \\ &= \frac{\lambda_{\min}(D)}{L \cdot \lambda_{\max}(D)} \langle D(\nabla f_g(x) - \nabla f_g(y)), GD(\nabla f_g(x) - \nabla f_g(y)) \rangle \\ &= \frac{\lambda_{\min}(D)}{L \cdot \lambda_{\max}(D)} \|\nabla f_D^g(x) - \nabla f_D^g(y)\|_G^2. \end{aligned}$$

Thus, the operator  $\nabla f_D^g$  is co-coercive. □

### 3 Main results

In this section, we propose several modified CQ algorithms with preconditioning techniques and prove the convergence.

#### 3.1 General preconditioning CQ algorithm

In this part, we have our first algorithm with fixed stepsize and preconditioner to solve GSP (1.2). The algorithm is as follows.

**Algorithm 3.1** Choose  $\forall x_0 \in H_1$  such that  $g(x_0) \in C$ , and let  $x_n \in H_1$  such that  $g(x_n) \in C$ , then we calculate  $x_{n+1}$  such that

$$g(x_{n+1}) = P_C[g(x_n) - \gamma \nabla f_D^g(x_n)], \quad n \geq 0, \tag{3.1}$$

where  $\gamma \in (0, \frac{2}{L \cdot L_D})$ ,  $L$  and  $L_D$  are the largest eigenvalues of  $A^T A$  and  $D$ , respectively.

Now we establish the weak convergence of Algorithm 3.1.

**Theorem 3.1** *Suppose that the operators  $g : H_1 \rightarrow C$  and  $g^{-1} : C \rightarrow H_1$  are continuous. If  $\Gamma \neq \emptyset$ , then the sequence  $\{x_n\}$  generated by Algorithm 3.1 converges to the solution of GSPF (1.2).*

*Proof* Firstly, for  $\forall x^* \in \Gamma$ , we have

$$g(x^*) = P_C[g(x^*) - \gamma \nabla f_D^g(x^*)].$$

From (3.1), (iii), (ii) and the definition of ism, we have

$$\begin{aligned} & \|g(x_{n+1}) - g(x^*)\|_G^2 \\ &= \|P_C[g(x_n) - \gamma \nabla f_D^g(x_n)] - P_C[g(x^*) - \gamma \nabla f_D^g(x^*)]\|_G^2 \\ &\leq \|g(x_n) - g(x^*) - \gamma [\nabla f_D^g(x_n) - \nabla f_D^g(x^*)]\|_G^2 \\ &\leq \|g(x_n) - g(x^*)\|_G^2 - 2\gamma \langle g(x_n) - g(x^*), \nabla f_g(x_n) - \nabla f_g(x^*) \rangle \\ &\quad + \gamma^2 \langle \nabla f_D^g(x_n) - \nabla f_D^g(x^*), \nabla f_g(x_n) - \nabla f_g(x^*) \rangle \\ &\leq \|g(x_n) - g(x^*)\|_G^2 - \left(\frac{2\gamma}{L} - \gamma^2 L_D\right) \|\nabla f_g(x_n) - \nabla f_g(x^*)\|^2, \end{aligned} \tag{3.2}$$

as  $\frac{2\gamma}{L} - \gamma^2 L_D > 0$ , which implies that the sequence  $\{\|g(x_n) - g(x^*)\|_G\}_{n \in \mathbb{N}}$  is monotonically decreasing, then we can obtain that the sequence  $\{\|g(x_n) - g(x^*)\|_G\}_{n \in \mathbb{N}}$  is also convergent, especially, the sequence  $\{g(x_n)\}_{n \in \mathbb{N}}$  is bounded. Consequently, we get from (3.2)

$$\lim_{n \rightarrow \infty} \|\nabla f_g(x_n) - \nabla f_g(x^*)\| = \lim_{n \rightarrow \infty} \|\nabla f_g(x_n)\| = 0. \tag{3.3}$$

Moreover, for each  $g(x_n) \in C$ , from (iii) and (2.1), we have

$$\begin{aligned} \|g(x_{n+1}) - g(x_n)\|_G &\leq \gamma \|\nabla f_D(x_n)\|_G \\ &\leq \gamma L_D \|\nabla f_g(x_n)\|_G \leq \gamma L_D \sqrt{\lambda_{\max}(G)} \|\nabla f_g(x_n)\|. \end{aligned}$$

Then by virtue of (3.3) we have

$$\lim_{n \rightarrow \infty} \|g(x_{n+1}) - g(x_n)\|_G = 0. \tag{3.4}$$

Hence, there exists a subsequence  $\{g(x_j)\}_{j \in \bar{\mathbb{N}}}$  of  $\{g(x_n)\}_{n \in \mathbb{N}}$  such that

$$\lim_{j \rightarrow \infty} \|g(x_{j+1}) - g(x_j)\|_G = 0.$$

Thus,  $\{g(x_j)\}_{j \in \bar{\mathbb{N}}}$  is also bounded.

Let  $\bar{x}$  be an accumulation point of  $\{x_n\}$ , then the subsequence of  $\{x_n\}$ ,  $\{x_j\}_{j \in \bar{\mathbb{N}}} \rightarrow \bar{x}$  as  $j \rightarrow \infty$ . Because of the continuity of  $g$ , there exists an accumulation point  $g(\bar{x}) \in C$  of the

sequence  $\{g(x_n)\}_{n \in \mathbb{N}}$ ; for the subsequence  $\{g(x_j)\}_{j \in \bar{\mathbb{N}}}$ , we have  $\{g(x_j)\}_{j \in \bar{\mathbb{N}}} \rightarrow g(\bar{x})$  as  $j \rightarrow \infty$ . After that, from (3.3) we obtain

$$\lim_{j \rightarrow \infty} \|\nabla f_g(x_{n_j})\| = \|\nabla f_g(\bar{x})\| = \|A^T A g(\bar{x}) - A^T P_Q A g(\bar{x})\| = 0,$$

that is,  $A g(\bar{x}) \in Q$ .

We use  $\bar{x}$  in place of  $x^*$  in (3.2) and obtain that  $\{\|g(x_n) - g(\bar{x})\|_G\}$  is convergent. Because its subsequence  $\{\|g(x_{n_j}) - g(\bar{x})\|_G\} \rightarrow 0$ , then we get that  $\{g(x_n)\}_{n \in \mathbb{N}}$  converges to  $g(\bar{x})$  as  $j \rightarrow \infty$ . As well as  $g^{-1}$  is continuous, we finally have

$$\lim_{n \rightarrow \infty} x_n = \bar{x} \in \Gamma.$$

Therefore,  $\bar{x}$  is a solution of GSFP (1.2). □

From Algorithm 3.1 and Theorem 3.1 we can deduce the following results easily.

**Corollary 3.1** *If  $g = I$ , then GSFP (1.2) reduces to SFP, Algorithm 3.1 also reduces to a preconditioning CQ (PCQ) algorithm for  $\forall x_0 \in H_1$ :*

$$x_{n+1} = P_C[x_n - \gamma \nabla f_D(x_n)], \quad n \geq 0, \tag{3.5}$$

where  $\gamma \in (0, \frac{2}{L-L_D})$ ,  $L$  and  $L_D$  are the largest eigenvalues of  $A^T A$  and  $D$ , respectively.  $P_C$  and  $P_Q$  are still the projection operators onto  $C$  and  $Q$  with respect to the norm  $\|\cdot\|_D$  and  $\|\cdot\|_{\tilde{D}}$ , respectively.

**Corollary 3.2** *If  $g = I$ ,  $D = I$ ,  $\tilde{D} = I$ , then GSFP reduces to SFP, then Algorithm 3.1 reduces to the CQ algorithm proposed in [25].*

**Corollary 3.3** *If  $g = I$ ,  $\tilde{D} = I$ ,  $P_C$  and  $P_Q$  are the projections onto  $C$  and  $Q$  with respect to the norm  $\|\cdot\|$ , set  $F(x_n) = DA^T(I - P_Q)ADx_n$ , (3.5) transforms into the algorithm in [26]*

$$x_{n+1} = P_C[x_n - \gamma F(x_n)], \quad n \geq 0,$$

where  $\gamma \in (0, \frac{2}{L})$ ,  $L = \|DA^T\|^2$ . Then the GSFP reduces to the extended split feasibility problem (ESFP) in [26].

### 3.2 An algorithm with variable projection metric

The algorithms above can speed the convergence of CQ algorithm, but the stepsize and preconditioner are fixed. In this subsection, we extend the results in [6] and construct an iterative scheme with variable stepsize and preconditioner  $D_n$  from one iteration to the next. As a key role,  $D_n$  will change arbitrarily or following some rules to achieve the convergence progress and better results.

Let  $D_n$  and  $\tilde{D}_n$  be two symmetric positive definite matrices for  $n = 0, 1, 2, \dots$ . Denote by  $P_C$  and  $P_Q$  the projections onto  $C$  and  $Q$  with respect to the norm  $\|\cdot\|_{D_n}$  and  $\|\cdot\|_{\tilde{D}_n}$ , respectively. Let  $\chi$  be a set of symmetric positive definite matrices, we have the following algorithm.

**Algorithm 3.2** Choose  $\forall x_0 \in H_1$  such that  $g(x_0) \in C$ , and let  $x_n \in H_1$  such that  $g(x_n) \in C$ ; for  $\forall D_n \in \chi$ , we compute  $x_{n+1}$  such that

$$g(x_{n+1}) = P_C[g(x_n) - \gamma_n D_n \nabla f_g(x_n)], \quad n \geq 0, \tag{3.6}$$

where  $\gamma_n \in (0, \frac{2}{L M_D})$ ,  $L$  is the largest eigenvalue of  $A^T A$ ,  $M_D$  is the minimum value of all the largest eigenvalue values  $L_{D_n}$  to matrices  $D_n$ .

**Remark 3.1** Define  $d_n = \|g(x_{n+1}) - g(x_n)\|_{G_n}$ , then for the next iteration,  $D_{n+1}$  is either chosen arbitrarily from  $\chi$  or equivalent to  $D_n$ . It is conditional on whether  $d_n$  has decreased or not. More particularly, we define a scalar  $\bar{d}_n$  with initial value  $\bar{d}_0 = \infty$ . Having chosen a scalar  $\alpha \in (0, 1)$  at the  $n$ th iteration, then  $\bar{d}_{n+1}$  is calculated by

$$\bar{d}_{n+1} = \begin{cases} \alpha \bar{d}_n, & \text{if } d_n \leq \bar{d}_n; \\ \bar{d}_n, & \text{if } d_n > \bar{d}_n, \end{cases}$$

then we select

$$D_{n+1} = \begin{cases} \forall D \in \chi, & \text{if } \bar{d}_{n+1} < \bar{d}_n; \\ D_n, & \text{if } \bar{d}_{n+1} = \bar{d}_n. \end{cases}$$

**Theorem 3.2** If  $\Gamma \neq \emptyset$ , then the sequence  $\{x_n\}$  generated by Algorithm 3.2 converges to the solution of GSFP (1.2).

*Proof* To obtain variable  $D_n$  at each iteration and keep the convergence of Algorithm 3.2,  $d_n$  must be a descending behavior for  $n = 0, 1, 2, \dots$ . We first show that

$$\liminf_{n \rightarrow \infty} d_n = 0. \tag{3.7}$$

Indeed if (3.7) is not true, we have  $\liminf_{n \rightarrow \infty} d_n > 0$ , then  $D_n$  must have changed a finite number of times, we set that this number is  $\kappa \in N$ . Therefore, let  $x^* \in \Gamma$  be a solution of GSFP, refer to (3.2) and for  $n > \kappa$ , we have

$$\begin{aligned} \|g(x_{n+1}) - g(x^*)\|_{G_\kappa}^2 &\leq \|g(x_n) - g(x^*)\|_{G_\kappa}^2 \\ &\quad - \left( \frac{2\gamma_\kappa}{L} - \gamma_\kappa^2 L_{D_\kappa} \right) \|\nabla f_g(x_n) - \nabla f_g(x^*)\|^2, \end{aligned}$$

as  $M_D = \min\{L_{D_n} \mid n = 0, 1, \dots, \kappa\}$ ,  $\frac{2\gamma_\kappa}{L} - \gamma_\kappa^2 L_{D_\kappa} > 0$ . Then, following the proof of Theorem 3.1, we also get

$$\lim_{n \rightarrow \infty} \|\nabla f_g(x_n)\| = 0 \tag{3.8}$$

and

$$\lim_{n \rightarrow \infty} d_n = \lim_{n \rightarrow \infty} \|g(x_{n+1}) - g(x_n)\|_{G_\kappa} = 0. \tag{3.9}$$

Equation (3.9) contradicts the above hypothesis, so (3.7) is true.

By using (3.6) and (iii), for the  $n$ th iteration, we have

$$\begin{aligned} d_n &= \|g(x_{n+1}) - g(x_n)\|_{G_n}^2 \\ &\geq \frac{1}{2} \|g(x_n) - P_C[g(x_n)]\|_{G_n}^2 - \|g(x_{n+1}) - P_C[g(x_n)]\|_{G_n}^2 \\ &= \frac{1}{2} \|g(x_n) - P_C[g(x_n)]\|_{G_n}^2 - \|P_C[g(x_n) - \gamma_n D_n \nabla f_g(x_n)] - P_C[g(x_n)]\|_{G_n}^2 \\ &\geq \frac{1}{2} \|g(x_n) - P_C[g(x_n)]\|_{G_n}^2 - \gamma_n L_{D_n} \|\nabla f_g(x_n)\|_{G_n}^2, \end{aligned}$$

where  $\gamma_n L_{D_n} > 0$ . Then from (2.1) and (3.8) we know that

$$\lim_{n \rightarrow \infty} \|\nabla f_g(x_n)\|_{G_n} = \lim_{n \rightarrow \infty} \|A^T(I - P_Q)Ag(x_n)\|_{G_n} = 0,$$

so  $Ag(x_n) \in Q$ . By virtue of (3.7), we also have

$$\lim_n \|g(x_n) - P_C[g(x_n)]\| = 0.$$

This means that at least a subsequence of  $\{g(x_n)\}_{n \in \mathbb{N}}$  converges to a solution of  $g(\bar{x}) \in \Gamma$ . Similar to the argumentation of accumulation in the proof of Theorem 3.1, we know that  $\{x_n\}$  converges to a solution of GSFP.  $\square$

### 3.3 Some methods for execution

In Algorithms 3.1 and 3.2, there still exists difficulty to implement the projections  $P_C$  and  $P_Q$  with respect to the defined norms, especially when  $C$  and  $Q$  are general closed convex sets. According to the relaxed method in [8, 27, 28], we consider the above algorithm in which the closed convex subsets  $C$  and  $Q$  are the following particular formula:

$$C = \{g(x) \in H_1 \mid c(g(x)) \leq 0\} \quad \text{and} \quad Q = \{Ag(x) \in H_2 \mid q(Ag(x)) \leq 0\},$$

where  $c: H_1 \rightarrow \mathbb{R}$  and  $q: H_2 \rightarrow \mathbb{R}$  are convex functions.  $C_n$  and  $Q_n$  are given as

$$\begin{aligned} C_n &= \{g(x) \in H_1 \mid c(g(x_n)) + \langle \xi_n, g(x) - g(x_n) \rangle \leq 0\}, \\ Q_n &= \{Ag(x) \in H_1 \mid q(Ag(x_n)) + \langle \eta_n, Ag(x) - Ag(x_n) \rangle \leq 0\}, \end{aligned}$$

where  $\xi_n \in \partial c(g(x_n))$ ,  $\eta_n \in \partial q(Ag(x_n))$ .

Here, we also replace  $P_C$  and  $P_Q$  by  $P_{C_n}$  and  $P_{Q_n}$ . However, in this paper, take Algorithm 3.2 for example, the projections are with respect to the norms corresponding to  $G_n$  and  $\tilde{D}_n$ , we should use the following methods to calculate them. For  $\forall z \in H_1$  and  $\forall y \in H_2$ ,

$$P_{C_n}(z) = \begin{cases} z - D_n \frac{c[g(x_n)] + \langle D_n \xi_n, z - g(x_n) \rangle}{\|\xi_n\|_{D_n}^2} \xi_n, & \text{if } c[g(x_n)] + \langle D_n \xi_n, z - g(x_n) \rangle > 0; \\ z, & \text{otherwise} \end{cases}$$

and

$$P_{Q_n}(y) = \begin{cases} y - \tilde{D}_n \frac{q[Ag(x_n)] + \langle \tilde{D}_n \eta_n, y - Ag(x_n) \rangle}{\|\eta_n\|_{\tilde{D}_n}^2} \eta_n, & \text{if } q[Ag(x_n)] + \langle \tilde{D}_n \eta_n, y - Ag(x_n) \rangle > 0; \\ y, & \text{otherwise.} \end{cases}$$



Set  $z = g(x_n) - \gamma_n D_n \nabla f_g(x_n)$ ,  $y = Ag(x_n)$ , let  $\bar{x} \in H_1$  be an accumulation of  $\{x_n\}_{n \in \mathbb{N}}$ . From the proof above, it is easy to deduce that

$$\lim_{n \rightarrow \infty} c[g(x_n)] + \langle D_n \xi_n, z - g(x_n) \rangle = c[g(\bar{x})] \leq 0,$$

$$\lim_{n \rightarrow \infty} q[Ag(x_n)] + \langle \tilde{D}_n \eta_n, y - Ag(x_n) \rangle = q[Ag(\bar{x})] \leq 0.$$

Therefore,  $g(\bar{x}) \in C \subseteq C_n$ ,  $Ag(\bar{x}) \in Q \subseteq Q_n$ , with the projections  $P_{C_n}$  and  $P_{Q_n}$ ,  $\bar{x}$  is a solution of GSFP.

Next, we present a new approximation method to estimate  $\gamma_n$  and  $D_n$  in Algorithm 3.2.

If  $\Gamma \neq \emptyset$ , for  $\forall x^* \in \Gamma$  and  $n \geq 0$ , such that

$$D_n \nabla f_g(x^*) = D_n A^T Ag(x^*) - D_n A^T P_{Q_n} Ag(x^*) = 0,$$

under the ideal condition, if  $D_n A^T A \approx I$ , the solution is done, but unfortunately,  $(A^T A)^{-1}$  cannot be calculated directly when  $A$  is a large matrix in practice. As

$$A^T Ag(x^*) = A^T P_{Q_n} Ag(x^*) = \lambda g(x^*),$$

where  $\lambda$  is an eigenvalue of  $A^T A$ . Let  $D_0 = I$ , for the  $n$ th iteration, we have the next  $j \times j$  approximation of  $D_{n+1}$

$$D_{n+1}^{jj} = \begin{cases} \frac{[g(x_n)]_j}{[A^T P_{Q_n} Ag(x_n)]_j}, & \text{if } [g(x_n)]_j \neq 0 \text{ and } [A^T P_{Q_n} Ag(x_n)]_j \neq 0; \\ D_n^{jj}, & \text{otherwise,} \end{cases}$$

where  $j = 1, 2, \dots$ . So, at the  $n$ th iteration, let  $l_{D_n}$  be the minimum eigenvalue of  $D_n$ , take  $M_{D_n} \approx \min\{L_{D_k} \mid k = 0, 1, \dots, n\}$ ,  $L_n \approx \max\{(l_{D_k})^{-1} \mid k = 0, 1, \dots, n\}$ , the variable stepsize is approximated by

$$\gamma_n = \frac{\rho_n}{L_n \cdot M_{D_n}}, \quad \rho_n \in (0, 2), n = 1, 2, \dots$$

#### 4 Numerical results

We consider the following problem from [29] in a finite dimensional Hilbert space:

Let  $C = \{x \in H_1 \mid c(x) \leq 0\}$ , where  $c(x) = -x_1 + x_2^2 + \dots + x_N^2$ , and  $Q = \{y \in H_2 \mid q(y) \leq 0\}$ , where  $q(y) = y_1 + y_2^2 + \dots + y_M^2 - 1$ .  $A_{M \times N}$  is a random matrix where every element of  $A$  is in  $(0, 1)$  satisfying  $\Gamma \neq \emptyset$ . Let  $x_0$  be a random vector in  $H_1$  where every element of  $x_0$  is in  $(0, 1)$ .

We set  $\|x_{n+1} - x_n\| \leq \varepsilon$  as the stop rule, and let  $N = 10$ ,  $M = 20$ ,  $g = I$ ,  $\tilde{D}_n = I$ , for  $n \geq 0$ . Using the methods in Section 3.3, we compare Algorithm 3.2 with the relaxed CQ algorithm (RCQ) in [30], with different  $\varepsilon$  and initial values. The results can be seen in Table 1. We see that the proposed methods in this paper behave better.

#### 5 Concluding remarks

In this paper, we have discussed a new general split feasibility problem, which is related to the general variational inequalities involving a co-coercive operator. By using the  $G$ -norm

**Table 1** The comparison between the results of preconditioning and relaxed CQ algorithms

$\varepsilon$	Algorithms	$n$	CPU (s)	$c(x)$	$q(y)$
0.01	RCQ	16	0.0112	-0.1665	57.1092
	3.2	22	0.0049	-0.1333	1.2636
0.001	RCQ	200	0.0216	-0.2684	1.2708
	3.2	13	0.0039	-0.0770	3.5300E-02
0.0001	RCQ	360	0.0295	-0.1039	1.0710E-01
	3.2	60	0.0081	-0.0479	1.4900E-02
0.00001	RCQ	464	0.0363	-0.2561	7.5000E-03
	3.2	21	0.0043	-0.0566	3.3809E-04

method, variable modulus method and relaxed method, two modified projection algorithms for solving the GSPF and some approximate methods for algorithm executing have been presented. The numerical results show that by preconditioning method, the convergence speed of CQ algorithm can be improved, but the way to obtain variable stepsize in the paper is inexact. To continue to improve it or combine it with the methods in [14] and [28] is another interesting subject.

#### Competing interests

The authors declare that they have no competing interests.

#### Authors' contributions

The authors take equal roles in deriving results and writing of this paper. All authors read and approved the final manuscript.

#### Author details

<sup>1</sup>The Second Training Base, Naval Aviation Institution, Huludao, 125001, China. <sup>2</sup>Department of Mathematics, Shijiazhuang Mechanical Engineering College, Shijiazhuang, 050003, China. <sup>3</sup>Department of Mathematics and Information, Hebei Normal University, Shijiazhuang, 050024, China.

#### Acknowledgements

The authors would like to thank the associate editor and the referees for their comments and suggestions. This research was supported by the National Natural Science Foundation of China (11071053).

Received: 14 February 2014 Accepted: 17 October 2014 Published: 31 Oct 2014

#### References

- Chen, K: Matrix Preconditioning Techniques and Applications. Cambridge University Press, New York (2005)
- Piana, M, Bertero, M: Projected Landweber method and preconditioning. *Inverse Probl.* **13**, 441-463 (1997)
- Strand, ON: Theory and methods related to the singular-function expansion and Landweber's iteration for integral equations of the first kind. *SIAM J. Numer. Anal.* **11**, 798-824 (1974)
- Auslender, A: Optimisation: Méthodes Numérique. Masson, Paris (1976)
- Dafermos, S: Traffic equilibrium and variational inequalities. *Transp. Sci.* **14**, 42-54 (1980)
- Bertsekas, PD, Gafni, EM: Projection methods for variational inequities with application to the traffic assignment problem. *Math. Program. Stud.* **17**, 139-159 (1982)
- Marcotte, P, Wu, JH: On the convergence of projection methods: application to the decomposition of affine variational inequalities. *J. Optim. Theory Appl.* **85**, 347-362 (1995)
- Fukushima, M: A relaxed projection method for variational inequalities. *Math. Program.* **35**, 58-70 (1986)
- Yang, Q: The revisit of a projection algorithm with variable steps for variational inequalities. *J. Ind. Manag. Optim.* **1**, 211-217 (2005)
- He, BS: Inexact implicit methods for monotone general variational inequalities. *Math. Program.* **86**, 199-217 (1999)
- Noor, MA, Wang, YJ, Xiu, N: Projection iterative schemes for general variational inequalities. *J. Inequal. Pure Appl. Math.* **3**, Article 34 (2002)
- Santos, PSM, Scheimberg, S: A projection algorithm for general variational inequalities with perturbed constraint sets. *Appl. Math. Comput.* **181**, 649-661 (2006)
- Muhammad, AN, Abdellah, B, Saleem, U: Self-adaptive methods for general variational inequalities. *Nonlinear Anal.* **71**, 3728-3738 (2009)
- Qu, B, Xiu, N: A note on the CQ algorithm for the split feasibility problem. *Inverse Probl.* **21**, 1655-1665 (2005)
- Byrne, C: A unified treatment of some iterative algorithms in signal processing and image reconstruction. *Inverse Probl.* **20**, 103-120 (2004)
- Dolidze, Z: Solution of variational inequalities associated with a class of monotone maps. *Èkon. Mat. Metody* **18**, 925-927 (1982)
- He, B, He, X, Liu, H, Wu, T: Self-adaptive projection method for co-coercive variational inequalities. *Eur. J. Oper. Res.* **196**, 43-48 (2009)

18. Facchinei, F, Pang, JS: *Finite-Dimensional Variational Inequality and Complementarity Problems*, vol. I. Springer, New York (2003)
19. Facchinei, F, Pang, JS: *Finite-Dimensional Variational Inequality and Complementarity Problems*, vol. II. Springer, New York (2003)
20. Yao, YH, Postolache, M, Liou, YC: Strong convergence of a self-adaptive method for the split feasibility problem. *Fixed Point Theory Appl.* **2013**, 201 (2013)
21. Yao, YH, Yang, PX, Kang, SM: Composite projection algorithms for the split feasibility problem. *Math. Comput. Model.* **57**, 693-700 (2013)
22. Yao, YH, Liou, YC, Shahzad, N: A strongly convergent method for the split feasibility problem. *Abstr. Appl. Anal.* **2012**, Article ID 125046 (2012)
23. Mohammad, E, Abdul, L: General split feasibility problems in Hilbert spaces. *Abstr. Appl. Anal.* **2013**, Article ID 805104 (2013)
24. Censor, Y, Elfving, T: A multiprojection algorithm using Bregman projections in a product space. *Numer. Algorithms* **8**, 221-239 (1994)
25. Byrne, C: Iterative oblique projection onto convex sets and the split feasibility problem. *Inverse Probl.* **18**, 441-453 (2002)
26. Wang, PY, Zhou, HY: A preconditioning method of the CQ algorithm for solving the extended split feasibility problem. *J. Inequal. Appl.* **2014**, 163 (2014). doi:10.1186/1029-242X-2014-163
27. Yang, Q: On variable-step relaxed projection algorithm for variational inequalities. *J. Math. Anal. Appl.* **302**, 166-179 (2005)
28. López, G, Martín-Márquez, M, Wang, F, Xu, H-K: Solving the split feasibility problem without prior knowledge of matrix norms. *Inverse Probl.* **28**, 085004 (2012)
29. Wang, Z, Yang, Q, Yang, Y: The relaxed inexact projection methods for the split feasibility problem. *Appl. Math. Comput.* **217**, 5347-5359 (2011)
30. Yang, Q: The relaxed CQ algorithm solving the split feasibility problem. *Inverse Probl.* **20**, 1261-1266 (2004)

10.1186/1029-242X-2014-435

**Cite this article as:** Wang et al.: Preconditioning methods for solving a general split feasibility problem. *Journal of Inequalities and Applications* 2014, **2014**:435

**Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:**

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

---

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)

---