

Technical University of Denmark



## Mass loss of the Greenland Ice Sheet since the Little Ice Age, implications on sea level

Kjeldsen, K. K.; Bjork, A. A.; Khan, Shfaqat Abbas; Korsgaard, N. J.; Larsen, N. K.; Long, A. J.; Woodroffe, S.; Milne, G. A.; Wahr, J. M.; Bamber, J. A.; van den Broeke, M. R.

*Publication date:*  
2013

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Kjeldsen, K. K., Bjork, A. A., Khan, S. A., Korsgaard, N. J., Larsen, N. K., Long, A. J., ... van den Broeke, M. R. (2013). Mass loss of the Greenland Ice Sheet since the Little Ice Age, implications on sea level. Abstract from AGU Fall Meeting 2013, San Francisco, United States.

## DTU Library

Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**CONTROL ID:** 1805858

**TITLE:** Mass loss of the Greenland Ice Sheet since the Little Ice Age, implications on sea level

**AUTHORS (FIRST NAME, LAST NAME):** Kristian K Kjeldsen<sup>1</sup>, Kurt Henrik Kjaer<sup>1</sup>, Anders A Bjork<sup>1</sup>, Shfaqat Abbas Khan<sup>2</sup>, Niels J Korsgaard<sup>1</sup>, Nicolaj K Larsen<sup>3</sup>, Antony J Long<sup>4</sup>, Sarah Woodroffe<sup>4</sup>, Glenn A. Milne<sup>5</sup>, John M Wahr<sup>6</sup>, Geruo A<sup>6</sup>, Jonathan L Bamber<sup>7</sup>, Michiel R van den Broeke<sup>8</sup>

**INSTITUTIONS (ALL):** 1. Centre for GeoGenetics, Natural History Museum of Denmark, Copenhagen, Denmark.  
2. DTU Space - Department of Geodesy, Technical University of Denmark, Kgs. Lyngby, Denmark.  
3. Department of Geoscience, Aarhus University, Aarhus, Denmark.  
4. Department of Geography, University of Durham, Durham, United Kingdom.  
5. Department of Earth Sciences, University of Ottawa, Ottawa, ON, Canada.  
6. CIRES, University of Boulder, Boulder, CO, United States.  
7. Bristol Glaciology Centre, University of Bristol, Bristol, United Kingdom.  
8. Institute for Marine and Atmospheric Research, Utrecht University, Utrecht, Netherlands.

**ABSTRACT BODY:** The impact of mass loss from the Greenland Ice Sheet (GrIS) on 20th Century sea level rise (SLR) has long been subject to intense discussions. While globally distributed tide gauges suggest a global mean SLR of 15-20 cm, quantifying the separate components is of great concern - in particular for modeling sea level projections into the 21st Century. Estimates of the past GrIS contribution to SLR have been derived using a number of different approaches, e.g. surface mass balance (SMB) calculations combined with estimates of ice discharge found by in correlating SMB anomalies and calving rates. Here, we adopt a novel geometric approach to determine the post-Little Ice Age (LIA) mass loss of the GrIS.

We use high quality aerial stereo photogrammetric imagery recorded between 1978 and 1987 to map morphological features such as trim lines (boundary between freshly eroded and non-eroded bedrock) and end moraines marking the ice extent of the LIA, which thereby enables us to obtain vertical point-based differences associated with changes in ice extent. These point measurements are combined with contemporary ice surface differences derived using NASA's Airborne Topographic Mapper (ATM) from 2002-2010, NASA's Ice, Cloud, and land Elevation Satellite (ICESat) from 2003-2009, and NASA's Land, Vegetation, and Ice Sensor (LVIS) from 2010, to estimate mass loss throughout the 20th and early 21st Century.

We present mass balance estimates of the GrIS since retreat commence from the maximum extent of the LIA to 2010 derived for three intervals, LIAMax (1900) - 1978/87, 1978/87 - 2002, and 2002 - 2010. Results suggest that despite highly spatially- and temporally variable post-LIA mass loss, the total mass loss and thus the contribution from the GrIS to global SLR has accelerated significantly during the 20th Century.

**KEYWORDS:** 0700 CRYOSPHERE, 0726 CRYOSPHERE Ice sheets, 0762 CRYOSPHERE Mass balance 0764 Energy balance, 0758 CRYOSPHERE Remote sensing.

(No Image Selected)

(No Table Selected)

### **Additional Details**

**Previously Presented Material:**

### **Contact Details**

**CONTACT (NAME ONLY):** Kristian Kjeldsen

**CONTACT (E-MAIL ONLY):** kkjeldsen@snm.ku.dk