

RESEARCH

Open Access

Existence results of a kind of Sturm-Liouville type singular boundary value problem with non-linear boundary conditions

Junfang Zhao^{1*} and Weigao Ge²

*Correspondence:
zhao_junfang@163.com
¹School of Science, China University of Geosciences, Beijing, 100083, P.R. China
Full list of author information is available at the end of the article

Abstract

By using the Leggett-Williams fixed-point theorem, a series of Sturm-Liouville type boundary value problems are studied. We impose sufficient conditions on f and get the existence of at least three positive solutions. The interesting point is that the boundary conditions are non-linear.

Keywords: Sturm-Liouville type; positive solutions; boundary value problem; fixed point theorem; p -Laplacian

1 Introduction

In this paper, we consider the existence of triple positive solutions of a kind of four-point boundary value problem

$$(\phi_p(u'(t)))' + h(t)f(t, u(t)) = 0, \quad 0 < t < 1, \quad (1.1)$$

subject to one of the following boundary conditions:

$$g_1(u'(0)) - u(\xi) = 0, \quad \beta u'(1) + u(\eta) = 0, \quad (1.2)$$

$$\alpha u'(0) - u(\xi) = 0, \quad g_2(u'(1)) + u(\eta) = 0, \quad (1.3)$$

where $\phi_p(s) = |s|^{p-2}s$, $p > 1$, $\alpha \geq \xi$, $\beta \geq 1 - \eta$, $0 < \xi < \eta < 1$, and g_1, g_2 satisfy

(H₁) (i) $g_1 \in C((-\infty, +\infty), (-\infty, +\infty))$, and $g_1(0) = 0$;

(ii) $g_1(x) \geq \xi x$ for $x \in [0, +\infty)$;

(iii) g_1 is increasing on x .

(H₂) (i) $g_2 \in C((-\infty, +\infty), (-\infty, +\infty))$ and g_2 is odd;

(ii) $g_2(x) \geq (1 - \eta)x$ for $x \in [0, +\infty)$;

(iii) g_2 is increasing on x .

We will assume throughout:

(C₁) $h(t) \in C((0, 1), (0, +\infty))$, which implies that $h(t)$ may be singular at $t = 0$ or/and $t = 1$.
 $0 < \int_0^1 h(t) dt < +\infty$, and $h(t)$ is not identically zero on any subinterval of $[0, 1]$.

(C₂) $f \in C([0, 1] \times [0, +\infty), [0, +\infty))$, $f \not\equiv 0$.

Boundary value problems (BVPs) for ordinary differential equations play a very important role in both theory and application. They are used to describe a large number of physical, biological and chemical phenomena. And in analyzing these non-linear phenomena, many mathematical models give rise to the problems for which only nonnegative solutions make sense. Therefore, since the publication of the monograph *Positive Solutions of Operator Equations* in 1964 by the academician Krasnosel'skii, many research articles on the theory of positive solutions of nonlinear problems have appeared [1–5]. In this vast field of research, particular attention has been focused on the BVPs with linear boundary conditions. However, BVPs with non-linear boundary conditions have not received much attention. Obviously, studying such kind of BVPs is more practical but more challenging. The main challenge is that it is not easy to construct an operator and, in turn, to get its solutions.

In [6], Kong *et al.* studied

$$(\Phi_p(u'(t)))' + q(t)f(u(t)) = 0, \quad 0 < t < 1 \tag{1.4}$$

subject to one of the following non-linear boundary conditions:

$$\begin{aligned} u(0) - g_1(u'(0)) &= 0, & u(1) &= 0, \\ u'(0) &= 0, & u(1) + g_2(u'(1)) &= 0, \end{aligned} \tag{1.5}$$

where $\Phi_p(v) = |v|^{p-2}v$, $p > 1$, $q(t)$ is a nonnegative measurable function, $f(u)$ is a nonnegative continuous function on $[0, +\infty)$, and $g_1(v)$, $g_2(v)$ are all continuous functions defined on $(-\infty, +\infty)$. By using the theory of fixed point index, the authors get the existence of at least two positive solutions. For more references about the classical Sturm-Liouville BVP, we refer the readers to [7–17].

BC ' $u'(0) - \alpha(u(\xi)) = 0$, $u'(1) + \beta(u(\eta)) = 0$ (*)' which is the special case of BC (1.2) and (1.3) can be called Sturm-Liouville type BC, the reason is that when $\xi = 0$, $\eta = 1$, BC (*) become the classic Sturm-Liouville BC. To the best knowledge of the authors, few papers have dealt with equation (1.1) subject to the non-linear BC (1.2) and (1.3). So to fill this gap, motivated by the results mentioned above, in this paper we consider BVP (1.1), (1.2) and (1.1), (1.3). At the end of this paper, we also give its generalization which is more universal.

On the other hand, the main tool used in this paper is the Leggett-Williams fixed point theorem, which is different from [6]. In order to apply this fixed point theorem to our paper, we need to make use of some techniques such as the definition of the cone and the zooming of inequalities *etc.* The interesting point in this paper is that the boundary condition is the non-linear Sturm-Liouville type boundary condition.

2 Background material and the fixed point theorem

In this section, for convenience, we present the main definitions and the theorem that will be used in this paper.

Definition 2.1 Let E be a real Banach space. A nonempty closed convex set $P \subset E$ is called a cone if it satisfies the following two conditions:

- (i) $\lambda u \in P$ for all $\lambda \geq 0$,
- (ii) $u, -u \in P$ implies $u = 0$.

Every cone $P \subset E$ induces an ordering in E given by $x \geq y$ if and only if $y - x \in P$.

Definition 2.2 The map α is said to be a nonnegative concave continuous function provided that $\alpha : P \rightarrow [0, \infty)$ is continuous and

$$\alpha(\lambda x + (1 - \lambda)y) \geq \lambda\alpha(x) + (1 - \lambda)\alpha(y)$$

for all $x, y \in P$ and $0 \leq \lambda \leq 1$.

Let a, b, r be constants, $P_r = \{u \in P : \|u\| < r\}$, $P(\alpha, b, d) = \{u \in P : a \leq \alpha(u), \|u\| \leq b\}$.

Theorem 2.1 (Leggett-Williams [18]) *Let $A : \bar{P}_c \rightarrow \bar{P}_c$ be a completely continuous map and α be a nonnegative continuous concave functional on P such that for $\forall u \in \bar{P}_c$, there holds $\alpha u \leq \|u\|$. Suppose there exist a, b, d with $0 < a < b < d \leq c$ such that*

- (i) $\{u \in P(\alpha, b, d) : \alpha u > b\} \neq \emptyset$ and $\alpha(Au) > b$ for all $u \in P(\alpha, b, d)$;
- (ii) $\|Au\| < a$ for all $u \in \bar{P}_a$;
- (iii) $\alpha(Au) > b$ for all $u \in P(\alpha, b, d)$ with $\|Au\| > d$.

Then A has at least three fixed points u_1, u_2, u_3 satisfying

$$\|u_1\| < a, \quad b < \alpha(u_2), \quad \|u_3\| > a, \quad \text{and} \quad \|u_3\| < b.$$

In what follows, by ϕ_q we denote the inverse to ϕ_p , where $\frac{1}{p} + \frac{1}{q} = 1$.

3 Existence results for BVP (1.1), (1.2)

3.1 Some useful lemmas

Lemma 3.1 *Suppose that $v \in C[0, 1]$, $v(t) \geq 0$, $v(t) \not\equiv 0$ on any subinterval of $[0, 1]$. Then BVP*

$$\begin{cases} (\phi_p(u'(t)))' + v(t) = 0, & 0 < t < 1, \\ g_1(u'(0)) - u(\xi) = 0, & \beta u'(1) + u(\eta) = 0, \end{cases} \tag{3.1}$$

has the unique solution

$$u(t) = \begin{cases} g_1(\phi_q(\int_0^\sigma v(s) ds)) + \int_\xi^t \phi_q(\int_s^\sigma v(\tau) d\tau) ds, & (\diamond) \quad t \in [0, \sigma], \\ \beta\phi_q(\int_\sigma^1 v(s) ds) + \int_t^\eta \phi_q(\int_t^s v(\tau) d\tau) ds, & (\heartsuit) \quad t \in [\sigma, 1], \end{cases} \tag{3.2}$$

where σ is a solution of the equation

$$A_1(t) - A_2(t) = 0, \quad t \in [0, 1]. \tag{3.3}$$

And $A_1(t), A_2(t)$ are defined as follows:

$$\begin{aligned} A_1(t) &= g_1\left(\phi_q\left(\int_0^t v(s) ds\right)\right) + \int_\xi^t \phi_q\left(\int_s^t v(\tau) d\tau\right) ds, \\ A_2(t) &= \beta\phi_q\left(\int_t^1 v(s) ds\right) + \int_t^\eta \phi_q\left(\int_t^s v(\tau) d\tau\right) ds. \end{aligned}$$

Proof At the beginning, we show that (3.3) has a unique solution.

$$\begin{aligned} &A_1(t) - A_2(t) \\ &= g_1\left(\phi_q\left(\int_0^t v(s) \, ds\right)\right) + \int_t^\xi \phi_q\left(\int_t^s v(\tau) \, d\tau\right) \, ds \\ &\quad - \beta\phi_q\left(\int_t^1 v(s) \, ds\right) - \int_t^\eta \phi_q\left(\int_t^s v(\tau) \, d\tau\right) \, ds \\ &= g_1\left(\phi_q\left(\int_0^t v(s) \, ds\right)\right) + \int_\xi^\eta \phi_q\left(\int_s^t v(\tau) \, d\tau\right) \, ds + \beta\phi_q\left(\int_1^t v(s) \, ds\right). \end{aligned}$$

Obviously, $A_1(t) - A_2(t)$ is strictly increasing on $t \in [0, 1]$. At the same time, we can notice that $A_1(0) - A_2(0) < 0$, and $A_1(1) - A_2(1) > 0$. Further, $A_1(t) - A_2(t)$ is continuous, thus there must exist one point $\sigma \in (0, 1)$ such that $A_1(t) - A_2(t) = 0$, which means that $A_1(t), A_2(t)$ must intersect at a unique point σ .

Next, we intend to prove that the solution to BVP (3.1) can be expressed as (3.2). Let u be a solution of BVP (3.1). $(\phi_p(u'(t)))' = -v(t) \leq 0$ implies $u''(t) \leq 0$, together with the boundary condition, we claim that there must exist one point $\sigma \in (0, 1)$ such that $u'(\sigma) = 0$. (It can be easily checked that σ cannot be 0 or 1. In fact, if $u'(0) = 0$, we have $u(\xi) = 0$, $u'(1) < 0$, $u(\eta) > 0$, which deduces a contradiction.) If not, without loss of generality, we suppose that $u'(t) < 0$ for all $t \in (0, 1)$. Then $u(t)$ must be decreasing on $t \in (0, 1)$, using the boundary condition in equation (3.1), we have $u(\xi) = g_1(u'(0)) < 0$, $u(\eta) = -\beta u'(1) > 0$, which is a contradiction

$$-(\phi_p(u'(t)))' = v(t). \tag{3.4}$$

Integrate both sides of (3.4) from t to σ to conclude

$$\phi_p(u'(t)) = \int_t^\sigma v(s) \, ds.$$

Then

$$u'(t) = \phi_q\left(\int_t^\sigma v(s) \, ds\right). \tag{3.5}$$

Integrate both sides of (3.5) from 0 to t to conclude

$$u(t) = u(0) + \int_0^t \phi_q\left(\int_s^\sigma v(\tau) \, d\tau\right) \, ds. \tag{3.6}$$

By (3.5) and (3.6), we have

$$\begin{aligned} u'(0) &= \phi_q\left(\int_0^\sigma v(s) \, ds\right), \\ u(\xi) &= u(0) + \int_0^\xi \phi_q\left(\int_s^\sigma v(\tau) \, d\tau\right) \, ds. \end{aligned}$$

Considering the BC in (3.1), we arrive at

$$u(0) = g_1 \left(\phi_q \left(\int_0^\sigma v(s) ds \right) \right) - \int_0^\xi \phi_q \left(\int_s^\sigma v(\tau) d\tau \right) ds. \tag{3.7}$$

Substitute (3.7) into (3.6), we get

$$u(t) = g_1 \left(\phi_q \left(\int_0^\sigma v(s) ds \right) \right) + \int_\xi^t \phi_q \left(\int_s^\sigma v(\tau) d\tau \right) ds. \tag{3.8}$$

By the similar argument, we can obtain

$$u(t) = \beta \phi_q \left(\int_\sigma^1 v(s) ds \right) + \int_\sigma^\eta \phi_q \left(\int_\sigma^s v(\tau) d\tau \right) ds. \tag{3.9}$$

Hence, for $t \in [0, 1]$, the solution of (3.1) can be expressed as (3.2), which completes our proof. \square

Remark 3.1 In fact, for $t \in [0, 1]$, the solution of (3.1) can be expressed both by (\diamond) and (\heartsuit) , but just for convenience, we write it into two parts.

Lemma 3.2 Let $v(t)$ satisfy all the conditions in Lemma 3.1, then the solution $u(t)$ of BVP (3.1) is concave on $t \in [0, 1]$. What is more, $u(t) \geq 0$.

Proof $(\phi_p(u'(t)))' = -v(t) \leq 0$ means $u''(t) \leq 0$, so $u(t)$ is concave on $t \in [0, 1]$.

Next, we try to prove that $u(t) \geq 0$. By Lemma 3.1, we know that $u(t)$ can be expressed as (3.2). When $t \in [0, \sigma]$, since $(H_1)(ii)$, we have

$$\begin{aligned} (\diamond) &= g_1 \left(\phi_q \left(\int_0^\sigma v(s) ds \right) \right) + \int_\xi^t \phi_q \left(\int_s^\sigma v(\tau) d\tau \right) ds \\ &\geq \xi \phi_q \left(\int_0^\sigma v(s) ds \right) - \int_0^\xi \phi_q \left(\int_s^\sigma v(\tau) d\tau \right) ds + \int_0^t \phi_q \left(\int_s^\sigma v(\tau) d\tau \right) ds \\ &\geq \int_0^\xi \phi_q \left(\int_0^\sigma v(\tau) d\tau \right) ds - \int_0^\xi \phi_q \left(\int_0^\sigma v(\tau) d\tau \right) ds + \int_0^t \phi_q \left(\int_s^\sigma v(\tau) d\tau \right) ds \\ &\geq \int_0^t \phi_q \left(\int_s^\sigma v(\tau) d\tau \right) ds \geq 0. \end{aligned}$$

Further, when $t \in [\sigma, 1]$, $\beta \geq 1 - \eta$, similarly, we can get $(\heartsuit) \geq 0$. Thus $u(t) \geq 0$ for all $t \in [0, 1]$. The proof is complete. \square

3.2 Existence of positive solutions to BVP (1.1), (1.2)

In this subsection, we try to impose some growth conditions on the non-linear term f which allows us to apply Theorem 2.1 to establish the existence of triple positive solutions to BVP (1.1), (1.2).

Let $E = C^+[0, 1] = \{u \in C[0, 1], u(t) \geq 0\}$, and let it be endowed with the maximum norm $\|u\| = \max_{0 \leq t \leq 1} |u(t)|$.

The cone $P \subset E$ is defined as

$$P = \{u \in E : u \text{ is concave on } t \in [0, 1], \text{ there exists one point } \sigma \in (0, 1) \text{ such that } u'(\sigma) = 0\}.$$

In our main results, we will make use of the following lemmas.

Lemma 3.3 For $u \in P$, then $u(t) \geq \min\{t, 1 - t\} \|u\|$.

Proof For $t \in [0, \sigma]$, we have

$$\frac{u(t)}{t} \geq \frac{u(\sigma)}{\sigma} \geq \frac{\max_{0 \leq t \leq 1} |u(t)|}{1} \Rightarrow u(t) \geq t \|u\|.$$

Similarly, $u(t) \geq (1 - t) \|u\|$ when $t \in [\sigma, 1]$. Thus $u(t) \geq \min\{t, 1 - t\} \|u\|$, which completes our proof. \square

Remark 3.2 From Lemma 3.3, we know that when $k \geq 3$, there holds $\min_{1/k \leq t \leq 1 - 1/k} u(t) \geq \frac{1}{k} \|u\|$.

Define $T_1 : P \rightarrow E$ as follows:

$$(T_1 u)(t) = \begin{cases} g_1(\phi_q(\int_0^\sigma h(s)f(s, u(s)) ds) + \int_\xi^t \phi_q(\int_s^\sigma h(\tau)f(\tau, u(\tau)) d\tau) ds), & t \in [0, \sigma], \\ \beta \phi_q(\int_\sigma^1 h(s)f(s, u(s)) ds) + \int_t^\eta \phi_q(\int_\sigma^s h(\tau)f(\tau, u(\tau)) d\tau) ds, & t \in [\sigma, 1]. \end{cases} \quad (3.10)$$

Lemma 3.4 Suppose that (C_1) and (C_2) hold, then $T_1 : P \rightarrow P$ is completely continuous.

Proof To justify this, we first show that $T_1 : P \rightarrow P$ is well defined. Let $u \in P$. Easily, we have $(T_1 u)(t) \geq 0$, $\beta(T_1 u)'(1) + (T_1 u)(\eta) = 0$. Next, we show that T_1 is concave on $t \in [0, 1]$. By (3.10), we have

$$(T_1 x)'(t) = \begin{cases} \phi_q(\int_t^\sigma h(s)f(s, u(s)u) ds), & t \in [0, \sigma], \\ -\phi_q(\int_\sigma^t h(s)f(s, u(s)) ds), & t \in [\sigma, 1]. \end{cases} \quad (3.11)$$

Obviously, $(T_1 x)''(t) \leq 0$, this implies that T is concave on $t \in [0, 1]$. Further, $(T_1 x)'(t) \geq 0$ on $t \in [0, \sigma]$, and $(T_1 x)'(t) \leq 0$ on $t \in [\sigma, 1]$, $(T_1 x)'(\sigma) = 0$. Thus $T_1 : P \rightarrow P$ is well defined.

It is easy to prove that T_1 is continuous with respect to x . For more details, see [19, p.6].

For any bounded subset Ω of P , it can be easily verified that $T_1(\Omega)$ is bounded and equicontinuous. Thus by the Arzela-Ascoli theorem, $T_1(\Omega)$ is relatively compact, therefore, T_1 is completely continuous. \square

Throughout, we suppose that $k \geq 3$ is big enough such that $\xi, \eta \in [\frac{1}{k}, 1 - \frac{1}{k}]$. And the nonnegative continuous concave functional α_1 is defined on the cone P by

$$\alpha_1(x) = \min_{\frac{1}{k} \leq t \leq 1 - \frac{1}{k}} |x(t)|, \quad \text{for } x \in P.$$

Let

$$\begin{aligned}
 M_1 &= \beta \phi_q \left(\int_0^1 h(s) \, ds \right) + \max \left\{ \int_0^\eta \phi_q \left(\int_0^s h(\tau) \, d\tau \right) \, ds, \int_\eta^1 \phi_q \left(\int_s^1 h(\tau) \, d\tau \right) \, ds \right\}, \\
 m_1 &= \frac{1}{k} \cdot \min \left\{ \xi \phi_q \left(\int_{\frac{1}{k}}^{\frac{1}{2}} h(s) \, ds \right) + \int_\xi^{\frac{1}{2}} \phi_q \left(\int_s^{\frac{1}{2}} h(\tau) \, d\tau \right) \, ds, \right. \\
 &\quad \left. \beta \phi_q \left(\int_{\frac{1}{2}}^{1-\frac{1}{k}} h(s) \, ds \right) + \int_{\frac{1}{2}}^\eta \phi_q \left(\int_{\frac{1}{2}}^s h(\tau) \, d\tau \right) \, ds \right\}.
 \end{aligned}$$

Theorem 3.5 *Assume that (H₁) holds, there exist constants such that $0 < a_1 < b_1 \leq \min\{\frac{1}{k}, \frac{m_1}{M_1}\}c_1$. Further suppose that*

- (S₁) $f(t, u) < \phi_p(a_1/M_1)$, for $(t, u) \in [0, 1] \times [0, a_1]$;
- (S₂) $f(t, u) \leq \phi_p(c_1/M_1)$, for $(t, u) \in [0, 1] \times [0, c_1]$;
- (S₃) $f(t, u) \geq \phi_p(b_1/m_1)$, for $(t, u) \in [\frac{1}{k}, 1 - \frac{1}{k}] \times [b_1, kb_1]$.

Then boundary value problem (1.1), (1.2) has at least three positive solutions u_1, u_2, u_3 such that

$$\|u_1\| < a_1, \quad \alpha_1(u_2) > b_1 \quad \text{and} \quad \|u_3\| > a_1, \quad \alpha_1(u_3) < b_1. \tag{3.12}$$

Proof Problem (1.1), (1.2) has a solution $u = u(t)$ if and only if it solves the operator equation $u = T_1 u$. Thus we set out to verify that the operator T_1 satisfies the Leggett-Williams fixed point Theorem 2.1, which will prove the existence of at least three fixed points of T_1 .

Now, we will prove the main theorem step by step.

Step 1. We firstly show that $T_1 : \bar{P}_{a_1} \rightarrow \bar{P}_{a_1}$. In view of (S₁), we have

$$\begin{aligned}
 \|T_1 u\| &= |(T_1 u)(\sigma)| = \beta \phi_q \left(\int_\sigma^1 h(s) f(s, u(s)) \, ds \right) + \int_\sigma^\eta \phi_q \left(\int_\sigma^s h(\tau) f(\tau, u(\tau)) \, d\tau \right) \, ds \\
 &\leq \max \left\{ \beta \phi_q \left(\int_0^1 h(s) f(s, u(s)) \, ds \right) + \int_0^\eta \phi_q \left(\int_0^s h(\tau) f(\tau, u(\tau)) \, d\tau \right) \, ds, \right. \\
 &\quad \left. \beta \phi_q \left(\int_0^1 h(s) f(s, u(s)) \, ds \right) + \int_\eta^1 \phi_q \left(\int_s^1 h(\tau) f(\tau, u(\tau)) \, d\tau \right) \, ds \right\} \\
 &\leq \frac{a_1}{M_1} \cdot M_1 = a_1.
 \end{aligned}$$

Similarly, (S₂) implies that $T_1 : \bar{P}_{c_1} \rightarrow \bar{P}_{c_1}$.

Step 2. We try to prove Theorem 2.1(i) is satisfied.

Choose $u(t) = kb_1$. Obviously, $u(t) \in P(\alpha_1, b_1, kb_1)$, thus $\{u(t) \in P(\alpha_1, b_1, kb_1)\} \neq \emptyset$.

If $u \in P(\alpha_1, b_1, kb_1)$, then for $t \in [\frac{1}{k}, 1 - \frac{1}{k}]$, $b_1 < u(t) \leq kb_1$. In view of (S₃), we have

$$\begin{aligned}
 \alpha_1(Tu) &= \min_{\frac{1}{k} \leq t \leq 1-\frac{1}{k}} |(Tu)(t)| \geq \min \left\{ \frac{1}{k}, 1 - \frac{1}{k} \right\} \max_{0 \leq t \leq 1} |(Tu)(t)| = \frac{1}{k} |(Tu)(\sigma)| \\
 &= \frac{1}{k} \left(g_1 \left(\phi_q \left(\int_0^\sigma h(s) f(s, u(s)) \, ds \right) \right) + \int_\xi^\sigma \phi_q \left(\int_s^\sigma h(\tau) f(\tau, u(\tau)) \, d\tau \right) \, ds \right) \\
 &\geq \frac{1}{k} \min \left\{ \xi \phi_q \left(\int_0^{\frac{1}{2}} h(s) f(s, u(s)) \, ds \right) + \int_\xi^{\frac{1}{2}} \phi_q \left(\int_s^{\frac{1}{2}} h(\tau) f(\tau, u(\tau)) \, d\tau \right) \, ds, \right.
 \end{aligned}$$

$$\begin{aligned} & \left. \beta \left(\phi_q \left(\int_{\frac{1}{2}}^1 h(s)f(s, u(s)) \, ds \right) \right) + \int_{\frac{1}{2}}^\eta \phi_q \left(\int_{\frac{1}{2}}^s h(\tau)f(\tau, u(\tau)) \, d\tau \right) \, ds \right\} \\ & \geq \frac{1}{k} \min \left\{ \xi \phi_q \left(\int_{\frac{1}{k}}^{\frac{1}{2}} h(s)f(s, u(s)) \, ds \right) + \int_{\xi}^{\frac{1}{2}} \phi_q \left(\int_s^{\frac{1}{2}} h(\tau)f(\tau, u(\tau)) \, d\tau \right) \, ds, \right. \\ & \left. \beta \left(\phi_q \left(\int_{\frac{1}{2}}^{1-\frac{1}{k}} h(s)f(s, u(s)) \, ds \right) \right) + \int_{\frac{1}{2}}^\eta \phi_q \left(\int_{\frac{1}{2}}^s h(\tau)f(\tau, u(\tau)) \, d\tau \right) \, ds \right\} \\ & \geq \frac{b_1}{m_1} \cdot m_1 = b_1. \end{aligned}$$

Thus, Theorem 2.1(i) holds.

Step 3. We try to prove that Theorem 2.1(iii) holds.

If $u \in P(\alpha_1, b_1, c_1)$ and $\|Tu\| > kb_1$, then we have

$$\alpha_1(Tu) = \min_{\frac{1}{k} \leq t \leq 1-\frac{1}{k}} (Tu)(t) \geq \frac{1}{k} \|Tu\| > b_1.$$

Hence, Theorem 2.1(iii) also holds. An application of Theorem 2.1 means that BVP (1.1), (1.2) has at least three positive solutions satisfied (3.12). The proof is complete. \square

4 Existence results for BVP(1.1), (1.3)

4.1 Some useful lemmas

Lemma 4.1 *Suppose that $v(t)$ satisfies all the conditions in Lemma 3.1. Then BVP*

$$\begin{cases} (\phi_p(u'(t)))' + v(t) = 0, & 0 < t < 1, \\ g_1(u'(0)) - u(\xi) = 0, & \beta u'(1) + u(\eta) = 0. \end{cases} \quad (4.1)$$

has the unique solution

$$u(t) = \begin{cases} \alpha \phi_q \left(\int_0^\sigma v(s) \, ds \right) + \int_\xi^t \phi_q \left(\int_s^\sigma v(\tau) \, d\tau \right) \, ds, & t \in [0, \sigma], \\ g_2 \left(\phi_q \left(\int_\sigma^1 v(s) \, ds \right) \right) + \int_t^\eta \phi_q \left(\int_\sigma^s v(\tau) \, d\tau \right) \, ds, & t \in [\sigma, 1], \end{cases} \quad (4.2)$$

where σ is a solution of the equation

$$B_1(t) - B_2(t) = 0, \quad t \in [0, 1]. \quad (4.3)$$

And $B_1(t), B_2(t)$ are defined as follows:

$$\begin{aligned} B_1(t) &= \alpha \phi_q \left(\int_0^t v(s) \, ds \right) + \int_\xi^t \phi_q \left(\int_s^t v(\tau) \, d\tau \right) \, ds \\ B_2(t) &= g_2 \left(\phi_q \left(\int_t^1 v(s) \, ds \right) \right) + \int_t^\eta \phi_q \left(\int_t^s v(\tau) \, d\tau \right) \, ds. \end{aligned}$$

Proof Essentially, we can prove it using the same discussion as in Lemma 3.1. Here, we omit it. \square

Lemma 4.2 *Let $v(t)$ satisfy all the conditions in Lemma 3.1, then the solution $u(t)$ of BVP (4.1) is concave on $t \in [0, 1]$. What is more, $u(t) \geq 0$.*

4.2 Existence of positive solutions to BVP (1.1), (1.3)

In this subsection, we try to impose some growth conditions on the non-linear term f which allow us to apply Theorem 2.1 to establish the existence of triple positive solutions to problem (1.1).

Let E , cone P and the norm be defined as in Section 3.

Let $k \geq 3$ be big enough so that $\xi, \eta \in [\frac{1}{k}, 1 - \frac{1}{k}]$; and the nonnegative continuous concave functional α_2 be defined on the cone P by

$$\alpha_2(x) = \min_{\frac{1}{k} \leq t \leq 1 - \frac{1}{k}} |x(t)|, \quad \text{for } x \in P.$$

Define $T_2 : P \rightarrow E$ as follows:

$$(T_2u)(t) = \begin{cases} \alpha \phi_q(\int_0^\sigma h(s)f(s, u(s)) ds) \\ \quad + \int_\xi^t \phi_q(\int_s^\sigma h(\tau)f(\tau, u(\tau)) d\tau) ds, & t \in [0, \sigma], \\ g_2(\phi_q(\int_\sigma^1 h(s)f(s, u(s)) ds) \\ \quad + \int_t^\eta \phi_q(\int_\sigma^s h(\tau)f(\tau, u(\tau)) d\tau) ds, & t \in [\sigma, 1]. \end{cases} \tag{4.4}$$

Lemma 4.3 *Suppose that (C_1) and (C_2) hold, then $T_2 : P \rightarrow P$ is completely continuous.*

Let

$$\begin{aligned} M_2 &= \alpha \phi_q\left(\int_0^1 h(s) ds\right) + \max\left\{\int_0^\xi \phi_q\left(\int_0^s h(\tau) d\tau\right) ds, \int_\xi^1 \phi_q\left(\int_s^1 h(\tau) d\tau\right) ds\right\} \\ m_2 &= \frac{1}{k} \cdot \min\left\{\alpha \phi_q\left(\int_{\frac{1}{k}}^{\frac{1}{2}} h(s) ds\right) + \int_\xi^{\frac{1}{2}} \phi_q\left(\int_s^{\frac{1}{2}} h(\tau) d\tau\right) ds, \right. \\ &\quad \left. (1 - \eta)\phi_q\left(\int_{\frac{1}{2}}^{1 - \frac{1}{k}} h(s) ds\right) + \int_{\frac{1}{2}}^\eta \phi_q\left(\int_{\frac{1}{2}}^s h(\tau) d\tau\right) ds\right\}. \end{aligned}$$

Theorem 4.4 *Assume that (H_2) holds, there exist constants such that $0 < a_2 < b_2 \leq \min\{\frac{1}{k}, \frac{m_2}{M_2}\}c_2$. Further suppose that*

- (S_4) $f(t, u) < \phi_p(a_2/M_2)$, for $(t, u) \in [0, 1] \times [0, a_2]$;
- (S_5) $f(t, u) \leq \phi_p(c_2/M_2)$, for $(t, u) \in [0, 1] \times [0, c_2]$;
- (S_6) $f(t, u) \geq \phi_p(b_2/m_2)$, for $(t, u) \in [\frac{1}{k}, 1 - \frac{1}{k}] \times [b_2, kb_2]$.

Then boundary value problem (1.1) (1.3) has at least three positive solutions u_1, u_2, u_3 such that

$$\|u_1\| < a_2, \quad \alpha_2(u_2) > b_2 \quad \text{and} \quad \|u_3\| > a_2, \quad \alpha_2(u_3) < b_2. \tag{4.5}$$

Proof By making use of Theorem 2.1, we can prove the result by essentially the same method. Here we omit it. □

5 A generalization result

In this section, we consider the existence results of (1.1) subject to the following non-linear four-point boundary condition:

$$g_1(u'(0)) - u(\xi) = 0, \quad g_2(u'(1)) + u(\eta) = 0. \tag{5.1}$$

In addition to (H_1) , (H_2) , we assume at least one of the following conditions holds:

(H_3) There exists a positive constant $B_1 \geq \xi$ such that $g_1(x) \leq B_1x$;

(H_4) There exists a positive constant $B_2 \geq 1 - \eta$ such that $g_1(x) \leq B_2x$.

Without loss of generality, we suppose (H_4) holds. Essentially by the same method, we can get the multiple existence results for BVP (1.1), (5.1). In what follows, we just list the main results but the proofs are all omitted.

Lemma 5.1 *Suppose that $v(t)$ satisfies all the conditions in Lemma 3.1, then BVP*

$$\begin{cases} (\phi_p(u'(t)))' + v(t) = 0, & 0 < t < 1, \\ g_1(u'(0)) - u(\xi) = 0, & g_2(u'(1)) + u(\eta) = 0 \end{cases} \quad (5.2)$$

has the unique solution

$$u(t) = \begin{cases} g_1(\phi_q(\int_0^\sigma v(s) ds)) + \int_\xi^t \phi_q(\int_s^\sigma v(\tau) d\tau) ds, & (i) \quad t \in [0, \sigma], \\ g_2(\phi_q(\int_\sigma^1 v(s) ds)) + \int_t^\eta \phi_q(\int_\sigma^s v(\tau) d\tau) ds, & (ii) \quad t \in [\sigma, 1], \end{cases} \quad (5.3)$$

where σ is a solution of the equation

$$C_1(t) - C_2(t) = 0, \quad t \in [0, 1], \quad (5.4)$$

and $C_1(t)$, $C_2(t)$ are defined as follows:

$$\begin{aligned} C_1(t) &= g_1\left(\phi_q\left(\int_0^t v(s) ds\right)\right) + \int_\xi^t \phi_q\left(\int_s^t v(\tau) d\tau\right) ds, \\ C_2(t) &= g_2\left(\phi_q\left(\int_t^1 v(s) ds\right)\right) + \int_t^\eta \phi_q\left(\int_t^s v(\tau) d\tau\right) ds. \end{aligned}$$

Lemma 5.2 *Let $v(t)$ satisfy all the conditions in Lemma 3.1, then the solution $u(t)$ of BVP (5.2) is concave on $t \in [0, 1]$. What is more, $u(t) \geq 0$.*

Let E , cone P and the norm be defined as in Section 3; and let the nonnegative continuous concave functional α_3 be defined on the cone P by

$$\alpha_3(x) = \min_{\frac{1}{k} \leq t \leq 1 - \frac{1}{k}} |x(t)|, \quad \text{for } x \in P.$$

Define $T_3 : P \rightarrow E$ as follows:

$$(T_3u)(t) = \begin{cases} g_1(\phi_q(\int_0^\sigma h(s)f(s, u(s)) ds)) \\ \quad + \int_\xi^t \phi_q(\int_s^\sigma h(\tau)f(\tau, u(\tau)) d\tau) ds, & t \in [0, \sigma], \\ g_2(\phi_q(\int_\sigma^1 h(s)f(s, u(s)) ds)) \\ \quad + \int_t^\eta \phi_q(\int_\sigma^s h(\tau)f(\tau, u(\tau)) d\tau) ds, & t \in [\sigma, 1]. \end{cases} \quad (5.5)$$

Lemma 5.3 *Suppose that (C_1) and (C_2) hold, then $T_3 : P \rightarrow P$ is completely continuous.*

Let

$$M_3 = B_1 \phi_q \left(\int_0^1 h(s) \, ds \right) + \max \left\{ \int_0^\xi \phi_q \left(\int_0^s h(\tau) \, d\tau \right) \, ds, \int_\xi^1 \phi_q \left(\int_s^1 h(\tau) \, d\tau \right) \, ds \right\}$$

$$m_3 = \frac{1}{k} \cdot \min \left\{ \xi \phi_q \left(\int_{\frac{1}{k}}^{\frac{1}{2}} h(s) \, ds \right) + \int_\xi^{\frac{1}{2}} \phi_q \left(\int_s^{\frac{1}{2}} h(\tau) \, d\tau \right) \, ds, \right.$$

$$\left. (1 - \eta) \phi_q \left(\int_{\frac{1}{2}}^{1 - \frac{1}{k}} h(s) \, ds \right) + \int_{\frac{1}{2}}^\eta \phi_q \left(\int_{\frac{1}{2}}^s h(\tau) \, d\tau \right) \, ds \right\}.$$

Theorem 5.4 Assume that (H_1) , (H_4) hold, there exist constants such that $0 < a_3 < b_3 \leq \min\{\frac{1}{k}, \frac{m_3}{M_3}\}c_3$. Further suppose that

- (S_7) $f(t, u) < \phi_p(a_3/M_3)$, for $(t, u) \in [0, 1] \times [0, a_3]$;
- (S_8) $f(t, u) \leq \phi_p(c_3/M_3)$, for $(t, u) \in [0, 1] \times [0, c_3]$;
- (S_9) $f(t, u) \geq \phi_p(b_3/m_3)$, for $(t, u) \in [\frac{1}{k}, 1 - \frac{1}{k}] \times [b_3, kb_3]$.

Then boundary value problem (1.1), (1.4) has at least three positive solutions u_1, u_2, u_3 such that

$$\|u_1\| < a_3, \quad \alpha_3(u_2) > b_3 \quad \text{and} \quad \|u_3\| > a_3, \quad \alpha_3(u_3) < b_3. \tag{5.6}$$

6 Example

In this section, we give examples to illustrate our main results.

Example 6.1 Consider the following non-linear four-point boundary value problem:

$$\begin{cases} (\frac{u'}{\sqrt{u'}})' + \frac{1}{t^{1/2}} f(t, u(t)) = 0, & 0 < t < 1, \\ u'^2(0) + u'(0) - u(1/2) = 0, & 2u'(1) + u(2/3) = 0, \end{cases} \tag{6.1}$$

where

$$f(t, u(t)) = \begin{cases} \sin t + (\frac{u}{15})^8, & 0 < u \leq 24.9, \\ \sin t + (\frac{24.9}{15})^8, & u > 24.9. \end{cases}$$

And we can see that $p = 3/2, g_1(x) = x^2 + x, \xi = 1/2, \eta = 2/3$. Choose $k = 5, a_1 = 80/9, b_1 = 23, c_1 = 32,000$. By direct calculation, we can get $M_1 = 80/9, m_1 = 0.027$. Also the following conditions hold.

- (i) $f(t, u) \leq 0.86 < 1 = \phi_p(a_1/M_1)$, for $(t, u) \in [0, 1] \times [0, 80/9]$;
- (ii) $f(t, u) \leq 58.51 < 60 = \phi_p(c_1/M_1)$, for $(t, u) \in [0, 1] \times [0, 32,000]$;
- (iii) $f(t, u, v) \geq 30.5 > 29.2 = \phi_p(b_1/m_1)$, for $(t, u) \in [\frac{1}{k}, 1 - \frac{1}{k}] \times [23, 115]$.

Thus, all the conditions in Theorem 3.5 are satisfied, so BVP (6.1) has at least three positive solutions u_1, u_2, u_3 satisfying

$$\|u_1\| < 80/9, \quad \alpha(u_2) > 23 \quad \text{and} \quad \|u_3\| > 80/9, \quad \alpha(u_3) < 23. \tag{6.2}$$

Example 6.2 Consider the following non-linear four-point boundary value problem

$$\begin{cases} u'' + \frac{1}{(1-t)^{1/2}}f(t, u(t)) = 0, & 0 < t < 1, \\ 3u'(0) - u(1/4) = 0, & g_2(u'(1)) + u(1/2) = 0, \end{cases} \quad (6.3)$$

where

$$f(t, u) = \begin{cases} -(t - \frac{1}{2})^2 + \frac{1}{4} + (\frac{u}{10})^3, & 0 \leq u \leq 3,400, \\ -(t - \frac{1}{2})^2 + \frac{1}{4} + (\frac{3400}{10})^3, & u \geq 3,400 \end{cases}$$

and

$$g_2(x) = x^3 + 1.$$

Choose $k = 4$, $a_2 = 6.866$, $b_2 = 150$, $c_2 = 2.75 \times 10^8$. Obviously, $p = 2$, $\alpha = 3$, $\xi = 1/4$, $\eta = 1/2$. By direct calculation, we can get that $q = 2$, $M_2 = 6.866$, $m_2 = 0.052$. Also the following conditions hold.

- (i) $f(t, u) \leq 0.5737 < 1 = \phi_p(a_2/M_2)$, for $(t, u) \in [0, 1] \times [0, 6.866]$;
- (ii) $f(t, u) \leq 3.9305 \times 10^7 < 4.005 \times 10^7 = \phi_p(c_2/M_2)$, for $(t, u) \in [0, 1] \times [0, 2.75 \times 10^8]$;
- (iii) $f(t, u, v) \geq 3,375 > 2,884.6 = \phi_p(b_2/m_2)$, for $(t, u) \in [\frac{1}{4}, \frac{3}{4}] \times [150, 600]$.

Thus, all the conditions in Theorem 4.4 are satisfied, so BVP (6.1) has at least three positive solutions u_1, u_2, u_3 satisfying

$$\|u_1\| < 6.866, \quad \alpha(u_2) > 150 \quad \text{and} \quad \|u_3\| > 6.866, \quad \alpha(u_3) < 150. \quad (6.4)$$

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

JZ and WG conceived of the study, and participated in its coordination. JZ drafted the manuscript. All authors read and approved the final manuscript.

Author details

¹School of Science, China University of Geosciences, Beijing, 100083, P.R. China. ²Department of Mathematics, Beijing Institute of Technology, Beijing, 100081, P.R. China.

Acknowledgements

The authors were very grateful to the anonymous referee whose careful reading of the manuscript and valuable comments enhanced presentation of the manuscript. The study was supported by Pre-research project and Excellent Teachers project of the Fundamental Research Funds for the Central Universities 2-9-2012-141.

Received: 27 January 2012 Accepted: 21 August 2012 Published: 5 September 2012

References

1. Agarwal, RP, O'Regan, D, Wong, PJY: Positive Solutions of Differential, Difference, and Integral Equations. Kluwer Academic, Boston (1999)
2. Avery, R: Multiple positive solutions of an nth order focal boundary value problem. *Panam. Math. J.* **8**, 39-55 (1998)
3. Lv, HS, O'Regan, D, Zhong, CK: Multiple positive solutions for the one-dimensional singular p -Laplacian. *Appl. Math. Comput.* **133**, 407-422 (2002)
4. Webb, JRL: Positive solutions of some three point boundary value problems via fixed point index theory. *Nonlinear Anal.* **47**, 4319-4332 (2001)
5. Wong, PJY, Agarwal, RP: Multiple positive solutions of two-point right focal boundary value problems. *Math. Comput. Model.* **28**, 41-49 (1998)
6. Kong, LB, Wang, JY: Multiple positive solutions for the one-dimensional p -Laplacian. *Nonlinear Anal.* **42**, 1327-1333 (2000)
7. Agarwal, RP, Hong, H, Yeh, C: The existence of positive solutions for the Sturm-Liouville boundary value problems. *Comput. Math. Appl.* **35**, 89-96 (1998)

8. Agarwal, RP, Bohner, M, Wong, PJY: Sturm-Liouville eigenvalue problems on time scales. *Appl. Math. Comput.* **99**, 153-166 (1999)
9. Bai, Z, Gui, Z, Ge, W: Multiple positive solutions for some p -Laplacian boundary value problems. *J. Math. Anal. Appl.* **300**, 477-490 (2004)
10. Davis, JM, Erbe, LH, Henderson, J: Multiplicity of positive solutions for higher order Sturm-Liouville problems. *Rocky Mt. J. Math.* **31**, 169-184 (2001)
11. Erbe, LH, Hu, SC, Wang, HY: Multiple positive solutions of some boundary value problems. *J. Math. Anal. Appl.* **184**, 640-648 (1994)
12. Ge, W, Ren, J: New existence theorem of positive solutions for Sturm-Liouville boundary value problems. *Appl. Math. Comput.* **148**, 631-644 (2004)
13. Wong, PJY: Solutions of constant signs of a system of Sturm-Liouville boundary value problems. *Math. Comput. Model.* **29**, 27-38 (1999)
14. Wong, PJY: Three fixed-sign solutions of system model with Sturm-Liouville type conditions. *J. Math. Anal. Appl.* **298**, 120-145 (2004)
15. Wong, PJY: Multiple fixed-sign solutions for a system of difference equations with Sturm-Liouville conditions. *J. Comput. Appl. Math.* **183**, 108-132 (2005)
16. Il'in, VA, Moiseev, EI: Nonlocal boundary value problem of the first kind for a Sturm-Liouville operator in its differential and finite difference aspects. *Differ. Equ.* **23**, 803-810 (1987)
17. Il'in, VA, Moiseev, EI: Nonlocal boundary value problem of the second kind for a Sturm-Liouville operator. *Differ. Equ.* **23**, 979-987 (1987)
18. Leggett, RW, Williams, LR: Multiple positive fixed points of nonlinear operators on ordered Banach spaces. *Indiana Univ. Math. J.* **28**, 673-688 (1979)
19. Lian, H, Ge, W: Positive solutions for a four-point boundary value problem with the p -Laplacian. *Nonlinear Anal.* **68**, 3493-3503 (2008)

doi:10.1186/1029-242X-2012-197

Cite this article as: Zhao and Ge: Existence results of a kind of Sturm-Liouville type singular boundary value problem with non-linear boundary conditions. *Journal of Inequalities and Applications* 2012 **2012**:197.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
