Falahi et al. Epigenetics & Chromatin 2013, 6(Suppl 1):P15 http://www.epigeneticsandchromatin.com/content/6/S1/P15

### **POSTER PRESENTATION**





# Gene specific overwriting of epigenetic signatures to modulate the expression of selected tumor-promoting genes in cancer

Fahimeh Falahi<sup>1\*</sup>, Christian Huisman<sup>1</sup>, Elisa Garcia Diaz<sup>1</sup>, Hinke G Kazemier<sup>1</sup>, Geke AP Hospers<sup>2</sup>, Marianne G Rots<sup>1</sup>

*From* Epigenetics and Chromatin: Interactions and processes Boston, MA, USA. 11-13 March 2013

### Background

Epidermal Growth Factor Receptor-2 (HER-2) and Estrogen receptor-alpha (ER- $\alpha$ ) have been found to be dysregulated in several types of cancer. Their dysregulation is associated with aberrant epigenetic modifications. Additionally, expression of ER- $\alpha$  and HER-2 is inversely correlated; their crosstalk has been shown to be involved in endocrine therapy resistance mechanisms [1]. Therapies targeting either receptor result in cell proliferation inhibition for some cancer types; the anti-cancer effect might be further optimized to be more efficient and applicable for more cancer types by co-targeting the genes at the DNA level. As epigenetic modifications provide a way to modify expression of genes in a sustained manner we aim to downregulate HER-2 and ER- $\alpha$  by inducing epigenetic silencing marks specifically onto their promoters (Epigenetic Editing [2]). We also aim to discover the possible crosstalk of ER-a with HER-2 and other important genes in cancer.

#### Methods and materials

Towards downregulation of HER-2 and ER- $\alpha$ , expression and epigenetic modification status of promoter of these genes were assessed in a panel of cancer cell lines (bisulfite sequencing and ChIP). Designed zinc finger proteins (ZFPs) targeting genes promoters were fused to a transcription repressor domain [SKD, histone methyltransferases (G9a, SUV39H1), or a DNA methyltransferase domain (C141S [3])] and expressed in cancer cells through viral transduction.

<sup>1</sup>Epigenetic Editing, Dept. Medical Biology and Pathology, 9713 GZ Groningen, The Netherlands

Full list of author information is available at the end of the article

## BioMed Central

### Results

HER2-ZFP fused to G9a or SUV39H1 induced the intended silencing histone methylation marks (H3K9me2, H3K9me3) on the HER-2 promoter. Up to 4-fold induced H3K9me2 mark was associated with up to  $54 \pm 2.9\%$  downregulation of HER-2 expression (P<0.0001). Of note, downregulation of HER-2 by induced H3K9 methylation mark was associated with inhibition of cell proliferation and clonogenicity. Additionally, up to 50% downregulation of ER- $\alpha$  was obtained by a ZFP specif c to ER- $\alpha$  fused to SKD, G9a, and C141S.

#### Conclusions

These results show that targeted induction of epigenetic marks is instructive in downregulation of HER-2 and ER- $\alpha$  expression. We conclude that Epigenetic Editing might provide a novel and synergistic approach to modulate (onco)genes. In addition, investigation of ER- $\alpha$  crosstalk with HER-2 and other genes might be promising to better interfere with pathways involved in cancer.

#### Author details

<sup>1</sup>Epigenetic Editing, Dept. Medical Biology and Pathology, 9713 GZ Groningen, The Netherlands. <sup>2</sup>Dept. Medical Oncology, 9713 GZ Groningen, The Netherlands.

Published: 18 March 2013

#### References

- Arpino G, Wiechmann L, Osborne CK, Schif R: Crosstalk between the estrogen receptor and the HER tyrosine kinase receptor family: molecular mechanism and clinical implications for endocrine therapy resistance. *Endocr Rev* 2008, 29(2):217-233.
- de Groote ML, Verschure PJ, Rots MG: Epigenetic editing: Targeted rewriting of epigenetic marks to Modulate Expression of selected target genes. Nucleic Acids Res 2012, 40(21):10596-10613.
- van der Gun BT, Maluszynska-Hofman M, Kiss A, Arendzen AJ, Ruiters MH, McLaughlin PM, Weinhold E, Rots MG: Targeted DNA methylation by a

© 2013 Falahi et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

DNA methyltransferase coupled to a triple helix forming oligonucleotide to down-regulate the epithelial cell adhesion molecule. *Bioconjug Chem* 2010, **21(7)**:1239-1245.

doi:10.1186/1756-8935-6-S1-P15

**Cite this article as:** Falahi *et al*:: Gene specific overwriting of epigenetic signatures to modulate the expression of selected tumor-promoting genes in cancer. *Epigenetics & Chromatin* 2013 6(Suppl 1):P15.

## Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar

**BioMed** Central

• Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit