

RESEARCH ARTICLE

Open Access



Evaluating emergency preparedness and impact of a hurricane sandy in pediatric patients with diabetes

Rubina Heptulla^{1*}, Rebecca Hashim², Doreen Newell Johnson³, Jeniece Trast Ilkowitz³, Gina DiNapoli³, Venkat Renukuntla⁴ and Jennifer Sivitz⁵

Abstract

Background: Natural disasters have always been associated with significant adverse events including medical and mental health problems. Children with chronic disease such as diabetes have also been believed to be affected to a greater extent by any natural disaster. The purpose of this study was to assess and compare emergency preparedness post-disaster and post-traumatic stress effects of Hurricane Sandy in affected and relatively unaffected populations.

Methods: The study was conducted between February and July 2013. A total of 142 families caring for children with Type 1 Diabetes Mellitus (T1DM) who attended clinics were recruited from hospitals in Bronx, NY (control) and in NJ (affected) by Hurricane Sandy. Subjects were recruited to participate in a survey 3–6 months after the hurricane. Data on demographics, glycemic control and insulin regimens were collected. Families were surveyed for socio-economic status (SES), using Hollingshead questionnaire, general and diabetes preparedness and the Hurricane Related Traumatic Experiences (HURTE) questionnaire was used to evaluate for symptoms of post-traumatic stress.

Results: Ninety-five percent of families reported to be generally well to moderately prepared for the hurricane and 83 % reported to be very well prepared with regards to their child's diabetes during the disaster. There was no difference between the sites for preparedness for the disaster, age or gender. There was a trend toward significance ($p < 0.06$) in New Jersey subjects as to a greater psychological impact from the hurricane. Poor glycemic control was significantly associated with lower SES ($p < 0.008$). Most importantly, SES was unrelated to preparedness for diabetes management during the hurricane.

Conclusions: Despite low SES, families were generally well to moderately prepared for hurricane. In children with diabetes, interventional studies should be designed and implemented so that glycemic control remains unaffected, following any major disaster.

Keywords: Disaster, Preparedness, Hurricane, Diabetes, Pediatrics

Background

The International Federation of Red Cross and Red Crescent Societies categorize a disaster as an event that causes more than ten deaths, affects more than 100 people or leads to an appeal by those affected for assistance [1]. Disasters can further be classified as natural or

manmade. Naturally occurring disasters include hurricanes, tornadoes, floods and earthquakes, whereas fires, mass transportation incidents, events involving environmental toxins and episodes of civil unrest are considered manmade disasters [2]. The Federal Emergency Management Agency (FEMA) recommends that families should be self-sufficient for greater than or equal to 3 days after a disaster occurs [3]. Disaster preparedness reduces fear, anxiety and losses that may occur as a result of disasters. The importance of disaster preparedness is highlighted by recent weather events, such as Hurricane Katrina and

*Correspondence: rheptull@montefiore.org

¹ Pediatric Endocrinology and Diabetes, Children's Hospital at Montefiore and Albert Einstein College of Medicine, 3411 Wayne Ave, Suite: 4 M, Bronx, NY 10467, USA

Full list of author information is available at the end of the article



the terrorist attacks of September 11, 2001. According to FEMA, the importance of disaster preparedness is only becoming more relevant as the frequency of natural disasters has increased over the past 50 years. On average, over the last 10 years there have been 65.1 declared disasters per year (2004–2013) compared to 14.7 declared disasters per year between 1954–1963 [3].

Management of chronic diseases, such as diabetes, is challenging during and after disasters. Furthermore, the psychological impact of a disaster may have negative consequences for patients as seen after Hurricane Katrina [4]. Diabetes affects approximately 3.4 million children in the United States [5]. Recent data suggests approximately one out of 500 children in the US have type 1 diabetes (T1DM) by 18 years of age [5]. Given the vulnerability of children to environmental factors (toxins/extreme temperatures) and the multiple resources needed to care for someone with T1DM (i.e. access to adequate food and water, insulin, testing supplies such as meters, glucose test strips, and lancets, etc.), disaster preparedness becomes absolutely essential. The Board of Directors of the American Academy of Pediatrics further emphasized this need when they identified disaster preparedness as one of seven priority issues requiring special attention and resources [2].

In recent years, there have been a number of disastrous events that have tested our readiness for such situations. Currently, we are reeling from the aftermath of Hurricane Sandy, the largest Atlantic Hurricane on record, which made landfall on October 29, 2012. The hurricane, also known as Super Storm Sandy, caused widespread destruction in the areas of New York City, Long Island, Westchester County of New York State, New Jersey, and Connecticut. Services for health care delivery, hospitals and pharmacies were significantly impacted during this time [6, 7]. The influence of disaster has effects that are more chronic and can have a negative impact on children. Furthermore, inadequate preparation prior to the storm can affect both the immediate and long-term consequences of diseases such as diabetes [8, 9]. Our previous research suggested that patients with T1DM who were surveyed before the disaster were inadequately prepared for a disaster [10]. In this trial, we examined the after-effects of a disaster in families that were directly affected and those that were unaffected and lived in the same geographic area hit by Hurricane Sandy.

Methods

The study was approved by the Institutional Review Board (IRB) at both participating sites. The protocol was expedited and the participants were consented before administering the surveys. After Hurricane Sandy in New York and New Jersey, a survey was designed to determine

the level of disaster preparedness and was distributed to families caring for children with T1DM who were attending the diabetes clinic at an urban hospital in Bronx, New York, which was minimally affected by Hurricane Sandy; and the diabetes clinic at a suburban hospital in Hackensack, New Jersey, which was severely affected by the hurricane. Severely affected areas were characterized by loss of services such as electricity, water and phone as well as destruction of homes, businesses and schools for many days, weeks or months. This includes damage caused by flooding, fires, and high winds. Severely affected areas required people to leave their homes before or after the storm to remain safe [11, 12]. Minimally affected areas had less destruction, for example trees falling, and had loss of services (such as electricity and transportation) for only a few hours or days [13]. Due to the media coverage, people were very aware of what areas were more and less affected. The survey was administered during routine clinic visits to English-speaking families. Adults 18 years of age and older did not need to be accompanied by a legal guardian to answer the questions in the survey and those under 18 filled out the survey with their legal guardian. The sample (subjects) was a convenience sample of families attending one of the two clinics described. Each family was approached individually in the clinic and was asked to complete the survey using paper and a pencil. On average, families took approximately 15–20 min to complete the survey. We collected a total of 142 surveys: 77 from hospital in the Bronx and 65 from the hospital in New Jersey. We approached 150 families and 142 agreed to participate. The eight families who choose not to participate did not specify the reason.

The survey included four parts: (1) preparation for a disaster in general, (2) preparation for caring for the child with diabetes in the event of a disaster, (3) demographic characteristics and (4) a measure to assess hurricane-related traumatic experiences. The questions in the first and second parts were graded on a scale of one to five points as described in our previous work [10], weighted according to their degree of importance in disaster preparedness. The general and diabetes preparedness raw scores were divided into tertiles, to indicate well, moderate and poor preparedness. The first set of questions on the survey elicited an estimation of level of general preparedness. A disaster supply kit, or a 72-hour emergency kit, consisted of 3-days supply of food, water and current prescriptions, and a first aid kit which includes non-prescription treatments for common illnesses and tools. The disaster supply kit and its different elements accounted for 44 of the 70 possible points awarded for the general preparedness portion of the survey. Further points were awarded for additional components of a family disaster plan such as identified emergency contact numbers,

household smoke alarms and fire extinguishers, in addition to communication and evacuation plans. A score of 0–23 reflected poor preparedness, 24–47 reflected moderate preparedness and 48–70 reflected well preparedness for an emergency or disastrous situation.

For the next part of the survey, questions were designed to evaluate diabetes preparedness in an emergency or disaster as previously described [10]. The survey addressed items and skills necessary to care for a child with diabetes including: wearing a medical alert bracelet, completion of diabetes education classes, maintenance of current blood glucose logs, possession of a glucagon emergency kit, at least 3 days supply of current prescriptions (insulin, insulin syringes/insulin pens and pen needles, glucose meter, glucose test strips, lancets and lancing device and insulin pump supplies if using an insulin pump) and up to date immunizations. One could earn up to 40 points on this portion of the survey. A total of zero to 13 points was considered poorly prepared, 14–27 indicated moderate preparation, and 28–40 points was considered well prepared to manage a child with diabetes in the setting of an emergency or disaster.

Demographic characteristics were elicited in the third section of the questionnaire. Data was collected on gender, race and the child's age. Hemoglobin A1C (HbA1C) level at the visit as well as the child's current insulin regimen was also obtained. HbA1C was further defined as well controlled if HbA1C was <7.5 %, moderately controlled if 7.6–8.9 % and poorly controlled if >9 %. The Hollingshead 4-Factor Index of Social Status was used to determine the estimated socioeconomic status (SES) of the families [14]. A calculation based on education and occupation levels was used to determine SES status. Education and occupation levels were self-reported by the families interviewed. A total score range of 8–66 could be earned in this section. A score of 48–66 was considered high SES, 28–47 was moderate and 8–27 was considered low SES.

Finally, Hurricane Related Traumatic Experiences (HURTE) was used to measure hurricane-related traumatic experiences as previously described [15]. In brief, HURTE elicited the children's reports of experiencing life threatening events and perceived loss and disruption [16]. The items on the HURTE were developed to identify potentially confirmed events and answered with a "yes" or "no." There was one question that addressed whether or not the child felt they were in a life threatening position during the event. The next six items inquired about witnessed events that were reflective of life threatening events and the sum of these questions was used to determine a score of *n* for life threatening events during a hurricane. Finally the last ten items regarded loss and disruption. These items were also totaled to obtain a score of

n for loss and disruption. Each "yes" response counted as one point and each "no" response counted as zero points. (See Appendix 1).

Statistics: The advanced model of SPSS 16.0 was used to perform statistical analyses (SPSS, Chicago, IL). Figures and graphs were obtained using both SPSS 16.0 and Graphpad Prism 6.0 (GraphPad, La Jolla, CA). Unless otherwise indicated, all data is expressed as means and SDs or proportions. The analyses included: cross-tabulation independent-sample *t* tests, Mann–Whitney test and analysis of variance and Pearson correlation. Significance was considered at $p < 0.05$ level.

Results

One hundred and forty-two subjects were surveyed from two centers (N = 77 from the Bronx, New York and N = 65 from Hackensack, New Jersey). The patient ages were well matched at the two sites (Table 1), 74 were females and 68 were males, mean HbA1c was 8.8 ± 1.9 % (24 % were well controlled, 40 % were moderately controlled and 36 % were poorly controlled). Subjects were primarily managed on insulin pumps (56 %) or multiple daily insulin injections with long acting and rapid acting analog at meals (40 %). The remaining was on two injections per day, using a combination of intermediate and rapid acting insulin analogs. Socioeconomic status (SES) of the cohort showed that 28 % were in the lower economic strata and 31 and 38 % were in the moderate to high economic strata, respectively. Two subjects did not divulge information on their SES. SES was significantly higher at the New Jersey site and the subjects at that site had better glycemic control (Table 1). Table 1 also shows marked differences in race at the two sites. We divided the race based on different socioeconomic status. For subjects between Bronx (New York) and New Jersey, we did see a significant difference between the race and the socioeconomic status. In New Jersey, there were significant number of subjects that belonged to the high and moderate socioeconomic status, when compared to the population in Bronx, New York (NJ vs. Bronx, $p = 0.001$). Compared to New Jersey, the diabetes preparedness was slightly better in subjects with low socioeconomic status in Bronx, New York ($p = 0.03$). There were no other differences that were noted regarding general preparedness ($p = 0.61$, NS) or diabetes preparedness ($p = 0.76$, NS) at either site despite socio-economic disparities. Further, there were no statistically significant differences between the HURTE scores (for post-traumatic stress events) and the different socioeconomic status at either site ($p = 0.36$).

In Fig. 1, the HURTE scores showed that the subjects in New Jersey had more symptoms of post-traumatic stress than the Bronx patients. There was a trend toward significance ($p < 0.06$).

Table 1 Patient characteristics at the two participating sites (Hackensack, NJ, and Bronx, NY)—patient characteristics include age in years, HbA1C %, race, gender, Hollingshead (SES status), past history of facing disaster, general preparedness and diabetes preparedness


	Hackensack, NJ (n = 65)	Bronx, NY (n = 77)	Total	Significance (p value)
Age (years)				
Mean ± SD	13.3 ± 2	13.3 ± 2	13.3 ± 2	NS
HbA1C %				
Mean ± SD	8.2 ± 1	9.3 ± 2.3	8.8 ± 1.9 % well controlled 24 % moderately controlled 40 % poorly controlled 36 %	<0.008
Frequencies (%) race				
Caucasian	75 (n = 49)	12 (n = 9)	41 (n = 58)	
African American	5 (n = 3)	22 (n = 17)	14 (n = 20)	<0.0001
Hispanic	14 (n = 9)	58 (n = 45)	38 (n = 54)	
Mixed	6 (n = 4)	8 (n = 6)	7 (n = 10)	
Gender				
Female	38 (49 %)	36 (55 %)	74 (52 %)	
Hollingshead (SES status)	45 ± 14	33 ± 16	39 ± 15	<0.0001
SES (frequency %)				
High	53 (n = 35)	27 (n = 21)	39 (n = 56)	<0.002
Moderate	36 (n = 23)	29 (n = 22)	31 (n = 45)	<0.001
Low	11 (n = 7)	44 (n = 34)	29 (n = 41)	0.31
Past history of facing disaster (frequency %)				
No history	1 (n = 1)	0	7 (n = 1)	
History of one	63 (n = 41)	66 (n = 51)	65 (n = 92)	NS
More than one	36 (n = 23)	34 (n = 26)	35 (n = 49)	
General preparedness (frequency %)				
High	39 (n = 25)	52 (n = 40)	46 (n = 65)	
Moderate	61 (n = 40)	42 (n = 32)	51.5 (n = 72)	NS
Low	0	6 (n = 5)	4 (n = 5)	
Diabetes preparedness (frequency %)				
High	85 (n = 55)	82 (n = 63)	83 (n = 118)	NS
Moderate	15 (n = 10)	14 (n = 11)	15 (n = 21)	NS
Low	0	4 (n = 3)	2 (n = 3)	0.03

Discussion

In this cross-sectional study, we compared two populations in the area where a major Hurricane occurred on October 29, 2012. Hackensack, NJ was more severely affected than the Bronx, NY. We showed that despite differences in SES and glycemic control in the two patient populations, general and diabetes emergency preparedness was similar for the two populations except for the diabetes preparedness in low SES population in Bronx, NY which was slightly better than the low SES population in Hackensack, NJ. There was a trend towards post-traumatic stress symptoms in the New Jersey population after the storm, perhaps because they were more affected by the disaster than the Bronx.

These study results were surprising because, of a previous study done in Texas, where we have shown that

poor glycemic control is associated with lower general and diabetes preparedness [10]. Those finding may have been due to storm lethargy, the idea that many storms are predicted in general and that people are asked to prepare for it. However, even in low SES populations particularly New York, emergency preparedness, especially for diabetes, was high. This is very reassuring that patients with diabetes are taking heed to warnings and have adequate supplies and report that they were prepared. The reason for storm preparedness may be due to community support, or the urban setting versus the rural setting in Texas. Emergency preparedness in the low SES population should be explored more. However, in concordance with our previous findings, general preparedness scores were lower than diabetes preparedness. For diabetes



patients it is not only important to have diabetes supplies but adequate supplies of food and water as well.

In a year with multiple disasters, our past research on disaster preparedness suggested hurricane fatigue and lower preparedness over time. In 2012, the only major storm to hit the Northeastern United States was Hurricane Sandy and perhaps it is possible that the low frequency of such incidents was the reason for good emergency preparedness, as compared to our survey in Texas when there were many powerful storms that were predicted but did not happen.

SES had no impact on diabetes preparedness and that is also contrary to our previous finding. SES continues to have an impact on overall glycemic control as previously reported within the US and outside [17–19]. Exploring why patients with low SES were prepared for storms may give insight into how to improve glycemic control in this population as well.

In this study, we also wanted to study the post-traumatic effects of the disaster and hence we compared two populations. New Jersey was more severely affected than the Bronx, and thus in looking at these two populations we could evaluate the psychological effects of the disaster. The study suggested that there was a trend toward more psychological symptoms in the New Jersey population compared to the Bronx. This is consistent with other disasters, such as the earthquakes in Japan [20, 21] and can affect both glycemic control and blood pressure management [22]. Additionally, the HURTE questionnaire could have been affected by the perceived fear of potential damage in the less affected group due to the exposure to media coverage of the disaster in the neighboring area.

The limitations of this study were that only two locations were surveyed and the most severely

affected areas were not surveyed. It is possible that if we had surveyed the more affected areas, we would have noted a more profound effect on post-traumatic stress related events. However, we note that even in moderately affected areas there is a psychological impact of disasters. Since this is a cross-sectional study it is not possible to have a cause and effect of disasters. There is also a recall bias associated with this kind of study since it is post-impact. In future, we will do a baseline evaluation of our patient population and compare it to an evaluation done after a disaster has occurred. Despite the limitations of the study, we have uncovered important factors that we previously thought affected disaster preparedness, like poor glycemic control and SES, and found them to not have an effect in this study.

Conclusions

In conclusion, preparing for a disaster is important for diabetes care and interventional studies to address the stress following disasters should be designed and ready to be implemented so that glycemic control remains unaffected and patients that are impacted by a major disaster are able to cope with the aftermath.

Abbreviations

FEMA: Federal Emergency Management Agency; SES: Socio-economic Status; T1DM: Type 1 Diabetes Mellitus; HURTE: Hurricane Related Traumatic Experiences.

Authors' contributions

RH participated in the conception, design and coordination of this study and revising it critically for important intellectual content. RH participated in the assignment of HURTE scores (Hurricane Related Traumatic Experiences) to evaluate the symptoms of post-traumatic stress. DNJ participated in the study design and acquisition of the data. JT participated in the data analysis and interpretation of the data. GD also participated in the design of the study, evaluation of HURTE scores (Hurricane Related Traumatic Experiences) for the symptoms of post-traumatic stress and study design. VR participated in the design of the study, performed the statistical analysis and drafted the manuscript. JS participated in the design and coordination of this study and helped to draft the manuscript. All authors read and approved the final manuscript.

Author details

¹ Pediatric Endocrinology and Diabetes, Children's Hospital at Montefiore and Albert Einstein College of Medicine, 3411 Wayne Ave, Suite: 4 M, Bronx, NY 10467, USA. ² Department of Psychiatry and Pediatrics, Children's Hospital at Montefiore, Bronx, NY, USA. ³ Pediatric Endocrinology and Diabetes, Children's Hospital at Montefiore, Bronx, NY, USA. ⁴ Pediatric Endocrinology and Diabetes, Albert Einstein College of Medicine, Bronx, NY, USA. ⁵ Pediatric Endocrinology and Diabetes, Hackensack University Medical Center, Hackensack, NJ, USA.

Acknowledgements

We would like to thank both the organizations, Children's Hospital at Montefiore/Albert Einstein College of Medicine at Bronx, NY and Hackensack Medical Center, Hackensack, NJ for participating in this study. There were no sponsors involved in this project.

Competing interests

The authors declare that they have no competing interests.

Appendix 1

DISASTER PREPAREDNESS SURVEY

Subject ID # _____

Date: _____

1) Do you live in an area which is prone to disasters? (Ex: a hurricane prone zone, area prone to floods, large city prone to terrorist attack)

Yes **No**

2) Have you faced any of the following disasters? (Check if you faced any one of the below)

Hurricane

Dam failure

Volcano

Tornado

Landslide

Fire (home fire, wild fire)

Flood

Earthquake

Nuclear emergency

(Nuclear power plant accident)

Tsunami

Winter storm

Terrorist attack

3) How many disasters have you faced in your lifetime? (any of the above mentioned)

None

One

More than one

Subject ID # _____

4) Do you have a family disaster plan to deal with disasters and emergencies? (a list of things to be done to be prepared in the event of a disaster)

Yes **No**

5) Do you have an emergency contact at a distant location in a case of evacuation due to a disaster/emergency? (Ex: A relative or friend's house in another city)

Yes **No**

6) Do you have an emergency communication plan in the event of a disaster? (Ex: how to contact family members located at different places in an emergency situation, through phone or any other source)

Yes **No**

17) Do you have an emergency evacuation plan to deal with disasters? (Like a place to meet, how to meet, a safer place to move in the event of a disaster)

Yes **No**

8) Do you know about the emergency helpline number (311) for information on evacuation routes, shelters and special needs transportation?

Yes **No**

9) Do you know about your community emergency plan for disasters?

Yes **No**

Subject ID # _____

10) Do you have a disaster supplies kit (72 Hour Emergency Kit)?

- At home
- At work place
- In your car/Truck
- I do not have a kit

11) Do you have a supply of the following items?

- i) A minimum of 3 day supply of food and water Yes No
- ii) At least a 3 day supply of current prescription Yes No
- iii) A first aid kit Yes No
- iv) Tools and emergency supplies Yes No
- v) Over the counter pills for aches and pains, nausea and vomiting, and diarrhea, antibiotic creams, bandages Yes No
- vi) Clothing and bedding Yes No
- vii) Copies of current drug prescriptions and lab-tests/procedures Yes No
- viii) Copies of Insurance or Medicare card Yes No

Subject ID # _____

ix) List of doctors and their phone numbers **Yes** **No**

x) Pharmacy phone number **Yes** **No**

12) Do you have smoke alarms in working condition installed in your house?

Yes **No**

13) Do you have a fire extinguisher in working condition?

Yes **No**

14) Do you know how to use a fire extinguisher?

Yes **No**

15) How many disasters/emergencies have you faced along with your child who has diabetes?

None

One

More than one

16) When was your child, first diagnosed with diabetes?

Less than one year

1-5 years

>5-10 years

More than 10 years

Subject ID # _____

17) Does your child with diabetes, always wear a Diabetes medical alert identification (ID)?

Yes **No**

18) Is your child up-to-date with all immunizations, including tetanus?

Yes **No**

19) Are you up-to-date with the diabetes education classes?

Yes **No**

20) Do you keep a supply of the following items with you always?

i) Prescription pills and/ insulin sufficient for at least 3 days **Yes** **No**

ii) Meter to measure blood sugar **Yes** **No**

iii) Lancets/ syringes/ pump supplies, if the child is on an insulin pump, meter supplies, sufficient for at least 3 days, cooler and cool pack for insulin **Yes** **No**

iv) Glucose tablets, soda with sugar, juice and hard candies **Yes** **No**

Subject ID # _____

v) Unexpired glucagon emergency kit Yes No

vi) Log book of blood sugars Yes No

vii) Copies of diabetes prescriptions and lab tests/ procedures Yes No

viii) Contact list of your child's Physician(s) Yes No

ix) Copy of your child's insurance card Yes No

21) Gender: Male Female

22) Ethnicity: Caucasian
 African American
 Non-White Latin American/ Spanish Ancestry
 Native American/ American Indian
 Asian/ Pacific Islander
 Other _____

23) Age of the child: _____

Subject ID # _____

24) Type of Diabetes Mellitus:

Type 1 Type 2

25) Hb-A1c %: _____

26) Insulin Regimen:

- NPH plus Humalog/ Novolog
- Lantus plus Humalog/Novolog
- Insulin Pump
- Insulin and Pills
- Only pills

27) Highest Level of Education of parent(s):

- Master's or Doctoral Degree
- Bachelor's Degree
- At least 1 year of college or any other specialized training
- Received high school diploma or GED
- Completed 10th or 11th grade
- Completed 7th, 8th, or 9th grade
- Completed less than 7th grade

28) Job description of parent(s): _____

29) Annual Household Income:

(According to the US Census Bureau, 2006)

- <\$22,500
- \$22,500-\$77,500
- >\$77,500

Subject ID # _____

Date: _____

Hurricane Related Traumatic Experiences (HURTE) Questionnaire

Please answer as **Y for Yes and N for No** beside the question

Perceived Life Threat:

1. At any time during the hurricane, did you think that you might die?

Life-Threatening Experiences:

1. Did windows or doors break in the place you stayed during the hurricane?
2. Did you get hurt during the hurricane?
3. Did you see anyone else get badly hurt during the hurricane?
4. Did you have to go outside during the hurricane because the building you were staying in was badly damaged?
5. Did a pet you like get hurt or die during the hurricane?
6. Did you get hit by anything falling or flying during the hurricane?

Loss-Disruption Experiences:

1. Was your home badly damaged or destroyed by the hurricane?
2. Were your clothes or toys ruined by the hurricane?
3. Has it been hard to see your friends since the hurricane because they moved or you moved?
4. Did you or your family have trouble getting enough food or water after the hurricane?
5. Did you move to a new place because of the hurricane?
6. Did you have to go to a new school because of the hurricane?

Subject ID # _____

7. Did you have to live away from your parents for a week or more because of the hurricane?
8. Has anyone stolen anything from your home since the hurricane?
9. Did one of your parents lose his or her job because of the hurricane?
10. Did your pet run away or have to be given away because of the hurricane?

Time 2:

Since the Hurricane:

1. Is one of your parents now out of a job?
2. Living in a house with a roof that leaks?
3. Have to travel longer to get to your school?
4. Living in the house where you lived before?
5. Has the damage to your house been fixed?
6. How many times have you moved?

For the questions below please answer as Not at all, a little, a lot or a whole lot:

7. How bothered are you by the way things look at home?
8. How bothered are you by the way things look in the neighborhood?
9. How bothered are you by family members not getting along?
10. How bothered are you by problems regarding spending time with friends?

Received: 22 October 2015 Accepted: 17 January 2016

Published online: 03 February 2016

References

1. The House D. The red cross and disaster planning. *J Ky Med Assoc.* 1964;62:40–1.
2. The pediatrician's role in disaster preparedness. American Academy of Pediatrics Committee on Pediatric Emergency Medicine. *Pediatrics.* 1997;99(1):130–3.
3. Cannon GM. Strengthening the Federal Emergency Management Agency's disaster response capabilities. *J Bus Contin Emer Plan.* 2008;2(3):281–93.
4. Fonseca VA, Smith H, Kuhadiya N, Leger SM, Yau CL, Reynolds K, et al. Impact of a natural disaster on diabetes: exacerbation of disparities and long-term consequences. *Diabetes Care.* 2009;32(9):1632–8.
5. Dabelea D, Mayer-Davis EJ, Saydah S, Imperatore G, Linder B, Divers J, et al. Prevalence of type 1 and type 2 diabetes among children and adolescents from 2001 to 2009. *JAMA.* 2014;311(17):1778–86. doi:10.1001/jama.2014.3201.
6. Lin CJ, Pierce LC, Roblin PM, Arquilla B. Impact of hurricane sandy on hospital emergency and dialysis services: a retrospective survey. *Prehosp Disaster Med.* 2014;29(4):374–9.
7. Lien C, Raimo J, Abramowitz J, Khanijo S, Kritharis A, Mason C, et al. Community healthcare delivery post-Hurricane Sandy: lessons from a mobile health unit. *J Community Health.* 2014;39(3):599–605.
8. Thethi TK, Yau CL, Shi L, Leger S, Nagireddy P, Waddadar J, et al. Time to recovery in diabetes and comorbidities following Hurricane Katrina. *Disaster Med Public Health Prep.* 2010;4(Suppl 1):S33–8. doi:10.1001/dmp.2010.10.
9. Kamps JL, Varela RE. Predictors of metabolic control in children with type 1 diabetes: the impact of Hurricane Katrina in a prospective study. *Diabetes Res Clin Pract.* 2010;88(3):234–41.
10. Renukuntla VS, Hassan K, Wheat S, Heptulla RA. Disaster preparedness in pediatric type 1 diabetes mellitus. *Pediatrics.* 2009;124(5):e973–7.
11. Bergen Co. Police: Possible levee breach could mean 1,000 in need of rescue, some from roofs of homes. CNN—Starting point; 2012.
12. Hurricane Sandy leaves N.J., but destruction, questions remain. *NJ.com;* 2012.
13. One year later Superstorm Sandy's impact still felt in the Bronx. *Bronx-Times;* 2013.
14. Mollica RF, Milic M. Social class and psychiatric practice: a revision of the Hollingshead and Redlich model. *Am J Psychiatry.* 1986;143(1):12–7.
15. Vigil JM, Geary DC. A preliminary investigation of family coping styles and psychological well-being among adolescent survivors of Hurricane Katrina. *J Fam Psychol.* 2008;22(1):176–80.
16. Ben-Ezra M, Palgi Y, Hamama-Raz Y, Rubin GJ, Goodwin R. The association of disaster-related experiences and self-reported recollections of national trauma with posttraumatic stress disorder symptoms following Hurricane Sandy. *J Clin Psychiatry.* 2013;74(12):1265–6.
17. Secrest AM, Costacou T, Gutelius B, Miller RG, Songer TJ, Orchard TJ. Association of socioeconomic status with mortality in type 1 diabetes: the Pittsburgh epidemiology of diabetes complications study. *Ann Epidemiol.* 2011;21(5):367–73.
18. Hatherly K, Smith L, Overland J, Johnston C, Brown-Singh L, Waller D, et al. Glycemic control and type 1 diabetes: the differential impact of model of care and income. *Pediatr Diabetes.* 2011;12(2):115–9.
19. Hassan K, Loar R, Anderson BJ, Heptulla RA. The role of socioeconomic status, depression, quality of life, and glycemic control in type 1 diabetes mellitus. *J Pediatr.* 2006;149(4):526–31.
20. Fujihara K, Saito A, Heianza Y, Gibo H, Suzuki H, Shimano H, et al. Impact of psychological stress caused by the Great East Japan Earthquake on glycemic control in patients with diabetes. *Exp Clin Endocrinol Diabetes.* 2012;120(9):560–3.
21. Kamoi K, Tanaka M, Ikarashi T, Miyakoshi M. Effect of the 2004 Mid Niigata Prefecture earthquake on glycemic control in type 1 diabetic patients. *Diabetes Res Clin Pract.* 2006;74(2):141–7.
22. Kamoi K, Tanaka M, Ikarashi T, Miyakoshi M. Effect of the 2004 Mid-Niigata Prefecture earthquake on home blood pressure measurement in the morning in type 2 diabetic patients. *Clin Exp Hypertens.* 2006;28(8):719–29.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

