

1 **Impact of birth weight and gender on early postnatal hypothalamic energy balance**  
2 **regulatory gene expression in the young lamb**

3  
4  
5  
6 3

7  
8  
9 4 C.L. Adam, T. Bake, P. A. Findlay, J. S. Milne, R. P. Aitken and J. M. Wallace

10  
11  
12 5 *Rowett Institute of Nutrition and Health, University of Aberdeen, Bucksburn, Aberdeen, AB21*

13  
14  
15 6 *9SB, UK*

16  
17  
18 7

19  
20  
21 8

22  
23  
24 9 **Sort title:** Birth weight, sex and hypothalamic programming

25  
26  
27  
28 10  
29  
30  
31 11  
32  
33 12  
34  
35 13 **Corresponding author:**

36  
37 14 Dr Clare L Adam

38  
39 15 Rowett Institute of Nutrition & Health

40  
41 16 University of Aberdeen

42  
43 17 Bucksburn

44  
45 18 Aberdeen

46  
47 19 AB21 9SB

48  
49 20 UK

50  
51 21 Tel: +44 1224 438658

52  
53 22 Fax: +44 1224 438699

54  
55 23 Email: [c.adam@abdn.ac.uk](mailto:c.adam@abdn.ac.uk)

56  
57 24

58  
59  
60  
61  
62  
63  
64  
65

**Abstract**

Intra-uterine growth restriction (IUGR) is involved in developmental metabolic programming and here we test the hypothesis that IUGR affects the developing hypothalamic energy balance regulatory pathways in a sex-specific manner. This experiment investigated early postnatal hypothalamic gene expression for six primary leptin- and insulin-sensitive neuropeptides and receptors in male and female IUGR (n = 8 and 9, respectively) and normal (N) birth weight lambs (n = 8 per gender) gestated and suckled by overnourished mothers. IUGR lambs were smaller at birth, had increased fractional growth rates (FGR), lower final body weight (11 weeks) and similar body fat content compared with N lambs, while males had higher final body weight and insulinemia but lower body fat and leptinemia than females. *In situ* hybridization revealed greater gene expression in the hypothalamic arcuate nucleus at 11 weeks for anorexigenic genes in females and orexigenic genes in males, with no effect of IUGR. Leptinemia correlated with gene expression for neuropeptide Y (NPY, negatively) in both sexes and pro-opiomelanocortin (POMC, positively) in females but with leptin receptor (negatively) only in males. Current FGR for girth correlated negatively with gene expression for NPY in males and POMC in females. Neither IUGR nor gender affected suckling activity (proxy for appetite) assessed at 3 weeks, but final NPY gene expression correlated with suckling weight gain in males. This study has revealed no effect of IUGR on early postnatal hypothalamic energy balance gene expression but a major effect of gender associated with major sex differences in adiposity and leptinemia.

**Key words:** developmental programming, intra-uterine growth restriction, adiposity, leptin

## 46 1. Introduction

1  
2  
3 47 The basic model of energy homeostasis in the mature animal includes peripheral  
4  
5 48 metabolic feedback hormones leptin and insulin regulating the activities of opposing  
6  
7  
8 49 orexigenic and anorexigenic circuits in the hypothalamus (Schwartz et al., 2000). These  
9  
10 50 neuronal circuits develop in the fetal brain, they are well established by birth in precocial  
11  
12  
13 51 species such as sheep and primates, and their development is influenced by the prenatal  
14  
15 52 nutritional environment (Grayson et al., 2010). Changes in the developing hypothalamic  
16  
17  
18 53 circuitry may underlie the apparent programming of a predisposition to obesity and altered  
19  
20 54 metabolic phenotype by intra-uterine growth restriction (IUGR), especially when such  
21  
22 55 offspring are born into an unrestricted nutritional environment (Gluckman and Hanson,  
23  
24  
25 56 2008). However, existing data largely come from studies of rodents in which hypothalamic  
26  
27  
28 57 neuroendocrine maturation occurs after birth (Breton, 2013); it is now appropriate to increase  
29  
30 58 our understanding of prenatal nutritional programming of the hypothalamus in larger,  
31  
32 59 precocial mammalian species.

34  
35 60 Previously we have demonstrated the presence of key anorexigenic and orexigenic  
36  
37 61 gene expression in the hypothalamic arcuate nucleus of the ovine fetus at mid as well as late  
38  
39  
40 62 gestation (Adam et al., 2008; Mühlhäusler et al., 2004). A consistent finding was the  
41  
42 63 sensitivity of anorexigenic neuropeptides in the arcuate nucleus to fetal nutrient (glucose)  
43  
44 64 supply, thus pro-opiomelanocortin (POMC) gene expression correlated with fetal glycemia at  
45  
46  
47 65 81 days (term = 147 days; Adam et al., 2008), intra-fetal glucose infusion increased POMC  
48  
49  
50 66 gene expression at 140 days (Mühlhäusler et al., 2005), and cocaine- and amphetamine-  
51  
52 67 regulated transcript (CART) gene expression correlated with fetal liver glycogen and was  
53  
54  
55 68 decreased in IUGR compared with normally-growing fetuses at 130 days (Adam et al.,  
56  
57 69 2011a). It remains to be determined to what extent prenatal changes in hypothalamic  
58  
59 70 neuropeptides persist in postnatal life when they may affect appetite and body weight  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

71 regulation in the free-living sheep. Findings from rodent offspring indicate early postnatal  
72 down-regulation of anorexigenic and up-regulation of orexigenic hypothalamic neuropeptides  
73 following prenatal fetal undernutrition caused by maternal undernutrition or placental  
74 insufficiency (Cripps et al., 2009; Desai et al., 2007; Huizinga et al., 2001). Earlier we  
75 reported in sheep that the presumed increase in fetal arcuate POMC gene expression induced  
76 in late gestation by maternal overnutrition and elevated glycemia was sustained in lambs at  
77 postnatal day 30 (Mühlhäusler et al., 2006). Now it is pertinent to investigate whether the  
78 presumed changes in anorexigenic hypothalamic gene expression in IUGR ovine fetuses are  
79 sustained postnatally.

80           Although insulin and leptin are present in the fetal circulation, there is little evidence  
81 for either hormone playing an adult-like nutritional signaling role in the hypothalamus.  
82 Precocial species like sheep lay down adipose tissue in the latter stages of gestation, which  
83 secretes leptin in proportional concentrations (Mühlhäusler et al., 2002), and the ovine fetal  
84 pancreas secretes insulin from mid gestation onwards (Aldoretta et al., 1998). Gene  
85 expression for receptors for both hormones is detected in the arcuate nucleus from mid  
86 gestation, but no correlation was found between circulating concentrations of either hormone  
87 and expression of key appetite regulatory genes in the fetal arcuate nucleus (Adam et al.,  
88 2008, 2011; Mühlhäusler et al., 2004). Circulating fetal insulin concentrations correlated  
89 negatively with arcuate insulin receptor (Ins-R) gene expression in an apparently adult-like  
90 ligand-receptor relationship, but circulating leptin concentrations were found to correlate  
91 positively with arcuate leptin receptor (OB-Rb) gene expression in late gestation (Adam et  
92 al., 2011). These latter findings were consistent with the substantial evidence for leptin  
93 playing a neurotrophic role in the neonate (Bouret, 2010). However by 5-6 months of age  
94 intracerebroventricular (ICV) leptin suppresses appetite in both male and female sheep  
95 indicating that a functional role for leptin signaling in the hypothalamus has developed by

96 this age (Adam et al., 2011b). Conversely, ICV-administered insulin had no effect on appetite  
97 in 5-6 month-old sheep (Adam et al., 2011b). In the present study, we examine relationships  
98 between hypothalamic arcuate neuropeptide and receptor gene expression, leptinemia and  
99 insulinemia at 3 months of age.

100 In addition to the effects of IUGR, we examine the influence of gender since there is  
101 evidence for sex differences in hypothalamic programming with respect to the hypothalamo-  
102 pituitary-adrenal ‘stress’ axis (Gardner et al., 2006; Wallace et al., 2011), and yet there is a  
103 lack of equivalent data with respect to the hypothalamic appetite regulatory axis.  
104 Furthermore, in the cohort of lambs used for the present study, whilst IUGR impacted  
105 postnatal fractional growth rates and glucose metabolism, gender had the overriding influence  
106 on body composition and metabolic hormone status (Wallace et al., 2013). Thus, in these  
107 lambs from our overnourished adolescent dam model of utero-placental insufficiency, IUGR  
108 led to increased fractional growth rates to 11 weeks of age and impaired glucose handling at 7  
109 weeks compared with normal birth weight lambs whereas females had increased adiposity  
110 and leptinemia compared with males (Wallace et al., 2013). Importantly, IUGR and normal  
111 birth weight lambs in this cohort were both born to dams that had high dietary intakes  
112 throughout pregnancy and lactation, thus allowing us to examine their early postnatal  
113 phenotype without the confounding effect of differences in maternal nutrition. In order to  
114 estimate appetite drive and voluntary food intake in the lambs, suckling activity was assessed  
115 at 3 weeks of age.

116 This experiment therefore investigated the influences of IUGR and gender on the  
117 early postnatal phenotype with respect to the developing hypothalamic appetite regulatory  
118 pathways. Specifically, we examined gene expression for six primary leptin- and insulin-  
119 sensitive hypothalamic neuropeptides and receptors at 11 weeks of age in low and normal  
120 birth weight male and female lambs born to overnourished adolescent mothers. We test the

121 hypothesis that IUGR affects the developing hypothalamic energy balance regulatory  
122 pathways in a sex-specific manner.

123

## 124 **2. Material and Methods**

### 125 *2.1 Animals*

126 All procedures were licensed under the UK Animals (Scientific Procedures) Act 1986  
127 and approved by local Ethical Review Committee. The derivation of the lambs is described in  
128 detail by Wallace et al. (2013). Briefly, growing adolescent recipient ewes (Dorset Horn x  
129 Mule) had been implanted with singleton embryos, derived from superovulated donors  
130 (Border Leicester x Scottish Blackface) and a single sire (Dorset Horn), and given a high  
131 quality complete diet *ad libitum* throughout pregnancy and lactation. The complete diet  
132 contained 12 MJ metabolizable energy and 140 g crude protein per kg dry matter and *ad*  
133 *libitum* intakes were calculated to promote rapid maternal growth during pregnancy leading  
134 to restricted placental growth, and hence restricted fetal growth in ~50% cases, followed by  
135 maximal milk yields during the 11-week lactation. In addition, lambs had access to their  
136 mothers' feed at all times. There was a continuous distribution of birth weights from which  
137 lambs were categorized as intra-uterine growth restricted (IUGR) or normal birth weight (N)  
138 (Wallace et al., 2013). The present study comprised 17 IUGR (n = 8 male, n = 9 female) and  
139 16 N lambs (n = 8 per gender), with most of the individual embryo donors represented in  
140 both categories.

141 The lambs were weighed, measured and blood sampled mid-morning at 5-day  
142 intervals up to ~68 days of age and just before euthanasia at 77 days (11 weeks). Plasma  
143 leptin and insulin were determined by in-house radioimmunoassays (Marie et al., 2001 and  
144 MacRae et al., 1991, respectively) with all inter and intra-assay coefficients of variation less  
145 than 10% (as reported in Wallace et al., 2013). They were euthanized by lethal injection of

146 sodium pentobarbitone (10-15 ml Euthesate; 200 mg pentobarbitone/ml; Willows Francis  
147 Veterinary, Crawley, UK). Whole brains were removed, immediately frozen in isopentane  
148 over dry ice and stored at -80°C. Perirenal and visceral (omental and mesenteric) fat depots  
149 were dissected out and weighed.

## 150 *2.2 Suckling activity assessment*

151 Suckling activity was determined at  $23 \pm 0.9$  days of age, coincident with the  
152 presumed peak lactation and prior to lambs showing any significant interest in eating their  
153 mothers' food. Ewes were milked by hand following intravenous oxytocin injection  
154 (Oxytocin-S<sup>®</sup> 10 i.u. per ewe; Intervet Ltd, Cambridge, UK) in order to empty the udder, the  
155 lamb was weighed and access to the udder was prevented using an udder cover. After 3  
156 hours, the lamb was reweighed to determine fasting weight loss, the udder cover was  
157 removed and the number and duration of suckling bouts were determined for a period of 60  
158 minutes. Lambs were weighed at 15-minute intervals throughout this observation period and  
159 again at 90 minutes.

## 160 *2.3 Hypothalamic gene expression*

161 The frozen brains were trimmed down to a mid-ventral block, mounted and sectioned  
162 by cryostat coronally through the hypothalamus from the mammillary body (caudal) to the  
163 optic chiasm (rostral). Sections (20  $\mu$ m) were thaw-mounted onto poly-L-lysine-coated slides  
164 and stored at -80°C. Gene expression for neuropeptide Y (NPY), agouti-related peptide  
165 (AGRP), POMC, CART, OB-Rb and Ins-R was measured by *in situ* hybridization, using  
166 techniques described in detail elsewhere (Adam et al., 1997). The NPY riboprobe was  
167 generated from a rat cDNA (Adam et al., 1997), the CART probe from a cloned sheep cDNA  
168 (Barrett et al., 2001), and AGRP and POMC probes were generated from cloned Siberian  
169 hamster cDNAs (Mercer et al., 2000). A riboprobe complementary to fragments of the  
170 intracellular domain of OB-Rb was generated from a cloned sheep cDNA (Mercer et al.,

1998), and the Ins-R riboprobe was generated from a partial ovine cDNA (Archer et al., 2005). All probes have previously been validated in sheep brain (Adam et al., 2002; Archer et al., 2005) and corresponding sense probes showed no hybridization. Briefly, sections were fixed, acetylated, and hybridized overnight at 58 °C using <sup>35</sup>S-labelled cRNA probes (1–1.5 x 10<sup>7</sup> cpm/ml). They were then treated with RNase A, desalted, with a final high stringency wash (30 min) in 0.5 X SSC at 60 °C (Ins-R at 75 °C), dried and apposed for 7-10 days to Hyperfilm β-max (Amersham Pharmacia Biotech UK Ltd, Little Chalfont, Bucks, UK). Intensity and total area of hybridization were quantified in the hypothalamic arcuate nucleus on each autoradiographic image, using the Image-Pro Plus system (Media Cybernetics, Silver Spring, MD, USA). Example images are shown in Fig. 1. The integrated intensity of the hybridization signal (i.e. the optical density integrated over the total hybridization area) was computed using standard curves generated from <sup>14</sup>C autoradiographic micro-scales (Amersham Pharmacia Biotech UK Ltd, Little Chalfont, Bucks, UK). For each probe, up to 6 sections spanning the medial hypothalamus (i.e. in the region midway between the mammillary body and optic chiasm, further identified by third ventricle morphology) were examined from each brain. All reagents were obtained from Sigma (Sigma UK, Poole, Dorset, UK) unless otherwise stated.

#### 2.4 Statistical analyses

Effects of birth weight category, gender and their interaction were examined by analysis of variance (General Linear Model; Minitab 16, Minitab Inc., State College, PA), and data are presented as means ± standard errors. Pearson product-moment correlation analyses and linear regression were used to explore relationships between variables where indicated (Minitab 16), and data are presented as correlation coefficients (r). Statistical significance was taken as P < 0.05, and a trend is indicated where P = 0.06 to 0.10.



### 196 3. Results

#### 197 3.1 Postnatal growth, body composition, insulinemia and leptinemia

198 Detailed growth and metabolic data have been reported elsewhere for these lambs (n  
199 = 38 in Wallace et al., 2013) and relevant values are shown here for the representative subset  
200 of animals used in the present study (n = 33, Table 1). Birth weight was lower for IUGR  
201 versus N ( $P < 0.001$ ) and for females versus males ( $P < 0.05$ ), shoulder height at birth was  
202 lower for IUGR versus N ( $P < 0.001$ ), but fractional growth rates (FGR, absolute growth rate  
203 relative to size at birth) for body weight, shoulder height, girth at the thorax and girth at the  
204 umbilicus were all higher in IUGR lambs ( $P < 0.05-0.001$ ), with an effect of gender on FGR  
205 for girth at the thorax (females  $>$  males,  $P < 0.001$ ) and a trend towards an effect of gender on  
206 FGR for girth at umbilicus (females  $>$  males,  $P < 0.08$ ). Nonetheless, IUGR lambs had lower  
207 final body weight at 11 weeks compared with N lambs ( $P < 0.001$ ), with no effect on internal  
208 body fat content, while males had higher final body weight than females ( $P < 0.001$ ) but  
209 lower perirenal fat mass ( $P < 0.01$ ) and lower proportional total internal fat mass ( $P < 0.01$ ).  
210 Plasma insulin during days 65-73 (males  $>$  females;  $P < 0.05$ ) and leptin both during days 65-  
211 73 (males  $<$  females;  $P < 0.01$ ) and on final day 77 (males  $<$  females;  $P < 0.001$ ) were  
212 influenced by gender but not IUGR (Table 1). For both males and females, plasma leptin  
213 concentrations in the days before euthanasia correlated positively with measures of internal  
214 fat mass ( $P < 0.05-0.001$ ; Table 2) but plasma insulin only correlated with perirenal fat mass  
215 in males ( $P < 0.01$ ).

#### 216 3.2 Suckling activity

217 Weight loss during the 3-hour fast prior to the suckling assessment at 23 days of age,  
218 the number and duration of suckling episodes, and weight gain during suckling were  
219 independent of IUGR status and gender (Table 3). However, for all animals together, birth  
220 weight correlated positively with absolute fasting weight loss ( $r = 0.407$ ,  $P < 0.05$ ), post

221 fasting weight gain to 60 ( $r = 0.576$ ,  $P < 0.001$ ) and 90 minutes ( $r = 0.445$ ,  $P < 0.01$ ) and  
 222 tended to correlate with the number of suckling episodes per 60 minutes ( $r = 0.335$ ,  $P < 0.06$ ).  
 223 Similar and slightly stronger relationships were evident between current weight and absolute  
 224 fasting weight loss ( $r = 0.540$ ,  $P < 0.001$ ), post fasting weight gain to 60 ( $r = 0.555$ ,  $P <$   
 225  $0.001$ ) and 90 ( $r = 0.503$ ,  $P < 0.005$ ) minutes. However, when expressed as relative (%)  
 226 weight changes as opposed to absolute weight changes, or when males and females were  
 227 examined separately, none of the foregoing correlations were significant.

### 228 3.3 Hypothalamic gene expression

229 *In situ* hybridization revealed greater gene expression in the hypothalamic arcuate  
 230 nucleus for CART and POMC in females than males ( $P < 0.01-0.05$ ), and greater gene  
 231 expression for OB-Rb, NPY and AGRP in males than females ( $P < 0.001-0.01$ ), but no  
 232 effects of IUGR, and there was no effect of gender or IUGR on Ins-R gene expression (Figs 1  
 233 and 2).

234 Given the clear gender differences in gene expression, significant interrelationships  
 235 and relationships with measures of growth, adiposity, metabolic hormones and appetite were  
 236 explored within each sex. NPY and AGRP gene expression were positively correlated in both  
 237 males ( $r = 0.807$ ,  $P < 0.001$ ) and females ( $r = 0.637$ ,  $P < 0.01$ ). In males, NPY and AGRP  
 238 correlated negatively with plasma leptin ( $P < 0.01-0.001$ ; Table 4 and Fig. 3A) and positively  
 239 with OB-Rb ( $P < 0.05$ ), and NPY correlated negatively with plasma insulin ( $P < 0.05$ ); NPY,  
 240 AGRP and OB-Rb all correlated negatively with perirenal fat mass ( $P < 0.05-0.001$ ; Table 4).  
 241 In females, POMC gene expression correlated positively with final plasma leptin ( $P < 0.05$ )  
 242 and NPY correlated negatively with plasma leptin during the last week ( $P < 0.05$ ), but none  
 243 of the neuropeptide genes correlated with OB-Rb; AGRP correlated negatively with perirenal  
 244 fat mass ( $P < 0.05$ ; Table 4 and Figs 3 A,B). OB-Rb correlated negatively with plasma leptin  
 245 in males but not females (Table 4 and Fig. 3C) and with plasma insulin in males but not

246 females (Table 4). Gene expression for ins-R and plasma insulin did not correlate in either  
247 sex.

248           None of the hypothalamic genes correlated with birth weight or final body weight  
249 within either sex but some correlations were found with final current FGR (CFGR, absolute  
250 growth rate relative to current weight at 68 days of age). Thus, within only males, NPY  
251 correlated negatively with thorax girth CFGR ( $P < 0.05$ ) and there was a trend towards  
252 similar correlation for AGRP ( $P < 0.07$ ; Table 4). Within only females, POMC correlated  
253 negatively with body weight CFGR ( $P < 0.05$ ) and umbilical girth CFGR ( $P < 0.05$ ; Table 4).  
254 Finally, in males only, NPY correlated positively with percentage weight gain to 90 minutes  
255 during the suckling assessment at 23 days of age ( $P < 0.05$ ; Table 4). No other significant  
256 correlations were found between any of the hypothalamic genes and weight loss during  
257 fasting, weight gain during suckling or suckling activity during the assessment.

258

#### 259 **4. Discussion**

260           This study has revealed no effect of birth weight but a major effect of gender on gene  
261 expression for hypothalamic energy balance regulatory pathways in young ovine offspring.  
262 Thus females had higher anorexigenic gene expression (POMC, CART) but males had higher  
263 orexigenic gene expression (NPY, AGRP, OB-Rb) in the arcuate nucleus. Gender also had a  
264 major effect on adiposity since females had higher levels of body fat and leptinemia, with no  
265 effect of birth weight. We therefore reject the hypothesis that IUGR affects the developing  
266 hypothalamic energy balance regulatory pathways in a sex-specific manner at this early  
267 postnatal life stage, since gender itself had the overriding influence irrespective of birth  
268 weight.

269 Growth and body composition data from the present subset of lambs was wholly  
1  
2 270 representative of the total number reported in Wallace et al. (2013) whereby low birth weight  
3  
4 271 (categorized as IUGR) was a direct reflection of reduced placental size. IUGR decreased birth  
5  
6  
7 272 weight and final body weight at 11 weeks and increased postnatal FGR, but absolute catch-up  
8  
9  
10 273 growth was not shown. However, gender had the major effect on final body weight (males >  
11  
12 274 females), adiposity (females > males), leptinemia (females > males) and insulinemia (males >  
13  
14 275 females). Importantly, plasma leptin was an accurate circulating indicator of adipose reserves  
15  
16  
17 276 in both sexes. By contrast, plasma insulin showed a weaker relationship with adiposity (only  
18  
19 277 with perirenal fat mass) and only in males. Despite birth weight and on-going body weight  
20  
21  
22 278 differences, there was no evidence for IUGR or gender affecting postnatal appetite drive  
23  
24 279 since there were no group effects on weight gain or frequency or duration of suckling bouts  
25  
26  
27 280 during the suckling assessment at 3 weeks of age. By contrast, De Blasio et al. (2007)  
28  
29 281 observed that IUGR lambs (obtained by pre-mating maternal carunclectomy) had increased  
30  
31  
32 282 suckling (feeding) activity at 15 days of age, indicative of increased appetite. The difference  
33  
34 283 may be attributable to the younger age and/or the more rapid catch-up growth in De Blasio's  
35  
36  
37 284 lambs compared with our model. Although correlation analysis of our data revealed that  
38  
39 285 heavier lambs (at birth and current) consumed more, with higher absolute weight gain during  
40  
41  
42 286 suckling, this relationship did not hold true for relative (percentage) weight gain, suggesting  
43  
44 287 that there was indeed no effect on appetite drive. On the other hand, these data are consistent  
45  
46  
47 288 with the original premise that milk yields (i.e. available food for the offspring) would be the  
48  
49 289 same between IUGR and N mothers receiving the same nutritional regime in our model.

50  
51 290 Given the lack of sex differences in postnatal appetite drive and body weight FGR,  
52  
53  
54 291 the clear sex differences in underlying hypothalamic appetite-regulatory neuropeptides were  
55  
56 292 perhaps surprising, albeit examined at a later age. However, a recent review of sex  
57  
58  
59 293 differences in developmental programming, based largely on phenotype and peripheral  
60  
61  
62  
63  
64  
65

294 molecular outcomes, identifies temporal, spatial and biochemical differences between male  
1  
2  
3 295 and female pre- and postnatal development (Aiken and Ozanne, 2013). Our data have now  
4  
5 296 additionally shown important sex differences in hypothalamic neuroendocrine development.  
6  
7 297 The reasons for these differences may include inherent (genetic) sex differences or  
8  
9 298 differences in timing of development, but also the influence of different steroid hormone  
10  
11 299 exposure and, in the present study, differences in adiposity. Gonadal steroid hormone  
12  
13 300 secretion is initiated early in ovine gestation (Quirke et al., 2001) and so development pre-  
14  
15 301 and postnatally occurs in an environment of high testosterone for males and high estrogen for  
16  
17 302 females. Consistent with the increased orexigenic/anorexigenic balance in gene expression in  
18  
19 303 the present males, testosterone is known to up-regulate NPY gene expression in adult male  
20  
21 304 sheep (Dobbins et al., 2004), increase AGRP immunoreactivity in prenatally androgenized  
22  
23 305 adult ewes (Sheppard et al., 2011) and decrease POMC gene expression in perinatally  
24  
25 306 androgenized female mice (Nohara et al., 2011). Consistent with the increased  
26  
27 307 anorexigenic/orexigenic balance in the females, estradiol stimulates POMC and inhibits NPY  
28  
29 308 and AGRP gene expression in young rats (Santollo et al., 2012). Exposure to the different  
30  
31 309 steroid milieux during development may program the different ratios of  
32  
33 310 orexigenic/anorexigenic expression between males and females to produce a similar appetite  
34  
35 311 phenotype, in other words producing a sex difference in hypothalamic 'set point' for a given  
36  
37 312 food intake.

313           Apart from the presumed difference in circulating sex steroids, the major significant  
314 gender difference observed in the present study was in body composition. The importance of  
315 the hypothalamus-adipose axis in developmental programming has recently been recognized  
316 (Breton, 2013), and the present sex differences in adiposity may have been in part causally  
317 related to the differences in hypothalamic gene expression. Thus females had greater amounts  
318 of adipose tissue, circulating leptin, and hypothalamic POMC and CART gene expression

1 319 whereas males had less adipose tissue and leptin but more NPY, AGRP and OB-Rb gene  
2 320 expression. Both scenarios fit the basic model of adult energy balance regulation (Schwartz et  
3  
4 321 al., 2000), but some additional sex differences were revealed by the correlation analyses.  
5  
6 322 Both NPY and AGRP gene expression correlated with OB-Rb gene expression (positively)  
7  
8 323 and plasma leptin (negatively), consistent with their adult-like regulation by leptin via its  
9  
10 324 receptor (Adam et al., 2002), but only in males. Meanwhile in females, although gene  
11  
12 325 expression for NPY (negatively) and POMC (positively) correlated with leptin in an adult-  
13  
14 326 like manner, there was no correlation with OB-Rb gene expression, which was inconsistent  
15  
16 327 with neuropeptide regulation by leptin via its receptor. Nonetheless, females had overall  
17  
18 328 lower OB-Rb expression and higher plasma leptin values and the absence of receptor-ligand  
19  
20 329 correlation may have been because OB-Rb levels were already baseline and unable to  
21  
22 330 respond to further increases in leptinemia. It would seem that changes in post-receptor  
23  
24 331 sensitivity may underlie the present association between higher leptin and higher POMC gene  
25  
26 332 expression in the female lambs.  
27  
28  
29  
30  
31  
32  
33

34 333 Wallace et al. (2013) found no evidence for IUGR leading to decreased neonatal  
35  
36 334 leptin in these lambs, unlike in rats (Coupé et al., 2010), perhaps reflecting a difference  
37  
38 335 between precocial and altricial species. All the lambs had similar (baseline) plasma leptin  
39  
40 336 concentrations at birth which increased rapidly for the first week; thereafter IUGR and N  
41  
42 337 groups had similar values throughout but the sexes diverged (females > males) with the  
43  
44 338 difference significant from 4 weeks (Wallace et al., 2013). Compared to males, the female  
45  
46 339 hypothalamus therefore had prolonged postnatal exposure to greater amounts of leptin as well  
47  
48 340 as the significantly greater leptinemia in the week before and on the day of euthanasia  
49  
50 341 reported here. Leptin concentrations in early life are important for the development of the  
51  
52 342 adult metabolic profile (Granado et al. 2012) and these authors concluded that reported  
53  
54 343 differences in the effects of postnatal hyperleptinemia may be attributed to critical time points  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

344 of sensitivity and/or prenatal influences; however, our present data suggest that gender could  
345 also be a major influencing factor.

346         The evidence for leptin's critical role in neonatal programming of metabolic  
347 neuroendocrine hypothalamic pathways comes from altricial rodent models (Bouret and  
348 Simerly, 2007), and it is likely that the temporal importance of leptinemia may be different in  
349 precocial species like sheep in which hypothalamic development and adipogenesis occur  
350 prenatally. Moreover, although sex differences in adult leptinemia (female > male) have long  
351 been recognized (Kennedy et al., 1997; Watanobe and Suda, 1999), the influence of gender  
352 on leptin's neurotrophic actions is unreported. There are few studies of neonatal leptin  
353 concentrations in lambs, with no gender comparisons (Bispham et al., 2002; Long et al.,  
354 2011; McFadin et al., 2002), but the sex difference in leptinemia in the present lambs was  
355 unequivocal (Wallace et al., 2013). Higher than normal leptin concentrations occurring earlier  
356 than normal in neonatal rodents decrease hypothalamic leptin receptors (Toste et al., 2006)  
357 and lead to leptin resistance in adulthood (Ahima and Hileman, 2000; Yura et al., 2005). The  
358 present female lambs, exposed to elevated leptinemia from soon after birth, may have been  
359 showing signs of leptin resistance (i.e. lack of correlation between leptin and its receptor and  
360 low expression of leptin receptor) whereas the males exposed to lower leptinemia exhibited  
361 adult-like hypothalamic neuropeptide and receptor responses and no sign of leptin resistance.  
362 Nonetheless, in spite of the observed sex differences in the developing hypothalamic leptin  
363 signaling system at 3 months, both sexes show a decreased food intake response to intra-  
364 hypothalamic leptin indicative of adult-like functionality at 5-6 months of age (Adam et al.,  
365 2011b).

366         In early postnatal life (11 weeks), the expression of hypothalamic energy balance  
367 genes was not affected by IUGR and none were correlated with birth weight or final weight  
368 in either sex. NPY gene expression correlated positively with the percentage weight gain to

369 90 minutes during the suckling assessment, indicative of increased orexigenic drive with  
1  
2 370 higher amounts of food intake, but only in the males. Since the suckling assessment was  
3  
4 371 made 8 weeks before brains were taken for gene expression analysis, current fractional  
5  
6 372 growth rates (CFGR) just prior to euthanasia may provide a more temporally relevant  
7  
8 373 indicator of overall energy balance in these lambs. However, the opposite relationship was  
9  
10 374 seen, with NPY gene expression in males correlating negatively with thorax girth CFGR,  
11  
12 375 suggesting that orexigenic drive decreased with increasing anabolic state (as in adult  
13  
14 376 mammals; Schwartz et al., 2000). Conversely, POMC gene expression in female lambs  
15  
16 377 correlated negatively with body weight CFGR and umbilical girth CFGR, suggesting that  
17  
18 378 increased body energy status was associated with decreased anorexigenic drive (unlike in  
19  
20 379 adults; Schwartz et al., 2000). In other words, the dominant orexigenic pathway in males  
21  
22 380 appears to down-regulate in order to limit positive energy balance whereas the dominant  
23  
24 381 anorexigenic pathway in females appears to down-regulate in order to facilitate positive  
25  
26 382 energy balance. These findings may reflect sex differences in the stage of maturity for the  
27  
28 383 developing neuroendocrine hypothalamus, as seen in other organ systems (Aiken and  
29  
30 384 Ozanne, 2013).

385         Unlike leptinemia, insulinemia was higher in males than females in this study. In  
39  
40 386 males, insulinemia correlated with perirenal fat mass, providing an additional adiposity signal  
41  
42 387 to the brain where it correlated negatively with NPY and OB-Rb gene expression, in  
43  
44 388 agreement with the basic model of adult energy balance regulation (Schwartz et al., 2000).  
45  
46 389 There was no evidence for an association between insulinemia and hypothalamic gene  
47  
48 390 expression in the females, and indeed studies in adult rats have shown that the male brain is  
49  
50 391 more sensitive to insulin than the female brain (Clegg et al., 2003). Conversely, the female  
51  
52 392 brain is more leptin-sensitive than that of the male, with estrogen increasing leptin sensitivity  
53  
54 393 and decreasing insulin sensitivity (Clegg et al., 2006). There was no evidence for insulin  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 394 signal regulation through its hypothalamic receptor in either sex in this study, in contrast to  
2 395 the late gestation fetus (Adam et al., 2011a), but a negative correlation with OB-Rb gene  
3  
4 396 expression in the males agrees with the inhibitory action of insulin on OB-Rb previously  
5  
6  
7 397 demonstrated in adult sheep (Daniel et al., 2000). Altogether, our data would tend to indicate  
8  
9 398 that insulin may play a significant hypothalamic signaling role in addition to leptin in the  
10  
11 399 young postnatal male but not in the female. However, in terms of the food intake response to  
12  
13 400 intrahypothalamic insulin, neither the male nor female brain appeared sensitive to insulin at  
14  
15 401 5-6 months of age (Adam et al., 2011b).  
16  
17  
18

19 402 Therefore, while the present data do not preclude an effect of IUGR in later life, this  
20  
21 403 study has revealed no effect of IUGR on early postnatal hypothalamic energy balance gene  
22  
23 404 expression but a major effect of gender which was associated with major sex differences in  
24  
25 405 adiposity and leptinemia. Furthermore, the data strongly indicate that the sexes should be  
26  
27 406 considered separately in studies of developmental hypothalamic programming.  
28  
29  
30

31 407

### 32 408 **Acknowledgements**

33  
34 409 Research funded by the Scottish Government's Rural and Environment Science and  
35  
36 410 Analytical Services Division, including the Strategic Partnership for Animal Science  
37  
38  
39 411 Excellence. No conflicts of interest are declared by the authors.  
40  
41  
42  
43  
44 412  
45  
46 413  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

414 **References**

- 1  
2 415 Adam, C.L., Archer, Z.A., Findlay, P.A., Thomas, L., Marie, M., 2002. Hypothalamic gene  
3  
4 416 expression in sheep for cocaine- and amphetamine-regulated transcript, pro-  
5  
6  
7 417 opiomelanocortin, neuropeptide Y, agouti-related peptide and leptin receptor, and  
8  
9  
10 418 responses to negative energy balance. *Neuroendocrinology* 75, 250–256.
- 11  
12 419 Adam, C.L., Bake, T., Findlay, P.A., Milne, J.S., Aitken, R.P., Wallace, J.M., 2011a. Effects  
13  
14 420 of altered glucose supply and adiposity on expression of hypothalamic energy balance  
15  
16  
17 421 regulatory genes in late gestation growth restricted ovine fetuses. *Int. J. Dev.*  
18  
19 422 *Neurosci.* 29, 775-781.
- 20  
21  
22 423 Adam, C.L., Findlay, P.A., Aitken, R.P., Milne, J.S., Wallace, J.M., 2011b. Influence of birth  
23  
24 424 weight, sex, age and adiposity on central leptin and insulin sensitivity in young  
25  
26  
27 425 growing sheep, as indicated by changes in voluntary food intake. *Proc. Nut. Soc.* 70  
28  
29 426 (OCE6), E382 (abstract).
- 30  
31  
32 427 Adam, C.L., Findlay, P.A., Chanet, A., Aitken, R.P., Milne, J.S., Wallace, J.M., 2008.  
33  
34 428 Expression of energy balance regulatory genes in the developing ovine fetal  
35  
36 429 hypothalamus at midgestation and the influence of hyperglycemia. *Am. J. Physiol.*  
37  
38  
39 430 *Regul. Integr. Comp. Physiol.* 294, R1895-1900.
- 40  
41 431 Adam, C.L., Findlay, P.A., Kyle, C.E., Young, P., Mercer, J.G., 1997. Effect of chronic food  
42  
43 432 restriction on pulsatile luteinizing hormone secretion and hypothalamic neuropeptide  
44  
45  
46 433 Y gene expression in castrate male sheep. *J. Endocrinol.* 152, 329–337.
- 47  
48  
49 434 Ahima, R.S., Hileman, S.M., 2000. Postnatal regulation of hypothalamic neuropeptide  
50  
51 435 expression by leptin: implications for energy balance and body weight regulation.  
52  
53 436 *Regul. Pept.* 92, 1-7.
- 54  
55  
56 437 Aiken, C.E., Ozanne, S.E., 2013. Sex differences in developmental programming models.  
57  
58 438 *Reproduction* 145, R1-R13.
- 59  
60  
61  
62  
63  
64  
65

- 1  
2  
3 440 Aldoretta, P.W., Carver, T.D., Hay, W.W. Jr., 1998. Maturation of glucose-stimulated insulin  
4  
5 441 secretion in fetal sheep. *Biol. Neonate* 73, 375-386.  
6  
7 442 Archer, Z.A., Rhind, S.M., Findlay, P.A., Kyle, C.E., Barber, M.C., Adam, C.L., 2005.  
8  
9 443 Hypothalamic responses to peripheral glucose infusion in food-restricted sheep are  
10  
11 444 influenced by photoperiod. *J. Endocrinol.* 184, 515-525.  
12  
13 445 Barrett, P., Morris, M.A., Moar, K.M., Mercer, J.G., Davidson, J.A., Findlay, P.A., Adam,  
14  
15 446 C.L., Morgan, P.J., 2001. The differential regulation of CART gene expression in a  
16  
17 447 pituitary cell line and primary cell cultures of ovine pars tuberalis cells. *J.*  
18  
19 448 *Neuroendocrinol.* 13, 347-352.  
20  
21 449 Bispham, J., Budge, H., Mostyn, A., Dandrea, J., Clarke, L., Keisler, D.H., Symonds, M.E.,  
22  
23 450 Stephenson, T., 2002. Ambient temperature, maternal dexamethasone, and postnatal  
24  
25 451 ontogeny of leptin in the neonatal lamb. *Pediatr. Res.* 52, 85-90.  
26  
27 452 Bouret, S.G., 2010. Neurodevelopmental actions of leptin. *Brain Res.* 1350, 2-9.  
28  
29 453 Bouret, S.G., Simerly, R.B., 2007. Development of leptin-sensitive circuits. *J.*  
30  
31 454 *Neuroendocrinol.* 19, 575-582.  
32  
33 455 Breton, C., 2013. The hypothalamus-adipose axis is a key target of developmental  
34  
35 456 programming by maternal nutritional manipulation. *J. Endocrinol.* 216, R19-31.  
36  
37 457 Clegg, D.J., Brown, L.M., Woods, S.C., Benoit, S.C., 2006. Gonadal hormones determine  
38  
39 458 sensitivity to central leptin and insulin. *Diabetes* 55, 978-987.  
40  
41 459 Clegg, D.J., Riedy, C.A., Smith, K.A., Benoit, S.C., Woods, S.C., 2003. Differential  
42  
43 460 sensitivity to central leptin and insulin in male and female rats. *Diabetes* 52, 682-687.  
44  
45 461 Coupé, B., Amarger, V., Grit, I., Benani, A., Parnet, P., 2010. Nutritional programming  
46  
47 462 affects hypothalamic organization and early response to leptin. *Endocrinology* 151,  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 463 Cripps, R.L., Martin-Gronert, M.S., Archer, Z.A., Hales, C.N., Mercer, J.G., Ozanne, S.E.,  
1  
2 464 2009. Programming of hypothalamic neuropeptide gene expression in rats by  
3  
4 465 maternal dietary protein content during pregnancy and lactation. *Clin. Sci. (Lond.)*  
5  
6  
7 466 117, 85-93.
- 8  
9 467 Daniel, J.A., Thomas, M.G., Hale, C.S., Simmons, J.M., Keisler, D.H., 2000. Effect of  
10  
11 468 cerebroventricular infusion of insulin and (or) glucose on hypothalamic expression of  
12  
13 469 leptin receptor and pituitary secretion of LH in diet-restricted ewes. *Domest. Anim.*  
14  
15 470 *Endocrinol.* 18, 177-185.
- 16  
17  
18 471 De Blasio, M.J., Gatford, K.L., Robinson, J.S., Owens, J.A., 2007. Placental restriction of  
19  
20 472 fetal growth reduces size at birth and alters postnatal growth, feeding activity, and  
21  
22 473 adiposity in the young lamb. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 292,  
23  
24 474 R875-R886.
- 25  
26  
27 475 Desai, M., Gayle, D., Han, G., Ross, M.G., 2007. Programmed hyperphagia due to reduced  
28  
29 476 anorexigenic mechanisms in intrauterine growth-restricted offspring. *Reprod. Sci.* 14,  
30  
31 477 329-337.
- 32  
33  
34 478 Dobbins, A., Lubbers, L.S., Jackson, G.L., Kuehl, D.E., Hileman, S.M., 2004. Neuropeptide  
35  
36 479 Y gene expression in male sheep: influence of photoperiod and testosterone.  
37  
38 480 *Neuroendocrinology* 792, 82-89.
- 39  
40  
41 481 Gardner, D.S., Van Bon, B.W., Dandrea, J., Goddard, P.J., May, S.F., Wilson, V.,  
42  
43 482 Stephenson, T., Symonds, M.E., 2006. Effect of periconceptional undernutrition and  
44  
45 483 gender on hypothalamic-pituitary-adrenal axis function in young adult sheep. *J.*  
46  
47 484 *Endocrinol.* 190, 203-212.
- 48  
49  
50 485 Gluckman, P.D., Hanson, M.A., 2008. Developmental and epigenetic pathways to obesity: an  
51  
52 486 evolutionary-developmental perspective. *Int. J. Obes. (Lond.)* 32 Suppl. 7, S62-71.
- 53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 487 Granado, M., Fuente-Martín, E., García-Cáceres, C., Argente, J., Chowen, J.A., 2012. Leptin  
1  
2 488 in early life: a key factor for the development of the adult metabolic profile. *Obes.*  
3  
4 489 *Facts* 5, 138-150.  
6
- 7 490 Grayson, B.E., Kievit, P., Smith, M.S., Grove, K.L., 2010. Critical determinants of  
8  
9 491 hypothalamic appetitive neuropeptide development and expression: species  
10  
11 492 considerations. *Front. Neuroendocrinol.* 31, 16-31.  
13
- 14 493 Huizinga, C.T., Oudejans, C.B., Delemarre-van de Waal, H.A., 2001 Persistent changes in  
16  
17 494 somatostatin and neuropeptide Y mRNA levels but not in growth hormone-releasing  
18  
19 495 hormone mRNA levels in adult rats after intrauterine growth retardation. *J.*  
20  
21 496 *Endocrinol.* 168, 273-281.  
23
- 24 497 Kennedy, A., Gettys, T.W., Watson, P., Wallace, P., Ganaway, E., Pan, Q., Garvey, W.T.,  
25  
26 498 1997. The metabolic significance of leptin in humans: gender-based differences in  
27  
28 499 relationship to adiposity, insulin sensitivity, and energy expenditure. *J. Clin.*  
30  
31 500 *Endocrinol. Metab.* 82, 1293-1300.  
33
- 34 501 Long, N.M., Ford, S.P., Nathanielsz, P.W., 2011. Maternal obesity eliminates the neonatal  
35  
36 502 lamb plasma leptin peak. *J. Physiol.* 589.6, 1455-1462.  
37  
38
- 39 503 Marie, M., Findlay, P.A., Thomas, L., Adam, C.L., 2001. Daily patterns of plasma leptin in  
40  
41 504 sheep: effects of photoperiod and food intake. *J. Endocrinol.* 170, 277-286.  
42
- 43 505 MacRae, J.C., Bruce, L.A., Hovell, F.D.B., Hart, I.C., Inkster, J., Atkinson, T., 1991.  
45  
46 506 Influence of protein nutrition on the response of growing lambs to exogenous bovine  
47  
48 507 growth hormone. *J. Endocrinol.* 130, 53-61.  
49  
50
- 51 508 McFadin, E.L., Morrison, C.D., Buff, P.R., Whitley, N.C., Keisler, D.H., 2002. Leptin  
52  
53 509 concentrations in periparturient ewes and their subsequent offspring. *J. Anim. Sci.* 80,  
54  
55 510 738-743.  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- 511 Mercer, J.G., Moar, K.M., Findlay, P.A., Hoggard, N., Adam, C.L., 1998. Association of  
512 leptin receptor (OB-Rb), NPY and GLP-1 gene expression in the ovine and murine  
513 brainstem. *Reg. Pept.* 75-76, 271-278.
- 514 Mercer, J.G., Moar, K.M., Ross, A.W., Hoggard, N., Morgan, P.J., 2000. Photoperiod  
515 regulates arcuate nucleus POMC, AGRP, and leptin receptor mRNA in Siberian  
516 hamster hypothalamus. *Am. J. Physiol.* 278, R271-281.
- 517 Mühlhäusler, B.S., Adam, C.L., Findlay, P.A., Duffield, J.A., McMillen, I.C., 2006.  
518 Increased maternal nutrition alters development of the appetite-regulating network in  
519 the brain. *FASEB J.* 20, 1257-1259.
- 520 Mühlhäusler, B.S., Adam, C.L., Marrocco, E.M., Findlay, P.A., Roberts, C.T., McFarlane,  
521 J.R., Kauter, K.G., McMillen, I.C., 2005. Impact of glucose infusion on the structural  
522 and functional characteristics of adipose tissue and on hypothalamic gene expression  
523 for appetite regulatory neuropeptides in the sheep fetus during late gestation. *J.*  
524 *Physiol.* 565.1, 185-195.
- 525 Mühlhäusler, B.S., McMillen, I.C., Rouzaud, G., Findlay, P.A., Marrocco, E.M., Rhind,  
526 S.M., Adam, C.L., 2004. Appetite regulatory neuropeptides are expressed in the sheep  
527 hypothalamus before birth. *J. Neuroendocrinol.* 16, 502-507.
- 528 Mühlhäusler, B.S., Roberts, C.T., McFarlane, J.R., Kauter, K.G., McMillen, I.C., 2002. Fetal  
529 leptin is a signal of fat mass independent of maternal nutrition in ewes fed at or above  
530 maintenance energy requirements. *Biol. Reprod.* 67, 493-499.
- 531 Nohara, K., Zhang, Y., Waraich, R.S., Laque, A., Tiano, J.P., Tong, J., Münzberg, H.,  
532 Mauvais-Jarvis, F., 2011. Early-life exposure to testosterone programs the  
533 hypothalamic melanocortin system. *Endocrinology* 152, 1661-1669.
- 534 Quirke, L.D., Juengel, J.L., Tisdall, D.J., Lun, S., Heath, D.A., McNatty, K.P., 2001.  
535 Ontogeny of steroidogenesis in the fetal sheep gonad. *Biol. Reprod.* 65, 216-228.

- 536 Santollo, J., Yao, D., Neal-Perry, G., Etgen, A.M., 2012. Middle-aged female rats retain  
1  
2 537 sensitivity to the anorexigenic effect of exogenous estradiol. *Behav. Brain Res.* 232,  
3  
4 538 159-164.  
5  
6  
7 539 Schwartz, M.W., Woods, S.C., Porte, D. Jr., Seeley, R.J., Baskin, D.G., 2000. Central  
8  
9 540 nervous system control of food intake. *Nature* 404, 661-671.  
10  
11 541 Sheppard, K.M., Padmanabhan, V., Coolen, L.M., Lehman, M.N., 2011. Prenatal  
12  
13 542 programming by testosterone of hypothalamic metabolic control neurones in the ewe.  
14  
15 543 *J. Neuroendocrinol.* 23, 401-411.  
16  
17 544 Toste, F.P., de Moura, E.G., Lisboa, P.C., Fagundes, A.T., de Oliveira, E., Passos, M.C.,  
18  
19 545 2006. Neonatal leptin treatment programmes leptin hypothalamic resistance and  
20  
21 546 intermediary metabolic parameters in adult rats. *Br. J. Nutr.* 95, 830-837.  
22  
23  
24 547 Wallace, J.M., Milne, J.S., Aitken, R.P., Adam, C.L., 2013. Impact of embryo donor  
25  
26 548 adiposity, birth weight and gender on early postnatal growth, glucose metabolism and  
27  
28 549 body composition in the young lamb. *Reprod. Fertil. Dev.* (in press).  
29  
30  
31 550 Wallace, J.M., Milne, J.S., Green, L.R., Aitken, R.P., 2011. Postnatal hypothalamic-pituitary-  
32  
33 551 adrenal function in sheep is influenced by age and sex, but not by prenatal growth  
34  
35 552 restriction. *Reprod. Fertil. Dev.* 23, 275-284.  
36  
37  
38 553 Watanobe, H., Suda, T., 1999. A detailed study on the role of sex steroid milieu in  
39  
40 554 determining plasma leptin concentrations in adult male and female rats. *Biochem.*  
41  
42 555 *Biophys. Res. Commun.* 259, 56-59.  
43  
44  
45 556 Yura, S., Itoh, H., Sagawa, N., Yamamoto, H., Masuzaki, H., Nakao, K., Kawamura, M.,  
46  
47 557 Takemura, M., Kakui, K., Ogawa, Y., Fujii, S., 2005. Role of premature leptin surge  
48  
49 558 in obesity resulting from intrauterine undernutrition. *Cell. Metab.* 1, 371-378.  
50  
51  
52  
53  
54  
55  
56  
57 559  
58  
59  
60  
61 560  
62  
63  
64  
65

561 **Fig. 1.** Example autoradiographic images of coronal hypothalamic sections hybridized to  
1  
2 562 radiolabelled riboprobes for POMC, CART, NPY, AGRP, OB-Rb and Ins-R in (A, C, E, G, I,  
3  
4 563 K) male and (B, D, F, H, J, L) female lambs. 3V = third ventricle. ARC = arcuate nucleus.  
5  
6  
7 564 Bar = 2 mm.  
8  
9

10 565 **Fig. 2.** Gene expression in the hypothalamic arcuate nucleus for POMC, CART, NPY,  
11  
12 566 AGRP, OB-Rb and Ins-R in (A) IUGR (dark grey bars; n = 17) versus normal birth weight  
13  
14 567 (N, pale grey bars; n = 16) lambs and (B) female (solid bars; n = 17) versus male (open bars;  
15  
16 568 n = 16) lambs. Results are shown as mean  $\pm$  SEM. \*P < 0.05; \*\*P < 0.01; \*\*\*P < 0.001.  
17  
18  
19  
20

21 569 **Fig. 3.** Relationships between plasma leptin concentration (final sample) and hypothalamic  
22  
23 570 arcuate nucleus gene expression for (A) NPY, (B) POMC and (C) OB-Rb in female (solid  
24  
25 571 symbols) and male (open symbols) lambs, irrespective of birth weight. Pearson correlation  
26  
27 572 analysis: males r = -0.776 (P < 0.001) and females r = -0.185 (not significant, NS) for A;  
28  
29 573 males r = 0.090 (NS) and females r = 0.494 (P < 0.05) for B; males r = -0.542 (P < 0.05) and  
30  
31 574 females r = 0.061 (NS) for C.  
32  
33  
34  
35  
36  
37 575  
38  
39  
40  
41 576  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



**Table 1.** Size at birth, fractional growth rates, final body weight, body fat depot weights and plasma leptin and insulin concentrations in lambs at 11 weeks of age.

	Females		Males		P value		
	N n = 8	IUGR n = 9	N n = 8	IUGR n = 8	N v. IUGR	Female v. male	Interaction
Birth weight (g)	4662 ± 137	3131 ± 221	5254 ± 198	3402 ± 167	<b>&lt;0.001</b>	<b>0.028</b>	0.399
Shoulder height at birth (cm)	22.8 ± 0.30	18.7 ± 0.76	23.9 ± 0.95	19.5 ± 1.04	<b>&lt;0.001</b>	0.253	0.862
FGR bodyweight (%/day)	8.42 ± 0.235	11.31 ± 0.699	8.53 ± 0.378	10.83 ± 0.220	<b>&lt;0.001</b>	0.688	0.518
FGR shoulder height (%/day)	1.20 ± 0.022	1.61 ± 0.080	1.22 ± 0.061	1.53 ± 0.076	<b>&lt;0.001</b>	0.647	0.489
FGR thorax girth (%/day)	1.37 ± 0.018	1.60 ± 0.052	1.25 ± 0.043	1.41 ± 0.040	<b>&lt;0.001</b>	<b>0.001</b>	0.423
FGR umbilical girth (%/day)	1.75 ± 0.072	1.82 ± 0.044	1.60 ± 0.054	1.76 ± 0.058	<b>0.043</b>	0.077	0.432
Final bodyweight (kg)	33.7 ± 0.53	29.6 ± 1.25	39.4 ± 0.628	32.8 ± 1.49	<b>&lt;0.001</b>	<b>&lt;0.001</b>	0.254
Perirenal fat (g)	462 ± 50.7	444 ± 48.9	338 ± 28.5	296.4 ± 48.5	0.514	<b>0.006</b>	0.797
Internal fat (g/kg) <sup>1</sup>	47.4 ± 2.24	50.3 ± 3.53	34.5 ± 3.25	35.4 ± 2.47	0.733	<b>&lt;0.001</b>	0.528
Leptin (ng/ml during final week) <sup>2</sup>	3.90 ± 0.640	4.03 ± 0.779	1.58 ± 0.551	1.76 ± 0.578	0.817	<b>0.002</b>	0.972
Leptin (ng/ml, final day) <sup>3</sup>	4.04 ± 0.484	2.93 ± 0.429	1.48 ± 0.422	1.68 ± 0.514	0.336	<b>&lt;0.001</b>	0.166
Insulin (ng/ml during final week) <sup>2</sup>	2.52 ± 0.209	2.78 ± 0.417	3.71 ± 0.609	3.70 ± 0.417	0.782	<b>0.022</b>	0.766

Values are means ± SEM. N, normal birth weight. IUGR, intra-uterine growth-restricted. FGR, fractional growth rate. <sup>1</sup>Combined omental and mesenteric fat depots, g/kg empty bodyweight. <sup>2</sup>Average of three samples taken on days 65-73. <sup>3</sup>Sample taken on day 77. Data from Wallace et al. (2013).

**Table 2.** Relationships between plasma leptin and insulin concentrations and body fat depot weights in male and female lambs at 11 weeks of age, irrespective of birth weight.

	Females		Males	
	n = 16		n = 17	
	Perirenal fat (g)	Internal fat (g/kg) <sup>3</sup>	Perirenal fat (g)	Internal fat (g/kg) <sup>3</sup>
Leptin (ng/ml during final week) <sup>1</sup>	<b>0.778***</b>	<b>0.723***</b>	<b>0.833***</b>	<b>0.783***</b>
Leptin (ng/ml, final day) <sup>2</sup>	<b>0.559*</b>	0.288	<b>0.704**</b>	<b>0.830***</b>
Insulin (ng/ml during final week) <sup>1</sup>	0.011	0.005	<b>0.658**</b>	0.197

Values are Pearson correlation coefficients. Significant correlations shown in bold. \*P<0.05; \*\*P<0.01; \*\*\*P<0.001. <sup>1</sup>Average of three samples taken on days 65-73. <sup>2</sup>Sample taken on day 77. <sup>3</sup>Combined omental and mesenteric fat depots (g/kg empty bodyweight). Data derived from Wallace et al. (2013).

**Table 3.** Lamb body weight changes and suckling activity during assessment at 23 days of age.

	Female		Male		<i>P</i> -value		
	N	IUGR	N	IUGR	N v. IUGR	Female v. male	Interaction
	n = 8	n = 9	n = 8	n = 8			
Weight loss during 3h fast (g)	171 ± 16.9	199 ± 22.8	147 ± 14.7	171 ± 13.7	0.143	0.143	0.933
Weight loss during fasting (%)	1.3 ± 0.11	1.4 ± 0.14	1.3 ± 0.12	1.6 ± 0.08	0.161	0.341	0.892
Suckling episodes per 60 min (n)	5.9 ± 0.83	6.5 ± 1.33	4.9 ± 1.13	5.4 ± 0.96	0.341	0.614	0.950
Suckling duration per 60min (s)	120 ± 21.5	154 ± 35.9	130 ± 26.0	129 ± 15.4	0.758	0.522	0.509
Weight gain to 60 min after fast (g)	298 ± 53.1	388 ± 39.9	217 ± 60.6	289 ± 24.7	0.069	0.103	0.866
Weight gain to 90 min after fast (g)	291 ± 46.0	349 ± 44.1	260 ± 45.4	278 ± 14.5	0.214	0.360	0.524
Weight gain to 90 min after fast (%)	2.7 ± 0.19	2.4 ± 0.23	2.5 ± 0.41	2.6 ± 0.13	0.469	0.654	0.902

Values are means ± SEM. N, normal birth weight. IUGR, intra-uterine growth-restricted.

**Table 4.** Relationships between arcuate nucleus expression of energy balance regulatory genes, body fat, leptinemia, insulinemia, weight gain during a suckling activity assessment, and final current fractional growth rates for body weight and girth in lambs at 11 weeks of age.

	Gene expression (arbitrary densitometry units)					
	POMC	CART	NPY	AGRP	OBRb	Ins-R
<b>Females</b>						
OB-Rb gene expression	-0.234	-0.420	0.346	0.350	-	-
Ins-R gene expression	0.214	-0.057	0.009	0.069	-	-
Perirenal fat (g)	0.405	0.171	-0.362	<b>-0.520*</b>	-0.110	0.331
Internal fat (g/kg) <sup>1</sup>	0.159	0.205	-0.313	-0.456	0.009	0.088
Leptin (ng/ml, final day) <sup>2</sup>	<b>0.494*</b>	0.128	-0.185	-0.364	0.061	0.317
Leptin (ng/ml, during last week) <sup>3</sup>	0.303	0.204	<b>-0.510*</b>	-0.440	-0.162	0.046
Insulin (ng/ml, during last week) <sup>3</sup>	-0.386	0.268	0.040	-0.001	-0.324	-0.396
Suckling % weight gain to 90min <sup>4</sup>	-0.232	-0.065	-0.263	0.263	0.061	-0.025
CFGR body weight (%/day) <sup>5</sup>	<b>-0.490*</b>	-0.123	-0.195	0.167	0.369	0.015
CFGR thorax girth (%/day) <sup>5</sup>	-0.290	-0.019	0.231	0.212	0.392	0.122
CFGR umbilical girth (%/day) <sup>5</sup>	<b>-0.547*</b>	-0.340	0.324	0.237	-0.002	-0.389
<b>Males</b>						
OB-Rb gene expression	-0.147	-0.242	<b>0.535*</b>	<b>0.507*</b>	-	-
Ins-R gene expression	-0.261	0.164	0.315	0.146	-	-
Perirenal fat (g)	0.257	-0.045	<b>-0.784***</b>	<b>-0.683**</b>	<b>-0.529*</b>	-0.324
Internal fat (g/kg) <sup>1</sup>	0.245	-0.295	<b>-0.751***</b>	<b>-0.596*</b>	-0.281	-0.481
Leptin (ng/ml, final day) <sup>2</sup>	0.090	-0.287	<b>-0.776***</b>	<b>-0.670**</b>	<b>-0.542*</b>	-0.383
Leptin (ng/ml, during last week) <sup>3</sup>	0.237	-0.180	<b>-0.798***</b>	<b>-0.632**</b>	-0.472	-0.344
Insulin (ng/ml, during last week) <sup>3</sup>	-0.128	0.127	<b>-0.501*</b>	-0.433	<b>-0.643**</b>	0.130
Suckling % weight gain to 90min <sup>4</sup>	0.195	-0.094	<b>0.623**</b>	0.264	0.329	0.114
CFGR body weight (%/day) <sup>5</sup>	-0.129	-0.277	-0.248	-0.287	0.216	0.423
CFGR thorax girth (%/day) <sup>5</sup>	-0.022	-0.236	<b>-0.564*</b>	-0.466	-0.089	0.199
CFGR umbilical girth (%/day) <sup>5</sup>	-0.013	-0.191	-0.393	-0.302	-0.106	0.385

Values are Pearson correlation coefficients. Significant correlations shown in bold. \*P < 0.05; \*\*P < 0.01; \*\*\*P < 0.001. <sup>1</sup>Combined omental and mesenteric fat depots (g/kg empty bodyweight); <sup>2</sup>Sample taken on day 77; <sup>3</sup>Average of three samples taken on days 65-73; <sup>4</sup>Suckling assessment at 23 days; <sup>5</sup>CFGR (current fractional growth rate) measurements at 68 days of age.

**Figure(s)**










Fig. 2

A



B






Fig. 3

A



B



C

