



# Microbial communities in sediment from *Zostera marina* patches, but not the *Z. marina* leaf or root microbiomes, vary in relation to distance from patch edge

Cassandra L. Ettinger<sup>1</sup>, Sofie E. Voerman<sup>2,3</sup>, Jenna M. Lang<sup>1,4</sup>,  
John J. Stachowicz<sup>5</sup> and Jonathan A. Eisen<sup>1,5,6</sup>

<sup>1</sup> Genome Center, University of California, Davis, CA, United States

<sup>2</sup> Climate Change Cluster, University of Technology Sydney, Sydney, Australia

<sup>3</sup> School of Life Sciences, University of Technology Sydney, Sydney, Australia

<sup>4</sup> Trace Genomics, San Francisco, CA, United States

<sup>5</sup> Department of Evolution and Ecology, University of California, Davis, CA, United States

<sup>6</sup> Department of Medical Microbiology and Immunology, University of California, Davis, Davis, CA, United States

## ABSTRACT

**Background.** *Zostera marina* (also known as eelgrass) is a foundation species in coastal and marine ecosystems worldwide and is a model for studies of seagrasses (a paraphyletic group in the order *Alismatales*) that include all the known fully submerged marine angiosperms. In recent years, there has been a growing appreciation of the potential importance of the microbial communities (i.e., microbiomes) associated with various plant species. Here we report a study of variation in *Z. marina* microbiomes from a field site in Bodega Bay, CA.

**Methods.** We characterized and then compared the microbial communities of root, leaf and sediment samples (using 16S ribosomal RNA gene PCR and sequencing) and associated environmental parameters from the inside, edge and outside of a single subtidal *Z. marina* patch. Multiple comparative approaches were used to examine associations between microbiome features (e.g., diversity, taxonomic composition) and environmental parameters and to compare sample types and sites.

**Results.** Microbial communities differed significantly between sample types (root, leaf and sediment) and in sediments from different sites (inside, edge, outside). Carbon:Nitrogen ratio and eelgrass density were both significantly correlated to sediment community composition. Enrichment of certain taxonomic groups in each sample type was detected and analyzed in regard to possible functional implications (especially regarding sulfur metabolism).

**Discussion.** Our results are mostly consistent with prior work on seagrass associated microbiomes with a few differences and additional findings. From a functional point of view, the most significant finding is that many of the taxa that differ significantly between sample types and sites are closely related to ones commonly associated with various aspects of sulfur and nitrogen metabolism. Though not a traditional model organism, we believe that *Z. marina* can become a model for studies of marine plant-microbiome interactions.

Submitted 4 January 2017

Accepted 29 March 2017

Published 27 April 2017

Corresponding author

Jonathan A. Eisen,  
jaeisen@ucdavis.edu,  
jonathan.eisen@gmail.com

Academic editor

Gerrit Beemster

Additional Information and  
Declarations can be found on  
page 16

DOI 10.7717/peerj.3246

 Copyright  
2017 Ettinger et al.

Distributed under  
Creative Commons CC-BY 4.0

OPEN ACCESS

**Subjects** Marine Biology, Microbiology, Plant Science

**Keywords** Seagrass, Eelgrass, Sulfur metabolism, Microbiome, Edge effect, Fragmentation, Rhizosphere

## INTRODUCTION

The seagrass, *Zostera marina*, is a foundation species in protected bays and estuaries throughout the temperate northern hemisphere. Seagrasses are fully submerged marine angiosperms and are a paraphyletic group comprised of three lineages in the order *Alismatales* that convergently adapted to the marine environment between 70 and 100 million years ago (Les, Cleland & Waycott, 1997). There are only approximately 60 species of seagrass compared to the 250,000 species of terrestrial angiosperms, a testament to the strict selective pressure posed by re-entry to the marine environment (Orth et al., 2006). Seagrass patches serve as habitat and nursery grounds for many marine species, play key roles in nutrient cycling and carbon sequestration, and serve to protect the coastline from erosion (Williams & Heck Jr, 2001). *Z. marina* populations, like those of many seagrass species, are negatively affected by climate change, pollution and habitat destruction and so far, restoration efforts have been costly and ineffective (Orth et al., 2006). As a result, *Z. marina* is vulnerable to habitat fragmentation and loss.

The work described here was originally focused on a phenomenon known as “edge effects” in which the border between habitats is intermediate in abiotic conditions from the center of either adjacent habitat and thus the biotic composition of the border habitat, or edge, may differ from that of interior, intact habitat. Edges often support a mixture of organisms from two adjacent habitats (Fox et al., 1997; Davies-Colley, Payne & Elswijk, 2000), but may be abiotically unsuitable for species found in the center of either habitat. Increased predation and invasion by non-native species can also be a common feature of edges (Paton, 1994; Fox et al., 1997; Harrison & Bruna, 1999; Flaspohler, Temple & Rosenfield, 2001). Prior work on seagrasses have shown edge effects on species abundances (Smith et al., 2008; Smith et al., 2011; Tanner, 2005) and abiotic conditions such as turbulence (Granata et al., 2001; Folkard, 2005), carbon stocks (Ricart et al., 2015) and organic matter deposition (Duarte & Sand-Jensen, 1990; Cebrián et al., 2000). Here we investigate whether such edge effects are evident in the microbiota found in, on and near *Z. marina* plants.

Our interest in the microbiota for this study is driven by our overarching goal of developing *Z. marina* as a model for studies of microbial communities associated with marine plants. Although we speculate that plant microbe interactions are important for seagrasses, little is known about how the roles of microbial communities associated with marine plants may affect plant health and what abiotic and biotic factors affect the composition of these communities. Terrestrial plants, like *Arabidopsis* (Lundberg et al., 2012), corn (Peiffer et al., 2013; Bouffaud et al., 2014), rice (Peiffer et al., 2013; Edwards et al., 2015) and poplar (Beckers et al., 2016) have been shown to have distinct microbial communities on the inside (endophytes) and on the surface (epiphytes) of plant leaves and roots, as well as in the surrounding soil or sediment (rhizosphere) (Lundberg et al., 2012).

These communities can vary across different stages of plant development (*Chaparro, Badri & Vivanco, 2014*) and with local environmental conditions. In terrestrial systems the main drivers of plant associated microbial community composition are considered to be environmental factors, like soil particle size, pH and moisture content, as well as host plant species (*Aleklett et al., 2015; Lakshmanan, 2015*). Thus, examining eelgrass microbiota across a known environmental gradient from the center to the outside of a patch has the potential to provide insights into factors that shape the eelgrass microbiome, the full community of microorganisms associated with eelgrass. Recently a few culture-independent surveys of seagrass microbiomes have been published and these provide good initial reference points for our work here (*Jiang et al., 2015; Sun et al., 2015; Cúcio et al., 2016; Mejia et al., 2016*). Although, these studies have similar big picture findings, there are small differences in microbiome composition between them and thus further teasing apart of the factors that shape seagrass microbiomes is necessary and important work.

In this study, we characterized and then compared the epiphytic and rhizospheric bacterial communities of eelgrass using root, leaf and sediment samples obtained from the inside, edge and outside of a single subtidal *Z. marina* patch. We focused on characterizing the bacterial and archaeal members of the microbiome in each of these samples using high throughput sequencing of 16S ribosomal gene PCR libraries. We focus in particular on the following questions: What is the general taxonomic composition of the *Z. marina* microbiome? Are there changes in sediment microbial community composition or in biodiversity at the patch edge and, if so, what factors are driving observed differences, environmental abiotic factors or presence/absence of *Z. marina*? This analysis reveals multiple novel insights into the general structure of the *Z. marina* microbiome and lays the groundwork for further studies.

## MATERIALS & METHODS

### Sample collection

We collected leaf, root and sediment samples for microbiome analysis from 0.25 m<sup>2</sup> quadrats ( $n = 4$ ) located in the interior (2.5 m from the edge), on the edge (but within the eelgrass habitat) and outside (2.5 m from the edge) of a single shallow subtidal eelgrass patch in Bodega Bay, CA (GPS: 38.319435, -123.053838) during the summer of 2013. Quadrats were positioned 2.5 m from each other parallel to the patch' edge. Samples were collected during low tide ( $\pm 0.5$  m water depth) at night (11 PM). For quadrats located at the center or edge of the eelgrass patch, one eelgrass shoot was sampled and directly separated into root and shoot tissue. The root tissue consisted of one entire root bundle sampled, the leaf tissue consisted of a clipped leaf of  $\pm 3$  cm in length positioned at about half way along the shoot length ( $\pm 20$  cm from the base). For each quadrat, sediment samples were collected at two sediment depths, 0.5 cm or less and 3 cm deep, from randomly selected locations within the quadrats. Microbial samples were directly stored on ice and transported to the laboratory within one hour where samples were frozen at  $-20$  °C until further analysis.

Environmental data and the samples used for microbiome analysis were collected simultaneously. For each quadrat, eelgrass density was estimated by direct count.

Temperature, pH, salinity and dissolved oxygen were measured at 20 cm above the sediment with a YSI 556 handheld multimeter (YSI Inc., Yellow Springs, OH, USA), at a similar height as the shoot tissue was sampled. Sediment chemical and physical properties were assessed by separately coring the top 4 cm of sediment (10 cm diameter, taken twice within a quadrat and combined for analysis), to correspond with the sediment layer most influenced by the eelgrass roots. Sediment was dried (three days at 40 °C), mixed, sieved (sieve sizes: 710, 500, 355, 250, 180, 90 and 30  $\mu\text{m}$ ) and particle size fractions were weighed to investigate particle size distribution. A portion of the mixed sediment samples ( $\pm 50$  g) was separately analyzed for total organic carbon (TOC), total inorganic carbon (TIC) and Carbon:Nitrogen (C:N) ratio by the UC Davis Analytical Laboratory.

### Molecular methods

DNA was extracted from leaf ( $n = 8$ ), root ( $n = 8$ ) and both shallow ( $n = 12$ ) and deep sediment ( $n = 12$ ) samples as well as from a kit control ( $n = 1$ ) with the PowerSoil DNA Isolation kit (MO BIO Laboratories, Inc., Carlsbad, CA, USA) according to the manufacturer's protocol. For the DNA extractions, root and leaf tissues were placed directly into PowerBead tubes from the freezer without grinding. Microbial 16S rRNA genes were amplified using a two-step protocol targeting the V4 region using the "universal" 515F and 806R primers (Caporaso *et al.*, 2012). The primer set was modified to include Illumina adapters and barcode sequences using a dual indexing approach as in Lang, Eisen & Zivkovic (2014). The 41 samples described in this paper were multiplexed with 103 samples from other experiments, for a total of 144 samples on the run. Libraries were sequenced by the UC Davis Genome Center Core Facilities on an Illumina MiSeq (Illumina, Inc., San Diego, CA, USA) to generate 250 bp paired-end reads.

### Sequence processing

A custom in-house script was used to demultiplex, quality check and merge paired reads ([https://github.com/gjospin/scripts/blob/master/Demul\\_trim\\_prep.pl](https://github.com/gjospin/scripts/blob/master/Demul_trim_prep.pl)). The resulting sequences were analyzed using the Quantitative Insights Into Microbial Ecology (QIIME) v. 1.9.0 workflow (Caporaso *et al.*, 2010).

For a detailed walkthrough of the following analysis using QIIME, see the IPython notebook (<http://nbviewer.jupyter.org/gist/casett/86da7fc8749d27574f183498df65134a>).

The sequencing run for this project included samples from other projects. In total, for the entire run, 14,163,470 reads passed quality filtering (Q20). Of these reads, 4,573,318 were associated with the 41 samples for this project. Of the 4,573,318 reads for this project, 4,212,549 merged successfully (92.11%). The sample with the lowest number of sequences after merging was the negative control with 444 sequences, the next lowest sample, BB039, had 22,897 sequences, approximately a fifty-fold increase. The most abundant sequence in the negative control was chloroplast DNA, and thus, we conclude that these 444 sequences were likely the result of contamination from other samples during sequencing or molecular analyses. We considered removing shared operational taxonomic units (OTUs) or 100 percent identical DNA sequences between the negative control and our environmental samples, but determined both of these actions to be too stringent on the dataset when

taking into account the abundance of the OTU's/DNA sequences in the samples and the number of reads in the negative control. Instead the negative control was simply removed from downstream analysis.

A total of 4,976 chimeras were identified using USEARCH v. 6.1 and were filtered out. The remaining sequences were clustered using the open reference approach into OTUs at 97 percent similarity using UCLUST ([Edgar, 2010](#)). Taxonomy was assigned using the `assign_taxonomy.py` QIIME script with the GreenGenes database (v.13\_8) ([DeSantis et al., 2006](#)) using UCLUST. Further filtering was performed using the QIIME scripts, `filter_taxa_from_otu_table.py` and `filter_otus_from_otu_table.py`, to remove chloroplast DNA, mitochondrial DNA and singletons. Reads classified as "Unassigned" at the domain level were also removed from downstream analysis. After these filtering steps, the lowest number of sequences in a sample dropped to 3,277. This reduction in the number of sequence reads can be largely attributed to the removal of *Z. marina* chloroplast DNA from the leaf and root samples.

To aid in statistical comparison between different sample types (leaf, root, sediment), we subset our 16S rDNA sequences to a minimum sequence count of 3,277 to retain the maximum number of samples. However, when comparing only sediment samples, the 16S rDNA sequences were randomly subset to 20,000 sequences using the `single_rarefaction.py` QIIME script.

### Data visualization and statistical analyses

Data visualization was performed exclusively in R and statistical analyses were performed using a combination of QIIME scripts and R ([R Core Team, 2016](#)). For analysis done in R, the rarefied OTU tables were converted to json format and exported for analysis using the `ggplot2` ([Wickham, 2009](#)), `vegan` ([Dixon, 2003](#)) and `phyloseq` ([McMurdie & Holmes, 2013](#)) packages. Initial analysis indicated no significant differences between the microbial communities associated with shallow (0.5 cm or less) and deep (3 cm) sediment samples, thus sediment depth was not considered further here. We describe the different types of analyses below.

- Intra-sample (alpha) diversity. We were interested in if significant differences existed between the intra-sample (alpha) diversities (richness, evenness) of the microbial communities associated with different sample types (leaf, root, sediment) and different sediment locations (inside, edge, outside). We calculated the following diversity metrics: Chao1 ([Chao, 1984](#)), Observed OTUs, Shannon ([Shannon & Weaver, 1949](#)) and Simpson Indices ([Simpson, 1949](#)) in R. To determine if there were significant differences between the alpha diversities of different sample types and different sediment locations, we first performed Kruskal–Wallis tests. We then implemented Bonferroni corrected post-hoc Dunn tests to identify which pairwise comparisons were driving differences.
- Inter-sample (beta) diversity. We assessed the inter-sample (beta) diversities of the microbial communities associated with different sample groupings (sample type, location, etc) and if there were any significant correlations between environmental variables and community dissimilarity. We used both Unifrac (weighted and unweighted) ([Lozupone et al., 2007](#); [Hamady, Lozupone & Knight, 2010](#)) and Bray–Curtis ([Bray](#)


(Curtis, 1957) dissimilarities calculated in R using phyloseq. These dissimilarities were then plotted using principal coordinate analysis (PCoA) and non-metric multidimensional scaling (NMDS) methods. Multiple tests were then performed on these beta-diversity results. To test for significant differences in centroids between different sample groupings (sample type, location, etc.) PERMANOVA tests were performed using the adonis function from the vegan package in R with 9,999 permutations (Anderson, 2001). PERMANOVA tests can be sensitive to differences in dispersion when using abundance-based distance matrices (Warton, Wright & Wang, 2012), but are more robust than other tests, especially for balanced designs (Anderson & Walsh, 2013). To test for differences in mean dispersions between different groupings, the betadisper and permutest functions from the vegan package in R were used with 999 permutations. To test for correlations between the Bray Curtis dissimilarities of our samples and the environmental factors (C:N ratio, pH, etc) measured, euclidean distances were calculated in R using vegan and Mantel tests were performed using 9,999 permutations. The supervised\_learning.py QIIME script was used to see if a random forest classifier could differentiate between sample type or sediment location using leave-one-out cross validation and 1,000 trees.

- Taxonomic variation. To determine if the mean relative abundance of taxonomic orders varied significantly between different sample types and sediment locations, we first used the summarize\_taxa.py QIIME script to remove rare OTUs (less than one percent of total abundance) and to collapse OTUs at the Order level. We then used the group\_significance.py QIIME script on the resulting OTU table to test for differences using Bonferroni corrected Kruskal–Wallis tests with 1000 permutations. We removed the rare OTUs, as suggested in the documentation for the groups\_significance.py QIIME script, to avoid spurious significance from very low abundance OTUs, to simplify analyses and to focus on abundant organisms and overall patterns.
- Environmental variation. To determine if environmental factors varied significantly between different locations in the eelgrass patch (inside, edge, outside), ANOVA tests were performed in R for each factor. The post-hoc Tukey’s Honest Significant Difference (HSD) test was performed in R for factors found significantly different by the ANOVA (Tukey, 1953; Kramer, 1956; Kramer, 1957).

## RESULTS & DISCUSSION

### Diversity metrics I: intra-sample variation between sample types and locations

Alpha diversity is greater in the sediment than in the leaves and roots ( $p < 0.001$ ) for a variety of metrics including observed number of OTUs, Chao1, Shannon and Simpson (Fig. 1). However, there is no difference in alpha diversity between leaf and root samples ( $p > 0.05$ ) (Table S1). This is not altogether unexpected as in terrestrial systems soil has been observed to have increased diversity compared to host associated sample types (Edwards et al., 2015). There is conflict between the diversity metrics when determining if the intra-sample diversity of sediment at different locations (inside, edge, outside) varies




**Figure 1** Alpha diversity across samples. Four alpha diversity metrics, observed number of OTUs, Chao1, the Shannon and Simpson diversity indices, are shown as boxplots for (A) different sample types (leaf, root, sediment) and for (B) sediment from different locations (inside, edge, outside).

(Table S2). Two of the metrics, observed number of OTUs and Chao1, indicate greater diversity outside compared to inside the patch ( $p < 0.05$ ). The non-significant metrics, the Shannon and Simpson indices, account for both richness and evenness and are less sensitive to rare taxa than richness only metrics (Bent & Forney, 2008). Thus, one possible explanation for the difference in diversity between the inside and outside sediment is an increased number of rare taxa in sediment from outside the patch. No significant differences were found between the alpha diversity of leaves and roots between the inside and edge of the eelgrass patch.

### Diversity metrics II: inter-sample variation between sample types and locations

Distinct microbial communities were detected in association with *Z. marina* leaves, roots and sediment (Fig. 2). PERMANOVA tests performed on three different beta diversity metrics, weighted UniFrac, unweighted UniFrac and Bray–Curtis Dissimilarity, found these communities to be significantly different from each other with  $p = 0.0001$  (Table 1). Root and leaf associated communities were found to have more within group variance, or dispersion, than sediment communities ( $p = 0.001$ ), which could indicate that stabilizing




**Figure 2** Principal coordinates analysis (PCoA) of microbial communities based on weighted UniFrac distances. Samples are colored by sample type (leaf, root, sediment) with different shapes for location (inside, edge, outside).

**Table 1** Permanova results. Comparing microbial community composition between different sample types (leaf, root, sediment) and locations (inside, edge, outside) using multiple beta diversity metrics.

| Category    | Diversity metric   | Pseudo-F | R2     | P (perm) |
|-------------|--------------------|----------|--------|----------|
| Location    | Weighted UniFrac   | 2.22     | 0.107  | 0.0213   |
|             | Unweighted UniFrac | 1.91     | 0.0938 | 0.0043   |
|             | Bray Curtis        | 2.82     | 0.133  | 0.0009   |
| Sample type | Weighted UniFrac   | 13.75    | 0.426  | 0.0001   |
|             | Unweighted UniFrac | 6.16     | 0.249  | 0.0001   |
|             | Bray Curtis        | 9.53     | 0.34   | 0.0001   |
| LocXType    | Weighted UniFrac   | 1.98     | 0.0541 | 0.0426   |
|             | Unweighted UniFrac | 1.19     | 0.0455 | 0.1586   |
|             | Bray Curtis        | 1.482    | 0.0458 | 0.0795   |

selection is acting on these sediment communities. Random forest analysis further validated the observed differences between leaves, roots and sediment microbial communities (Table S3). The classifier had an estimated error of 5% (versus a baseline error of 40%) and correctly identified all leaf samples ( $n = 8$ ) and all sediment samples ( $n = 24$ ). The classifier did misclassify two of the root samples ( $n = 8$ ) as leaves, but this is not unexpected as these two samples also appear to cluster more closely with the leaf samples when visualized using Principal Coordinates Analysis (PCoA) (Fig. 2). The reason that these root samples cluster more closely with the leaf samples may be due to which root bundles were sampled; preliminary results indicate that the microbiota associated with the roots can vary depending on the proximity of the root to the base of the leaf, with roots closer to the base appearing more “leaf-like” (HE Holland-Moritz et al., 2017, unpublished data).






**Figure 3** Principal coordinates analysis (PCoA) of microbial communities in sediment based on weighted Unifrac distances. Samples are colored by location (inside, edge, outside).

**Table 2** Sediment PERMANOVA results. Comparing sediment microbial community composition between different locations (inside, edge, outside) and eelgrass densities using multiple beta diversity metrics.

| Category | Diversity metric   | Pseudo-F | R <sup>2</sup> | P (perm) |
|----------|--------------------|----------|----------------|----------|
| Location | Weighted UniFrac   | 8.69     | 0.453          | 0.0001   |
|          | Unweighted UniFrac | 2.92     | 0.217          | 0.0001   |
|          | Bray Curtis        | 8.01     | 0.433          | 0.0001   |
| Density  | Weighted UniFrac   | 2.81     | 0.551          | 0.0002   |
|          | Unweighted UniFrac | 1.51     | 0.398          | 0.0001   |
|          | Bray Curtis        | 2.86     | 0.555          | 0.0001   |

To determine if there was a difference in community composition at the eelgrass patch edge relative to the inside or outside of the patch, beta diversity metrics were calculated for the sediment microbial communities. As can be seen in Fig. 3, these diversity metrics show the communities clustering by sampling location (inside, edge, outside). PERMANOVA tests indicate that these clusters are significantly different between locations ( $p = 0.0001$ ) and also for eelgrass shoot densities ( $p < 0.0002$ ) (Table 2). However, leaf and root microbial communities do not differ significantly based on sampling location, possibly indicating that these plant tissue associated communities are more stable than the sediment communities in regards to location. Whereas sediment communities, although distinct when associated with eelgrass, may be under less selection from the host plant. One possible explanation for the correlation between the sediment communities and eelgrass shoot density may be the release of exudates and oxygen by the roots of the eelgrass, which would increase in concentration with eelgrass density.



**Figure 4** Average relative abundance of taxonomic groups associated with each sample type (leaf, root, sediment). OTUs are shown grouped by taxonomic order and colored by taxonomic class. Only orders with a mean abundance of at least one percent are shown here. The bars represent the standard error of the mean.

Random forest analysis confirmed differences between sediment microbial communities taken from the inside of the patch, the edge and unvegetated sediment from outside the patch (Table S4). The classifier had an estimated error of 12.5% (versus a baseline error of 66.7%) and correctly identified all of the unvegetated sediment ( $n = 8$ ). The classifier did mistakenly classify one sample from the edge ( $n = 8$ ) as being from the inside of the patch and two samples from the inside of the patch ( $n = 8$ ) as being from the edge. In Fig. 3, there is some overlap in the clustering of sediment from the inside and edges of patches which might account for these misclassifications.

### Major patterns in community composition of the leaves, roots and rhizosphere sediment

The analysis of diversity metrics presented above shows that there are distinct communities associated with leaves and roots, and these both differ from the sediment, whereas location effects are weaker. We therefore examined in more detail the taxonomic composition and possible functional roles of the microbes on *Z. marina* leaves, roots and rhizosphere sediment (sediment from the inside and edge of the eelgrass patch). We summarize our findings regarding this here.

Figure 4 shows the average relative abundance of different orders of bacteria for leaves, roots and sediment. On leaves, the most abundant orders were *Clostridiales*, *Bacteroidales*, *Rhodobacterales*, *Flavobacteriales*, *Saprospirales*, *Thiotrichales* and Unidentified *Gammaproteobacteria*. On roots, the most abundant orders were *Campylobacterales*, *Bacteroidales*, *Clostridiales*, *Desulfobacterales*, *Flavobacteriales* and *Desulfuromonadales*. In the rhizosphere sediment, the most abundant orders were *Bacteroidales*, *Flavobacteriales*, *Desulfobacterales*, *Thiotrichales*, *Clostridiales* and *Alteromonadales*.

We also examined the overall patterns in our results at the class level (Table S5). For leaves, the most abundant class of epiphytes observed was *Gammaproteobacteria* ( $20.5 \pm 7.3\%$ ). Other abundant classes included *Clostridia* ( $16.5 \pm 12\%$ ), *Bacteroidia* ( $12.6 \pm 8.6\%$ ), *Alphaproteobacteria* ( $11.4 \pm 8.5\%$ ), *Flavobacteria* ( $7.8 \pm 4.7\%$ ) and *Saprospirae* ( $7.7 \pm 4.2\%$ ). For roots, the dominant class associated with the roots were *Epsilonproteobacteria* ( $17.9 \pm 15.5\%$ ). Other abundant classes observed on the roots include *Deltaproteobacteria* ( $13.4 \pm 11.3\%$ ), *Bacteroidia* ( $12.8 \pm 8.8\%$ ), *Gammaproteobacteria* ( $12.6 \pm 11\%$ ), *Clostridia* ( $8 \pm 8.4\%$ ), *Flavobacteriia* ( $6.1 \pm 6.1\%$ ) and *Alphaproteobacteria* ( $4.8 \pm 6.3\%$ ). In the rhizosphere sediment, the dominant class was *Gammaproteobacteria* ( $18.2 \pm 3.4\%$ ), as it was on the leaves. Other abundant classes found in the rhizosphere sediment include *Deltaproteobacteria* ( $14.9 \pm 2.6\%$ ), *Bacteroidia* ( $13.3 \pm 2.3\%$ ), *Flavobacteriia* ( $9.3 \pm 3.4\%$ ), *Clostridia* ( $5.1 \pm 3.3\%$ ) and *Anaerolineae* ( $3.9 \pm 1.6\%$ ).

The summary results above allow a comparison to findings from a recent study on the rhizosphere sediment microbiomes of three seagrass species, including *Z. marina*, Cúcio et al., 2016. We chose to focus our comparison on the Cúcio et al. study because it is one of the more comprehensive culture independent studies of seagrasses. Overall, there are general similarities and differences when comparing the class-level patterns between the studies. The authors reported that the most abundant classes were *Gammaproteobacteria* (32–38% depending on the species sampled), *Deltaproteobacteria* (23–26%), and *Bacteroidia* (6–7%). These were the three most abundant classes in our sediment samples as well, but at different relative abundances (see above). These differences could be due to true differences in microbiomes in the sediments sampled, or due to the use of different primer sets, extraction methods, and sample collection strategies (among many other differences).

When examined at higher taxonomic ranks, the microbiome of the leaves of *Z. marina* shares some similarities with the microbiomes of various marine algae (e.g., kelp and seaweeds), with *Gammaproteobacteria* being the most abundant class in both cases (Hollants et al., 2013). However, these similarities are not seen at lower ranks (e.g., order, family, genus). This finding is similar to what has been observed between different marine algal microbiomes, with similarities observed at higher, but not lower taxonomic levels (Hollants et al., 2013; Egan et al., 2013). This is further supported by a recent study, which focused on surface-associated communities, that observed that the microbiomes of seagrass and seaweed species were host specific, but had broad-scale functional similarities (Roth-Schulze et al., 2016).

### **Differences in microbial communities between sample types (leaves, roots and rhizosphere sediment) and possible functional implications**

We used a Bonferroni corrected Kruskal–Wallis test to test for differences in relative abundance of the orders between sample types. This showed that *Saprospirales*, *Thiotrichales*, *Rhodobacterales*, *Desulfobacterales*, *Desulfuromonadales*, *Marinicellales*, *Spirochaetales*, *Chromatiales* and *Campylobacterales* are significantly different between sample types ( $p < 0.05$ ). *Campylobacterales*, *Desulfobacterales*, *Spirochaetales* and *Desulfuromonadales* were enriched on *Z. marina* roots. *Thiotrichales*, *Rhodobacterales* and *Saprospirales* were enriched on the leaves. *Thiotrichales*, *Marinicellales*, *Chromatiales*, *Desulfobacterales* and


*Spirochaetales* were enriched in the rhizosphere. We note that many of the taxa that differ significantly between communities are closely related to ones commonly associated with various aspects of sulfur and nitrogen metabolism. This is interesting because prior studies have suggested that nitrogen and sulfur metabolism are critical functions for the seagrass associated microbiome (Lovell, 2002). For example, acquisition of nitrogen (in its many forms) is frequently a limiting factor for the health of plants, including seagrasses (Short, 1987; Elser et al., 2007) and associations with microbes are frequently critical for such acquisition (Welsh, 2000; Nielsen et al., 2001). In addition, since the reduced sulfur compounds that accumulate in aquatic sediments are known phytotoxins (Lamers et al., 2013), it is thought that sulfur metabolizing microbes could play important roles in aiding seagrass survival in such sediments (Barber & Carlson, 1993; Terrados et al., 1999; Erskine & Koch, 2000; Van der Heide et al., 2012). Sulfur and nitrogen metabolism are not necessarily independent—it has been postulated that sulfate-reducing bacteria may be responsible for most of the nitrogen fixation that occurs in seagrass sediments (Capone, 1982).

Given this context, we discuss several of the specific taxa that differ between samples and their possible connection to nitrogen and/or sulfur metabolism below. For example, *Campylobacterales*, specifically *Sulfurimonas* species, from the class *Epsilonproteobacteria*, were enriched on *Z. marina* roots. Previous studies of *Spartina* (Thomas et al., 2014) and *Z. marina* (Jensen, Kühl & Priemé, 2007) also found enrichment of *Epsilonproteobacteria* on roots relative to the surrounding sediment. All known *Sulfurimonas* species are sulfur-oxidizing chemolithoautotrophs, can perform denitrification and are postulated to play significant roles in biogeochemical cycling in marine sediments (Campbell et al., 2006). Members of *Campylobacterales* have previously been identified as nitrogen fixers when isolated from *Spartina* roots (McClung & Patriquin, 1980). Additionally, *Campylobacterales* and *Desulfobacterales*, known sulfur-reducing bacteria, have been previously found to be abundant in association with plants from brackish habitats (e.g., mangroves—Gomes et al., 2010). *Rhodobacterales* which are enriched on the *Z. marina* leaves in our study, are purple nonsulfur bacteria, that have been identified as primary surface colonizers in marine habitats and have been shown to have the ability to fix nitrogen (Palacios & Newton, 2005; Dang et al., 2008). *Desulfobacterales* and *Rhodobacterales* species have been previously found in association with the tropical seagrass *Thalassia hemprichii* (Jiang et al., 2015). *Thiotrichales*, which are enriched in the sediment, are generally filamentous sulfur-oxidizing bacteria (Garrity, Bell & Lilburn, 2005) and are postulated to be dominant sulfur-oxidizers in salt marsh sediments (Thomas et al., 2014).

### Variation in sediment microbial communities between locations

The analysis of diversity metrics reported above also showed that there are significant differences in the sediment microbial communities from different locations (inside a eelgrass patch, the edge of a patch and outside of a patch). We therefore examined in more detail the taxonomic groups that differ significantly between sediment locations and their potential functional roles (Fig. 5).

*Bacteroidales*, *Myxococcales*, *Thiotrichales* and *Chromatiales* are significantly different between locations with a Bonferroni corrected Kruskal–Wallis test ( $p < 0.01$ ). *Thiotrichales*




**Figure 5** Average relative abundance of taxonomic groups associated with sediment from each location (inside, edge, outside). Operational taxonomic units (OTUs) are shown grouped by taxonomic order and are colored by taxonomic class. Only orders with a mean relative abundance of at least one percent are shown. Bars represent the standard error of the mean.

and *Chromatiales* are enriched outside of *Z. marina* patches in the unvegetated sediment compared to the inside or edge of patches. In contrast, *Bacteroidales* and *Myxococcales* are enriched in the rhizosphere sediment inside and at the edge of eelgrass patches compared to the outside. The functional significance of these differences is unclear but we note a few things here. First, *Thiotrichales* and *Chromatiales* are common taxa in other marine and brackish sediments including those associated with various plants (e.g., [Thomas et al., 2014](#)). This is thought to be reflective of a role in sulfur oxidation (see discussion above). Some studies have indicated that these taxa are associated with plants (e.g., seagrasses in Portugal [Cúcio et al., 2016](#)). However, other studies have indicated that these are found more in the sediment near plants but not specifically associated with plants ([Thomas et al., 2014](#)). *Myxococcales*, commonly found in freshwater and marsh sediments, includes microorganisms known to be involved in organic matter degradation ([Bowen et al., 2012](#); [Kou et al., 2016](#); [Cleary et al., 2016](#)). The abundance of *Myxococcales* inside the eelgrass patch aligns with the expectation of higher prevalence of organic matter degradation inside the patch as opposed to surrounding unvegetated sediment.

### Environmental drivers of sediment communities

In addition to investigating the taxonomic composition of the microbial communities of sediment collected from the inside, edge and outside of eelgrass patches, we decided to test for correlations between observed community differences and environmental factors to elucidate key factors that may be driving the microbial communities in eelgrass patches.

A variety of abiotic factors were significantly different between locations including C:N ratio, TIC, dissolved oxygen, pH and sediment size fractions 710  $\mu\text{m}$  and 63  $\mu\text{m}$  (ANOVA,  $p < 0.05$ ) ([Tables S6](#) and [S7](#)). Unsurprisingly, eelgrass shoot density was significantly



**Figure 6** Relationship between environmental data and microbial communities. Non-metric multi-dimensional scaling (NMDS) of Bray Curtis dissimilarities of microbial communities found in sediment samples are shown here colored by location (inside, edge, outside). Environmental factors ( $p < 0.055$ , ANOVA) were overlaid as vectors onto the NMDS using the `envfit` function in `vegan`.

different between locations (ANOVA,  $p < 0.05$ ). To determine which pair-wise locations were driving the significant differences between location overall, we performed Tukey's HSD tests (Tables S8 and S9). We also performed Tukey's HSD tests on percent TOC which was marginally non significantly different (ANOVA,  $p = 0.0519$ ). All pair-wise location comparisons of eelgrass density and dissolved oxygen were significantly different (Tukey's HSD,  $p < 0.05$ ). The C:N ratio and sediment fraction  $63 \mu\text{m}$  were significantly different for the outside-inside comparison (Tukey's HSD,  $p < 0.05$ ). Percent TIC and TOC as well as sediment fraction  $710 \mu\text{m}$  were significantly different for the outside-edge comparison and pH was significantly different for the inside-edge comparisons (Tukey's HSD,  $p < 0.05$ ).

To test if there was a correlation between environmental measures and microbial community composition, Mantel tests were performed on Euclidean distances of environmental measures and the Bray-Curtis dissimilarities of sediment communities. A combined dataset including C:N ratio, TIC, TOC, dissolved oxygen, pH and eelgrass density was found to be significantly positively correlated with the sediment microbial community data ( $r = 0.1122$ ,  $p = 0.0474$ ) (Fig. 6). However, when measures were tested individually only the C:N ratio ( $r = 0.1701$ ,  $p = 0.016$ ) and eelgrass density ( $r = 0.1292$ ,  $p = 0.0381$ ) were significantly correlated with microbial community composition.

The significant correlation between the sediment communities and the C:N ratio may indicate a change in ecosystem nutrient cycling at the patch edge. As Carbon (TIC and TOC) was not correlated with the sediment microbial communities, the correlation with the C:N ratio may hint at the importance of nitrogen, which was not measured in this study, to sediment community composition. Nitrogen is often a limiting terrestrial plant nutrient

and N-limitation has also been observed in several seagrass studies, more frequently in temperate habitats (De Boer, 2007). Terrestrial plants overcome N-limitation by having beneficial interactions with nitrogen fixing bacteria and these bacteria have previously been observed to form associations with eelgrasses (Capone & Budin, 1982; Welsh, 2000; Bagwell et al., 2002; Adhitya, Thomas & Ward, 2007; Sun et al., 2015). Nitrogen fixation can account for 5–10% of plant nitrogen requirements in temperate eelgrass patch sediments, and up to 50% in tropical patch sediments, indicating an important role for nitrogen fixation in overall patch health (Welsh, 2000).

A previous study looking at forest soil microbial communities found that microbial biomass and activity were significantly lower at forest edges due to decreased litter decomposition in the edge habitat and thus, changes in nutrient cycling (Malmivaara-Lämsä et al., 2008). In seagrass patches, on average, vegetated sediments are significantly enriched in organic matter compared to unvegetated sediments, with carbon stocks generally higher on the inside of patches (Duarte, Holmer & Marbà, 2005; Ricart et al., 2015). It is possible that the observed community structure changes in the sediment between locations and the correlations with C:N ratio and eelgrass density here are indicative of a similar trend of location based nutrient cycling resulting from differing nutrient deposition and decomposition rates.

Eelgrass density may have direct or indirect effects on sediment microbial communities as a result of the role eelgrass plays in its environment as a foundation species and an ecosystem engineer (Koch, 2001). Seagrasses are known to modify their surrounding habitat in a variety of ways including enhancing the input and retention of carbon and other nutrients in the sediment (Gacia et al., 2002; Duarte et al., 2005; Duarte & Cebrián, 1996), altering flow velocity and turbulence in the water column above patches (Fonseca et al., 1982; Granata et al., 2001; Folkard, 2005) which can increase sedimentation (Short & Short, 1984; Dauby et al., 1995; Gacia et al., 2002) and oxygenating the sediment using their roots (Caffrey & Kemp, 1991; Pedersen et al., 1998; Connell, Colmer & Walker, 1999).

Other factors at play in the observed differences between locations as a result of eelgrass density may be the development stage of the eelgrass at the edge (if the patch is expanding or contracting) and the variable use of eelgrass as a habitat by macroorganisms. From terrestrial systems, it is known that microbial communities can vary across different stages of plant development (Chaparro, Badri & Vivanco, 2014). Seagrasses at earlier stages in development are known to have different carbon deposition rates than later stages, so if seagrass patch was in the process of expanding this may change the abiotic conditions at the patch edge, and thus might be reflected in distinct microbial communities at the edge of a patch compared to the inside (Duarte & Sand-Jensen, 1990; Cebrián et al., 2000). Additionally, seagrass patches are habitats for a large number of macroorganisms with variable abundance across seagrass patch landscapes (seagrass densities) (Tanner, 2005; Smith et al., 2008; Smith et al., 2011).

Ultimately, although we see differences between locations in environmental abiotic measurements, we are unable, given the limitations of this study, to decouple these measurements from the eelgrass itself (eelgrass density), which is highly correlated with sediment community composition.

## CONCLUSIONS

This study provides new insights into the composition and assembly of the *Z. marina* microbiome. Major findings include that distinct microbial communities are associated with the leaves and roots of the plant, sediment associated communities are correlated with host plant density, and specific microbial taxa are found to have high relative abundances on particular tissues. Differences in the rhizosphere sediment community composition at the patch edge were observed and correlated with variation in environmental measurements. However, we were unable to disentangle these measures from eelgrass density, with the strongest correlated factor with community differences being presence/absence of the host plant. This is perhaps not unexpected from a field study, as eelgrass species are ecosystem engineers that actively change the sediment chemistry and landscape (Orth et al., 2006; Bos et al., 2007).

Overall, we believe that the results of this study hint at a network of complex interactions between *Z. marina*, the microbes associated with *Z. marina* and biogeochemical cycling. Untangling such networks is difficult but increasingly feasible. Although *Z. marina* is not a model organism in the sense of *Arabidopsis* or poplar, we believe it can nevertheless become a model for host-microbiome-environment interaction studies. Advantages of working on this species include that there is a genome now available (Olsen et al., 2016), that there is a large network of collaborating labs focusing on this species (Zostera Experimental Network; <http://zenscience.org>), and that it can be used in common garden and reciprocal transplant experiments. Along these lines we have been building a library of cultured isolates associated with this species and sequencing the genomes of many of these (Lee et al., 2015a; Lee et al., 2015b; Lee et al., 2016a; Lee et al., 2016b; Alexiev et al., 2016a; Alexiev et al., 2016b). There are still areas in need of improvement (e.g., there a limited amount of full length 16S and 18S other reference data; only limited information on the in situ functions of microbes are available, there is a need for more genetic tools for the host), but given the importance of coastal marine systems and seagrasses generally, we believe continued efforts to study the host-microbiome-environment interactions in this and related species are important.

## ACKNOWLEDGEMENTS

Illumina sequencing was performed at the DNA Technologies Core facility in the Genome Center at UC Davis, Davis, California. We thank Qingyi “John” Zhang for his help with the Illumina library preparation and Hannah Holland-Moritz for her help with DNA extractions. We thank Ted Grosholz and Susan Williams for hosting Sofie Voerman during her visit and for help with the sampling design. We thank Daniel Bradley for his help with sample collection.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This work was supported by a grant from the Gordon and Betty Moore Foundation (GBMF333) “Investigating the co-evolutionary relationships between seagrasses and their



microbial symbionts.” The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

Gordon and Betty Moore Foundation: GBMF333.

### Competing Interests

Jonathan A. Eisen is an Academic Editor for PeerJ. Jenna M. Lang is an employee of Trace Genomics, Inc.

### Author Contributions

- Cassandra L. Ettinger analyzed the data, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Sofie E. Voerman conceived and designed the experiments, reviewed drafts of the paper, performed sampling.
- Jenna M. Lang conceived and designed the experiments, reviewed drafts of the paper.
- John J. Stachowicz reviewed drafts of the paper, advised on experimental design, edited drafts of paper.
- Jonathan A. Eisen contributed reagents/materials/analysis tools, reviewed drafts of the paper, advised on data analysis, edited drafts of paper.

### DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:

This 16S rRNA sequencing project has been deposited at GenBank under the accession no. [PRJNA350006](https://doi.org/10.6084/m9.figshare.1014334.v1).

### Data Availability

The following information was supplied regarding data availability:

Coil, David; Eisen, Jonathan; Stachowicz, Jay; Green, Jessica; Holland-Moritz, Hannah; Lang, Jenna (2014): The Seagrass Microbiome. figshare.

<https://doi.org/10.6084/m9.figshare.1014334.v1>.

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.3246#supplemental-information>.

## REFERENCES

- Adhitya A, Thomas FIM, Ward BB. 2007.** Diversity of assimilatory nitrate reductase genes from plankton and epiphytes associated with a seagrass bed. *Microbial Ecology* 54:587–597 DOI [10.1007/s00248-006-9175-0](https://doi.org/10.1007/s00248-006-9175-0).
- Aleklett K, Leff JW, Fierer N, Hart M. 2015.** Wild plant species growing closely connected in a subalpine meadow host distinct root-associated bacterial communities. *PeerJ* 3:e804 DOI [10.7717/peerj.804](https://doi.org/10.7717/peerj.804).

- Alexiev A, Krusor ML, Jospin G, Lang JM, Eisen JA, Coil DA. 2016a.** Draft genome sequences of two *Pseudoalteromonas* strains isolated from roots and leaf blades of the seagrass *Zostera marina*. *Genome Announcements* **4**(1):e00010-16 DOI [10.1128/genomeA.00010-16](https://doi.org/10.1128/genomeA.00010-16).
- Alexiev A, Krusor ML, Jospin G, Lang JM, Eisen JA, Coil DA. 2016b.** Draft genome sequence of *Cobetia* sp. UCD-24C, isolated from roots and leaves of the seagrass *Zostera marina*. *Genome Announcements* **4**(2):e00116-16 DOI [10.1128/genomeA.00116-16](https://doi.org/10.1128/genomeA.00116-16).
- Anderson MJ. 2001.** A new method for non-parametric multivariate analysis of variance. *Austral Ecology* **26**:32–46.
- Anderson MJ, Walsh DC. 2013.** PERMANOVA, ANOSIM, and the mantel test in the face of heterogeneous dispersions: what null hypothesis are you testing? *Ecological Monographs* **83**:557–574 DOI [10.1890/12-2010.1](https://doi.org/10.1890/12-2010.1).
- Bagwell CE, Rocque JR, Smith GW, Polson SW, Friez MJ, Longshore JW, Lovell CR. 2002.** Molecular diversity of diazotrophs in oligotrophic tropical seagrass bed communities. *FEMS Microbiology Ecology* **39**:113–119 DOI [10.1111/j.1574-6941.2002.tb00912.x](https://doi.org/10.1111/j.1574-6941.2002.tb00912.x).
- Barber TR, Carlson PR. 1993.** Effects of seagrass die-off on benthic fluxes and porewater concentrations of  $\sum\text{CO}_2$ ,  $\sum\text{H}_2\text{S}$ , and  $\text{CH}_4$  in Florida Bay Sediments. In: Oremland RS, ed. *Biogeochemistry of global change: radiatively active trace gases*. New York: Chapman and Hall, 530–550.
- Beckers B, OP De Beeck M, Weyens N, Van Acker R, Van Montagu M, Boerjan W, Vangronsveld J. 2016.** Lignin engineering in field-grown poplar trees affects the endosphere bacterial microbiome. *Proceedings of the National Academy of Sciences of the United States of America* **113**:2312–2317 DOI [10.1073/pnas.1523264113](https://doi.org/10.1073/pnas.1523264113).
- Bent SJ, Forney LJ. 2008.** The tragedy of the uncommon: understanding limitations in the analysis of microbial diversity. *The ISME Journal* **2**:689–695 DOI [10.1038/ismej.2008.44](https://doi.org/10.1038/ismej.2008.44).
- Bos AR, Bouma TJ, De Kort GLJ, Van Katwijk MM. 2007.** Ecosystem engineering by annual intertidal seagrass beds: sediment accretion and modification. *Estuarine, Coastal and Shelf Science* **74**:344–348 DOI [10.1016/j.ecss.2007.04.006](https://doi.org/10.1016/j.ecss.2007.04.006).
- Bouffaud ML, Poirier MA, Muller D, Moënné-Loccoz Y. 2014.** Root microbiome relates to plant host evolution in maize and other Poaceae. *Environmental Microbiology* **16**:2804–2814 DOI [10.1111/1462-2920.12442](https://doi.org/10.1111/1462-2920.12442).
- Bowen JL, Morrison HG, Hobbie JE, Sogin ML. 2012.** Salt marsh sediment diversity: a test of the variability of the rare biosphere among environmental replicates. *The ISME Journal* **6**:2014–2023 DOI [10.1038/ismej.2012.47](https://doi.org/10.1038/ismej.2012.47).
- Bray JR, Curtis JT. 1957.** An ordination of the upland forest communities of Southern Wisconsin. *Ecological Monographs* **27**:325–349 DOI [10.2307/1942268](https://doi.org/10.2307/1942268).
- Caffrey JM, Kemp WM. 1991.** Seasonal and spatial patterns of oxygen production, respiration and root-rhizome release in *Potamogeton perfoliatus* L. and *Zostera marina* L. *Aquatic Botany* **40**:109–128 DOI [10.1016/0304-3770\(91\)90090-R](https://doi.org/10.1016/0304-3770(91)90090-R).

- Campbell BJ, Engel AS, Porter ML, Takai K. 2006.** The versatile epsilon-proteobacteria: key players in sulphidic habitats. *Nature Reviews. Microbiology* **4**:458–468 DOI [10.1038/nrmicro1414](https://doi.org/10.1038/nrmicro1414).
- Capone DG. 1982.** Nitrogen fixation (Acetylene Reduction) by rhizosphere sediments of the eelgrass *Zostera marina*. *Marine Ecology Progress Series* **10**:67–75 DOI [10.3354/meps010067](https://doi.org/10.3354/meps010067).
- Capone DG, Budin JM. 1982.** Nitrogen fixation associated with rinsed roots and rhizomes of the eelgrass *Zostera marina*. *Plant Physiology* **70**:1601–1604 DOI [10.1104/pp.70.6.1601](https://doi.org/10.1104/pp.70.6.1601).
- Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Peña AG, Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig JE, Ley RE, Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR, Turnbaugh PJ, Walters WA, Widmann J, Yatsunencko T, Zaneveld J, Knight R. 2010.** QIIME allows analysis of high-throughput community sequencing data. *Nature Methods* **7**:335–336 DOI [10.1038/nmeth.f.303](https://doi.org/10.1038/nmeth.f.303).
- Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, Owens SM, Betley J, Fraser L, Bauer M, Gormley N, Gilbert JA, Smith G, Knight R. 2012.** Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *The ISME Journal* **6**:1621–1624.
- Cebrián J, Pedersen MF, Kroeger KD, Valiela I. 2000.** Fate of production of the seagrass *Cymodocea nodosa* in different stages of meadow formation. *Marine Ecology Progress Series* **204**:119–130 DOI [10.3354/meps204119](https://doi.org/10.3354/meps204119).
- Chao A. 1984.** Non-parametric estimation of the number of classes in a population. *Scandinavian Journal of Statistics, Theory and Applications* **11**:265–270.
- Chaparro JM, Badri DV, Vivanco JM. 2014.** Rhizosphere microbiome assemblage is affected by plant development. *The ISME Journal* **8**:790–803 DOI [10.1038/ismej.2013.196](https://doi.org/10.1038/ismej.2013.196).
- Cleary DFR, Polónia ARM, Sousa AI, Lillebø AI, Queiroga H, Gomes NCM. 2016.** Temporal dynamics of sediment bacterial communities in monospecific stands of *Juncus maritimus* and *Spartina maritima*. *Plant Biology* **18**:824–834 DOI [10.1111/plb.12459](https://doi.org/10.1111/plb.12459).
- Connell EL, Colmer TD, Walker DI. 1999.** Radial oxygen loss from intact roots of *Halophila ovalis* as a function of distance behind the root tip and shoot illumination. *Aquatic Botany* **63**:219–228 DOI [10.1016/S0304-3770\(98\)00126-0](https://doi.org/10.1016/S0304-3770(98)00126-0).
- Cúcio C, Engelen AH, Costa R, Muyzer G. 2016.** Rhizosphere microbiomes of European seagrasses are selected by the plant, but are not species specific. *Frontiers in Microbiology* **7**:440 DOI [10.3389/fmicb.2016.00440](https://doi.org/10.3389/fmicb.2016.00440).
- Dang H, Li T, Chen M, Huang G. 2008.** Cross-ocean distribution of Rhodobacterales bacteria as primary surface colonizers in temperate coastal marine waters. *Applied and Environmental Microbiology* **74**:52–60 DOI [10.1128/AEM.01400-07](https://doi.org/10.1128/AEM.01400-07).
- Dauby P, Bale AJ, Bloomer N, Canon C, Ling RD, Norro A, Robertson JE, Simon A, Théate JM, Watson AJ, Frankignoulle M. 1995.** Particle fluxes over a Mediterranean seagrass bed: a one year case study. *Marine Ecology Progress Series* **126**:233–246 DOI [10.3354/meps126233](https://doi.org/10.3354/meps126233).

- Davies-Colley RJ, Payne GW, Van Elswijk M. 2000.** Microclimate gradients across a forest edge. *New Zealand Journal of Ecology* **11**:1–121.
- De Boer WF. 2007.** Seagrass–sediment interactions, positive feedbacks and critical thresholds for occurrence: a review. *Hydrobiologia* **591**:5–24  
DOI [10.1007/s10750-007-0780-9](https://doi.org/10.1007/s10750-007-0780-9).
- DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, Huber T, Dalevi D, Hu P, Andersen GL. 2006.** Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Applied and Environmental Microbiology* **72**:5069–5072 DOI [10.1128/AEM.03006-05](https://doi.org/10.1128/AEM.03006-05).
- Dixon P. 2003.** VEGAN, a package of R functions for community ecology. *Journal of Vegetation Science: Official Organ of the International Association for Vegetation Science* **14**:927–930 DOI [10.1111/j.1654-1103.2003.tb02228.x](https://doi.org/10.1111/j.1654-1103.2003.tb02228.x).
- Duarte CM, Borum K, Short FT, Walker DI. 2005.** Seagrass ecosystems: their global status and prospects. In: Polunin NVC, ed. *Aquatic ecosystems: trends and global prospects*. Cambridge Univ., 281–294.
- Duarte CM, Cebrián J. 1996.** The fate of marine autotrophic production. *Limnology and Oceanography* **41**:1758–1766 DOI [10.4319/lo.1996.41.8.1758](https://doi.org/10.4319/lo.1996.41.8.1758).
- Duarte CM, Holmer M, Marbà N. 2005.** Plant-microbe interactions in seagrass meadows. *Coastal and Estuarine Studies* Epub ahead of print Mar 23 2013  
DOI [10.1029/CE060p0031](https://doi.org/10.1029/CE060p0031).
- Duarte CM, Sand-Jensen K. 1990.** Seagrass colonization: patch formation and patch growth in *Cymodocea nodosa*. *Marine Ecology Progress Series* **65**:193–200  
DOI [10.3354/meps065193](https://doi.org/10.3354/meps065193).
- Edgar RC. 2010.** Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**:2460–2461 DOI [10.1093/bioinformatics/btq461](https://doi.org/10.1093/bioinformatics/btq461).
- Edwards J, Johnson C, Santos-Medellín C, Lurie E, Podishetty NK, Bhatnagar S, Eisen JA, Sundaresan V. 2015.** Structure, variation, and assembly of the root-associated microbiomes of rice. *Proceedings of the National Academy of Sciences of the United States of America* **112**:E911–E920 DOI [10.1073/pnas.1414592112](https://doi.org/10.1073/pnas.1414592112).
- Egan S, Harder T, Burke C, Steinberg P, Kjelleberg S, Thomas T. 2013.** The seaweed holobiont: understanding seaweed—bacteria interactions. *FEMS Microbiology Reviews* **37**:462–476 DOI [10.1111/1574-6976.12011](https://doi.org/10.1111/1574-6976.12011).
- Elser JJ, Bracken MES, Clel EE, Gruner DS, Harpole WS, Hillebr H, Ngai JT, Seabloom EW, Shurin JB, Smith JE. 2007.** Global analysis of nitrogen and phosphorus limitation of primary producers in freshwater, marine and terrestrial ecosystems. *Ecology Letters* **10**:1135–1142 DOI [10.1111/j.1461-0248.2007.01113.x](https://doi.org/10.1111/j.1461-0248.2007.01113.x).
- Erskine JM, Koch MS. 2000.** Sulfide effects on *Thalassia testudinum* carbon balance and adenylate energy charge. *Aquatic Botany* **67**:275–285  
DOI [10.1016/S0304-3770\(00\)00099-1](https://doi.org/10.1016/S0304-3770(00)00099-1).
- Flaspohler DJ, Temple SA, Rosenfield RN. 2001.** Species-specific edge effects on nest success and breeding bird density in a forested landscape. *Ecological Applications* **11**:32–46 DOI [10.2307/3061053](https://doi.org/10.2307/3061053).

- Folkard AM. 2005.** Hydrodynamics of model *Posidonia oceanica* patches in shallow water. *Limnology and Oceanography* **50**:1592–1600 DOI [10.4319/lo.2005.50.5.1592](https://doi.org/10.4319/lo.2005.50.5.1592).
- Fonseca MS, Fisher JS, Zieman JC, Thayer GW. 1982.** Influence of the seagrass, *Zostera marina* L. on current flow. *Estuarine, Coastal and Shelf Science* **15**:351–364 DOI [10.1016/0272-7714\(82\)90046-4](https://doi.org/10.1016/0272-7714(82)90046-4).
- Fox BJ, Taylor JE, Fox MD, Williams C. 1997.** Vegetation changes across edges of rainforest remnants. *Biological Conservation* **82**:1–13 DOI [10.1016/S0006-3207\(97\)00011-6](https://doi.org/10.1016/S0006-3207(97)00011-6).
- Gacia E, Esperança G, Duarte CM, Middelburg JJ. 2002.** Carbon and nutrient deposition in a Mediterranean seagrass (*Posidonia oceanica*) meadow. *Limnology and Oceanography* **47**:23–32 DOI [10.4319/lo.2002.47.1.0023](https://doi.org/10.4319/lo.2002.47.1.0023).
- Garrity GM, Bell JA, Lilburn T. 2005.** Order V. Thiotrichales ord nov. In: Brenner DJ, Krieg NR, Staley JT, Garrity GM, eds. *Bergey's manual of systematic bacteriology (The Proteobacteria), Part B (The Gammaproteobacteria)*. Vol. 2. New York: Springer, 131.
- Gomes NCM, Cleary DFR, Pinto FN, Egas C, Almeida A, Cunha A, Mendonça-Hagler LCS, Smalla K. 2010.** Taking root: enduring effect of rhizosphere bacterial colonization in mangroves. *PLOS ONE* **5**(11):e14065 DOI [10.1371/journal.pone.0014065](https://doi.org/10.1371/journal.pone.0014065).
- Granata TC, Serra T, Colomer J, Casamitjana X, Duarte CM, Gacia E. 2001.** Flow and particle distributions in a nearshore seagrass meadow before and after a storm. *Marine Ecology Progress Series* **218**:95–106 DOI [10.3354/meps218095](https://doi.org/10.3354/meps218095).
- Hamady M, Lozupone C, Knight R. 2010.** Fast UniFrac: facilitating high-throughput phylogenetic analyses of microbial communities including analysis of pyrosequencing and PhyloChip data. *The ISME Journal* **4**:17–27 DOI [10.1038/ismej.2009.97](https://doi.org/10.1038/ismej.2009.97).
- Harrison S, Bruna E. 1999.** Habitat fragmentation and large-scale conservation: what do we know for sure? *Ecography* **22**:225–232 DOI [10.1111/j.1600-0587.1999.tb00496.x](https://doi.org/10.1111/j.1600-0587.1999.tb00496.x).
- Hollants J, Leliaert F, De Clerck O, Willems A. 2013.** What we can learn from sushi: a review on seaweed–bacterial associations. *FEMS Microbiology Ecology* **83**:1–16 DOI [10.1111/j.1574-6941.2012.01446.x](https://doi.org/10.1111/j.1574-6941.2012.01446.x).
- Jensen SI, Kühl M, Priemé A. 2007.** Different bacterial communities associated with the roots and bulk sediment of the seagrass *Zostera marina*. *FEMS Microbiology Ecology* **6**:108–117.
- Jiang YF, Ling J, Dong JD, Chen B, Zhang YY, Zhang YZ, Wang YS. 2015.** Illumina-based analysis the microbial diversity associated with *Thalassia hemprichii* in Xincun Bay, South China Sea. *Ecotoxicology* **24**:1548–1556 DOI [10.1007/s10646-015-1511-z](https://doi.org/10.1007/s10646-015-1511-z).
- Koch EW. 2001.** Beyond light: physical, geological, and geochemical parameters as possible submersed aquatic vegetation habitat requirements. *Estuaries* **24**:1–17 DOI [10.2307/1352808](https://doi.org/10.2307/1352808).
- Kou W, Zhang J, Lu X, Ma Y, Mou X, Wu L. 2016.** Identification of bacterial communities in sediments of Poyang Lake, the largest freshwater lake in China. *SpringerPlus* **5**:401 DOI [10.1186/s40064-016-2026-7](https://doi.org/10.1186/s40064-016-2026-7).
- Kramer CY. 1956.** Extension of multiple range tests to group means with unequal numbers of replications. *Biometrics* **12**:307–310 DOI [10.2307/3001469](https://doi.org/10.2307/3001469).

- Kramer CY. 1957.** Extension of multiple range tests to group correlated adjusted means. *Biometrics* 13:13–18 DOI 10.2307/3001898.
- Lakshmanan V. 2015.** Root microbiome assemblage is modulated by plant host factors. In: Bais H, Sherrier J, eds. *Plant microbe interactions. Advances in botanical research*. Vol. 75, 57–79.
- Lamers LPM, Govers LL, Janssen ICJM, Geurts JJM, Van der Welle MRW, Van Katwijk MM, Van der Heide T, Roelofs JGM, Smolders AJP. 2013.** Sulfide as a soil phytotoxin—a review. *Frontiers in Plant Science* 4:Article 268 DOI 10.3389/fpls.2013.00268.
- Lang JM, Eisen JA, Zivkovic AM. 2014.** The microbes we eat: abundance and taxonomy of microbes consumed in a day’s worth of meals for three diet types. *PeerJ* 2:e659 DOI 10.7717/peerj.659.
- Lee RD, Jospin G, Lang JM, Eisen JA, Coil DA. 2015a.** Draft genome sequence of *Pseudoalteromonas tetraodonis* Strain UCD-SED8 (Phylum *Gammaproteobacteria*). *Genome Announcements* 3(6):e01276-15 DOI 10.1128/genomeA.01276-15.
- Lee RD, Jospin G, Lang JM, Eisen JA, Coil DA. 2015b.** Draft genome sequence of *Bacillus vietnamensis* Strain UCD-SED5 (Phylum *Firmicutes*). *Genome Announcements* 3(6):e01376-15 DOI 10.1128/genomeA.01376-15.
- Lee RD, Jospin G, Lang JM, Eisen JA, Coil DA. 2016a.** Draft genome sequences of two *Vibrio splendidus* strains, isolated from seagrass sediment. *Genome Announcements* 4(1):e01769-15 DOI 10.1128/genomeA.01769-15.
- Lee RD, Jospin G, Lang JM, Eisen JA, Coil DA. 2016b.** Draft genome sequences of two *Pseudoalteromonas porphyrae* strains isolated from seagrass sediment. *Genome Announcements* 4(2):e00092-16 DOI 10.1128/genomeA.00092-16.
- Les DH, Cleland MA, Waycott M. 1997.** Phylogenetic studies in alismatidae, II: evolution of marine angiosperms (seagrasses) and hydrophily. *Systematic Botany* 22:443–463 DOI 10.2307/2419820.
- Lovell CR. 2002.** Plant-microbe interactions in the marine environment. In: Bitton G, ed. *Encyclopedia of environmental microbiology*. Vol. 5. Hoboken: Wiley, 2539–2554.
- Lozupone CA, Hamady M, Kelley ST, Knight R. 2007.** Quantitative and qualitative diversity measures lead to different insights into factors that structure microbial communities. *Applied and Environmental Microbiology* 73:1576–1585 DOI 10.1128/AEM.01996-06.
- Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S, Tremblay J, Engelbrektson A, Kunin V, Del Rio TG, Edgar RC, Eickhorst T, Ley RE, Hugenholtz P, Tringe SG, Dangl JL. 2012.** Defining the core *Arabidopsis thaliana* root microbiome. *Nature* 488:86–90 DOI 10.1038/nature11237.
- Malmivaara-Lämsä M, Hamberg L, Haapamäki E, Liski J, Kotze DJ, Lehvävirta S, Fritze H. 2008.** Edge effects and trampling in boreal urban forest fragments—impacts on the soil microbial community. *Soil Biology & Biochemistry* 40:1612–1621 DOI 10.1016/j.soilbio.2008.01.013.

- McClung CR, Patriquin DG. 1980.** Isolation of a nitrogen-fixing *Campylobacter* species from the roots of *Spartina alterniflora* Loisel. *Canadian Journal of Microbiology* 26:881–886 DOI 10.1139/m80-153.
- McMurdie PJ, Holmes S. 2013.** phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLOS ONE* 8(4):e61217 DOI 10.1371/journal.pone.0061217.
- Mejia AY, Rotini A, Lacasella F, Bookman R, Thaller MC, Shem-Tov R, Winters G, Migliore L. 2016.** Assessing the ecological status of seagrasses using morphology, biochemical descriptors and microbial community analyses. A study in Halophila Stipulacea (*Forsk.*) *Aschers meadows in the northern Red Sea. Ecological Indicators* 60:1150–1163.
- Nielsen LB, Finster K, Welsh DT, Donnelly A, Herbert RA, De Wit R, Lomstein BA. 2001.** Sulphate reduction and nitrogen fixation rates associated with roots, rhizomes and sediments from *Zostera noltii* and *Spartina maritima* meadows. *Environmental Microbiology* 3:63–71 DOI 10.1046/j.1462-2920.2001.00160.x.
- Olsen JL, Rouzé P, Verhelst B, Lin Y-C, Bayer T, Collen J, Dattolo E, De Paoli E, Dittami S, Maumus F, Michel G, Kersting A, Lauritano C, Lohaus R, Töpel M, Tonon T, Vanneste K, Amirebrahimi M, Brakel J, Boström C, Chovatia M, Grimwood J, Jenkins JW, Jueterbock A, Mraz A, Stam WT, Tice H, Bornberg-Bauer E, Green PJ, Pearson GA, Procaccini G, Duarte CM, Schmutz J, Reusch TBH, Van de Peer Y. 2016.** The genome of the seagrass *Zostera marina* reveals angiosperm adaptation to the sea. *Nature* 530:331–335 DOI 10.1038/nature16548.
- Orth RJ, Carruthers TJB, Dennison WC, Duarte CM, Fourqurean JW, Heck KL, Hughes RA, Kendrick GA, Kenworthy WJ, Olyarnik S, Short FT, Waycott M, Williams SL. 2006.** A global crisis for seagrass ecosystems. *Bioscience* 56:987–996 DOI 10.1641/0006-3568(2006)56[987:AGCFSE]2.0.CO;2.
- Palacios R, Newton WE. 2005.** *Genomes and genomics of nitrogen-fixing organisms*. Dordrecht: Springer.
- Paton PWC. 1994.** The effect of edge on avian nest success: how strong is the evidence? *Conservation Biology: the Journal of the Society for Conservation Biology* 8:17–26 DOI 10.1046/j.1523-1739.1994.08010017.x.
- Pedersen O, Borum J, Duarte CM, Fortes MD. 1998.** Oxygen dynamics in the rhizosphere of *Cymodocea rotundata*. *Marine Ecology Progress Series* 169:283–288 DOI 10.3354/meps169283.
- Peiffer JA, Spor A, Koren O, Jin Z, Tringe SG, Dangl JL, Buckler ES, Ley RE. 2013.** Diversity and heritability of the maize rhizosphere microbiome under field conditions. *Proceedings of the National Academy of Sciences of the United States of America* 110:6548–6553 DOI 10.1073/pnas.1302837110.
- R Core Team. 2016.** R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing. Available at <https://www.r-project.org>.
- Ricart AM, York PH, Rasheed MA, Pérez M, Romero J, Bryant CV, Macreadie PI. 2015.** Variability of sedimentary organic carbon in patchy seagrass landscapes. *Marine Pollution Bulletin* 100:476–482 DOI 10.1016/j.marpolbul.2015.09.032.

- Roth-Schulze AJ, Zozaya-Valdés E, Steinberg PD, Thomas T. 2016.** Partitioning of functional and taxonomic diversity in surface-associated microbial communities. *Environmental Microbiology* **18**:4391–4402 DOI [10.1111/1462-2920.13325](https://doi.org/10.1111/1462-2920.13325).
- Shannon CE, Weaver W. 1949.** *The mathematical theory of communication*. Urbana: University of Illinois Press, 29.
- Short FT. 1987.** Effects of sediment nutrients on seagrasses: literature review and mesocosm experiment. *Aquatic Botany* **27**:41–57 DOI [10.1016/0304-3770\(87\)90085-4](https://doi.org/10.1016/0304-3770(87)90085-4).
- Short FT, Short CA. 1984.** The seagrass filter: purification of estuarine and coastal waters. In: Kennedy VS, ed. *The estuary as a filter*. Orlando: Academic Press, 395–413.
- Simpson EH. 1949.** Measurement of diversity. *Nature* **163**:688–688 DOI [10.1038/163688a0](https://doi.org/10.1038/163688a0).
- Smith TM, Hindell JS, Jenkins GP, Connolly RM. 2008.** Edge effects on fish associated with seagrass and sand patches. *Marine Ecology Progress Series* **359**:203–213 DOI [10.3354/meps07348](https://doi.org/10.3354/meps07348).
- Smith TM, Hindell JS, Jenkins GP, Connolly RM, Keough MJ. 2011.** Edge effects in patchy seagrass landscapes: the role of predation in determining fish distribution. *Journal of Experimental Marine Biology and Ecology* **399**:8–16 DOI [10.1016/j.jembe.2011.01.010](https://doi.org/10.1016/j.jembe.2011.01.010).
- Sun F, Zhang X, Zhang Q, Liu F, Zhang J, Gong J. 2015.** Seagrass (*Zostera marina*) colonization promotes the accumulation of diazotrophic bacteria and alters the relative abundances of specific bacterial lineages involved in benthic carbon and sulfur cycling. *Applied and Environmental Microbiology* **81**:6901–6914 DOI [10.1128/AEM.01382-15](https://doi.org/10.1128/AEM.01382-15).
- Tanner JE. 2005.** Edge effects on fauna in fragmented seagrass meadows. *Austral Ecology* **30**:210–218 DOI [10.1111/j.1442-9993.2005.01438.x](https://doi.org/10.1111/j.1442-9993.2005.01438.x).
- Terrados J, Duarte CM, Kamp-Nielsen L, Agawin NSR, Gacia E, Lacap D, Fortes MD, Borum J, Lubanski M, Greve T. 1999.** Are seagrass growth and survival constrained by the reducing conditions of the sediment? *Aquatic Botany* **65**:175–197 DOI [10.1016/S0304-3770\(99\)00039-X](https://doi.org/10.1016/S0304-3770(99)00039-X).
- Thomas F, Giblin AE, Cardon ZG, Sievert SM. 2014.** Rhizosphere heterogeneity shapes abundance and activity of sulfur-oxidizing bacteria in vegetated salt marsh sediments. *Frontiers in Microbiology* **5**:Article 309.
- Tukey JW. 1953.** *The problem of multiple comparisons. The collected works of John W. Tukey*, vol. 8. New York: Chapman & Hall.
- Van der Heide T, Govers LL, De Fouw J, Olf H, Van Der Geest M, Van Katwijk MM, Piersma T, Van de Koppel J, Silliman BR, Smolders AJP, Van Gils JA. 2012.** A three-stage symbiosis forms the foundation of seagrass ecosystems. *Science* **336**:1432–1434 DOI [10.1126/science.1219973](https://doi.org/10.1126/science.1219973).
- Warton DI, Wright ST, Wang Y. 2012.** Distance-based multivariate analyses confound location and dispersion effects. *Methods in Ecology and Evolution* **3**:89–101 DOI [10.1111/j.2041-210X.2011.00127.x](https://doi.org/10.1111/j.2041-210X.2011.00127.x).



**Welsh DT. 2000.** Nitrogen fixation in seagrass meadows: regulation, plant-bacteria interactions and significance to primary productivity. *Ecology Letters* 3:58–71  
DOI [10.1046/j.1461-0248.2000.00111.x](https://doi.org/10.1046/j.1461-0248.2000.00111.x).

**Wickham H. 2009.** *ggplot2: elegant graphics for data analysis*. New York: Springer-Verlag.

**Williams SL, Heck Jr KL. 2001.** Seagrass communities ecology. In: Bertness MD, Gaines SD, Hay ME, eds. *Marine community ecology*. Sunderland: Sinauer Associates, 317–337.