



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Quantifying particulate matter accumulated on leaves by 17 species of urban trees in Beijing, China

Citation for published version:

Xu, Y, Xu, W, Mo, L, Heal, MR, Xu, X & Yu, X 2018, 'Quantifying particulate matter accumulated on leaves by 17 species of urban trees in Beijing, China' *Environmental science and pollution research*, vol. 25, no. 13, pp. 12545-12556. DOI: 10.1007/s11356-018-1478-4

Digital Object Identifier (DOI):

[10.1007/s11356-018-1478-4](https://doi.org/10.1007/s11356-018-1478-4)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Environmental science and pollution research

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Quantifying Particulate Matter Accumulated on Leaves by 17

Species of Urban Trees in Beijing, China

Yansen Xu^{1,2}, Wen Xu², Li Mo^{1,3}, Mathew R. Heal⁴, Xiaowu Xu¹, Xinxiao Yu^{1*}

¹Beijing Laboratory of Urban and Rural Ecological Environment, Beijing Forestry University, No.35 Tsinghua East Road, Beijing, China.

²State key laboratory of Urban and Regional Ecology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, No. 18 Shuangqing Road, Beijing, China.

³China-ASEAN Environmental Cooperation Center, No. 5 Houyingfang Hutong, Beijing, China

⁴School of Chemistry, University of Edinburgh, David Brewster Road, Edinburgh, UK

Abstract

Airborne particulate matter (PM) has become a serious environmental problem and harms human health worldwide. Trees can effectively remove particles from the atmosphere and improve the air quality. In this study, a washing and weigh method was used to quantify accumulation of water-soluble ions and insoluble PM on the leaf surfaces and within the wax of the leaves for 17 urban plant species (including 4 shrubs and 13 trees). The deposited PM was determined in three size-fractions: fine (0.2–2.5 μm), coarse (2.5–10 μm), and large ($> 10 \mu\text{m}$). Significant differences in the accumulation of PM were detected among various species. The leaves of *Platycladus orientalis* and *Pinus armandi* were the most effective in capturing PM. Across the species, 65% and 35% of PM on average deposited on the leaf surface and in the wax, respectively. The greatest PM accumulation by mass on leaves was in the largest PM size fraction, while accumulation of coarse and fine particles size fractions was smaller. Water-soluble ions accumulated on leaf surfaces contributed 28% to the total PM mass on average. This study demonstrated that leaves of woody plants accumulate PM differently, and the most effective plant species should be selected in urban areas for attenuating ambient PM.

Key words Particulate matter; Urban trees; Leaf deposition; Wax; Water-soluble ions.

* Corresponding author.

E-mail address: yuxinxiao111@126.com

33 **Introduction**

34

35 Airborne particulate matter (PM), consisting of particles with aerodynamic diameters in the
36 range of 0.001-100 μm , is a major atmospheric pollutant (Pope III and Dockery 2006; WHO
37 2006). Anthropogenic sources contributing to PM include vehicle exhausts, road dust, domestic
38 and large-scale coal burning, and cement and other industrial processes (Bosco et al. 2005;
39 Dzierzanowski et al. 2011). Due to rapid urbanization and industrialization, the Beijing-Tianjin-
40 Hebei region is undergoing serious PM pollution. For example, the annual average $\text{PM}_{2.5}$ (PM
41 with diameter $<2.5 \mu\text{m}$) concentration in Beijing was $86 \mu\text{g m}^{-3}$ in 2014 (Chen et al. 2015), which
42 was about 2.5 times higher than the WHO Air Quality Guideline value of $35 \mu\text{g m}^{-3}$ (WHO
43 2006). Positive associations have been found between $\text{PM}_{2.5}$ and health impacts (e.g. lower
44 respiratory infections, trachea bronchus, lung cancers and cerebrovascular disease) (Lelieveld et
45 al. 2015; Liu et al. 2016). It has been estimated that $\text{PM}_{2.5}$ pollution in urban areas of China led to
46 763,595 premature deaths in 2013 (Song et al. 2016). Beijing was worst affected with estimates
47 of 5.2, 9.0, 2.3, and 1.6 thousand premature mortalities from ischemic heart disease (IHD),
48 cerebrovascular disease, chronic obstructive pulmonary disease (COPD) and lung cancer (LC),
49 respectively (Liu et al. 2016).

50 Numerous studies have identified that urban greening filters accumulate atmospheric particles
51 more effectively than other land surfaces (Chen et al. 2016a; Janhäll 2015; Tallis et al. 2011;

52 Thithanhthao et al. 2015). Enhanced the deposition flux to the surface is the benefit for reducing
53 pollutant concentrations near the ground and thus exposure of people in urban areas. This is one
54 of the recognized ecosystem services of urban vegetation (Salmond et al. 2016). Compared with
55 other urban surfaces, vegetation enhances deposition of particles because of the finely divided
56 structure of many leaves, especially conifers. They have a larger collecting surface per unit
57 ground area, and reduce the laminar boundary layer that limits the particles uptake. Using the i-
58 tree model, Nowak et al. (2013) modeled PM_{2.5} removal by trees in ten American cities and
59 estimated that annual masses of particles removal ranged from 4.7 t in Syracuse to 64.5 t in
60 Atlanta. Schaubroeck et al. (2014) used the canopy interception and PM removal multilayered
61 model (CIPAM) to estimate PM accumulated on a forestry canopy. For a case study in 2010 they
62 estimated that a Scots pine stand in Belgium accumulated about 31 kg PM_{2.5} ha⁻¹ yr⁻¹. The plain
63 afforestation project in Beijing has been estimated to decrease annual PM_{2.5} concentrations by
64 0.57 µg m⁻³, or 2% of the target set by the “Beijing Clean Air Action Plan (2013-2017)” (Chen et
65 al. 2014).

66 At a leaf scale, particles deposited on leaf surface and ultrafine PM (particles with diameter <
67 100 nm) may enter the leaf stomata. Previous studies have examined the effectiveness of urban
68 trees around the world for accumulation of PM_{2.5} and PM₁₀ (Chen et al. 2016b; Sgrigna et al.
69 2015) and demonstrated that PM accumulated on leaves differed significantly among plant

70 species (Beckett et al. 2000; Przybysz et al. 2014). Conifer species have greater potential in
71 capturing PM than broad-leaved species and evergreen conifers accumulate PM throughout the
72 year (Sæbø et al. 2012). However, broad-leaved species with large amounts of pubescence and
73 rougher surfaces also accumulate greater masses of particles (Mo et al. 2015). Particles mainly
74 deposit on the adaxial leaf surfaces, and accumulated masses about six times higher than on
75 abaxial leaf surfaces (Wang et al. 2006). The importance of leaf microstructure characteristics on
76 the effectiveness of capturing PM was evidenced by using scanning electron microscopy (SEM)
77 and energy dispersive X-ray (EDX) analysis (Yan et al. 2016).

78 Given limited greening space in urban Beijing, the most effective plant species in removing
79 PM should be selected for urban greening. Although many studies have focused on quantifying
80 the amount of PM deposited on trees (Yin et al. 2011; Chen et al. 2015; Wang et al. 2015;
81 Leonard et al. 2016), there is still a knowledge gap in terms of accumulation of PM in
82 epicuticular waxes between different species. Xu et al. (2017) used an artificial rainfall
83 simulation system to investigate the influence of rainfall duration and intensity on PM removal
84 from four broad-leaved species, and they detected that final wash off rates in the rainfall events
85 was only 51-70% of initial deposition. In other words, most PM captured by leaves is stored
86 temporarily, since deposited particles on leaf surfaces may be subsequently resuspended from the
87 leaves to the air by wind and rain (Schaubroeck et al. 2014). However, PM accumulation in the

88 wax fraction is very important because these particles are trapped permanently (Song et al. 2015).
89 Most studies collect leaves only once at the beginning or in the middle of the season (Binze et al.
90 2014; Cheng et al. 2016; Mo et al. 2015). However, the characteristics of leaf structure are
91 different during the growing season, which may impact their ability to accumulate PM. The leaf-
92 washing methodology is the most common experimental approach. A large proportion of PM is
93 water-soluble ions, which comprised approximately 40% of PM_{2.5} in Beijing (Han et al. 2016).
94 However, the rinse and weigh method may underestimate the mass of particles on leaves.
95 Therefore, water solutions ions need to be quantified in order to accurately assess the
96 effectiveness of plant species in capturing PM. Analyzing water-soluble ions in deposited PM on
97 leaf surface allows to identify pollution source of PM. The aim of this study was to investigate
98 the effectiveness of 17 plant species in accumulating PM during the growth season in Beijing for
99 the year 2014. Two key features of this study were investigation of: 1) particles deposited onto
100 the leaf surface and into the wax as a function of three particles size fractions; and (2) the
101 accumulation of water-soluble ions by the leaf surface.

102

103 **Material and methods**

104

105 **Study Area**

106 All of the plant materials were collected from the campus of Beijing Forestry University (BFU)
107 (40°0'29.64"N, 116°20'46.04"E), Beijing City, China. The BFU campus is located in Haidian

108 District outside the Fourth Ring Road, which is the transition from urban to suburban areas.
109 There are no polluting industries or power plants within 5 km space range. All plant species were
110 located within 600 m each other, in the center of the BFU campus and 50 m away from
111 surrounding streets. It is a reasonable assumption that all species were exposed to the same
112 background PM concentrations. The average background PM₁₀ and PM_{2.5} concentrations were
113 139 and 84 $\mu\text{g m}^{-3}$ from April to September 2014, respectively, in Olympic Sport Center which
114 was the nearest air quality monitoring station (about 4.4 km apart) set by the Ministry of
115 Environmental Protection of China (<http://113.108.142.147:20035/emcpublish/>).

116 **Plant material and sample collection**

117 Vegetation samples were collected from 17 plant species (listed in Table 1). All are common
118 plants in northern China and widely used in urban areas. For each species three similar
119 individuals were used as replicates. Leaves were collected on 16 April, 20 May, 30 June, 27 July,
120 11 September and 14 October 2014. Criteria for the weather on a sampling day were sunny, wind
121 speed less than 5 m s^{-1} , and 9-10 days since precipitation amount of the last rain event $>5 \text{ mm}$.
122 Leaves were assumed to be washed clean by rainfall, so the average mass of daily deposited PM
123 was measured for the same exposure days. The study of Liu et al. (2013) investigated the dust-
124 retaining capacity of the four urban species in Guangzhou, and their results showed that the
125 amounts of PM deposited on leaf surface may approach saturation around 24 days. In the present
126 study, the average mass of daily deposited PM was measured for the same exposure days without

127 saturation, and leaves were assumed to be fully washed by rainfall. Intact leaves were in good
128 growth conditions, namely, with little or no disease and/or pests. They were sampled from four
129 directions in the canopy with the same height for each species. Between 200 and 300 cm² of leaf
130 area (about 5 - 30 leaves) were cut off with scissors and stored in plastic bags. Great care was
131 taken to ensure that particles did not dislodge from the leaves. The samples were then transported
132 to the laboratory and stored at 4°C freezer prior to analysis. Although the detailed collection
133 method of leaves samples has described in our previous studies (Chen et al. 2016b; Mo et al.
134 2015), it is repeated here for the reader's convenience.

135 **Quantitative analysis**

136 PM was washed and quantified from the leaf surfaces and in leaf waxes using the methods
137 described by Dzierzanowski et al. (2011). The methods for rinsing water-soluble ions from the
138 leaves were as described by Freer-Smith et al. (2005). Leaves were first washed with 250 mL of
139 deionized water. The rinse water was passed through a 100 µm metal sieve to remove particles
140 >100 µm and then filtered sequentially through 10 µm, 2.5 µm and 0.2 µm filters (EMD
141 Millipore Corp., Billerica, Massachusetts, USA) to separate the PM into the following three
142 particles size fractions: fine (0.2–2.5 µm), coarse (2.5–10 µm), and large (> 10 µm). The aqueous
143 sample after filtering was stored in the freezer (–18°C) prior to determination of the water-soluble
144 ions (see below). Before filtering, a deionizer gate (AP-BC2451, AP&T, Shanghai, China) was be
145 use to avoid an electrostatic charge on the filters. Before and after analysis, filters were dried in

146 an oven for 30 minutes at 50°C and stabilized in an 80 cm × 80 cm × 80 cm
147 polytetrafluoroethylene (PTFE) balance box at 25°C and 40% relative humidity for 24 hours,
148 which was controlled by a balance and a humidity controller (WHD48-11, ACREL Co., Ltd.,
149 Jiangsu, China). Filters were weighed using a BT125D balance (Sartorius, precision: 10 µg). PM
150 accumulated in waxes was also measured similarly except that the deionized water was replaced
151 with chloroform in order to dissolve the wax on the leaves. Also since nitrocellulose filters are
152 damaged by chloroform, PTFE filters were used in the filtering for in-wax PM.

153 Each washed broad-leaved sample was scanned using a scanner (HP Scanjet 4850, China
154 Hewlett-Packard Co., Ltd., Beijing, China), then the area was measured by Image J (1.50i,
155 National Institutes of Health, Bethesda, US). For needle leaves, leaf area was measured by
156 measuring the water displacement leaf volume and converting to leaf area using the following
157 equation:

158

159
$$S = 2L\left(1 + \frac{\pi}{n}\right)\sqrt{\frac{nV}{\pi L}}$$

160

161 where S represents leaf area; L is the average length of the leaves; V is the needle volume, and
162 n is the number of needle leaves.

163 To analyze the concentrations of water-soluble inorganic ions, the volumes of post-filtering
164 aqueous extracts were measured (V_{water}) and stored at $-18\text{ }^{\circ}\text{C}$ until analysis within four weeks.
165 Three anions (Cl^- , SO_4^{2-} , NO_3^-) and five cations (NH_4^+ , Ca^{2+} , Na^+ , Mg^{2+} , K^+), were determined
166 using a Dionex model ICS-120 ion chromatograph equipped with a conductivity detector (ASRS-
167 ULTRA) following the method of Lun et al. (2003) and Tang et al. (2016). The mass of these
168 ions obtained in each filtrate was calculated by multiplying V_{water} by the ion concentration, and
169 was expressed in per unit area per leaf.

170 **Statistical analysis**

171 One-way analysis of variance (ANOVA) was used to test for significant differences in PM
172 accumulation between species. Post-Hoc analysis (Duncan's test) was performed when multiple
173 comparisons among species were necessary. Spearman's correlation analysis was used to assess
174 the linear correlations between surface PM, in-wax PM and water-soluble ions. All statistical
175 analyses were carried out using SPSS 17.0 (SPSS 17.0 for Windows, SPSS Inc., IL, USA).

176

177 **Results and discussion**

178

179 **PM accumulation on leaves**

180 Fig. 1 shows the mass of accumulated surface and in-wax PM on the 17 plant species. The PM
181 accumulation on leaves varied significantly between plant species. Several studies have also
182 identified that different height of trees may affect the capacity for accumulating PM

183 (Dzierżanowski et al. 2011; Sæbø et al. 2012; Mo et al. 2015). In this study, we divided the
184 species into shrubs and trees (Table 1). The average PM accumulation for four shrubs was 49 μg
185 cm^{-2} . *E. japonicus* showed the highest PM accumulation (56 $\mu\text{g cm}^{-2}$) of the four shrubs.
186 Although the difference was not significant. Xie et al. (2014) and Wang et al. (2006) reported that
187 *E. japonicas* captured larger amounts of PM than other shrubs such as *Buxus sinica*, *Syringa*
188 *oblata* and *Lagerstroemia indica*. The leaf surface structure of *E. japonicus* is unique to those
189 shrubs. Tomentose pubescence distributes on the abaxial leaf surface, which can efficiently
190 capture and accumulate PM (Mo et al. 2015).

191 PM accumulation on *P. orientalis* and *P. armandi* were significantly higher than for other
192 trees. Accumulations on *F. pennsylvanica*, *P. tomentosa*, *A. altissima* and *S. japonica* were
193 significantly lower than for other trees (Fig. 1). The average PM accumulation on *P. orientalis*
194 and *P. armandi* (156 $\mu\text{g cm}^{-2}$) was 4.5 times greater than the average accumulation on the latter
195 four species (28.4 $\mu\text{g cm}^{-2}$). The PM accumulation on *R. typhina* (76 $\mu\text{g cm}^{-2}$) was second only to
196 *P. orientalis* and *P. armandi*. Trees can be divided into conifer or broad-leaved species. *P.*
197 *tabulaeformis*, *P. orientalis* and *P. armandi* are three common evergreen conifer species, which
198 accumulated more PM on leaves than other species except *R. typhina*. The PM accumulation
199 difference between the adaxial and abaxial broad-leaved surfaces is due to wind turbulence.
200 Wang et al (2006) investigated 11 broad-leaved species such as *P. tomentosa*, *S. Japonica* and

201 *Ailanthus altissima* using a scanning electron microscope (SEM) in Beijing urban area. They
202 found that on average only 17% of PM accumulated on the abaxial surface. However, the needles
203 of conifer species can accumulate PM over the entire leaf surface (Ottelé et al. 2010). The leaves
204 of conifer species also have unique microstructure, such as mucus oils, a thicker epicuticular wax
205 layer and a grooved ridge protuberance, which can help leaves to accumulate large particles
206 (Sabin et al. 2006). Previous studies showed that the leaves of *Platycladus* were more rough than
207 *Pinus* (Wang et al. 2007). This structure retained more particles on leaves of *P. orientalis*. The
208 average PM accumulation of broad-leaved trees was less than broad-leaved shrubs. The meta-
209 analysis results showed that PM leaf deposition on shrubs was significantly higher than that of
210 trees (Cai et al. 2017).

211 Across all species, PM was found both on leaf surface and in waxes. Fig. 1 shows that particles
212 distribution between surface and waxes is similar for all shrub species, with PM mass in waxes
213 about 4 times lower than on surface. The average surface and in-wax PM deposition on the shrub
214 leaves were 39 and 10 $\mu\text{g cm}^{-2}$. This corresponds to 21% of PM deposition in waxes, on average.
215 There were significant differences between tree species in the accumulation of PM on leaf
216 surfaces and in waxes. The lowest and highest surface PM deposition were found in *P. tomentosa*
217 (12.5 $\mu\text{g cm}^{-2}$) and *P. armandi* (56.9 $\mu\text{g cm}^{-2}$). The in-wax PM accumulation of *P. orientalis*
218 (101.9 $\mu\text{g cm}^{-2}$) and *P. armandi* (96.6 $\mu\text{g cm}^{-2}$) was significantly higher than other tree species. A

219 positive relationship between different PM fractions accumulation and the quantity of leaf waxes
220 was detected (Sæbø et al. 2012). Popek et al (2013) analyzed all tested species and also detected a
221 positive correlation between the amount of waxes and coarse PM. But the model had a low
222 partial fit ($r=0.54$). By analyzing for each species separately, a significant correlation was found
223 in leaves of *Tilia cordata* (Dzieranowski et al. 2011) and *Corylus colurna* (Popek et al. 2013). In
224 some cases, only weak, no relationship or negative correlation between mass of in-wax PM and
225 mass of wax were reported across some plant species (Jouraeva et al. 2002; Dzieranowski et al.
226 2011). The relationship showed different results among species. Song (2015) found that the
227 waxes in conifer species was a factor of about 2.5 times higher than broadleaves. *P. orientalis*
228 and *P. armandi*, of which the waxes can accumulate large fraction of the PM. The composition
229 and structure of the waxes may significantly affect the capacity of leaves in accumulating PM
230 (Kaupp et al. 2000; Jouraeva et al. 200; Bukhardt et al. 2010). On the other hand, *P.*
231 *tabulaeformis*, *P. orientalis* and *P. armandi* which accumulated greater in-wax PM than other
232 plant species are evergreen conifer species. The leaf growth cycle of these evergreen conifer
233 species of more than 12 months is considerably longer than for deciduous species. Song et al
234 (2015) investigated the mass of PM deposited on five evergreen species in Beijing. The PM
235 accumulated on the leaf surface were range from 72.31 to 231.84 $\mu\text{g cm}^{-2}$. Cai et al. (2017)
236 reviewed 150 studies and used a meta-analysis and also found that the weekly PM leaf deposition

237 of conifer species was significantly higher than broad leaves, by approximately 31.9%. These
238 results showed that evergreen conifer species had better performance in accumulating PM both
239 on the leaf surface and in the surface wax, which was accounted for by the structural
240 characteristics and habits of conifer species.

241 In our study we found significant positive correlation between in-wax PM accumulation and
242 surface PM accumulation ($R^2 = 0.43$, $p = 0.002$). Based on collection and analysis of existing raw
243 data of 56 plant species in 3 different urban areas (Popek et al. 2013; A. Przybysz et al. 2014; Mo
244 et al. 2015; and Song et al. 2015), a positive correlation relationship between total amount of
245 surface PM and in-wax PM accumulated on foliage was also detected ($R^2=0.64$, $P<0.0001$, In-
246 wax PM= $0.24 \times$ Surface PM+2.35). The leaf surface PM deposition amounts was 3.57 times
247 higher than that of in-wax. Actually, PM in the wax layer can account for a significant amount, about
248 22% in the present study. The epicuticular wax layer and releasing PM were dissolved by
249 chloroform, which has environmental health concern. In the previous studies, PM accumulated on
250 leaf surface was assessed (Freer-Smith et al. 2005; Chen et al. 2015). Lack of the examination of
251 PM encapsulated in the wax layer may lead to an underestimation of the present results. However,
252 a significant correlation was found between surface and in-wax PM, which can improve our
253 knowledge on this issue. Because leaves with greater amounts of surface PM may also

254 accumulate high mass of PM in wax layer, and these species should be attractive options for
255 urban greening.

256 The average mass of surface PM accumulation in trees ($34 \mu\text{g cm}^{-2}$) was lower than that in
257 shrubs ($39 \mu\text{g cm}^{-2}$). But trees have the higher average of in-wax PM accumulation ($28 \mu\text{g cm}^{-2}$)
258 relative to shrubs ($10 \mu\text{g cm}^{-2}$) but not statistical significant. Contributions of in-wax depositions
259 to leaf total (surface plus in-wax) depositions were $> 50\%$ for *P. tomentosa* (54%), *P. orientalis*
260 (65%) and *P. armandi* (63%). In the remaining 10 tree species, the contributions of in-wax to
261 total insoluble PM were on average 32%. In Norway and Poland, *Fagus sylvatica* and
262 *Stephanandra incisa* also accumulated about 25% and 28% of PM in the waxes and on the
263 surface, respectively. *Betula pendula* accumulated 82.6% of PM in the wax fraction (Sæbø et al.
264 2012), which is significantly higher than the value of coniferous species obtained in this study.
265 Popek et al. (2013) reported that more PM was deposited on the leaf surface. In their study, the
266 waxes in 5 of 39 woody species accumulated greater than 50% of insoluble PM (range 53%-63%),
267 consistent with our findings.

268 This study (Fig. 6 and 7) showed that PM leaf accumulation varied greatly among months,
269 which was different between leaf surface and waxes. Leaf surface PM retention amounts in trees
270 and shrubs were significantly higher in April and May, respectively, compared with those in other
271 months. The lowest amounts were measured in September. Masses of PM accumulated on

272 surfaces of trees and shrubs showed inconsistent relations among different months. Atmospheric
273 PM concentration, meteorological condition and leaf characteristics varied among months, which
274 may influence PM leaf deposition. Basis on analysis of our dataset, monthly ambient average
275 concentrations of PM₁₀ and PM_{2.5} did not correlate with deposited amounts of total surface
276 particle. However, with increasing atmospheric PM concentrations, particles accumulated on leaf
277 surface raised slightly. By quantitatively analyzing in four districts of the city in Italy, leaf
278 surface PM deposition didn't correlate with local atmospheric PM₁₀ concentration (G. Sgrigna et
279 al. 2015). On the other hand, a meta-analysis result showed that PM accumulation is generally
280 highest in winter compared to other seasons and thought this variable is affected by atmospheric
281 PM concentration (Cai et al. 2017). In this study, monthly data was converted to seasons and
282 showed highest total PM deposited was in spring for trees and shrubs. Leaf PM depositions were
283 all most equally between summer and autumn. But background PM concentrations in summer
284 were lower than spring and autumn. On the other hand, monthly change of PM accumulation in
285 waxes is increased firstly and then decreased during growing season, which in trees and shrubs
286 were highest in July and September, respectively. Vegetable characteristics and ambient PM
287 concentrations show the interaction in leaf PM deposition. In winter, most studies focused on
288 evergreen species such as coniferous plants, which show high capacity to capture particles. So,
289 the conclusion that PM leaf deposition is highest in winter is one-sided. Across all growing

290 months, trees show higher PM accumulation in wax than shrubs. The PM accumulation on leaves
291 varied significantly among months. Leaves only once at the begin or the middle of the seasons to
292 assess the capacity of PM capturing may underestimate or overestimate it.

293 Overall, results suggested that leaf surface accumulated more PM than waxes. Although for
294 most species there was lower mass of PM deposited in epicuticular waxes. However, PM
295 accumulation in the wax fraction is very important because these particles are trapped
296 permanently (Song et al. 2015).

297 **Different Size-fraction Particles**

298 Different size fractions of airborne PM have different chemical composition and posed adverse
299 effects on human health (Luo et al. 2011; Li et al. 2017). For example, particulate matters were
300 responsible for respiratory, cardiovascular and others (Cai et al. 2017), especially fine particles
301 (Englert et al. 2004). To assess the capacity of plant leaves to capture different size fractions, the
302 present study separated quantification of PM into: fine (0.2–2.5 μm), coarse (2.5–10 μm), and
303 large ($> 10 \mu\text{m}$). The masses in each of these size fractions accumulated on leaf surface and in
304 waxes are presented in Fig. 2 and 3, respectively.

305 PM of different size fractions accumulated on leaf surface and in waxes differed among the
306 species. Non-significant differences were observed among the different size-fraction surface PM
307 accumulations on shrubs (Fig. 2). The averages of the three size fractions for surface PM

308 accumulations on shrubs were $31.5 \mu\text{g cm}^{-2}$ (81%), $4.7 \mu\text{g cm}^{-2}$ (12%) and $2.5 \mu\text{g cm}^{-2}$ (7%) for
309 large, coarse and fine PM, respectively.

310 Across all trees, *P. orientalis* ($44.4 \mu\text{g cm}^{-2}$) and *P. armandi* ($43.7 \mu\text{g cm}^{-2}$) showed the greatest
311 large particles fraction accumulation on the leaf surface. Large particles fraction accumulation on
312 *P. tomentosa* ($9.8 \mu\text{g cm}^{-2}$), *A. altissima* ($15.4 \mu\text{g cm}^{-2}$) and *S. japonica* ($12.7 \mu\text{g cm}^{-2}$) leaves
313 were significantly lower than the greatest accumulators. The average surface accumulation of
314 large particles fraction was $25.0 \mu\text{g cm}^{-2}$, which accounted for 75% of the total insoluble PM.

315 The average coarse fraction PM accumulated on leaf surfaces ($5.1 \mu\text{g cm}^{-2}$) comprised 15% of
316 the total insoluble PM. The greatest accumulation of coarse PM was found on *P. armandi* leaves
317 ($8.9 \mu\text{g cm}^{-2}$), which was weakly significant difference from accumulation by other trees. The
318 species with the lowest surface accumulation of the coarse fraction was on *P. tomentosa* ($1.8 \mu\text{g}$
319 cm^{-2}).

320 The greatest accumulation of fine fraction PM on the leaf surface was on *R. typhina* ($9.30 \mu\text{g}$
321 cm^{-2}), which was significantly greater than for other trees. *P. occidentalis* ($2.0 \mu\text{g cm}^{-2}$), *F.*
322 *pennsylvanica* ($1.2 \mu\text{g cm}^{-2}$), *P. tomentosa* ($1.0 \mu\text{g cm}^{-2}$), *G. biloba* ($2.1 \mu\text{g cm}^{-2}$) and *S. japonica*
323 ($1.6 \mu\text{g cm}^{-2}$) had considerably lower accumulation of fine fraction PM. The average fine PM
324 fraction surface accumulation was $3.5 \mu\text{g cm}^{-2}$, or 10% of the total insoluble PM deposition.

325 These present results showed that shrubs accumulated more mass of large size fraction PM on
326 leaf surface than most of the trees, except that *P. orientalis* and *P. amandi* which showed slightly
327 higher accumulation than that of shrubs. However, trees were more efficient than shrubs in
328 accumulating coarse and fine PM on the leaf surface.

329 There were no significant differences between the three particles size fraction accumulations in
330 the wax of shrub leaves. On average, the PM accumulation in wax on the shrub leaves was 6.8 μg
331 cm^{-2} (65%) for the large particles size fraction, 2.3 $\mu\text{g cm}^{-2}$ (21%) for the coarse size fraction and
332 1.5 $\mu\text{g cm}^{-2}$ (14%) for the fine particles size fractions. For the trees, *P. orientalis* (84.1 $\mu\text{g cm}^{-2}$)
333 and *P. armandi* (85.3 $\mu\text{g cm}^{-2}$) showed significantly greater accumulation of the large particles
334 size fraction in wax. The average accumulation of the large particles size fraction in wax on trees
335 was 22.0 $\mu\text{g cm}^{-2}$, which was 77% of the total accumulated PM. The highest coarse particles
336 accumulation in wax was for *P. orientalis* (10.9 $\mu\text{g cm}^{-2}$). The coarse particles in-wax deposition
337 on *P. armandi* (6.5 $\mu\text{g cm}^{-2}$) and *S. matsudana* (6.6 $\mu\text{g cm}^{-2}$) were marginally higher than for the
338 remaining trees. *F. pennsylvanica* (1.1 $\mu\text{g cm}^{-2}$) and *A. altissima* (0.9 $\mu\text{g cm}^{-2}$) had low deposition
339 of fine particles in waxes. The average coarse and fine particles size fraction accumulations in
340 wax were 3.8 $\mu\text{g cm}^{-2}$ and 2.5 $\mu\text{g cm}^{-2}$, which corresponded to 13% and 9% of the total in-wax
341 particles deposition. Particle size fraction is very similar for surface and in wax PM accumulated
342 by trees, but differs for shrubs showing amount of large PM on surface about 5 more than in wax,

343 which led to less fine and coarse PM fractions on surface. Dzierzanowski et al (2011) detected
344 a small shrub accumulated the largest amounts of PM as compared to trees, and more large PM
345 also deposited on surface than wax. As we explained above, shrubs growing low to the ground
346 were presumably more exposed to soil splash and traffic dust on the leaves than trees with an
347 upright growth habit, which were mostly large particles. However, these large PM deposited on
348 surface cannot be fixed by epicuticular wax in shrubs, which was significantly different with
349 trees.

350 Overall, PM depositions by mass for the three size fraction was in the order: large > coarse >
351 fine. The average proportions of large, coarse and fine PM were 73%, 16% and 11%. Previous
352 studies reported slightly different proportions of these three size fractions (Popek et al. 2013;
353 Song et al. 2015; Zhang et al. 2014), which may be due to the local ambient PM composition.
354 The proportions of the three size fractions were slightly different between species. This might be
355 explained by the leaf and structural characteristics of species (Song et al. 2015). However, as we
356 introduced that the average background PM₁₀ and PM_{2.5} concentrations in this area were 139 and
357 84 $\mu\text{g m}^{-3}$, respectively. The concentration of fine PM was higher than that of coarse particles,
358 which was different with fine and coarse PM deposited in leaf surface and waxes. Freer-smith et
359 al. (2005) found that sedimentation under gravity principally leads to large PM deposited on leaf
360 surfaces. Impaction and interception affects the deposition of coarse and fine PM on leaf surfaces.

361 Therefore, particles sedimentation is the main process by which PM deposits to foliage. The mass
362 of fine PM deposited by impaction and interception was $5.5 \mu\text{g cm}^{-2}$ on average, which
363 contributed only 11% of the total. However, the number of fine particles on leaves was large.
364 Previous studies proposed that the number of particles in the fine fraction contributed over 90%
365 of total insoluble PM (Li et al. 2015; Zhao et al. 2014). Therefore, plant also showed high
366 efficiently accumulation of fine PM in urban areas. The amount of coarse and fine PM
367 contributed 18% and 12% of total in-wax PM respectively. This ratio was higher than for
368 deposition to leaf surfaces. Shrub leaf surfaces accumulated greater mass of the three size
369 fractions PM than most of trees. Shrub and tree leaves grow at different heights, which may have
370 an effect on PM accumulations on leaf surface because there is more dust near the ground
371 (Dzierżanowski et al. 2011; A. Sæbø et al. 2012; Mo et al. 2015). The significant correlation
372 between the mass of PM size fraction was detected (R^2 for large, coarse and fine PM are 0.99 ($p <$
373 0.0001), 0.78 ($p < 0.0001$) and 0.52 ($p = 0.001$), respectively). So, species that accumulated more
374 total particles also have the high capability to capture greater amounts of fine particles, which is
375 the most dangerous for human health. The most efficient species of fine PM accumulation should
376 be used for urban greening, such as *R. typhina*, *P. orientalis*, *P. armandi*, and *M. denudate*. Tree-
377 dimensional configuration of the most efficient shrub and tree species can reduce PM pollution

378 on different spatial scales. We also suggest that conifer species should be priority in urban
379 greening, which show high capability in capturing PM during winter.

380 **Water-soluble ions on leaf surfaces**

381 The foliage analyzed for water-soluble ions was that sampled on July 27, 2014. Fig. 4 shows
382 the results for ions and total insoluble suspended particulate (TSP). The ratio in Fig. 4 represents
383 the percentage of water-soluble ions and the total of soluble and insoluble PM. Across all species,
384 the mass of ions on leaf surfaces ranged from 3.7 $\mu\text{g cm}^{-2}$ (in *M. denudata*) to 31.6 $\mu\text{g cm}^{-2}$ (in *P.*
385 *armandi*). There were significant differences between plant species, *U. pumila*, *S. matsudana*, *P.*
386 *orientalis* and *P. armandi* accumulating more ions on leaf surfaces ($p < 0.05$). For other species,
387 accumulations for all ions were low to 10 $\mu\text{g cm}^{-2}$. The mean value for ions deposition was 12.9
388 $\mu\text{g cm}^{-2}$. Significant positive correlations were observed between the amounts of ions and TSP
389 across species ($R^2=0.48$, $p=0.001$). The highest and lowest percentages of water-soluble ions to
390 total PM on leaves were 50% and 7% respectively, with an average of 28%. The proportions of
391 water-soluble ions in total PM varied greatly among different plant species, especially for trees.
392 Variations of proportion between species were also detected in the study of Freer-Smith et al.
393 (2005). The data show that the contributions of ions to total PM was in the order $\text{NO}_3^- > \text{Ca}^{2+} >$
394 $\text{SO}_4^{2-} > \text{Mg}^{2+} > \text{Cl}^- > \text{Na}^+ > \text{K}^+ > \text{NH}_4^+$. The proportion of NH_4^+ in the ions was lowest across all
395 species, which is surprising considering the ambient levels of NH_4^+ in Beijing. However, the
396 findings are consistent with Freer-Smith et al. (2005) who also showed that NO_3^- was the main

397 ion component and that the proportion of NH_4^+ was low. Cheng et al. (2016) reported average
398 concentrations of SO_4^{2-} , Cl^- and NO_3^- on plant leaves of $0.9098 \mu\text{g cm}^{-2}$, $0.7298 \mu\text{g cm}^{-2}$ and
399 $0.0878 \mu\text{g cm}^{-2}$, respectively, which are slightly different from this study. The difference may be
400 accounted for by the composition of inorganic compounds in the air. Shen et al. (2011) showed
401 ions comprised 33% PM_{10} at the Dongbeiwang experiment site, near to this study site, which was
402 relatively high compared with this study. The contribution of NH_4^+ to ions was 16%. Liu et al.
403 (2015) used ^{15}N tracer techniques to show that water-soluble ions (NH_4^+ and NO_3^-) in $\text{PM}_{2.5}$ can
404 be absorbed effectively by *P. euramericana* seedlings. Given that plant leaves absorb ultrafine
405 PM through their stomata (Bell and Treshow 2002), we postulate that NH_4^+ is translocated into
406 stomata which leads to the low contribution of NH_4^+ to ions on the leaf surface. The
407 concentration of NO_3^- was in the range of previous studies. In summary this study has shown that
408 water-soluble ions on leaf surfaces are also an important component of total PM deposited onto
409 plant leaves. As many past studies did not analyze the soluble PM components this means that the
410 interception capacity of plant species for PM has previously been undervalued. According to
411 studies on water soluble ions in atmosphere particles, without considering the mass of soluble
412 ions was slightly undervalued the effects of plants on PM deposition and may be a deviation in
413 the estimation of the PM accumulated by the whole trees (Song et al. 2015). In this study, the
414 percentages of water-soluble ions to total PM on coniferous leaves ranged from 24% to 29%.

415 Based on analysis of their data (Song et al. 2015), undervalued mass of water ions on conifer
416 species was about $38.59 \mu\text{g cm}^{-2}$ on average. This may have more serious impact on the whole
417 plant assessment. The elemental composition of the ions helps the identification the source of the
418 particles, which provides further important new information. In previous studies, the water-
419 soluble ions in PM in Beijing were classified into 4 sources (Sun et al. 2004; Wang et al. 2005):
420 NO_3^- , SO_4^{2-} and NH_4^+ were assigned to type 1, which comes from secondary inorganic aerosol;
421 Ca^{2+} and Mg^{2+} were to type 2, which comes from road and construction dusts; Na^+ and Cl^- is
422 assigned to type 3, which mainly come from waste incineration; and K^+ is assigned to type 4,
423 which mainly comes from biomass combustion. Figure 5 shows the average concentrations of
424 ions in each type. The greatest contribution to was from type 1 secondary aerosol; NO_2 and SO_2
425 from coal combustion and vehicle exhausts are converted to NO_3^- and SO_4^{2-} by gas-to-particles
426 reactions, which are mainly of anthropogenic origin. The contribution of Ca^{2+} and Mg^{2+} to total
427 PM were the next greatest and was significantly higher than contribution from type 3 and 4. Type
428 3 and 4 were low in our study site, which indicated the burning of waste and biomass. The
429 important feature of the data reported is that traffic was the main source of pollution for the
430 background air at this study site.

431
432 **Conclusion**
433

434 Seventeen urban plant species showed significantly different capacity in accumulating PM.
435 The average accumulation of PM on leaves was $58.9 \mu\text{g cm}^{-2}$, of which 65% was deposited on the
436 leaf surface and 35% in the leaf wax. The greatest PM accumulations on leaves occurred in *P.*
437 *orientalis* and *P. armandi*, both of which are evergreen conifer species. *F. pennsylvanica*, *P.*
438 *tomentosa*, *A. altissima* and *S. japonica* was less effective for PM deposition. *P. tomentosa*, *P.*
439 *orientalis* and *P. armandi* were the only three of seventeen species, which accumulated greater
440 than 50% of total insoluble PM in waxes. PM larger than $10 \mu\text{m}$ comprised 73% of PM deposited
441 on leaves, with the coarse and fine particles size fractions comprising 16% and 11% of the
442 deposited PM, respectively. Water-soluble ions comprised 28% of total PM on leaves on average.
443 Lack of knowledge about ions captured by leaves obviously leads to underestimation of the
444 ability of plant species to intercept PM. This study has shown that trees and shrubs should be
445 considered as an effective approach to remove aerial PM in urban areas.

446

447 **Acknowledgments**

448

449 This research was supported by the National Natural Science Foundation of China (41705130),
450 the National Key R&D Program of China (2017YFC0210106), the Forestry Public Welfare
451 Project of China (201304301), the Beijing Municipal Education Commission (CEFF-
452 PXM2017_014207_000043), and Beijing Laboratory Project (PXM2015-014207-000014).

453

454 **References**

455
456 Beckett KP, Freer-Smith PH, Taylor G (2000) Particulate pollution capture by urban trees: effect
457 of species and windspeed. *Glob. Change Biol* 6: 995-1003
458 Bell JNB, Treshow M (2002) *Air Pollution and Plant Life*. second ed. John Wiley, Chichester
459 Bosco ML, Varrica D, Dongarrà G (2005) Case study: Inorganic pollutants associated with
460 particulate matter from an area near a petrochemical plant. *Environ. Res* 99: 18-30
461 Cai MF, Xin ZB, Yu XX (2017) Spatio-temporal variations in PM leaf deposition: A meta-
462 analysis. *Environ. Pollut* 231: 207-218
463 Chen B, Li SN, Yang XB, Lu SW, Wang B, Niu X (2016a) Characteristics of atmospheric PM_{2.5}
464 in stands and non-forest cover sites across urban-rural areas in Beijing, China. *Urban*
465 *Ecosystems* 19: 867-883
466 Chen L, Liu C, Zou R, Yang M, Zhang Z (2016b) Experimental examination of effectiveness of
467 vegetation as bio-filter of particulate matters in the urban environment. *Environ. Pollut* 208:
468 198-208
469 Chen L, Liu CL, Pan T, Chen CC, Li Z, Wang HH, Pei X, Sun L (2014) Assessment of the effect
470 of PM_{2.5} reduction by plain afforestation project in Beijing based on dry deposition model.
471 *Chin. J. Appl. Ecol* 33: 2897-2904
472 Chen W, Tang H, Zhao H (2015) Diurnal, weekly and monthly spatial variations of air pollutants
473 and air quality of Beijing. *Atmos. Environ* 119: 21-34

- 474 Cheng Y, Wang Y, Wang Y, Zhang H, Liang D (2016) Particulate matter adsorption and the main
475 influencing factors of five typical plants in Beijing. *Environ. Chem* 35: 1690-1697
- 476 Dzierzanowski K, Popek R, Gawronska H, Saebo A, Gawronski SW (2011) Deposition of
477 particulate matter of different size fractions on leaf surfaces and in waxes of urban forest
478 species. *Int. J. Phytoremediat* 13: 1037-1046
- 479 Enqlert N (2004) Fine particles and human health--a review of epidemiological studies. *Toxicol.*
480 *Lett* 149: 235-242
- 481 Freer-Smith PH, Beckett KP, Taylor G (2005) Deposition velocities to *Sorbus aria*, *Acer*
482 *campestre*, *Populus deltoides* × *trichocarpa* ‘Beaupré’, *Pinus nigra* and × *Cupressocyparis*
483 *leylandii* for coarse, fine and ultra-fine particles in the urban environment. *Environ. Pollut* 133:
484 157-167
- 485 Gao G, Xu YS, Mo L, Yu XX (2016) The Ability of Plant Leaves on Depositing Size-fractionated
486 Particles. *Ecol. Environ. Sci* 25: 260-265
- 487 Han X, Guo QJ, Liu CQ, Strauss H, Yang JX, Hu J, Wei RF, Tian LY, Kong J (2016) Effect of
488 the pollution control measures on PM_{2.5} during the 2015 China Victory Day Parade:
489 Implication from water-soluble ions and sulfur isotope. *Environ. Pollut* 218: 230-241
- 490 Janhäll S (2015) Review on urban vegetation and particle air pollution – Deposition and
491 dispersion. *Atmos. Environ* 105: 130-137

- 492 Jun L, Han YQ, Tang X, Zhu J, Zhu T (2016) Estimating adult mortality attributable to PM2.5
493 exposure in China with assimilated PM2.5 concentrations based on a ground monitoring
494 network. *Sci. Total Environ* 568: 1253-1262
- 495 Jouraeva VA, Johnson DL, Hassett JP, Nowak DJ (2002) Differences in accumulation of PAHs
496 and metals on the leaves of *Tilia×euchlora* and *Pyrus calleryana*. *Environ. Pollut* 120: 331-338
- 497 Lelieveld J, Evans JS, Fnais M, Giannadaki D, Pozzer A (2015) The contribution of outdoor air
498 pollution sources to premature mortality on a global scale. *Nature* 525: 367-371
- 499 Leonard R J, Mcarthur C, Hochuli D F (2016) Particulate matter deposition on roadside plants
500 and the importance of leaf trait combinations. *Urban For. Urban Gree* 20: 249-253
- 501 Li D, Xu YX, Wang BT (2015) The Deposition of Particulate Matters on Climbing Plant. *Ecol.*
502 *Environ. Sci* 24: 1486-1492
- 503 Li HX, Ji HB, Shi CJ, Yang G, Yan Z, Xu XY, Ding HJ, Tang L, Xing YX (2017) Distribution of
504 heavy metals and metalloids in bulk and particle size fractions of soils from coal-mine
505 brownfield and implications on human health. *Chemosphere* 172: 505-515
- 506 Liu J, Han Y, Tang X, Zhu J, Zhu T (2016) Estimating adult mortality attributable to PM2.5
507 exposure in China with assimilated PM2.5 concentrations based on a ground monitoring
508 network. *Sci. Total Environ* 568: 1253-1262
- 509 Liu Q, Shi J, An HL, Cao XH, Liu C, Ying WL, Xia XL, Guo HH (2015) Absorption and

510 Distribution of PM_{2.5} NH₄⁺ and NO₃⁻ in *Populus euramericana* Neva.. Acta Ecol. Sin 35: 6541-
511 6548

512 Lun X, Zhang X, Mu Y, Nang A, Jiang G (2003) Size fractionated speciation of sulfate and nitrate
513 in airborne particulates in Beijing, China. Atmos. Environ 37: 2581-2588

514 Luo XS, Yu S, Li XD (2011) Distribution, availability, and sources of trace metals in different
515 particle size fractions of urban soils in Hong Kong: Implications for assessing the risk to
516 human health. Environ. Pollut 159: 1317-1326

517 Mo L, Ma ZY, Xu YS, Sun FB, Lun XX, Liu XH, Chen J, Yu XX (2015) Assessing the Capacity
518 of Plant Species to Accumulate Particulate Matter in Beijing, China. Plos One 10: e0140664

519 Nowak DJ, Hirabayashi S, Bodine A, Hoehn R (2013) Modeled PM_{2.5} removal by trees in ten
520 U.S. cities and associated health effects. Environ. Pollut 178: 395-402

521 Ottelé M, van Bohemen HD, Fraaij ALA (2010) Quantifying the deposition of particulate matter
522 on climber vegetation on living walls. Ecol. Eng 36: 154-162

523 Pope III CA, Dockery DW (2006) Health effects of fine particulate air pollution: lines that
524 connect. J. Air Waste Manage 56: 709-742

525 Popek R, Gawronska H, Wrochna M, Gawronski SW, Sæbø A (2013) Particulate matter on
526 foliage of 13 woody species: deposition on surfaces and phytostabilisation in waxes--a 3-year
527 study. Int. J. Phytoremediat 15: 245-256

528 Przybysz A, Sæbø A, Hanslin HM, Gawronski SW (2014) Accumulation of particulate matter and
529 trace elements on vegetation as affected by pollution level, rainfall and the passage of time.
530 Sci. Total Environ 481: 360-369

531 Sabin LD, Hee Lim J, Teresa Venezia M, Winer AM, Schiff KC, Stolzenbach KD (2006) Dry
532 deposition and resuspension of particle-associated metals near a freeway in Los Angeles.
533 Atmos. Environ 40: 7528–7538

534 Sæbø A, Popek R, Nawrot B, Hanslin HM, Gawronska H, Gawronski SW (2012) Plant species
535 differences in particulate matter accumulation on leaf surfaces. Sci. Total Environ 427–428:
536 347-354

537 Salmond JA, Tadaki M, Vardoulakis S, Arbuthnott K, Coutts A, Demuzere M, Dirks KN,
538 Heaviside C, Lim S, Macintyre H, McInnes RN, Wheeler BW (2016) Health and climate
539 related ecosystem services provided by street trees in the urban environment. Environ. Health
540 15: S36

541 Schaubroeck T, Deckmyn G, Neiryneck J, Staelens J, Adriaenssens S, Dewulf J, Muys B, Verheyen
542 K (2014) Multilayered modeling of particulate matter removal by a growing forest over time,
543 from plant surface deposition to washoff via rainfall. Environ. Sci. Technol 48: 10785-10794

544 Sgrigna G, Sæbø A, Gawronski S, Popek R, Calfapietra C (2015) Particulate Matter deposition on
545 Quercus ilex leaves in an industrial city of central Italy. Environ. Pollut 197: 187-194

- 546 Shen J, Tang A, Liu X, Kopsch J, Fangmeier A, Goulding K, Zhang F (2011) Impacts of Pollution
547 Controls on Air Quality in Beijing during the 2008 Olympic Games. *J. Environ. Qual* 40: 37-45
- 548 Song YS (2015) The study on fine particulate air pollution characteristics and the dust detention
549 effect of greening trees in Beijing. Dissertation, Research Center for Eco-Environmental
550 Sciences, Chinese Academy of Sciences.
- 551 Song YS, Maher B, Li F, Wang X, Sun X, Zhang H (2015) Particulate matter deposited on leaf of
552 five evergreen species in Beijing, China: Source identification and size distribution. *Atmos.*
553 *Environ* 105: 53-60
- 554 Song YS, Wang XK, Maher B, Li F, Xu C, Liu X, Sun X, Zhang Z (2016) The spatial-temporal
555 characteristics and health impacts of ambient fine particulate matter in China. *J. Clean. Prod*
556 112: 1312-1318
- 557 Sun Y, Zhuang G, Wang Y, Han L, Guo J, Dan M, Zhang W, Wang Z, Hao Z (2004) The air-borne
558 particulate pollution in Beijing—concentration, composition, distribution and sources. *Atmos.*
559 *Environ* 38: 5991-6004
- 560 Tallis M, Taylor G, Sinnott D, Freer-Smith P (2011) Estimating the removal of atmospheric
561 particulate pollution by the urban tree canopy of London, under current and future
562 environments. *Landscape Urban Plan* 103: 129-138
- 563 Tang X, Zhang X, Wang Z, Ci Z (2016) Water-soluble organic carbon (WSOC) and its

564 temperature-resolved carbon fractions in atmospheric aerosols in Beijing. *Atmos. Res* 181:
565 200-210

566 Thithanhthao N, Yu X, Zhang Z, Liu M, Liu X (2015) Relationship between types of urban forest
567 and PM_{2.5} capture at three growth stages of leaves. *J. Environ. Sci* 27: 33-41

568 Wang L, Gao S, Liu L, Ha S (2006) Atmospheric particle -retaining capability of eleven garden
569 plant species in Beijing. *Chin . J. Appl . Ecol* 17, 597-601

570 Wang L, Gong HL, Liao WB, Wang Z (2015) Accumulation of particles on the surface of leaves
571 during leaf expansion. *Sci. Total Environ* 532: 420-434 Wang L, Hasi E, Liu L, Gao S (2007)
572 Physico-chemical characteristics of ambient particles settling upon leaf surface of six conifers
573 in Beijing. *Chin. J. Appl. Ecol* 18: 487-492

574 Wang Y, Zhuang G, Tang A, Yuan H, Sun Y, Chen S, Zheng A (2005) The ion chemistry and the
575 source of PM_{2.5} aerosol in Beijing. *Atmos. Environ* 39: 3771-3784

576 WHO (2006) WHO air quality guidelines for particulate matter, ozone, nitrogen dioxide and
577 sulfur dioxide. Global Update 2005. Summary of Risk Assessment World Health Organization.
578 Geneva

579 Xie BZ, Wang HX, Yang J (2014) Retention capability of PM 2.5 and it's explanation by leaf
580 surface micro-structure of common broad-leaved plant species in Beijing. *Acta Bot. Boreal. -*
581 *Occident. Sin* 34: 2432-2438

- 582 Yan J, Lin L, Zhou W, Ma K, Pickett STA (2016) A novel approach for quantifying particulate
583 matter distribution on leaf surface by combining SEM and object-based image analysis.
584 Remote Sens. Environ 173: 156-161
- 585 Yin S, Shen Z, Zhou P, Zou X, Che S, Wang WH (2011) Quantifying air pollution attenuation
586 within urban parks: An experimental approach in Shanghai, China. Environ. Pollut 159: 2155-
587 2163
- 588 Zhang ZD, Xi BY, Cao ZG, Jia LM (2014) Exploration of a quantitative methodology to
589 characterize the retention of PM_{2.5} and other atmospheric particulate matter by plant leaves:
590 Taking *Populus tomentosa* leaves as an example. Chin. J. Appl. Ecol 25: 2238-2242
- 591 Zhao ST, Li XY, Li YM (2014) The Characteristics of deposition of airborne particulate matters
592 with different size on certain plants. Ecol. Environ. Sci 23: 271-276
593

594

595 **Table 1.** Species and allometric data for the trees sampled. (mean \pm SD, n = 3)

Classification	Species	Height (m)	Diameter at breast height (cm)	Crown diameter (m)
Shrubs	<i>Amygdalus triloba</i> (Lindl.) Ricker	2.1 \pm 0.4	7.7 \pm 1.4	2.8 \pm 0.2
	<i>Euonymus japonicus</i> Thunb.	1.9 \pm 0.2	5.0 \pm 1.1	2.4 \pm 0.2
	<i>Lonicera maackii</i> (Rupr.) Maxim.	3.2 \pm 0.3	10.8 \pm 3.4	5.6 \pm 0.4
	<i>Prunus Cerasifera</i> Ehrh	2.4 \pm 0.3	23.0 \pm 2.4	5.4 \pm 1.2
Trees	<i>Magnolia denudata</i> Desr.	10.5 \pm 0.4	23.6 \pm 3	6.8 \pm 1.7
	<i>Rhus typhina</i> Nutt.	3.6 \pm 0.2	8.8 \pm 1.3	2.3 \pm 1.1
	<i>Platanus occidentalis</i> L.	14.4 \pm 0.6	53.7 \pm 6.2	6.6 \pm 2.1
	<i>Fraxinus pennsylvanica</i> Marsh.	9.4 \pm 0.6	25.0 \pm 2.1	7.9 \pm 2.6
	<i>Populus tomentosa</i> Carr.	13.4 \pm 0.7	51.9 \pm 3.6	5.3 \pm 0.9
	<i>Ginkgo biloba</i> L.	14.5 \pm 0.5	53.5 \pm 5.6	5.8 \pm 1.4
	<i>Ulmus pumila</i> L.	8.4 \pm 0.3	77.1 \pm 7.4	6.8 \pm 2.3
	<i>Salix matsudana</i> Koidz.	12.6 \pm 0.4	34.4 \pm 5.8	6.9 \pm 2.3
	<i>Pinus tabulaeformis</i> Carr.	10.0 \pm 0.8	29.4 \pm 5.3	5.6 \pm 0.8
	<i>Platyclusus orientalis</i> (L.) Franco	6.1 \pm 0.1	16.1 \pm 2.3	1.7 \pm 0.6
	<i>Pinus armandi</i> Franch.	5.4 \pm 0.6	16.5 \pm 1.6	3.8 \pm 0.8
	<i>Ailanthus altissima</i> (Mill.) Swingle	9.8 \pm 0.5	27.8 \pm 3.5	4.6 \pm 0.7
<i>Sophora japonica</i> L.	10.7 \pm 0.3	24.5 \pm 0.3	3.4 \pm 0.5	

596

597

Figure Captions

598

599 **Fig. 1.** Total amount of PM accumulated on leaf surfaces and in-wax. Data are mean \pm SE, n = 18.

600 Bars marked with different letters are significantly different ($p < 0.05$).

601 **Fig. 2.** The sum of PM accumulation of three size fractions on leaf surfaces. Data are mean \pm SE,

602 n=18. Bars marked with different letters are significantly different ($p < 0.05$).

603 **Fig. 3.** The sum of PM accumulation of three size fractions in waxes. Data are mean \pm SE, n=18.

604 Bars marked with different letters are significantly different ($p < 0.05$).

605 **Fig. 4.** Total insoluble suspended particulate and water-soluble ions accumulated on leaf

606 surfaces. The ratio of ions to TSP. Data are mean \pm SE, n =3.

607 **Fig. 5.** The concentrations of dissolvable inorganic ions presented as the sum mass of four types

608 ions. Data are mean \pm SE, n=3.

609 **Fig. 6.** Monthly change of PM accumulation on leaf surface between shrubs and trees. Data are

610 mean \pm SE, n=3. Bars marked with different letters are significantly different ($p < 0.05$).

611 **Fig. 7.** Monthly change of PM accumulation in waxes between shrubs and trees. Data are mean

612 \pm SE, n=3. Bars marked with different letters are significantly different ($p < 0.05$).

613

614


615

616

617

618

619



620

621

622

623

624

625

Fig. 1.


626

627

628

629

630



631

632

633

634

635

636

Fig. 2.


637

638

639

640

641



642

643

644

645

646

647

Fig. 3.


648

649

650

651

652



653

654

655

656

657

658

Fig. 4.


659

660

661

662

663



664

665

666

667

668

669

670

Fig. 5.


671

672

673

674

675



676

677

678

679

680

681

Fig. 6.


682

683

684

685

686



687

688

689

690

691

692

Fig. 7.