



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Army imposters

Citation for published version:

Perez-Espona, S, Goodall-Copestake, WP, Berghoff, SM, Edwards, KJ & Franks, NR 2017, 'Army imposters: diversification of army ant-mimicking beetles with their Eciton hosts' *Insectes sociaux*, vol. 65, no. 1, pp. 59-75. DOI: 10.1007/s00040-017-0588-1

Digital Object Identifier (DOI):

[10.1007/s00040-017-0588-1](https://doi.org/10.1007/s00040-017-0588-1)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Insectes sociaux

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1 **Army imposters: diversification of army ant-mimicking beetles with their *Eciton* hosts**

2 Sílvia Pérez-Espona^{1,2*}, William P. Goodall-Copetake³, Stefanie M. Berghoff^{1,4}, Keith J. Edwards¹, Nigel R.
3 Franks¹

4 1. The University of Bristol, School of Biological Sciences, 24 Tyndall Avenue, Bristol BS8 1TH, United
5 Kingdom

6 2. Estación Biológica de Doñana, CSIC, Isla de La Cartuja, Américo Vespucio s/n, Sevilla E41092, Spain

7 3. British Antarctic Survey, High Cross, Madingley Road, Cambridge CB3 0ET, United Kingdom

8 4. Max Planck Institute of Neurobiology, Am Kopferspitz 18, 82152 Martinsried, Germany

9 * Corresponding author: Sílvia Pérez-Espona, e-mail: s.perezespona@gmail.com

10 Short running title: Diversification of myrmecophiles with hosts

11 **Abstract**

12 Colonies of neotropical army ants of the genus *Eciton* Latreille offer some of the most captivating examples of
13 intricate interactions between species, with hundreds of associated species already described in colonies of *Eciton*
14 *burchellii* Westwood. Among this plethora of species found with *Eciton* colonies, two genera of staphylinid
15 beetles, *Ecitomorpha* Wasmann, and *Ecitophya* Wasmann, have evolved to mimic the appearance and parallel the
16 colouration of the most abundant ant worker cast. Here, we study for the first time the association of these ant-
17 mimicking beetles with their ant host in an evolutionary and population genetics framework. The central emphasis
18 is on colonies of *E. burchellii*, the only *Eciton* species that harbours both genera of ant-mimicking beetles.
19 Phylogenetic and population structure analyses using the same mtDNA COI region (802bp) for ants and beetles
20 indicated that speciation patterns of the myrmecophiles were congruent with specialization to a particular *Eciton*
21 (sub)species. Therefore, current taxonomic treatments of *Eciton* and its *Ecitomorpha* and *Ecitophya* associates
22 need revision. Molecular clock analyses suggested that diversification of the *Eciton* hosts pre-date that of their
23 guests, with a possible earlier association of *Ecitophya* (found with a large number of *Eciton* species) than with
24 *Ecitomorpha* (found only with *E. burchellii* colonies). Population-level analyses revealed that patterns of
25 diversification for the myrmecophiles are also consistent with specialisation to a particular host across broad
26 geographical areas but not at small geographical scales, with gene flow within each species found between host
27 colonies even across landscape features that are strong barriers for *Eciton* female-mediated gene flow.

28 **Keywords**

29 Gene flow; mimicry; mitochondrial DNA; myrmecophily; myrmecophory; population structure; speciation;
30 taxonomy

31 **Introduction**

32 The study of associations between species is vital if we are to understand the evolution of biological diversity
33 (Thompson, 2013). This was recognised by Darwin in the closing paragraph of *On the Origin of Species by Means*
34 *of Natural Selection* (1859) when he used the term ‘entangled bank’ to refer to the interaction between species that
35 form biological communities, as highlighted by Thompson (1994). Nature is full of examples of intricate and
36 intimate associations between different species but it is in the world of ants where some of the most numerous and
37 astonishing associations can be found (Hölldobler and Wilson, 1990). Among the ants, army ants harbour the most
38 extensive array of species associations, with multiple vertebrate and invertebrate associate species exploiting the
39 ants and the different environments and homeostatic conditions that their colony life creates (Gotwald Jr., 1995;
40 Hughes et al., 2008). Of all army ants, those of the genus *Eciton*, inhabiting the tropics of the New World, are the
41 ones exhibiting the most captivating display of associates (Gotwald Jr., 1995), with 557 species already recorded
42 with *Eciton burchellii* and many more still to be described (Ivens et al., 2016; Rettenmeyer et al., 2011). Associates
43 found with *Eciton* army ants include, among others, mites that feed on secretions and hemolymph of the ants; flies
44 that feed on the middens’ refuse; beetles that steal prey from the ants or predate on the ants or their brood; the
45 iconic army-ant-following birds that feed on arthropod prey flushed out by the ants during their raids; and
46 butterflies that feed on droppings from ant-following birds (Gotwald Jr., 1995; Kistner, 1979; Rettenmeyer et al.,
47 2011; Schneirla, 1971)

48
49 As part of this plethora of associates, some have evolved to conceal their presence among the army ants through
50 chemical, tactile and morphological mimicry (Gotwald Jr., 1995; Kistner, 1979). These ‘imposters’ have evolved
51 different strategies to associate with army ants, with staphylinid beetles having mastered the art of blending in
52 with the ants by evolving to resemble the appearance of their hosts (Hölldobler and Wilson, 1990; Kistner and
53 Jacobson, 1990; Maruyama and Parker, 2017). Two genera of myrmecomorph (ant-like) staphylinid beetles,
54 *Ecitomorpha* and *Ecitophya*, both in the subfamily Aleocharinae (tribe Athetini; Elven, Bachmann, & Gusarov,
55 2012), are found with neotropical army ants of the genus *Eciton*. These two genera of beetles are highly specialized
56 to the epigeaic patterns and nomadic life of *Eciton* army ants and both mimic the most abundant worker cast in
57 *Eciton* colonies (Seevers, 1965); the media workers (Franks, 1985). Both genera present similar morphological
58 modifications that confer resemblance to their host: similarity in surface sculpturing, subpetiolate and ellipsoidal
59 abdomens, slender heads and pronota, and long appendages (Seevers, 1965). Their appearance is considered
60 moderately ant-like as they are not a perfect mimic of the ants, but their colour parallels that of the species or

61 subspecies of *Eciton* with which they are associated (Akre and Rettenmeyer, 1966; Kistner and Jacobson, 1990;
62 Seevers, 1965). The two genera are mainly distinguished by *Ecitophya* presenting a much slender body and longer
63 appendages than *Ecitomorpha*, with a head more than twice as long as wide, a bilobed mentum, slender gula with
64 sutures not converging in the front, longer antennae (more than six times as long as the head width) but with the
65 terminal segments of the antennae not much wider than the preceding segments (terminal segments in *Ecitomorpha*
66 are more club-shaped) (Seevers, 1965).

67
68 The mimicry of *Ecitomorpha* and *Ecitophya* to *Eciton* media workers is likely to be both an adaptive response to
69 avoid predators such as ant-following birds (Batesian mimicry), as well as an adaptation for integration into the
70 ant colony and avoidance of host aggression - Wasmannian mimicry (Parker, 2016). The presence of *Ecitomorpha*
71 and *Ecitophya* in *Eciton* colonies is rare (and sometimes absent), with many colonies presenting less than one of
72 these beetles per 1,000 worker ants. These two myrmecophile genera are considered hunting guests of *Eciton* army
73 ants, as they are found running among ants in raiding columns where they feed on dropped prey or at booty caches
74 (Kistner and Jacobson, 1990). These beetles are also found in emigration columns (Akre and Rettenmeyer, 1966;
75 Kistner and Jacobson, 1990) when the conspicuously nomadic *Eciton* colonies move to another location to set up
76 their new bivouac (temporary nest). During *Eciton* colonies emigration, *Ecitomorpha* and *Ecitophya* individuals
77 run in the centre of the columns or ride on prey captured by the ants or on ant pupae (Kistner and Jacobson, 1990).
78 Therefore, they are adapted to the movement and life cycle of their host (Akre and Rettenmeyer, 1966), as has
79 been shown for other *Eciton* myrmecophiles (Berghoff et al., 2009; Von Beeren et al., 2016a, 2016b).

80
81 Taxonomically, the genus *Ecitophya* was initially divided into five species acknowledging the colour parallel
82 between this myrmecophile and its *Eciton* hosts (Reichensperger, 1933). *Ecitophya rapaxae* Mann found
83 associated with the ant *Eciton rapax* Smith, *Ecitophya consecta* Mann associated with *Eciton vagans* Olivier,
84 *Ecitophya gracillima* Mann that is associated with *Eciton hamatum* Fabricius, and *Ecitophya simulans* Wasmann
85 and *Ecitophya bicolor* Reichensperger associated with *E. burchellii*. The latter two species were later grouped into
86 a single species, *E. simulans*, as it was considered that specimens of *Ecitophya* collected with *E. burchellii* colonies
87 did not differ sufficiently to be considered as separate species (Kistner and Jacobson, 1990; Seevers, 1965).
88 Another species found associated with *Eciton lucanoides* Emery was later described as *Ecitophya rettenmeyeri*
89 (Kistner and Jacobson, 1990).

90 *Ecitomorpha* beetles have only been found with the army ant *Eciton burchellii*, with the first specimens of
91 *Ecitomorpha* described by Wasmann in 1889 as *Ecitomorpha arachnoides* (Akre and Rettenmeyer, 1966).
92 Reichensperger divided this genus into four species taking into account the colour polymorphism within *Eciton*
93 *burchellii*: *Ecitomorpha arachnoides* Wasmann, *Ecitomorpha nevermanni* Reichensperger, *Ecitomorpha*
94 *breviceps* Reichensperger and *Ecitomorpha melanotica* Mann (Reichensperger, 1935, 1933). However, due to the
95 difficulty in finding consistent morphological characters (besides colouration) supporting the separation of these
96 species, they were subsequently lumped back into a single species, *Em. arachnoides* (Kistner and Jacobson, 1990;
97 Seevers, 1965).

98
99 In this study, we investigated for the first time in an evolutionary and population genetics framework the interaction
100 of *Ecitomorpha* and *Ecitophya* ant-mimicking beetles with their *Eciton* hosts; in particular, *E. burchellii*, as it is
101 the only *Eciton* species known to host both genera of beetles. Genetic analyses of ants and beetles collected in
102 Panama, a geographical area where many different *Eciton* species have overlapping ranges (Watkins, 1976), were
103 conducted to test the following hypotheses: (i) considering the strong level of association of *Ecitophya* and
104 *Ecitomorpha* with *Eciton*, phylogenetic patterns of the myrmecophiles will mirror that of their host, (ii) due to
105 the dependence of these two genera on *Eciton*'s hunted prey and the pedestrian dispersal capability of the queen
106 and workers, the level of specificity between host and myrmecophile will be observable at broad geographical
107 scales, (iii) if myrmecophiles are truly host-specific and have evolved and diverged via increased specification on
108 a particular *Eciton* host, molecular patterns should support earlier taxonomic classifications of *Ecitophya* and
109 *Ecitomorpha* by Reichensperger (i.e. each *Eciton* species will host a particular *Ecitophya* and *Ecitomorpha*
110 species).

111

112 **Methods**

113 **Study area and sampling**

114 Sampling for this study was targeted on colonies of *E. burchellii* ssp. *foreli* Mayr and *E. b.* ssp. *parvispinum* Forel,
115 the two most-studied *E. burchellii* subspecies. These two subspecies are highly epigaeic and their distribution
116 ranges overlap in Panama, Costa Rica, and Honduras (Watkins, 1976). Descriptions of these species highlight their
117 morphological similarity (Borgmeier, 1955; Santschi, 1925), with main differences reported being the colouration
118 of media workers, *E. b. foreli* having black head and mesosoma but reddish metasoma, and *E. b. parvispinum*'s
119 media workers having complete black bodies. However, studies assessing the genetic differences between these

120 two subspecies in view of their current taxonomic treatment have yet to be conducted. This study focused on
121 sampling of colonies in Panama, an area where the geographical range of both subspecies partly overlaps. Three
122 main areas of Panama were sampled (Fig. 1); in West Panama the Bosque Protector Palo Seco (BPPS) and the
123 adjacent Reserva Forestal Fortuna (RFF), and in Central Panama the Área Protegida San Lorenzo and its buffer
124 zone (APSL). As the Chagres River was found to be a barrier for *E. b. foreli* when gene flow was estimated with
125 mtDNA markers (Pérez-Espona et al., 2012), this area was divided into two (APSLA and APSLB) to group
126 colonies from each side of the Chagres River. In total, 13 colonies of *E. b. foreli* (4 in BPPS, 6 in APSLA and 3
127 in APSLB) and 12 colonies of *E. b. parvispinum* (all in RFF) for which we found associated *Ecitophya* and/or
128 *Ecitomorpha* beetles were sampled (Fig. 1; Table 1). In addition, four colonies of *E. hamatum* and the associated
129 *Ep. gracillima*: 1 colony in RFF, 2 colonies in BPPS, and 1 in Soberanía National Park (SOB), and one colony of
130 *E. lucanoides* and its associated *Ep. rettenmeyeri* collected in RFF were opportunistically sampled. The number
131 of *Ecitophya* and *Ecitomorpha* beetles sampled and sequenced from *E. b. foreli* and *E. b. parvispinum* colonies
132 are summarized in Table 1. Although the number of beetles collected is not directly comparable between colonies,
133 as effort spent searching and collecting the beetles was constrained by the time of the day a colony was
134 encountered, the maximum number of beetles collected in a colony was 78 *Em. arachnoides* (colony E114
135 collected in RFF with *E. b. parvispinum*) and 40 *Ep. simulans* (colony E89 collected in BPPS with *E. b. foreli*).

136

137 The sampling protocol for our study consisted of walking all available trails and adjacent less accessible areas
138 (off trails), through daily extensive walks (9 a.m. until dusk), to collect individuals from as many colonies of *E. b.*
139 *foreli* and *E. b. parvispinum* as possible. Once an *E. burchellii* colony was encountered, workers from all castes
140 were sampled from raid or emigration columns by removing them with the help of long forceps. Ant columns were
141 then carefully observed at several points of the raid or emigration columns to sample *Ecitophya* and *Ecitomorpha*
142 beetles with straight tube aspirators. Ant column observations per colony lasted several hours or until dusk,
143 depending on the time of the day when a colony was encountered. Collections were simultaneously conducted by
144 two people in order to maximise sampling of the beetles. All samples were preserved in 99% ethanol for further
145 examination and subsequent genetic studies.

146

147 **Laboratory procedures**

148 *DNA extraction*

149 Abdomens of ants and beetles were carefully dissected before DNA extractions to avoid any contamination from
150 consumed prey. To facilitate DNA extraction, the tissue samples were deposited in a 1.5mL Eppendorf tube and
151 briefly immersed in liquid nitrogen prior to extraction procedures. Genomic DNA was extracted using the DNeasy
152 tissue kit (QIAGEN) following the manufacturer's instructions.

153

154 *Sequencing of the mitochondrial marker Cytochrome Oxidase subunit I*

155 Ants and ant-mimicking beetles from both genera, *Ecitomorpha* and *Ecitophya*, were sequenced for the same
156 region of the mitochondrial cytochrome oxidase subunit I gene (COI, *coxI*). Mitochondrial DNA markers are the
157 most widely used genetic markers in species-level phylogenies and DNA barcoding studies. Due to their haploid
158 nature, maternal inheritance, and smaller population size, sequences derived from mtDNA coalesce over a shorter
159 time scale than those derived from nuclear DNA (Simon et al., 2006), with reciprocal monophyly at the species
160 level reached faster after speciation in mtDNA phylogenies than in nuclear DNA phylogenies (Sunnucks, 2000).
161 The faster mutation rate of mtDNA markers has been shown to offer more powerful resolutions of relationships
162 between closely related taxa in phylogeographic and population-level studies (Avise, 2000; Zhang and Hewitt,
163 2003). In insects, mutation rates of mtDNA markers have been estimated to be 2 to 9 times faster than nuclear
164 protein-coding genes (Moriyama and Powell, 1997) making them more suitable for the study of closely related
165 species that have diverged recently (Lin and Danforth, 2004). COI fragments were amplified from the *Eciton*
166 samples using a modified version of the primer pairs CI13/CI14 (Hasegawa et al., 2002) and Ben/Jerry (Simon et
167 al., 1994). Details of the modified primers and Polymerase Chain Reactions (PCR) conditions can be found in
168 Pérez-Espona *et al.* (2012). Fragments of COI were amplified from *Ecitomorpha* and *Ecitophya* using the primer
169 pairs C1-J-1634/C1-N-2317 and C1-J-2216/C2-N-3431 (Maus et al., 2001). These beetle PCR amplifications were
170 conducted in a total volume of 25µL, using 10-15ng of template DNA, 1X NH4 buffer, 2.5mM MgCl₂, 0.6µM of
171 each primer, 1 unit of BIOTAQ polymerase (Bioline, London) and double processed tissue culture distilled water
172 (Sigma-Aldrich, Buchs, Switzerland) to bring the volume up to 25µL. The PCR cycling protocol included an initial
173 denaturation step of 94°C for 3 min, a three-step cycling consisting of a denaturing step of 94°C for 30 s, annealing
174 at 51°C for 30 s and ramping at 0.3°C/s to an extension step of 72°C for 1 min. The cycle was repeated 29 times
175 and was followed by a final extension of 72°C for 10 min. PCR products were run on a 1.5% agarose gel and
176 visualised using ethidium bromide staining. Successful amplifications were purified using EXOSAP (GE
177 Healthcare), and forward and reverse strands for each of the fragments sequenced in two reactions using 6µL of
178 purified PCR product, 4µL of the reaction mix DYEnamic ET Terminator Cycle Sequence Kit (Amersham, GE

179 Healthcare) and 0.5 μ L of primer. Cycle sequencing consisted of 25 cycles including a denaturation step of 95°C
180 for 20 s, an annealing step of 50°C for 15 s and an extension step of 60°C for 1 min. Sequences were run on a
181 MegaBACE™ 1000 capillary sequencer (Amersham GE Healthcare) at The University of Bristol.
182 Electropherograms from the forward and reverse sequencing reads were edited and assembled into contigs using
183 the software Geneious version 10 (Biomatters: <http://www.geneious.com>). The resulting consensus reads from
184 each individual were sorted into unique haplotypes and subsequently manually aligned in Geneious, together with
185 some additional sequences obtained from four individuals of *Eciton dulcium* Forel (collected in RFF and APSL)
186 and sequences obtained from GenBank (Accession numbers: AY233691-4, AY233696, GQ980948). These
187 additional sequences were selected as ingroup placeholders and outgroups based on the army ant phylogeny of
188 Brady (2003) and the army ant-mimicking beetle phylogeny in Maruyama and Parker (2017). The resulting
189 alignments were all trimmed to include the same region COI fragment (802bp). All unique sequences were
190 submitted to DDBJ under the accession numbers LC258007-LC258019 for the *Eciton* sequences, and LC258020-
191 LC258064 for the *Ecitophya* and *Ecitomorpha* sequences.

192

193 *Phylogenetic and molecular clock analyses*

194 Unique haplotype alignments for *Eciton* and *Ecitomorpha* with *Ecitophya* were initially evaluated for nucleotide
195 compositional heterogeneity using the Chi-square test in PAUP version 4.0b10 (Swofford, 2002), and using
196 tetrahedral plots and matched-pairs tests for symmetry implemented in SeqVis version 1.5 (Ho et al., 2006). The
197 more conservative Chi-square test provided no significant evidence for compositional heterogeneity in either of
198 the alignments. The more sensitive tetrahedral plots and matched-pairs tests for symmetry also provided no strong
199 evidence for compositional heterogeneity in the *Ecitophya-Ecitomorpha* alignment; however, there was evidence
200 for some heterogeneity in the *Eciton* alignment: tetrahedral plots contained co-dispersed clusters of data points
201 and the number of the matched-pairs tests for symmetry was >5% at P = 0.05 and >1% at P = 0.01. To explore
202 whether data re-coding could reduce the level of compositional heterogeneity in the *Eciton* alignment, C and T
203 nucleotides were re-coded as Ys. SeqVis analysis of this re-coded DNA alignment revealed no significant evidence
204 for heterogeneity. Therefore, in addition to analyzing the full *Eciton* DNA alignment, we also analyzed a data
205 reduced AGY form of the *Eciton* alignment to account for artefacts that may arise from compositional
206 heterogeneity.

207

208 Phylogenies were generated from the *Eciton* and *Ecitophya-Ecitomorpha* DNA alignments using Maximum
209 Parsimony and Bayesian methods. For the Maximum Parsimony analysis, heuristic searches were conducted using
210 PAUP, with the full *Eciton* and *Ecitophya-Ecitomorpha* alignments and the AGY re-coded *Eciton* alignment. Each
211 heuristic search started from a random tree with 50 random addition replicates, one tree was held per step, saved
212 trees set to a maximum of 10,000 and other settings left at default values. Confidence values were generated using
213 non-parametric bootstrapping (10,000 replicates). For the Bayesian inference, the DNA alignments were analysed
214 as a single partition using a mixed nucleotide substitution model and gamma corrected rate heterogeneity across
215 sites with the software MrBayes version 3.2.6 (Ronquist et al., 2012). Parameter distributions were approximated
216 using reversible jump Metropolis-coupled Markov chain Monte Carlo methods, with three chains of 10,000,000
217 generations, chain heating at 0.05, sampling frequency at 1,000 and other settings at default values. Posterior
218 samples of parameter estimates were assessed using generation plots, distribution plots, the potential scale
219 reduction convergence diagnostic and estimated sample sizes as recommended in the MrBayes manual. These
220 revealed that the total number of generations and default burnin of 25% appeared to be sufficient to acquire final
221 parameter estimates from a stationary distribution.

222

223 To analyse the *Eciton* alignment under AGY coding using Bayesian inference, we used the three-state 'AGY'
224 model implemented in the software mcmphase within the PHASE version 3.0 software package
225 (<https://github.com/james-monkeyshines/rna-phase-3>). Heterogeneity across sites was modelled using a gamma
226 correction, and a chain length of 10,000,000 iterations, sampling period at 1,000 and burnin of 25% were used, as
227 in the MrBayes analyses. Perturbation proposal priorities were 10 for the tree and 1 for the substitution model.
228 Within these tree and model components, the proposal priorities were 1 for topology changes, 10 for branch lengths
229 (with an exponential (10) prior), 1 for frequencies, 1 for rate ratios, and 1 for the gamma parameter. Generation
230 versus log probability plots, parameter distribution plots, and repeated analyses starting from different random
231 seeds indicated that these settings generated final posterior estimates from a stationary distribution. Tree files
232 generated using mcmphase were analysed using the associated program mcmsummarize and also Geneious.

233

234 Chronological estimates for key diversification events were obtained with additional Bayesian analyses conducted
235 using MrBayes. For these analyses, we reduced the level of haplotype sampling in order to include only the most
236 abundant haplotypes and key biogeographic placeholders. This was to ameliorate the impact of intra-specific
237 differences in haplotype sampling between the *Eciton* and *Ecitomorpha-Ecitophya* datasets that could have a

238 negative impact on the date estimation procedure. The molecular clock analyses used the same underlying model
239 and chain settings as described for MrBayes above with an outgroup-ingroup division enforced as a strong prior
240 topological constraint. Two different methods of clock calibration were explored: a standard 1% per million years'
241 rate applied to both the *Eciton* and *Ecitomorpha-Ecitophya* datasets, and fixed date calibrations of 26 million years
242 on the most recent common ancestor (MRCA) of *Eciton* following the results obtained by Brady (2003), and a
243 date of 25 million years on the *Ecitomorpha-Ecitophya* divergence following the results in Maruyama and Parker
244 (2017). Based on the data in Brady (2003) and Maruyama and Parker (2017), a uniform tree age prior of 10-100
245 million years was used. Clock model options were explored using stepping-stone sampling estimates of the
246 marginal likelihood. The Thorne-Kishino 2002 'TK02' relaxed clock had the smallest log likelihood but this was
247 less than 5 units better than the independent gamma rates 'IGR' relaxed clock and strict clock in both the *Eciton*
248 and *Ecitophya-Ecitomorpha* datasets. This preliminary analysis thus provided no strong evidence in favor of either
249 of these alternative clock models and, consequently, all three were used to generate date estimates. Posterior
250 samples of parameter estimates were assessed for stationarity as described above.

251

252 *Population-level analyses of E. burchellii and associates*

253 Genetic diversity analyses for *E. b. foreli* and *E. b. parvispinum* colonies and their associated *Ecitophya* and
254 *Ecitomorpha* beetles were conducted using the softwares Arlequin version 3.1 (Excoffier et al., 2005) and DnaSP
255 version 5 (Librado & Rozas, 2009). Genetic diversity was estimated in terms of number of haplotypes, segregating
256 sites (S) and average number of nucleotide differences (k). In order to compare divergence within and between
257 species, S and k were also calculated for the different *Eciton*, *Ecitophya* and *Ecitomorpha* species included in our
258 study. Population structure for the host and each of the associates was estimated using a hierarchical analysis of
259 molecular variance (AMOVA) with the software popART (<http://popart.otago.ac.nz>). The partitioning of genetic
260 variation was assessed within and among two main geographical areas, West Panama (BPPS) and Central Panama
261 (APSLA, APSLB), and significance values obtained after 1,000 permutations. The software popART was also
262 used to build Median Joining haplotype networks (epsilon = 0) to assess haplotype relationships and identify
263 patterns of haplotype structure for *E. burchellii* and the myrmecophile beetles at different geographical scales.

264

265 **Results**

266 *Diversification and species relationships*

267 Phylogenetic and molecular clock analyses provided further insights into the evolution and diversification of the
268 *Eciton* species and their *Ecitophya* and *Ecitomorpha* beetle associates (Figs. 2 & 3, Table 2 & 3). Analyses using
269 the full and reduced datasets using different methods (i.e. standard and AGY coded, Parsimony and Bayesian
270 analyses) resolved the same suites of well-supported haplotype relationships; this indicated that these were robust
271 to the nucleotide compositional heterogeneity found in the *Eciton* DNA alignment. With the *Eciton* dataset,
272 haplotypes of *E. dulcium* and *E. hamatum*, and the subspecies *E. b. foreli*, were grouped according to these
273 taxonomic categories with strong statistical support ($\geq 99\%$ Parsimony bootstrap support, Bayesian posterior
274 probabilities of 1.0). The subspecies *E. b. foreli* and *E. b. parvispinum* were also resolved as a single group but
275 with lower support ($\geq 60\%$ bootstrap, ≥ 0.91 posterior probability). Other basal relationships between *Eciton* taxa
276 included in this study were poorly supported in all phylogenetic analyses. In the *Ecitophya-Ecitomorpha* analyses,
277 haplotypes assigned to *Ecitomorpha* and *Ecitophya* were clearly separated into these genera level categories (\geq
278 99% bootstrap, posterior probability of 1.0). The *Em. arachnoides* haplotypes formed two well-supported groups
279 that corresponded to their *E. burchellii* host ($\geq 98\%$ bootstrap, posterior probability of 1.0); *E. b. foreli* and *E. b.*
280 *parvispinum*. The *Ep. simulans* haplotypes also consisted of two well-supported groups following their *E.*
281 *burchellii* host (100% bootstrap, posterior probabilities of 1.0). However, these two *Ep. simulans* groups formed
282 a complex with two groups of *Ep. gracillima* (associated with *E. hamatum*). Within this complex, *Ep. simulans*
283 (associated with *E. b. foreli*) was clearly grouped with *Ep. gracillima* (87% bootstrap, posterior probability of
284 0.95). *Ep. rettenmeyeri* (associated with *E. lucanoides*) was separated from the other *Ecitophya* species with 92%
285 parsimony bootstrap and 0.78 posterior probability support. Lineage relationships within *Ecitophya*, therefore, did
286 not mirror that of their *Eciton* host. The number of segregating sites (S) and the average pairwise nucleotide
287 differences (*k*) between species revealed further insights into the taxonomy and species relationships of the
288 myrmecophiles and their hosts (Table 3). Levels of divergence estimated as S and *k* between *E. b. foreli* and *E. b.*
289 *parvispinum* were similar (or slightly higher) to those between taxa currently recognised as separate *Eciton* species.
290 Although speciation patterns of *Ecitophya* did not mirror those of the hosts, a strong divergence was observed
291 between *Ecitomorpha* or *Ecitophya* associated with each subspecies of *E. burchellii*. Divergence of *Ecitomorpha*
292 was only assessed for two *E. burchellii* subspecies so further subspecies would need to be studied to elucidate
293 further speciation patterns.
294
295 The molecular clock analyses recovered identical sets of *Eciton* and *Ecitophya-Ecitomorpha* haplotype
296 relationships but with different associated date estimates depending on the clock model and calibration method

297 used (Table 2). Analyses calibrated using fixed dates taken from the literature were consistently older than date
298 estimates calibrated using a 1% rate, while differences arising from the use of different clock models were subtler.
299 Regardless of the clock model or calibration method, the relative age differences between *Eciton* and *Ecitophya*-
300 *Ecitomorpha* lineage divergences were similar in all molecular clock analyses (Table 2, Figure 3). Although the
301 confidence intervals for the divergence estimates were wide and therefore we should be cautious with
302 interpretations, median estimates of divergences indicated that the diversification of the genus *Eciton* is likely to
303 be older than that of the associated myrmecophile genera *Ecitophya* and *Ecitomorpha*. This was the case when
304 considering date estimates for the MRCA of *Eciton* species taken from literature (used as a fixed 26 MYA
305 calibration date) and when considering independently obtained date estimates derived from the 1% calibration rate
306 (Table 2, Fig 3). Median estimates for diversification of the main *Eciton* species (or subspecies) also appears to
307 pre-date the diversification of the main myrmecophile lineages (Table 2, Fig 3). Poor branch support for most of
308 the basal *Eciton* relationships (Fig 2, Fig 3) may, in part, be due to the rapid diversification of these primary *Eciton*
309 lineages following the MRCA, as other studies using nuclear markers and phylogenomics have also failed to
310 unambiguously resolve these species relationships (see Discussion). Key among the main *Eciton* diversification
311 events was the MRCA of *E. b. parvispinum* and *E. b. foreli*, which pre-dated the diversification of the associated
312 myrmecophiles *Em. arachnoides* and the *Ep. simulans* – *Ep. gracillima* complex (Table 2, Fig 2, Fig 3). Among
313 the two genera of myrmecophiles, median estimates for the diversification of *Ecitophya*, which is found with more
314 species of *Eciton*, appear to be older than those for *Ecitomorpha*, although confidence intervals did overlap for
315 these estimates (Table 2, Fig 3). The genus *Ecitophya* has been reported in colonies of *E. rapax*, *E. vagans*, *E.*
316 *burchellii*, *E. hamatum* and *E. lucanoides* while *Ecitomorpha* has only been reported for *E. burchellii*.

317

318 *Population-level analyses of E. burchellii and its Ecitophya and Ecitomorpha associates*

319 The diversification of the ant-mimicking beetles *Ecitophya* and *Ecitomorpha* with *E. burchellii* was further
320 confirmed by the haplotype networks (Fig. 4). The networks clearly indicated a strong divergence between *E. b.*
321 *foreli* and *E. b. parvispinum* haplotypes, with 90 segregating sites among haplotypes of both subspecies (Fig. 4).
322 This divergence was mirrored in the associated beetles; however, the divergence between haplotypes in the beetles
323 was smaller than that found for the host (55 segregating sites in *Ep. simulans* and 50 segregating sites in *Em.*
324 *arachnoides*).

325

326 The haplotype networks confirmed that the diversification of *Ecitophya* and *Ecitomorpha* with *E. b. foreli* was
327 also prevalent over broad geographical areas. Haplotypes of ants and beetles collected in West Panama (BPPS)
328 markedly differed from those collected in Central Panama (APSLA and APSLB). Divergence time estimates for
329 the separation of haplotype lineages between West and Central Panama were older for *E. b. foreli* than for the
330 associate *Ep. simulans* (Fig. 3, Table 2). Although the haplotypes of *Em. arachnoides* found in the two main
331 geographical study areas were clearly distinct (Fig. 4), with haplotypes derived from two distinct lineages found
332 in West Panama (BPPS), the phylogenetic analyses did not support a strong grouping of haplotypes according to
333 these geographical areas (Fig. 2). In the Central Panama study area (APSL) the Chagres River was a strong gene
334 flow barrier for *E. b. foreli* females, with median estimates of divergence of *E. b. foreli* haplotypes either side of
335 the river estimated at 0.4-1.6 MYA (Fig. 3, Table 2). The Chagres River, however, was not a gene flow barrier for
336 either *Ecitomorpha* or *Ecitophya*, as haplotypes characteristic of a particular species of these myrmecophiles were
337 shared between their *Eciton* host colonies either side of the river (Fig. 4).

338

339 Estimates of population structure were higher for *E. b. foreli* than for the associated *Ecitophya* and *Ecitomorpha*
340 beetles (Table 4). In the host, genetic differentiation was found at all levels of analyses, with the majority of the
341 genetic variation explained by differences among groups (76.31%), and hardly any variation explained due to
342 differences within populations (0.20%). In *Ecitophya* and *Ecitomorpha*, most of the genetic variation was also
343 explained by differences among groups (84.62% and 79.57%, respectively) but genetic variation due to differences
344 within populations (15.51% and 22.03%, respectively) was much larger than that found for the host. Negative
345 values in variation among populations within groups found in *Ecitophya* and *Ecitomorpha* indicate the large
346 haplotype variation within populations and the sharing of haplotypes between these. No genetic variation was
347 attributed to differences among populations collected on either side of the river within each of the beetle species,
348 further corroborating that the Chagres River was not a gene flow barrier for these myrmecophiles.

349

350 **Discussion**

351 *Taxonomy of Eciton and associated Ecitophya and Ecitomorpha*

352 The phylogenetic analyses and haplotype networks generated for this study revealed that current taxonomic
353 treatments of *E. burchellii* and of its *Ecitophya simulans* and *Ecitomorpha arachnoides* associates need further
354 revision to take into account further speciation within these taxa. Morphological descriptions of *E. b. parvispinum*
355 have highlighted its black colouration, reduced spines in the metanotum and barely protruding epinotum teeth

356 (Borgmeier, 1955; Santschi, 1925), but these characteristics were considered not to warrant species status for *E.*
357 *b. parvispinum*. However, our results clearly indicate that genetic divergence between *E. b. foreli* and *E. b.*
358 *parvispinum* is as high as (or higher than) between taxa currently considered separate *Eciton* species. Divergence
359 estimates calculated in terms of *S* and *k* further suggest this speciation, as values between *E. b. foreli* and *E. b.*
360 *parvispinum* were higher than the those obtained between *E. hamatum* – *E. dulcium* and *E. hamatum* – *E.*
361 *lucanoides*; therefore, indicating that taxonomic treatments of *E. burchellii* need to be revised. Our results coupled
362 with those from a recent study phylogeographic study of this species in Central America (Winston et al., 2017),
363 provide strong evidence that speciation within this taxon has long been underestimated.

364
365 Although divergence between the *Eciton* species, and within *E. burchellii*, is clear from our genetic analyses,
366 phylogenetic relationships between *Eciton* species were not fully resolved in our study, despite the suitability and
367 widespread use of COI for resolving relationships between closely related insect taxa (Lin and Danforth, 2004;
368 Sunnucks, 2000). Weak statistical support for relationships between *Eciton* species does not appear solely reflect
369 our choice of COI as a molecular marker, because combined analyses of COI and nuclear markers (Brady, 2003),
370 and even a recent phylogenomic approach (Winston et al., 2017), also failed to unambiguously resolve these *Eciton*
371 species relationships. Rather, the lack of resolution in phylogenetic trees reported here is more likely indicative of
372 a rapid diversification of lineages within *Eciton* (Whitfield and Lockhart, 2007).

373
374 Important insights into the taxonomy of *Ecitophya* and *Ecitomorpha* were also revealed by our genetic analyses.
375 The taxonomy of these myrmecophiles has been challenging due to the subtle differences observed in
376 morphological characters, besides colouration, within each of these genera (Kistner and Jacobson, 1990; Seevers,
377 1965). The *Ecitomorpha* species key by Reichensperger was based on head dimensions, thoracic plates
378 sculpturing, length of the hind tibiae and depressions behind the eyes (Reichensperger, 1933). Further differences
379 between the species were highlighted in each of the species descriptions; these included different antennae
380 formation, elytra sculpturing, and colouration in different parts of the body of the specimens. Species
381 characterisation according to colouration was also indicated in previous descriptions such as that of *Em.*
382 *melanotica* found with *E. b. parvispinum* (Mann, 1926). Kistner and Jacobson (1990), after analysing a large series
383 of specimens, concluded that variation in colour was not consistent; however, they reported that black specimens
384 of *Ecitomorpha* were more frequently found in geographical areas where *E. b. parvispinum* was present, and rarer
385 in areas populated by *E. b. foreli*. In terms of other morphological features, Kistner and Jacobson (1990) did not

386 find consistency either in the shape of the groove of the pronotum or in spermathecas in the male genitalia that
387 would allow species differentiation. They, therefore, considered that early classifications of the *Ecitomorpha*
388 species by Reichensperger could not be supported. For *Ecitophya*, early species classification by Reichensperger
389 (1935, 1933) were based on the general appearance of specimens, measurements and characteristics of the head
390 and legs, differences in antennal segments, elytra sculpturing, presence and characteristics of abdominal bristles,
391 and the colouration of different body parts. In their analysis of *Ecitophya* specimens, that included dissections of
392 male genitalia, Kistner and Jacobson (1990) concluded that only some characteristics of the abdominal bristles,
393 the relative measurements of size of body parts and the relative length of antennal segments were reliable
394 characters for species identification. This reduction in the number of reliable characters to distinguish species led
395 Kistner and Jacobson (1990) to lump some of the earlier species described by Reichensperger into a single species,
396 *E. simulans*.

397

398 Our study has shown the importance of the use of genetic studies for resolving taxonomic challenges and has
399 provided strong evidence of speciation of *Ecitomorpha* and *Ecitophya* as a result of their specialisation to their
400 *Eciton* hosts. The speciation patterns found in our study for both myrmecophile genera (i.e. specialisation to an
401 *Eciton* host) would further support that *E. b. foreli* and *E. b. parvispinum* may be considered separate species. The
402 diversification patterns obtained for *Ecitophya* and *Ecitomorpha* were more concordant with initial species
403 classifications by Reichensperger (1935, 1933) . Divergence estimated as S and k further supported the speciation
404 of *Ecitomorpha* and *Ecitophya* according to their host, with values between *Ep. simulans* found with *E. b. foreli*
405 or *E. b. parvispinum* differing by a similar number of segregating sites than between *Ep. simulans* found with *E.*
406 *b. foreli* and those found with *Ep. gracillima* (found with *E. hamatum*). Therefore, the current taxonomic treatment
407 of these *Eciton burchellii* associates as single species, i.e. *Ep. simulans* and *Em. arachnoides*, merits revision to
408 take into account the speciation patterns revealed by our genetic data. For *Ecitomorpha* beetles found associated
409 with *E. b. parvispinum* we suggest the adoption of the previous name of *Em. melanotica* Mann. Given our results
410 and those from Winston et al. (2017) supporting speciation of *E. burchellii* in Central America, we expect that
411 further species of *E. burchellii* and its *Ecitomorpha* and *Ecitophya* associates will be reported, as morphological
412 diversification of *E. burchellii* within its broad distributional range has long been acknowledged (Kistner and
413 Jacobson, 1990; Reichensperger, 1935, 1933; Seevers, 1965).

414

415 The lack of consistent and reliable morphological characters to differentiate species within *Ecitomorpha* and
416 *Ecitophya* reported by Kistner and Jacobson (1990) cannot be explained simply in terms of the recent and rapid
417 diversification within these genera, as indicated by our genetic analyses. In myrmecophiles, it is expected that
418 morphological variation between species may be reduced or absent due to strong selection on morphological,
419 behavioural and physiological characters as adaptations to exploit their hosts (Schonrogge et al., 2002). Often,
420 species are determined by visual inspection of morphological features; however, chemical and acoustic characters
421 can be more important for species recognition in arthropods. Future studies of *Ecitophya* and *Ecitomorpha*
422 diversification, therefore, will benefit from thorough analyses of chemical and acoustic characters, as these cues
423 are likely to be very important for their adaptation to their *Eciton* hosts (Lenoir et al., 2001).

424

425 *Diversification of Ecitophya and Ecitomorpha with Eciton species*

426 Phylogenetic patterns of *Ecitophya* and *Ecitomorpha* indicated specialization of these myrmecophiles with their
427 *Eciton* hosts, confirming previous observations of ant-resemblance and colouration parallels (Akre and
428 Rettenmeyer, 1966; Kistner and Jacobson, 1990; Reichensperger, 1933), and behavioural observations such as the
429 preference of following trails of the host species (Akre and Rettenmeyer, 1968). However, the patterns of
430 speciation of the myrmecophiles did not mirror those of the host revealing that the beetles' phylogenies were not
431 an 'evolutionary print' of the host (Thomas et al., 1996). Furthermore, phylogenetic analyses and estimates of S
432 and *k* disagreed with previously reported evolutionary relationships between *Ecitophya* species based on
433 morphological characters (Kistner and Jacobson, 1990). *Ep. simulans* found with colonies of *E. b. foreli* were more
434 closely related to *Ep. gracillima* (associated with *E. hamatum*) than to the *Ep. simulans* found with *E. b.*
435 *parvispinum*. Additionally, *Ep. rettenmeyeri* (associated with *E. lucanoides*) was not closely related to *Ep.*
436 *gracillima* as previously inferred by the similar colouration between the ant hosts (Kistner and Jacobson, 1990);
437 in fact, *Ep. rettenmeyeri* was sister to the other *Ecitophya*.

438

439 Confidence intervals for time divergence estimates overlapped slightly between the ants and the myrmecophiles
440 and, therefore, we cannot discard with certainty potential coevolutionary processes between hosts and guests.
441 However, median time divergence estimates indicated that diversification patterns of the ants probably pre-dates
442 that of the associated beetles. We acknowledge that our divergence estimates are derived from a single genetic
443 marker and therefore we focus our discussion on *Eciton* and myrmecophile divergences on relative (rather than
444 absolute) date estimates. Comparisons of median time divergence estimates from the molecular clock analyses for

445 both beetle genera suggest that diversification of *Ecitophya* with *Eciton* ants is likely to have occurred earlier than
446 for *Ecitomorpha*. A possible earlier diversification of *Ecitophya*, and subsequently a longer time frame to fine-
447 tune their interaction with their host, could explain the association of *Ecitophya* with a larger number of *Eciton*
448 species.

449

450 Myrmecophily in Staphylinidae beetles is an ancient phenomenon, with a fossil of *Protoclaviger trichodens* gen.
451 et sp. nov. (Clavigeritae) in amber being dated to the Early Eocene (c. 52 MYA; Parker and Grimaldi, 2014). This
452 early association of Clavigeritae beetles with ants can explain the remarkable diversity of myrmecophilous species
453 within this supertribe (Parker and Grimaldi, 2014). High species diversification due to myrmecophily has also
454 been reported for ant-nest beetles of the genus *Paussus* L. (subfamily Paussinae; Moore and Robertson, 2014).
455 Although new species of *Ecitophya* and *Ecitomorpha* are likely to be described, due to underestimated speciation
456 within *E. burchellii* (and maybe in other *Eciton* species), the diversification of these two Aleocharinae genera is
457 not as exceptional as that found for other myrmecophile Staphylinidae such as *Paussus*. This difference probably
458 reflects the high specialisation of *Ecitophya* and *Ecitomorpha* to the genus *Eciton*, in contrast to *Paussus* beetles
459 that are found associated with different, and sometimes distantly unrelated, ant genera (Moore and Robertson,
460 2014). Furthermore, our results suggest that contrary to other myrmecophile beetles found associated with *Eciton*
461 colonies (e.g. *Vatesus*), *Ecitophya* and *Ecitomorpha* species have evolved more host-specific adaptations, probably
462 as a result of stronger selection pressures because they are hunting guests of day-time raiding epigeagic army ants.

463

464 *Geographical patterns of diversification of Ecitophya and Ecitomorpha with E. b. foreli*

465 Broad phylogeographic patterns of *Ecitophya* and *Ecitomorpha* were concordant with their *E. b. foreli* host, with
466 a clear separation of haplotypes between the West and Central Panama study areas. However, patterns of
467 diversification of the host and these two myrmecophiles differed at smaller geographical scales, indicating that
468 local differences might be the result of the spatial distribution of the host and the capability of dispersal of the
469 myrmecophiles (Tack and Roslin, 2010; Thompson, 2005). The haplotype networks revealed that *Ecitophya* and
470 *Ecitomorpha* beetles are not colony- or ant-mtDNA lineage-specific even though vertical transmission of
471 myrmecophiles is likely to occur during colony fission (Schneirla, 1971). Ant colonies with different mtDNA
472 haplotypes shared myrmecophile haplotypes, indicating that horizontal transmission of *Ecitophya* and
473 *Ecitomorpha* mtDNA lineages occurs between colonies. Horizontal transmission of myrmecophiles between
474 *Eciton* colonies has been previously reported for *Vatesus* (Akre and Torgerson, 1969), a beetle that does not mimic

475 the physical appearance of the ants but most of its life cycle is tightly linked to that of its host (Von Beeren et al.,
476 2016a). Horizontal transmission of *Ecitophya* and *Ecitomorpha* individuals, as a potential strategy to avoid
477 inbreeding within a colony, might occur by dispersal through flight between colonies such as shown for *Vatesus*
478 (Chatzimanolis et al., 2004; Von Beeren et al., 2016a), or whenever colony fusion - the aggregation of workers to
479 another colony after losing their queen (Schneirla, 1940; Schneirla & Brown, 1950) - takes place (Kronauer et al.,
480 2010). In our study, gene flow between the myrmecophile populations of each species was also found across the
481 Chagres River despite this being a major gene flow barrier for *E. b. foreli* females (Pérez-Espona et al., 2012).
482 Gene flow across the Chagres River, therefore, indicates a higher dispersal capability of the beetles in contrast to
483 the obligate pedestrian dispersal of *Eciton* queen and worker ants. Reports of flight in these beetles have only been
484 anecdotal and limited to observations of hovering of *Ep. consecta* over a colony of *E. vagans* when this colony
485 was spreading to attack the observer (Mann, 1921), and an individual of *Ep. gracillima* found with a colony of *E.*
486 *hamatum* (Pérez-Espona pers. obs.). In the latter case, hovering was observed when trying to aspirate one
487 individual from a raiding column. This specimen hovered to seek refuge under some fallen leaves but after a few
488 minutes tried to follow the ant trail (when it was successfully collected and included in this study). Fully developed
489 wings with venation characteristic of staphylinids have been described for both *Ecitophya* and *Ecitomorpha*
490 (Kistner and Jacobson, 1990). However, due to the close dependence of these beetles with their specific host and
491 the relatively low density of colonies, at least for *E. burchellii* (Franks, 1982), dispersal of the myrmecophiles
492 between colonies, in particular those located at further distance or separated by landscape features that act as
493 barriers for *Eciton* female dispersal, might be challenging. In such scenarios, it is likely that dispersal between
494 colonies is mediated by large and alate *Eciton* males when they leave their natal colony in search of conspecific
495 colonies to find a queen for mating (Gotwald Jr., 1995; Schneirla, 1971). Males of *E. b. foreli* are produced in
496 large numbers (c. 3,000) as part of sexual broods (including a small number of queens) when colony fission is
497 imminent (Franks and Hölldobler, 1987; Gotwald Jr., 1995; Schneirla, 1971). Males have been shown to be strong
498 fliers and able to disperse over 1km distances (Jaffé et al., 2009). They are therefore responsible for the majority
499 of gene flow between colonies (Berghoff et al., 2008; Jaffé et al., 2009; Pérez-Espona et al., 2012; Soare et al.,
500 2014). Dispersal of myrmecophiles by ant alate reproductives have been shown in *Atta* leaf-cutting ants, with
501 *Attafila* cockroaches observed on ant queens departing for nuptial flights (Moser, 1967), and similar strategies
502 have been suggested for the movement of myrmecophiles between *Eciton* colonies (Kronauer et al., 2010).
503

504 Ecological and evolutionary studies of myrmecophiles of army ants at the population-level are still in their infancy
505 due to the difficulty of keeping the host and myrmecophiles in laboratory conditions (Kistner and Jacobson, 1990).
506 However, as shown in this and recent studies in *Vatesus* and *Tetradonia* beetles (Von Beeren et al., 2016a, 2016b),
507 genetic approaches can shed light on the evolution and levels of specificity of these army ant imposters. Our study
508 was based on variation in mtDNA therefore future studies aiming to further elucidate the level of association of
509 *Ecitomorpha* and *Ecitophya*, in particular at small geographical scales, would benefit through the use of highly
510 variable markers such as microsatellites. Using a combination of mitochondrial and microsatellite data in previous
511 studies has been demonstrated that deforestation has a major impact on the connectivity of *E. burchellii*
512 populations (Pérez-Espona et al., 2012; Soare et al., 2014), threatening the long-term persistence not only of these
513 top neotropical predators but also the multitudes of species associated with them. Further research on *Eciton* and
514 their associates is crucial if we are to provide conservation solutions that would guarantee the maintenance of this
515 manifestation of Darwin's 'entangled bank'.
516

517 **Acknowledgements**

518 Autoridad Nacional del Ambiente (ANAM), Smithsonian Tropical Research Institute (STRI) and Centro de
519 Estudios de Acción Social Panameña (CEASPA) are thanked for arranging permits and fieldwork activities. We
520 thank J. Domínguez and F. García for assistance in the field, Carles Molina Rubio for laboratory assistance, John
521 N. Thompson for discussion of results, and two anonymous referees for comments on a previous version of the
522 manuscript. Cambridge Conservation Forum and Cambridge Conservation Initiative are thanked for allowing
523 Sílvia Pérez-Espona to use their office space in the David Attenborough Building while writing this manuscript.
524 Samples for this study were collected and exported under the permits SEX/A-8-08 and SEX/A-24-08, respectively.
525 This study was funded by The Leverhulme Trust (project grant F/00 182/AI).

526 **References**

- 527 Akre, R.D., Rettenmeyer, C.W., 1968. Trail-following by guests of army ants (Hymenoptera: Formicidae:
528 Ecitonini). *J. Kansas Entomol. Soc.* 41, 165–174.
529 Akre, R.D., Rettenmeyer, C.W., 1966. Behavior of Staphylinidae associated with army ants (Formicidae:
530 Ecitonini). *J. Kansas Entomol. Soc.* 39(4), 745–782.
531 Akre, R.D., Torgerson, R.L., 1969. Behavior of *Vatesus* beetles associated with army ants (Coleoptera:
532 Staphylinidae). *Pan-Pac. Entomol.* 45, 269–281.
533 Avise, J.C., 2000. *Phylogeography: The History and Formation of Species*. Cambridge, MA: Harvard University
534 Press, Cambridge, MA.
535 Berghoff, S.M., Kronauer, D.J.C., Edwards, K.J., Franks, N.R., 2008. Dispersal and population structure of a
536 New World predator, the army ant *Eciton burchellii*. *J. Evol. Biol.* 21(4), 1125–1132.
537 doi:<http://dx.doi.org/10.1111/j.1420-9101.2008.01531.x>
538 Berghoff, S.M., Wurst, E., Ebermann, E., Sendova-Franks, A.B., Rettenmeyer, C.W., Franks, N.R., 2009.
539 Symbionts of societies that fission: mites as guests or parasites of army ants. *Ecol. Entomol.* 34, 684–695.
540 doi:[10.1111/j.1365-2311.2009.01125.x](http://dx.doi.org/10.1111/j.1365-2311.2009.01125.x)

541 Borgmeier, T., 1955. Die Wanderameisen der neotropischen Region (Hym. Formicidae). Stud. Entomol. 3, 1–
542 716.

543 Brady, S.G., 2003. Evolution of the army ant syndrome: the origin and long-term evolutionary stasis of a
544 complex of behavioral and reproductive adaptations. Proc. Natl. Acad. Sci. U. S. A. 100, 6575–6579.
545 doi:10.1073/pnas.1137809100

546 Brady, S.G., Ward, P.S., 2005. Morphological phylogeny of army ants and other dorylomorphs (Hymenoptera:
547 Formicidae). Syst. Entomol. 30, 593–618. doi:10.1111/j.1365-3113.2005.00290.x

548 Chatzimanolis, S., Ashe, J.S., Hanley, R.S., 2004. Diurnal/nocturnal activity of rove beetles (Coleoptera:
549 Staphylinidae) on Barro Colorado Island, Panama assayed by flight intercept trap. Coleopt. Bull. 58, 569–
550 577. doi:10.1649/689.1

551 Darwin, C., 1859. On the Origin of the Species by Means of Natural Selection, or the Preservation of Favoured
552 Races in the Struggle for Life. John Murray, London. doi:10.1016/S0262-4079(09)60380-8

553 Elven, H., Bachmann, L., Gusarov, V.I., 2012. Molecular phylogeny of the Athetini-Lomechusini-Ecitocharini
554 clade of aleocharine rove beetles (Insecta). Zool. Scr. 41, 617–636. doi:10.1111/j.1463-6409.2012.00553.x

555 Excoffier, L., Laval, G., Schneider, S., 2005. Arlequin (version 3.0): an integrated software package for
556 population genetics data analysis. Evol. Bioinform. Online 1, 47–50. doi:10.1111/j.1755-
557 0998.2010.02847.x

558 Franks, N.R., 1985. Reproduction, foraging efficiency and worker polymorphism in army ants, in: Hölldobler,
559 B., Lindauer, M., Franks, N.R. (Eds.), Experimental Behavioral Ecology and Sociobiology : In Memoriam
560 Karl von Frisch, 1886-1982. pp. 91–107. doi:citeulike-article-id:7111831

561 Franks, N.R., 1982. A new method for censusing animal populations: The number of *Eciton burchelli* army ant
562 colonies on Barro Colorado Island, Panama. Oecologia 52, 266–268. doi:10.1007/BF00363847

563 Franks, N.R., Hölldobler, B., 1987. Sexual competition during colony reproduction in army ants. Biol. J. Linn.
564 Soc. 30, 229–243. doi:10.1111/j.1095-8312.1987.tb00298.x

565 Gotwald Jr., W.H., 1995. Army Ants: the Biology of Social Predation. Cornell University Press.

566 Hasegawa, E., Tinaut, A., Ruano, F., 2002. Molecular phylogeny of two slave-making ants: *Rossomyrmex* and
567 *Polyergus* (Hymenoptera : Formicidae). Ann. Zool. Fennici 39, 267–271.

568 Ho, J.W.K., Adams, C.E., Lew, J. Bin, Matthews, T.J., Ng, C.C., Shahabi-Sirjani, A., Tan, L.H., Zhao, Y.,
569 Eastal, S., Wilson, S.R., Jermini, L.S., 2006. SeqVis: Visualization of compositional heterogeneity in
570 large alignments of nucleotides. Bioinformatics 22, 2162–2163. doi:10.1093/bioinformatics/btl283

571 Hölldobler, B., Wilson, E.O., 1990. The Ants. Harvard University Press.

572 Hughes, D.P., Pierce, N.E., Boomsma, J.J., 2008. Social insect symbionts: evolution in homeostatic fortresses.
573 Trends Ecol. Evol. 23, 672–677. doi:10.1016/j.tree.2008.07.011

574 Ivens, A.B.F., von Beeren, C., Blüthgen, N., Kronauer, D.J.C., 2016. Studying the complex communities of ants
575 and their symbionts using ecological network analysis. Annu. Rev. Entomol. 61, 353–371.
576 doi:10.1146/annurev-ento-010715-023719

577 Jaffé, R., Moritz, R.F.A., Kraus, F.B., 2009. Gene flow is maintained by polyandry and male dispersal in the
578 army ant *Eciton burchellii*. Popul. Ecol. 51, 227–236. doi:10.1007/s10144-008-0133-1

579 Kistner, D.H., 1979. Social and evolutionary significance of social insect symbionts, in: Hermann, H.R. (Ed.),
580 Social Insects. Academic Press, New York, p. 1:339-413.

581 Kistner, D.H., Jacobson, H.R., 1990. Cladistic analysis and taxonomic revision of the ecitophilous tribe
582 Ecitocharini with studies of their behavior and evolution (Coleoptera, Staphylinidae, Aleocharinae).
583 Sociobiology 17, 333–480.

584 Kronauer, D.J.C., Schöning, C., D’Ettorre, P., Boomsma, J.J., 2010. Colony fusion and worker reproduction
585 after queen loss in army ants. Proc. R. Soc. B Biol. Sci. 277, 755–763. doi:10.1098/rspb.2009.1591

586 Lenoir, A., Ettorre, P.D., Errard, C., 2001. Chemical ecology and social parasitism in ants. Annu. Rev. Entomol.
587 46, 573–599.

588 Lin, C.-P., Danforth, B.N., 2004. How do insect nuclear and mitochondrial gene substitution patterns differ?
589 Insights from Bayesian analyses of combined datasets. Mol. Phylogenet. Evol. 30, 686–702.

590 Mann, W.M., 1926. New neotropical myrmecophiles. Jour. Wash. Acad. Sci. 16, 448–455.

591 Mann, W.M., 1921. Three new myrmecophilous beetles. Proc. United States Natl. Museum 59, 547–552.

592 Maruyama, M., Parker, J., 2017. Deep-time convergence in rove beetle symbionts of army ants. Curr. Biol. 27,
593 920–926.

594 Maus, C., Peschke, K., Dobler, S., 2001. Phylogeny of the genus *Aleochara* inferred from mitochondrial
595 cytochrome oxidase sequences (Coleoptera: Staphylinidae). Mol. Phylogenet. Evol. 18, 202–216.
596 doi:10.1006/mpev.2000.0874

597 Moore, W., Robertson, J.A., 2014. Report explosive adaptive radiation and extreme phenotypic diversity within
598 ant-nest beetles. Curr. Biol. 24, 2435–2439. doi:10.1016/j.cub.2014.09.022

599 Moriyama, E.N., Powell, J.R., 1997. Synonymous substitutions rates in *Drosophila* mitochondrial versus nuclear
600 genes. J. Mol. Evol. 45, 378–391.

601 Moser, J.C., 1967. Mating activities of *Atta texana* (Hymenoptera, Formicidae). *Insectes Soc.* 14, 295–312.
602 doi:10.1007/BF02252831

603 P, L., J, R., 2009. DnaSP v5: A software for comprehensive analysis of DNA polymorphism data.
604 *Bioinformatics* 25, 1451–1452.

605 Parker, J., 2016. Myrmecophily in beetles (Coleoptera): Evolutionary patterns and biological mechanisms.
606 *Myrmecological News* 22, 65–108.

607 Parker, J., Grimaldi, D.A., 2014. Specialized myrmecophily at the ecological dawn of modern ants. *Curr. Biol.*
608 24, 2428–2434. doi:10.1016/j.cub.2014.08.068

609 Pérez-Espona, S., McLeod, J.E., Franks, N.R., 2012. Landscape genetics of a top neotropical predator. *Mol.*
610 *Ecol.* 21, 5969–5985. doi:10.1111/mec.12088

611 Reichensperger, A., 1935. Beitrag zur Kenntnis der Myrmecophilenfauna Brasiliens und Costa Ricas III. (Col.
612 Staphyl. Hist.). *Arb. iiber Morphol. Taxon. Entomol. aus Berlin-Dahlem* 2, 188–218.

613 Reichensperger, A., 1933. Ecitophilen aus Costa Rica (II), Brasilien und Peru (Staph. Hist. Clavig.). *Rev.*
614 *Entomol.* 3, 179–194.

615 Rettenmeyer, C.W., Rettenmeyer, M.E., Joseph, J., Berghoff, S.M., 2011. The largest animal association
616 centered on one species: The army ant *Eciton burchellii* and its more than 300 associates. *Insectes Soc.* 58,
617 281–292. doi:10.1007/s00040-010-0128-8

618 Ronquist, F., Teslenko, M., Van Der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B., Liu, L., Suchard,
619 M.A., Huelsenbeck, J.P., 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice
620 across a large model space. *Syst. Biol.* 61, 539–542. doi:10.1093/sysbio/sys029

621 Santschi, F., 1925. Nouveaux Formicides brésiliens et autres. *Extr. des Ann. Bull. la Soc. Entomol. Belgique* 65,
622 221–247.

623 Schneirla, T., 1971. *Army ants: a study in social organization.* Freeman, WH, San Francisco.

624 Schneirla, T., 1940. Further studies on the army-ant behavior patterns: mass organization in the swarm-raiders. *J.*
625 *Comp. Psychol.* 29, 401–460.

626 Schneirla, T., Brown, R.Z., 1950. Army ant life and behavior under dry season conditions. 4. Further
627 investigation of cyclic processes in behavioral and reproductive functions. *Bull. Am. Museum Nat. Hist.*
628 95, 263–354.

629 Schonrogge, K., Barr, B., Wardlaw, J.C., Naf, E., Gardner, M.G., Breen, J., Elmes, G.W., Thomas, J.A., Olg,
630 P.H., Bassett, S., East, C., 2002. When rare species become endangered: cryptic speciation in
631 myrmecophilous hoverflies. *Biol. J. Linn. Soc.* 75, 291–300.

632 Seevers, C.H., 1965. The systematics, evolution and zoogeography of staphylinid beetles associated with army
633 ants (Coleoptera, Staphylinidae). *Fieldiana Zool.* 47, 137–351.

634 Simon, C., Buckley, T.R., Frati, F., Stewart, J.B., Beckenbach, A.T., 2006. Incorporating molecular evolution
635 into phylogenetic analysis, and a new compilation of conserved polymerase chain reaction primers for
636 animal mitochondrial DNA. *Annu. Rev. Ecol. Evol. Syst.* 37, 545–579.

637 Simon, C., Frati, F., Beckenbach, A., Crespi, B., Liu, H., Flook, P., 1994. Evolution, weighting, and
638 phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain
639 reaction primers. *Ann. Entomol. Soc. Am.* 87, 651–701. doi:10.1093/aesa/87.6.651

640 Soare, T.W., Kumar, A., Naish, K.A., O'Donnell, S., 2014. Genetic evidence for landscape effects on dispersal
641 in the army ant *Eciton burchellii*. *Mol. Ecol.* 23, 96–109. doi:10.1111/mec.12573

642 Sunnucks, P., 2000. Efficient genetic markers for population biology. *Trends Ecol. Evol.* 15, 199–203.
643 doi:10.1016/S0169-5347(00)01825-5

644 Swofford, D.L., 2002. PAUP*. *Phylogenetic Analysis Using Parsimony (* and Other Methods).* Version 4b10.
645 Sinauer Assoc. Sunderland, MA., Sunderland, MA.

646 Tack, A.J.M., Roslin, T., 2010. Overrun by the neighbors: Landscape context affects strength and sign of local
647 adaptation. *Ecology* 91, 2253–2260. doi:10.1890/09-0080.1

648 Thomas, F., Verneau, O., De Meeûs, T., Renaud, F., 1996. Parasites as to host evolutionary prints: Insights into
649 host evolution from parasitological data. *Int. J. Parasitol.* 26, 677–686. doi:10.1016/0020-7519(96)00023-9

650 Thompson, J.N., 2013. *Relentless Evolution.* University of Chicago Press, Chicago.

651 Thompson, J.N., 2005. *The Geographic Mosaic of Coevolution.* University of Chicago Press, Chicago.

652 Von Beeren, C., Maruyama, M., Kronauer, D.J.C., 2016a. Cryptic diversity, high host specificity and
653 reproductive synchronization in army ant-associated *Vatesus* beetles. *Mol. Ecol.* 25, 990–1005.
654 doi:10.1111/mec.13500

655 Von Beeren, C., Maruyama, M., Kronauer, D.J.C., 2016b. Community sampling and integrative taxonomy
656 reveal new species and host specificity in the army ant-associated beetle genus *Tetradonia* (Coleoptera,
657 Staphylinidae, Aleocharinae). *PLoS One* 11, e0165056.

658 Watkins, J.F., 1976. *The Identification and Distribution of New World Army Ants.* Baylor University Press,
659 Waco, Texas.

660 Whitfield, J.B., Lockhart, P.J., 2007. Deciphering ancient rapid radiations. *Trends Ecol. Evol.* 22, 258–265.

661 doi:10.1016/j.tree.2007.01.012
662 Winston, M.E., Kronauer, D.J.C., Moreau, C., 2017. Early and dynamic colonization of Central America drives
663 speciation in Neotropical army ants. *Mol. Ecol.* 26, 859–870.
664 Zhang, D.-X., Hewitt, G.M., 2003. Nuclear DNA analyses in genetic studies of populations: practice, problems
665 and prospects. *Mol. Ecol.* 12, 563–584.

666
667 **Figure legends**

668
669 **Figure 1.** Map of Panama indicating the main study areas. RFF = Reserva Forestal de Fortuna, BPPS = Bosque
670 Protector de Palo Seco, APSL = Area Protegida de San Lorenzo and its buffer zone; SOB = Soberanía National
671 Park. Colonies collected in APSL and its buffer zone were grouped in the analyses as APSLA and APSLB to
672 reflect colonies collected on either side of the Chagres River.

673
674 **Figure 2.** Majority-rule consensus phylograms for unique haplotypes derived from *Eciton* species (a) and the
675 associated myrmecophiles *Ecitophya* and *Ecitomorpha* (b), generated with MrBayes using a mixed nucleotide
676 model and gamma corrected rate heterogeneity. Support values are shown for nodes (marked by filled circles)
677 found in $\geq 50\%$ of parsimony bootstrap and posterior probability samples. Coloured bars indicate differences in
678 abdomen (or whole body) colour of different taxa.

679
680 **Figure 3.** All compatible groups consensus chronograms for the selected *Eciton* (a) and associated myrmecophiles
681 *Ecitophya* and *Ecitomorpha* (b) haplotypes, generated with MrBayes using a mixed nucleotide model, gamma
682 corrected rate heterogeneity, tk02 relaxed clock and fixed node calibration of 26 (*Eciton*) or 25 (associated
683 myrmecophiles) MYA (indicated by a star). Nodes are scaled to median date estimates with the 95% Highest
684 Posterior Density indicated by a translucent blue bar. Nodes with date estimates are labelled with roman numerals
685 and correspond to values shown in Table 2. Node support values are given for the presented chronograms, filled
686 circles indicate nodes found in $\geq 50\%$ of parsimony bootstrap and posterior probability samples in the main
687 phylogenetic analyses. Coloured bars indicate the abdomen (or whole body)'s colour of different taxa.

688
689 **Figure 4.** Haplotype networks of *Eciton burchellii* (a) and associated *Ecitophya* (b) and *Ecitomorpha* (c) in the
690 main study areas in Panama (RFF, BPPS, APSLA, APSLB) constructed using a median-joining approach. Study
691 areas are indicated with different colours. The size of the circles is proportional to the number of individuals
692 representing a particular haplotype. Missing intermediated haplotypes are indicated with black dots, nucleotide
693 substitutions between haplotypes are indicated by small lines over the haplotype connecting branches.

694

695
696
697

Table 1. Details of collection and sequencing of *Ecitophya* and *Ecitomorpha* myrmecophiles found with different *Eciton* army ant species in the study areas in Panama

Ant species	Location	Ant colony	Individuals collected		Individuals sequenced	
			<i>Ecitomorpha</i>	<i>Ecitophya</i>	<i>Ecitomorpha</i>	<i>Ecitophya</i>
<i>E. b. parvispinum</i>	RFF	E85	6	2	3	2
<i>E. b. parvispinum</i>	RFF	E86	4	12	2	0
<i>E. b. parvispinum</i>	RFF	E87	1	0	1	0
<i>E. b. parvispinum</i>	RFF	E94	3	8	3	4
<i>E. b. parvispinum</i>	RFF	E95	23	8	4	2
<i>E. b. parvispinum</i>	RFF	E96	2	0	2	0
<i>E. b. parvispinum</i>	RFF	E97	12	5	3	2
<i>E. b. parvispinum</i>	RFF	E99	8	20	2	6
<i>E. b. parvispinum</i>	RFF	E100	13	10	4	6
<i>E. b. parvispinum</i>	RFF	E108	15	4	2	1
<i>E. b. parvispinum</i>	RFF	E109	58	28	1	5
<i>E. b. parvispinum</i>	RFF	E114	78	32	8	8
<i>E. b. foreli</i>	BPPS	E89	15	40	4	2
<i>E. b. foreli</i>	BPPS	E101	21	2	2	2
<i>E. b. foreli</i>	BPPS	E103	5	13	2	2
<i>E. b. foreli</i>	BPPS	E104	7	12	2	2
<i>E. b. foreli</i>	APSLA	E126	8	6	6	2
<i>E. b. foreli</i>	APSLA	E127	9	3	2	2
<i>E. b. foreli</i>	APSLA	E132	28	17	14	8
<i>E. b. foreli</i>	APSLA	E143	14	6	13	5
<i>E. b. foreli</i>	APSLA	E154	4	0	2	2
<i>E. b. foreli</i>	APSLA	E156	8	4	2	1
<i>E. b. foreli</i>	APSLB	E162	3	5	2	3
<i>E. b. foreli</i>	APSLB	E165	2	3	2	2
<i>E. b. foreli</i>	APSLB	E166	8	4	2	2
<i>E. hamatum</i>	BPPS	E88	0	69	0	6
<i>E. hamatum</i>	RFF	E105	0	13	0	3
<i>E. hamatum</i>	BPPS	E110	0	8	0	2
<i>E. hamatum</i>	SOB	E169	0	1	0	1
<i>E. lucanoides</i>	RFF	E107	0	6	0	1

698

699

700
701
702
703

Table 2. Date estimates for selected nodes in the *Ecitophya* and myrmecophiles' chronograms generated using MrBayes with TK02, IGR and strict clock models and either a fixed date calibration according to Brady (2003) and Maruyama and Parker (2017) or 1% rate calibration.

Code	Dataset	Method	Median	95% Lower	95% Upper
I	Eciton	TK02 clock, 26 MYA date calibration	26.0	n.a.	n.a.
I	Eciton	TK02 clock, 1% rate calibration	13.7	8.8	21.9
I	Eciton	IGR clock, 26 MYA date calibration	26.0	n.a.	n.a.
I	Eciton	IGR clock, 1% rate calibration	12.1	8.9	16.7
I	Eciton	Strict clock, 26 MYA date calibration	26.0	n.a.	n.a.
I	Eciton	Strict clock, 1% rate calibration	11.5	9.2	14.5
II	Eciton	TK02 clock, 26 MYA date calibration	18.0	12.3	24.0
II	Eciton	TK02 clock, 1% rate calibration	9.5	5.0	16.4
II	Eciton	IGR clock, 26 MYA date calibration	16.8	10.2	23.6
II	Eciton	IGR clock, 1% rate calibration	7.8	4.3	11.8
II	Eciton	Strict clock, 26 MYA date calibration	17.9	13.4	22.8
II	Eciton	Strict clock, 1% rate calibration	7.8	5.7	10.3
III	Eciton	TK02 clock, 26 MYA date calibration	3.5	1.3	7.5
III	Eciton	TK02 clock, 1% rate calibration	1.8	0.6	5.3
III	Eciton	IGR clock, 26 MYA date calibration	3.9	1.2	8.3
III	Eciton	IGR clock, 1% rate calibration	1.8	0.5	4.0
III	Eciton	Strict clock, 26 MYA date calibration	2.4	1.3	3.8
III	Eciton	Strict clock, 1% rate calibration	1.1	0.6	1.6
IV	Eciton	TK02 clock, 26 MYA date calibration	1.3	0.3	3.5
IV	Eciton	TK02 clock, 1% rate calibration	0.7	0.1	2.3
IV	Eciton	IGR clock, 26 MYA date calibration	1.6	0.2	4.1
IV	Eciton	IGR clock, 1% rate calibration	0.7	0.1	1.9
IV	Eciton	Strict clock, 26 MYA date calibration	0.9	0.4	1.7
IV	Eciton	Strict clock, 1% rate calibration	0.4	0.2	0.7
IX	Myrmecophiles	TK02 clock, 25 MYA date calibration	0.8	0.2	2.0
IX	Myrmecophiles	TK02 clock, 1% rate calibration	0.5	0.1	1.4
IX	Myrmecophiles	IGR clock, 25 MYA date calibration	1.1	0.1	2.9
IX	Myrmecophiles	IGR clock, 1% rate calibration	0.5	0.1	1.2
IX	Myrmecophiles	Strict clock, 25 MYA date calibration	0.7	0.2	1.4
IX	Myrmecophiles	Strict clock, 1% rate calibration	0.4	0.1	0.7
V	Myrmecophiles	TK02 clock, 25 MYA date calibration	25.0	n.a.	n.a.
V	Myrmecophiles	TK02 clock, 1% rate calibration	14.3	9.5	23.1
V	Myrmecophiles	IGR clock, 25 MYA date calibration	25.0	n.a.	n.a.
V	Myrmecophiles	IGR clock, 1% rate calibration	12.3	9.3	16.1
V	Myrmecophiles	Strict clock, 25 MYA date calibration	25.0	n.a.	n.a.
V	Myrmecophiles	Strict clock, 1% rate calibration	12.3	9.8	15.4
VI	Myrmecophiles	TK02 clock, 25 MYA date calibration	8.0	4.2	14.1
VI	Myrmecophiles	TK02 clock, 1% rate calibration	4.5	1.9	11.2
VI	Myrmecophiles	IGR clock, 25 MYA date calibration	6.7	3.4	11.7
VI	Myrmecophiles	IGR clock, 1% rate calibration	3.2	1.7	5.2
VI	Myrmecophiles	Strict clock, 25 MYA date calibration	5.9	3.8	8.1
VI	Myrmecophiles	Strict clock, 1% rate calibration	2.9	2.0	4.0
VII	Myrmecophiles	TK02 clock, 25 MYA date calibration	13.8	9.9	18.0
VII	Myrmecophiles	TK02 clock, 1% rate calibration	7.8	4.4	14.2
VII	Myrmecophiles	IGR clock, 25 MYA date calibration	14.7	10.4	19.9
VII	Myrmecophiles	IGR clock, 1% rate calibration	7.1	4.8	9.7
VII	Myrmecophiles	Strict clock, 25 MYA date calibration	14.2	11.2	17.4
VII	Myrmecophiles	Strict clock, 1% rate calibration	6.9	5.3	8.8
VIII	Myrmecophiles	TK02 clock, 25 MYA date calibration	7.8	4.6	12.0
VIII	Myrmecophiles	TK02 clock, 1% rate calibration	4.4	2.1	9.5
VIII	Myrmecophiles	IGR clock, 25 MYA date calibration	8.4	5.1	13.3
VIII	Myrmecophiles	IGR clock, 1% rate calibration	4.0	2.6	6.0
VIII	Myrmecophiles	Strict clock, 25 MYA date calibration	7.4	5.4	9.6
VIII	Myrmecophiles	Strict clock, 1% rate calibration	3.7	2.7	4.8

704

705 **Table 3.** Number of segregating sites (S) and average number of nucleotide differences (*k*) between *Eciton* species,
 706 between the *Ecitophya* and *Ecitomorpha*, and between species within each of these genera; based on unique
 707 haplotypes (802bp mtDNA COI).
 708

Comparison		S	<i>k</i>
Ants			
Eciton	<i>b. parvispinum</i> – <i>dulcium</i>	104	103.5
	<i>b. parvispinum</i> – <i>hamatum</i>	100	94.33
	<i>b. foreli</i> – <i>dulcium</i>	100	91.67
	<i>b. parvispinum</i> – <i>lucanoides</i>	88	88.00
	<i>lucanoides</i> – <i>dulcium</i>	86	85.50
	<i>b. foreli</i> – <i>lucanoides</i>	90	81.83
	<i>b. foreli</i> – <i>b. parvispinum</i>	90	81.00
	<i>b. foreli</i> – <i>hamatum</i>	92	79.28
	<i>hamatum</i> – <i>dulcium</i>	81	74.33
	<i>hamatum</i> – <i>lucanoides</i>	80	73.33
Beetles	<i>Ecitomorpha</i> – <i>Ecitophya</i>	185	103.98
Ecitomorpha	<i>arachnoides</i> (<i>foreli</i>) – <i>arachnoides</i> (<i>parvispinum</i>)	19	33.90
Ecitophya	<i>simulans</i> (<i>foreli</i>) – <i>rettenmeyeri</i>	82	77.30
	<i>gracillima</i> – <i>rettenmeyeri</i>	97	74.67
	<i>simulans</i> (<i>parvispinum</i>) – <i>rettenmeyeri</i>	79	73.71
	<i>simulans</i> (<i>parvispinum</i>) – <i>gracillima</i>	72	42.43
	<i>simulans</i> (<i>foreli</i>) – <i>simulans</i> (<i>parvispinum</i>)	54	41.81
	<i>simulans</i> (<i>foreli</i>) – <i>gracillima</i>	55	24.83

709
 710
 711
 712
 713
 714
 715
 716
 717
 718
 719
 720
 721
 722
 723
 724
 725
 726
 727
 728
 729
 730
 731
 732
 733
 734
 735
 736
 737
 738

739
740
741
742

Table 4. Population structure estimates derived from haplotypic data from *E. burchellii foreli* and their associated populations of *Ecitophya* and *Ecitomorpha* myrmecophile beetles between the areas studied in West (BPPS) and Central Panama (APSLA, APSLB). * P <0.001, NS non-significant

Source of variation	df	SSQ	Variance	% variation	Fixation index
<i>Eciton burchellii foreli</i>					
Among groups	1	1139.658	76.888	76.31	$\Phi_{ST} = 0.9979^*$
Among populations within groups	1	210.061	23.660	23.48	$\Phi_{SC} = 0.9914^*$
Within populations	28	5.765	0.206	0.20	$\Phi_{CT} = 0.7631$
Total	30	1355.484	100.574		
<i>Ecitophya simulans</i>					
Among groups	1	107.796	8.859	84.62	$\Phi_{ST} = 0.8449^*$
Among populations within groups	1	1.522	-0.014	-0.13	$\Phi_{SC} = -0.0088NS$
Within populations	24	38.978	1.624	15.51	$\Phi_{CT} = 0.8462^*$
Total	26	148.296	10.469		
<i>Ecitomorpha arachnoides</i>					
Among groups	1	218.331	14.713	79.57	$\Phi_{ST} = 0.7797^*$
Among populations within groups	1	1.032	-0.296	-1.59	$\Phi_{SC} = -0.0783NS$
Within populations	48	195.5	4.073	22.03	$\Phi_{CT} = 0.7957^*$
Total	50	414.863	18.491		

743