



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Dietary and physical activity recommendations to prevent type 2 diabetes in South Asian adults

Citation for published version:

Muilwijk, M, Nicolaou, M, Qureshi, SA, Celis-Morales, C, Gill, JMR, Sheikh, A, Sattar, N, Beune, E, Jennum, AK, Stronks, K & van Valkengoed, IGM 2018, 'Dietary and physical activity recommendations to prevent type 2 diabetes in South Asian adults: A systematic review' PLoS ONE, vol. 13, no. 7, pp. e0200681. DOI: 10.1371/journal.pone.0200681

Digital Object Identifier (DOI):

[10.1371/journal.pone.0200681](https://doi.org/10.1371/journal.pone.0200681)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

PLoS ONE

Publisher Rights Statement:

This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the Creative Commons CC0 public domain dedication.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



RESEARCH ARTICLE

Dietary and physical activity recommendations to prevent type 2 diabetes in South Asian adults: A systematic review

Mirthe Muilwijk^{1*}, Mary Nicolaou¹, Samera A. Qureshi², Carlos Celis-Morales³, Jason M. R. Gill³, Aziz Sheikh⁴, Naveed Sattar³, Erik Beune¹, Anne Karen Jenum⁵, Karien Stronks¹, Irene G. M. van Valkengoed¹

1 Department of Public Health, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands, **2** The Norwegian Centre for Migrant and Minority Health Research, Oslo, Norway, **3** Institute of Cardiovascular and Medical Sciences, University of Glasgow, Glasgow, United Kingdom, **4** Usher Institute of Population Health Sciences and Informatics, The University of Edinburgh, Edinburgh, United Kingdom, **5** Department of General Practice, Faculty of Medicine, Institute of Health and Society, Blindern, Oslo, Norway

* m.muilwijk@amc.uva.nl



OPEN ACCESS

Citation: Muilwijk M, Nicolaou M, Qureshi SA, Celis-Morales C, Gill JMR, Sheikh A, et al. (2018) Dietary and physical activity recommendations to prevent type 2 diabetes in South Asian adults: A systematic review. *PLoS ONE* 13(7): e0200681. <https://doi.org/10.1371/journal.pone.0200681>

Editor: Noël C. Barengo, Florida International University Herbert Wertheim College of Medicine, UNITED STATES

Received: May 19, 2017

Accepted: July 2, 2018

Published: July 16, 2018

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](https://creativecommons.org/licenses/by/4.0/) public domain dedication.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was sponsored by the Health Programme 2014-2020 from the European Union, grant number 664609 HP-PJ-2014 (https://ec.europa.eu/health/programme/policy/2014-2020_en). KS and IV received the funding. The funders had no role in study design, data collection and

Abstract

Intervention trials and guidelines for the prevention of type 2 diabetes (T2D) in populations of South Asian origin often include strategies to improve diet and physical activity that are based on those developed for other populations. These may be suboptimal for the South Asian target populations. We aimed to provide an overview of included recommended dietary and physical activity components, and to identify whether these were supported by evidence of their effectiveness. Databases were searched until September 2017 for intervention studies and guidelines with an adult South Asian population without T2D. The protocol was registered in PROSPERO, registration number: CRD42015207067. The quality of included studies and guidelines was assessed. Dietary and physical activity components, and effects on T2D incidence, glycemic status and adiposity measures, were summarized in tabular format and evaluated narratively. Eighteen intervention studies and four guidelines were identified. Dietary and physical activity components were similar to recommendations for the general population. Intervention studies and guidelines did not reference evidence to support the effectiveness of components included in the intervention for South Asian populations in particular. Moreover, we were unable to assess patterns of components to determine the effects of specific components. Evaluation of current and emerging components among South Asian populations and subgroups seems necessary to formulate more specific recommendations in future intervention studies and guidelines.

Introduction

Populations of South Asian origin have a high risk for type 2 diabetes (T2D) and its complications, both in the country of origin and after migration [1–3]. In comparison with populations

analysis, decision to publish, or preparation of the manuscript.

Competing interests: We have read the journal's policy and the authors of this manuscript have the following competing interests: Some authors contributed to studies that were included in this systematic review. JG and NS were co-authors on the PODOSA trial but were not involved in its evaluation in the present paper. MN, IV and KS contributed to DHIAAN, and were therefore not involved in the quality assessment in the present paper. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

of European origin, South Asians have been shown to develop T2D at a younger age and at lower levels of adiposity [4, 5]. In the South Asian subcontinent, the diabetes epidemic has been fuelled by changes in the diet and patterns of physical activity due to economic transitions, especially in urban populations [6, 7]. As diet and physical activity are key modifiable risk factors for T2D [8], intervention studies [9–11] and guidelines [12] that aim to prevent or delay the onset of T2D include strategies to improve dietary intake and physical activity. For instance, the T2D prevention 'Population and community-level interventions guideline' issued by the National Institute for Health and Care Excellence (NICE) in the United Kingdom (UK), is aimed at dietary and physical activity components such as 'adopt a low-fat diet' and 'use physically active forms of travel' [12].

Recently, interventions targeting diet and physical activity have been developed to reduce the risk of T2D among populations of South Asian origin [13–17]. However, the trials evaluating these interventions only show small to moderate effects. We speculate that this may be related to the lack of targeting of recommended dietary and physical activity intervention components (hereinafter referred to as components) to the needs of South Asian populations. While many interventions reported cultural adaptation of the mode of delivery, and inclusion or exclusion of components based on the prevalence of certain health behaviours in the target population it is uncertain whether the included components were sufficiently targeted to specific characteristics of South Asian populations. In our own work, the components were often selected based on proven effectiveness of intervention studies conducted in other populations, mainly populations of European origin [13, 14]. Examples of intervention trials that followed this approach are Prevention of Diabetes & Obesity in South Asians (PODOSA) [17] and Diabetes prevention study in Hindustani Surinamese (DH!AAN) [13]. Using similar components may be appropriate in case these are also effective in South Asian populations, however it is also possible that the effects of specific changes in physical activity or diet is different across populations. There is some evidence to suggest that the effects of specific changes in physical activity or diet may be different in South Asian origin populations than in, for instance, those of European origin [18, 19]. For instance, a difference in the effect of a high-calorie, high-fat diet on insulin sensitivity was found between South Asian and White European origin men [18]. Other reports have suggested that the association between duration of physical activity and cardio-metabolic risk factors differs between South Asian and European origin men and women [19, 20]. While previous reviews have discussed the effects of interventions on the prevention of T2D, for instance by estimating the effects on weight loss, none have examined the specific components that were included in the interventions [8, 15, 21, 22].

Therefore, this systematic review set out to examine the dietary and physical activity components in intervention studies and in guidelines for the prevention of T2D in adult South Asian populations worldwide. Our broad aim was to identify gaps and, ultimately, stimulate the development of intervention studies and guidelines that are more targeted to the South Asian population. While public health messages are generally more consistent across populations, clinical guidelines may allow more specific recommendations for prevention at the individual level. The specific aims were to, first, describe the current dietary and physical activity components in intervention studies and guidelines. We included guidelines as these specify recommendations for South Asians based on graded evidence. Secondly, we determined whether the components are supported by population specific evidence. This evidence may include direct references to prior observational and experimental studies for each component listed in the intervention studies and guidelines. Additionally, intervention studies may provide insight in the effects of the included components.

Materials and methods

The methods of the review were registered with the International Prospective Register of Systematic reviews (PROSPERO registration no CRD42015207067). A review protocol was published [23], and the methods are summarised here. The review was written in line with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) reporting guidelines ([S1 PRISMA Checklist](#)) [24].

We systematically searched PUBMED, Embase, Cochrane library and Web of Science from the start of the databases until September 12th, 2017. To the best of our knowledge no relevant studies were published before the start of these databases. All types of experimental, quasi-experimental and before-after studies and guidelines on dietary and/or physical activity components to prevent T2D in a population consisting of at least 75% South Asian adults (≥ 18 years) were considered. The search results were supplemented with reference list tracing of four key reviews [8, 15, 21, 22] and included studies [13, 16, 17, 25–39]. Study authors were contacted in case protocols or clarifications were desired, for instance to obtain the detailed recommended dietary and physical activity components. Corresponding authors of eleven of the studies were contacted to obtain more detailed information on the dietary components that were included in the intervention studies. Authors were additionally queried on which evidence components were based. Eight authors were reached, and provided additional information [13, 16, 17, 25–27, 33, 36]. Unpublished and in-progress studies were identified by searching English language trial registries. The applied search strategies are provided in the supplementary materials [S1 Text](#). Additional studies were identified from experts in the field.

Guidelines were identified by searching online databases and searches of grey literature in all South Asian countries (India, Bangladesh, Pakistan, Sri Lanka, Nepal and Bhutan) and the top five countries to which South Asians have migrated with at least one million migrants a year (Saudi Arabia, United Kingdom, Kuwait, Oman, United States, United Arab Emirates, Canada and Qatar) [40]. The last search was conducted on November 8th, 2017. The applied search strategies are provided in [S2 Text](#).

Two reviewers (MM, SAQ or IGMV) independently screened titles and abstracts for eligibility, based on the pre-set inclusion criteria. Thereafter full-texts were screened for eligibility by two reviewers (MM, SAQ or IGMV). For both processes a discussion was held in case of discrepancies and, if needed, a third reviewer was consulted to reach consensus (MN). A piloted data extraction form was developed in Microsoft Excel.

For each identified study, we reference the first identified manuscript reporting effects. However, information was extracted from all published manuscripts, study protocols and personal communication with authors. Only dietary and physical activity components that were explicitly mentioned in the written study documents or guidelines were extracted. Some studies referred to existing guidelines, for instance the National Institute of Nutrition (NIN) (India) guideline [41] and this reference was noted. However, components listed in referenced guidelines were not included in the data extraction. Data were extracted in duplicate by two reviewers (MM, SAQ, or IGMV) independently from each other. A discussion was held in case of discrepancies, and a third reviewer was consulted if needed (MN). In case data on dietary and / or physical activity strategy components was missing, published protocols were obtained and additional unpublished protocols were requested.

For intervention studies, the risk of bias was assessed at the study level using the Quality Assessment Tool for Quantitative Studies [42]. The quality assessment of studies was done by three reviewers (MM, SAQ, CC or IGMV) instead of two reviewers as we perceived the quality assessment tool to be vulnerable to differences in interpretation. The general quality of guidelines was assessed by the Appraisal of Guidelines for Research & Evaluation II (AGREE II)

instrument [43]. Assessment of the guidelines was done by two reviewers (MM, SAQ), subsections were rated between zero and seven which resulted in section scores between zero and 100, if needed a third reviewer (MN) was consulted. Diet and physical activity components were extracted from each identified study or guideline. Results were summarised in tabular form and evaluated narratively. The identified components were compared to the most recent NICE guideline for diabetes prevention [12] a general guideline listing the current components to prevent T2D in the U.K. to examine whether components in interventions and guidelines for South Asians were similar to those recommended for the general population or possibly targeted towards the South Asian population. The NICE guideline was selected for comparison as it is a recent and well regarded guideline for the general population in a European country with a large South Asian population. For each identified components, we examined whether the studies or guidelines supported the chosen component with specific evidence showing it to be relevant or effective for South Asians. Such evidence may include references to previous observational and experimental studies.

We explored whether certain components were recommended more frequently for pre-specified subgroups. Subgroups analysed were migrant / non-migrant groups (non-migrants were defined by populations living in South Asia), geographical setting (Europe / U.S. / Asia / Oceania) and ethnicity as described by the included study or guideline. We also considered differences by age and sex.

Patterns of effects of studies by their included intervention components were evaluated for studies with at least a moderate quality rating. We, first, described intervention effects on reported outcomes, and explored the consistency of effects. We then evaluated narratively whether differences existed between the effects of studies when certain components were included. Study outcomes were considered to be significantly improved if the p-value was <0.05 . To achieve comparability across measures, the primary effect sizes of outcome measures per study were calculated by Cohen's d, which was calculated by the mean difference or the difference in mean change divided by the pooled standard deviation. Cohen's d values >0.8 were considered as large effect sizes, 0.5–0.79 as moderate and 0.2–0.49 as small [44]. Initially, comparisons of effects by components were only made if an outcome was reported by at least four studies that did and at least four studies that did not include a specific component. If a pattern was recognized, we also assessed the consistency across other outcomes of interest (separately listed in S1 Table). The cumulative evidence may be affected by publication bias, as it is conceivable that non-effective intervention studies were underreported. We checked the possibility of publication bias by listing all registered studies in the trial register. Bias was examined by distinguishing between studies that were yet ongoing and studies that had been completed for at least one year, but had not yet published results.

Results

Characteristics of identified studies

Our searches resulted in 3577 study records (Fig 1). No additional records were identified through other sources, although key reviews, reference lists of included papers and experts were consulted. After removing duplicates, 2533 records were left for title and abstract screening. From the 50 full-text records that were assessed for eligibility, 18 records were included in the review.

The majority ($n = 14$) of included studies were randomized controlled trials (RCTs; Table 1) with a study duration varying from 1.5 to 36 months [13, 16, 17, 28–33, 36–39]. Most studies ($n = 11$) were conducted in India [16, 25–29, 32, 35, 36, 38, 39]. In addition, four were conducted in Europe [13, 17, 31, 33], one in Bangladesh [37], one in New Zealand [30] and




Fig 1. Study flow.

<https://doi.org/10.1371/journal.pone.0200681.g001>

one in the U.S. [34] Most interventions combined dietary and physical activity components [13, 16, 17, 25–27, 32–34, 36, 38], but there were also interventions that focused solely on physical activity (n = 3) [29, 31, 35] or on diet alone (n = 4) [28, 30, 37, 39]. Most studies included South Asian populations of Indian ethnicity (n = 12) [16, 25–29, 31, 32, 35–38], had a wide age range (n = 11) [13, 17, 26–34, 37, 38], and focused specifically on those with pre-diabetes (n = 10) (impaired fasting glucose (IFG) or impaired glucose tolerance (IGT)) [13, 16, 17, 25, 28, 29, 32, 36, 37, 39].

Characteristics of identified guidelines

No guidelines were identified by searching the predefined guideline databases, but four [45–47] were identified in the grey literature. In addition, one was excluded after initially being identified as a guideline, because it was a preliminary protocol for a planned intervention study [26]. Three of the identified guidelines targeted the general South Asian population in

Table 1. Study characteristics.

Study ID	Study design	Country	Ethnicity	Age (years)	% Male	Target group	Intervention size (n)	Study duration (months)	Behaviour change objective	Theory for behaviour change
DH!AAN 2013 [13]	RCT	Netherlands	South Asian Surinamese	18–60	51	IFG, IGT, glycated haemoglobin 42–46 mmol/l, HOMA-IR ≥ 2.39	536	12	Diet, PA	Theory of planned behaviour
DPM 2008 [26]	BAS	India	Indian	>10	41	All inhabitants	850	7	Diet, PA, Stress coping	Community based health promotion
DPP 2012 [27]	BAS	India	Indian	>18	46	All inhabitants	1681	6	Diet, PA, Stress coping	Community based health promotion
Dutta et al. 2014 [28]	RCT	India	Indian	30–80	41	IFG and/or IGT in 2 OGTTs	170	28	Diet	-
Hegde et al. 2013 [29]	RCT	India	Indian	30–75	48	Pre-diabetes, non-alcoholic, non-smoker	29	3	PA	Community based health promotion
Hurst et al. 2010 [30]	RCT	New Zealand	South Asian	23–68	0	Vitamin D deficiency	106	6	Diet	-
IDPP-1 2006 [16]	RCT	India	Indian	35–55	79	IGT on 2 OGTT	531	30	Diet, PA	Community based health promotion
IDPP-2 2009 [36]	RCT*	India	Indian	35–55	87	IGT on 2 OGTT	203	36	Diet, PA	-
InnvaDiab-DE-PLAN 2012 [33]	RCT	Norway	Pakistani	25–62	0	Not specified	198	7	Diet, PA	-
Islam et al. 2016 [37]	RCT	Bangladesh	Bangladeshi	30–65	51	Pre-diabetes	28	6	Diet	-
Madsen et al. 2015 [35]	LBW vs NBW	India	Indian	18–22	100	LBW	117	1.5	PA	-
McDermott et al. 2014 [32]	RCT	India	Indian	30–65	39	FBG ≥ 5.6 mmol/l 1 st degree relative T2D	41	2	Diet**, PA	-
PAMH 2013 [31]	RCT	Norway	Pakistani	25–60	100	Not physically active	150	5	PA	Social cognitive theory
Patel et al. 2017 [38]	RCT	US	Indian	>18	46	High risk T2D	36	6	Diet, PA	Community based health promotion
PODOS 2014 [17]	RCT	UK	South Asian	35–80	46	WC ≥ 90 men ≥ 80 women, IGT or IFG	171	36	Diet, PA	Stages of change model
Ramachandran et al. 2013 [25]	RCT	India	Indian	35–55	100	IGT on 2 OGTT, mobile phone, able to read	537	20	Diet, PA	Stages of change
RICE 2014 [34]	Quasi-experimental	US	Indian	18–75	19	High risk T2D, adapted risk assessment tool ADA	126	6	Diet, PA, Stress coping	Community based health promotion
Thirunavukkarasu et al. 2017 [39]	RCT	India	Indian	40–60	0	Prediabetes, prehypertension	75	3	Diet	-
AAPI Guide [46]	Guideline	U.S.	South Asian	-	-	Not specified	-	-	Diet, PA	-
Apnee Sehat [45]	Guideline Community organization	UK	South Asian	-	-	Not specified	-	-	Diet, PA	-
FHC [47]	Guideline Health Centre	Canada	South Asian	-	-	Not specified	-	-	Diet, PA, Stress coping	-

(Continued)

Table 1. (Continued)

Study ID	Study design	Country	Ethnicity	Age (years)	% Male	Target group	Intervention size (n)	Study duration (months)	Behaviour change objective	Theory for behaviour change
Misra et al. 2011 [48]	Consensus guideline	India	South Asian	-		Not specified	-	-	Diet	-

* The control group in this study received lifestyle intervention and was therefore included in this systematic review.

**The focus of this study was a physical activity intervention, but dietary advice was provided as well.

<https://doi.org/10.1371/journal.pone.0200681.t001>

the UK or North America [45–47], the other guideline focused on India [48]. Three guidelines combined dietary and physical activity components [45–47], and one was solely focused on a dietary strategy [48] (Table 1).

Risk of bias: quality assessment of the studies

In total, seven studies were rated strong [13, 16, 17, 25, 30, 31, 36], seven moderate [17, 26–29, 33, 34, 37] and four weak [32, 35, 38, 39] (Table 2). Overall moderate scores seemed mostly related to low ratings for blinding, and overall weak scores with low ratings for selection bias, design and confounders.

Risk of bias: General quality assessment of the guidelines

In general, guidelines scored high on stakeholder involvement, but low on rigour of development and editorial independence. Scores for the scope and purpose section varied from 44 to 100 points (Table 3). Examples of issues encountered in this section were a lack of attention to

Table 2. Quality assessment intervention studies.

Study ID	Selection Bias	Design	Confounders	Blinding	Data Collection	Drop-Outs	Overall Rating
DH!AAN [13]	2	1	1	2	1	2	Strong
DPM [26]	1	2	1	3	1	2	Moderate
DPP [27]	1	2	1	3	1	1	Moderate
Dutta et al. 2014 [28]	1	1	1	3	1	1	Moderate
Hegde et al. 2013 [29]	1	1	1	3	1	1	Moderate
Hurst et al. 2010 [30]	1	1	1	1	1	2	Strong
IDPP-1 [16]	2	1	1	2	1	1	Strong
IDPP-2 [36]	2	1	1	1	1	1	Strong
InnvaDiab-DE-PLAN [33]	1	1	1	3	1	1	Moderate
Islam et al. 2016 [37]	3	1	1	1	1	1	Moderate
Madsen et al. 2015 [35]	3	3	3	2	1	1	Weak
McDermott et al. 2014 [32]	3	1	3	2	1	1	Weak
PAMH [31]	1	1	1	2	1	1	Strong
Patel et al. 2017 [38]	3	1	1	2	1	3	Weak
PODOSA [17]	1	1	1	2	1	1	Strong
Ramachandran et al. 2013 [25]	1	1	1	2	1	1	Strong
RICE [34]	2	2	2	3	1	1	Moderate
Thirunavakkurasu et al. 2017 [39]	3	1	3	2	1	3	Weak

1 = strong, 2 = moderate, 3 = weak. Studies without weak ratings are overall rated as strong, studies with one weak rating obtain an overall moderate rating and studies with two or more weak ratings are overall rated as weak.

<https://doi.org/10.1371/journal.pone.0200681.t002>

Table 3. Quality assessment guidelines.

Guideline	Scope & Purpose	Stakeholder involvement	Rigour of development	Clarity of presentation	Applicability	Editorial independence
AAPI Guide [46]	67	94	25	94	38	50
Apnee Sehat [45]	44	100	8	72	88	25
FHC [47]	100	89	27	100	88	0
Misra et al. 2011 [48]	89	89	50	100	13	73

Guideline domains were rated on a scale from 0 to 100. According to the Agree II quality assessment tool, domain scores are independent and should therefore not be aggregated into a single quality score. Please note that an overall assessment is not given, as the overall guideline assessment does not include the domain scores.

<https://doi.org/10.1371/journal.pone.0200681.t003>

prevention of T2D (AAPI Guide [46]) and a lack of a clear description of the target group and no consideration of age groups (Apnee Sehat [45]). Further, rigour of development could be improved by the inclusion of references citing work that support the recommended components. Nevertheless, based on the overall assessment in the AGREE II tool, the guidelines could be qualified as acceptable for use.

Dietary components included in intervention studies

In total, 15 of the 18 intervention studies used dietary components (Table 4). Of these, two studies focused solely on vitamin D supplementation [28, 30] one on zinc supplementation [37] and one on using soy flour [39]. The PODOSA study had a very detailed protocol and, therefore, listed the most dietary components (n = 25) [17]. We observed that most other studies listed some specific components in combination with a reference to existing general dietary guidelines or previous intervention studies [13, 16, 17, 25–27, 34, 36, 38].

The recommended dietary components in intervention studies were compared to the NICE guideline for T2D prevention, published May 2011 [12], as was explicated in the study protocol [23]. The NICE guideline mainly focused on starchy foods, fibre, low-fat diets and intake of fruits and vegetables, and further recommended to avoid fried products or products high in sugar, to minimise alcohol intake, to avoid increasing calorie intake, watch portion size and to use breakfast (S2 Table) [12].

The intervention studies were partly in line with the NICE guideline [12]. For instance, many specific components included in these studies were: fat or saturated fat (including fried products; n = 10) [13, 16, 17, 25–27, 33, 34, 36, 38], vegetables (n = 10) [13, 16, 17, 25–27, 33, 34, 36, 38], sugar (n = 8) [16, 17, 25–27, 33, 34, 36] whole grains (n = 8) [16, 17, 25–27, 33, 34, 36], fibre (n = 7) [13, 16, 17, 25–27, 36], legumes (n = 7) [16, 17, 25–27, 33, 36], fruits (n = 7) [13, 16, 17, 25, 34, 36, 38] and portion sizes (n = 4) [17, 26, 27, 34]. Only one study (DH!AAN [13]) recommended the consumption of breakfast, and one study recommended limiting alcohol and calorie intake (PODOSA [17]). Finally, some interventions also included advice on protein intake [17, 26, 27, 34, 38] (n = 5) and balanced meals [16, 17, 25, 34] (n = 4), while the NICE guideline did not [12]. We explored whether there were differences in patterns of recommended dietary components according to population subgroups over the various intervention studies, but did not find clear differences by migration status, ethnicity and geographical location (S1 Table). We were not able to explore differences in patterns according to sex and age due to a lack in variation of these variables across the studies.

Dietary components included in guidelines

The identified South Asian guidelines also recommended components that were in line with the NICE guideline. However, the NICE guideline used more general descriptions, whereas

Table 4. Dietary components in studies and guidelines on prevention of T2D in South Asian adults.

Studies ¹ :	DH! AAN [13] ²	DPM [26] DPP [27] ²	Dutta et al. 2014 [28] Hurst et al. 2010 [30]	IDPP-1[16] Ramachandran et al. 2013 [25] IDPP-2 [36] ²	InnvaDiab-DE-PLAN [33]	Islam et al. 2016 [37]	Patel et al. 2017 [38]	PODOSA [17] ²	RICE [34]	Thirunavukkarasu et al. 2017 [39]	AAPI Guide [46]	Apnee Sehat [45]	FHC [47]	Misra et al. 2011 [48]
Other trials or guidelines referred to	[49–51]	[41]		[41]			[11, 52]	[11, 53, 54]	[11, 55, 56]					
Any nutrients and foods components specified	N,F	N,F	N,F	N,F	N,F	N	N, F	N,F	N,F	F	N,F	N,F	N,F	N,F
Specific nutrients and foods:														
Protein		N,F						N,F	N					N,F
Fat	N	F		N	N,F		N	N,F	F		N	N,F	N,F	N,F
Carbohydrates				N	N			N						N,F
Sugar		F		N	N,F			N,F	N,F		F	N	N,F	N,F
Salt				N	N			N	N,F			N	N,F	N,F
Cholesterol									N					N
Vitamins and minerals			N			N		N				N		N
Fibre	N	N		N				N			N		N	N
Fruits	F			F			F	F	F		F	F	F	F
Vegetables	F	F		F	F		F	F	F		F	F	F	F
Whole grains		F		F	F			F	F			F	F	F
Cereal and grains				F								F	F	F
Legumes		F		F	F			F		F		F		F
Low GI foods														F
Alcohol								F						F
Nuts								F				F	F	F
Meat								F					F	
Fish								F				F		F
Dairy								F					F	
Dilute juices & choose fresh												F		
Cucumber instead of boondi raita								F						
Pickles and char alternatives								F						
Replace chevda with popcorn								F						
Any patterns specified	P	P		P				P	P			P	P	P
Specific patterns:														

(Continued)

Table 4. (Continued)

Studies ¹ :	DH! AAN [13] ²	DPM [26] DPP [27] ²	Dutta et al. 2014 [28] Hurst et al. 2010 [30]	IDPP-1 [16] Ramachandran et al. 2013 [25] IDPP-2 [36] ²	InnvaDiab-DE-PLAN [33]	Islam et al. 2016 [37]	Patel et al. 2017 [38]	PODOSA [17] ²	RICE [34]	Thirunavukkarasu et al. 2017 [39]	AAPI Guide [46]	Apnee Sehat [45]	FHC [47]	Misra et al. 2011 [48]
Portion sizes		P						P	P			P	P	P
Eat breakfast	P											P		P
Cooking methods												P		P
Vegetarian diet									P		P			
Salt									P					P
Iron absorption												P		
Balanced meals				P				P	P			P	P	
Timing of meals												P	P	P
Limit eating out									P				P	
Energy deficit (500-600kcal/day)								P						

N, F or P indicates that a component was formulated for the nutrient, food (a certain product) or pattern respectively. A more extended table is provided in the supplementary data (S4 Table).

¹Studies in which the same components were recommended are shown in a combined column for reasons of space.

²Study leaders provided additional information on components.

<https://doi.org/10.1371/journal.pone.0200681.t004>

the identified guidelines provided more detailed recommendations. Guidelines for instance recommend to “replace chevda with popcorn” while the NICE guideline recommends to “consume as little as possible fried food; drinks and confectionery high in added sugar; and other food high in fat and sugar”. In line with the NICE guideline, fat, sugar, fruits and vegetables were mentioned in all guidelines [45–48], while fibre [46–48], whole grains [45, 47, 48], cereal intake [45, 47, 48] and portion sizes [45, 47, 48] were listed in three out of four guidelines, only two out of four guidelines included legume intake [46, 48]. In contrast to the NICE guideline, three out of four identified guidelines included components on salt [45, 47, 48], nuts [45, 47, 48] and timing of meals [45, 47, 48].

Physical activity components in intervention studies

Table 5 summarizes the physical activity components in the studies and guidelines. All studies, except the ones that focused on a single dietary component [28, 30, 37, 39], included physical activity (Table 5). The NICE guideline recommends to be physically active for at least 30 minutes per day on at least five days per week at least at a moderate-intensity (S2 Table) [12]. Most components in the intervention studies were in line with the recommendation to be active for at least 30 minutes per day for at least five days a week. However, there were three studies that recommended a longer duration of activity per day, varying from 45 to 90 minutes [29, 33, 35], and one recommended to be active on two days per week [33]. The preferred type of

Table 5. Physical activity components in studies and guidelines on prevention of T2D in South Asian adults.

Study ID	' / day	Days / week	Activity	Intensity
DH!AAN [13]	30	7		
DPM [26]	150 / week			Moderate
DPP [27]	30	7		
Hegde et al. 2013 [29]	75–90	5	Yoga	
IDPP-1 [16]	30	7	Brisk walking	Moderate
IDPP-2 [36]	30	7	Brisk walking	Moderate
InnvaDiab-DE-PLAN [33]	60	2	Walking 5000 steps	Low
Madsen et al. 2015 [35]	45	7	Cycling	
McDermott et al. 2014 [32]	32	3 to 6	Yoga	Moderate
PAMH [31]	30	7		Moderate
Patel et al. 2017 [38]	150 / week		Or walking 10.000 steps	Moderate
PODOSA [17]	30	7	Brisk walking	Moderate
Ramachandran et al. 2013 [25]	30	7	Brisk walking	Moderate
RICE [34]			Domestic activities	
AAPI Guide [36]	30	Most or all	Walking or other sports	Moderate
Apnee Sehat [35]	30	7	What makes happy	
FHC [37]	150 / week			Moderate

<https://doi.org/10.1371/journal.pone.0200681.t005>

activity was specified in nine of the studies; walking was the most recommended activity [16, 17, 25, 33, 36]. Two studies were solely based on yoga [29, 32]. We explored whether there were differences in patterns of recommended physical activity components across subgroups (Table 5). Recommendations for duration, frequency and intensity did not systematically vary across studies among migrant versus non-migrant populations or other subgroups. The only observation that could be made was that studies conducted among non-migrants [16, 25–29, 36, 38] seemed to recommend specific activities more frequently, such as walking or yoga, than studies conducted among migrants.

Physical activity components in guidelines

Three out of four guidelines included physical activity (Table 5). A similar pattern of recommended physical activity duration and frequency as mentioned in the intervention studies was observed [45–47].

Targeting of components to the characteristics of the South Asian population

We found that components used in studies were often based on prior T2D intervention studies in the general population, or on general population guidelines from the geographical area where the study was conducted (Tables 4 and 5). For instance, the dietary components that were recommended in the IDPP-1 study [16] were based on the National Institute of Nutrition (India) dietary guidelines, whereas the components in the DH!AAN study were based on the Dutch study on lifestyle intervention and impaired glucose tolerance Maastricht (SLIM) [51] and guidelines that were developed for the Dutch population [49, 50]. The majority of studies reported having adapted the recommendations to the cultural context of the population, in particular the mode of delivery of the intervention for instance by using South Asian dietitians [13, 17, 26, 27, 33, 34]. However, the studies did not state that components were selected or adapted based on specific evidence on the effects of components on T2D risk in the target

population (e.g. people of Indian descent with IGT). We did not identify any referenced articles that state components to be effective within the South Asian population in either intervention studies or guidelines [13, 16, 17, 25–39, 45–48], nor did we identify evidence in original interventions or guidelines on which some intervention studies were based [11, 41, 49–51, 53–56].

Patterns of effects of intervention studies

We included the 12 studies with a moderate to strong rating in the evaluation of the effects on T2D prevalence and its underlying measures (S1 Table), four studies were excluded due to a weak rating [32, 35, 38, 39]. The studies reported the effects with regard to a variety of different glucose metabolism and adiposity measures [13, 16, 17, 25–31, 33, 34]. Yet, only five studies reported effects on T2D incidence [16, 17, 25, 28, 36]. These were the studies with the longest study durations (20–36 months). In one study the intervention group received the lifestyle intervention combined with medication, while the control group received the lifestyle intervention [36], we considered the study arm that only received lifestyle intervention as a before-fer study. All studies reported lower progression rates to T2D in the intervention group versus the control group, although the difference in one study was not statistically significant [17]. Results for fasting glucose and other measures of glucose metabolism were less consistent. We could include the effect size as determined by Cohen's *d* for fasting glucose in nine studies; effect sizes varied from -0.14 to 2.78 standard deviations [13, 16, 17, 26–29, 33, 34]. We could not include the other three studies as standard deviations were not reported. All, but one study [30] reported on adiposity measures. We found effect sizes of -0.05 to 0.34 standard deviations [13, 16, 17, 26–29, 31, 33, 34] for BMI, which was included as an outcome measure in twelve studies of which for ten studies Cohen's *d* could be calculated. Although not all results significantly improved, most results were in the direction of an improvement.

For some studies, strength of the effects were different for different types of outcomes [17, 31, 33, 34]. For example, PAMH [31] and PODOSA [17] found significant improved results on adiposity parameters, but not on glucose measures. In contrast, RICE [34] and InnvaDiab-DE-PLAN [33] found significant differences for glucose, but not for adiposity measures. However, the directionality of the effects was similar for the different outcomes.

The effects of studies according to the components protein, portion sizes, carbohydrates, fibre, fruits, legumes, portion sizes and balanced meals were evaluated for glucose, BMI and waist circumference measures (Table 6 and S3 Table). However, no clear conclusion could be drawn from these comparisons, as the number of studies was limited and components overlapped.

Risk of publication bias

We identified 13 additional studies from trial registers that had not been published (Table 7) [57]. Based on the reported end date, two studies are still ongoing. Three studies ended recently (in 2016 or 2017). The remaining eight studies have not made their results public, although the studies had an estimated end date between 2009 and 2015. This suggests the possibility of a publication bias. Although the reason for this underreporting is not clear, it may be related to unsuccessful inclusion, discontinuation or failure to publish negative results.

Discussion

This review found that most intervention studies and guidelines to prevent T2D in South Asians recommended a wide variety of dietary and physical activity components that were underpinned by evidence from studies in non-South Asian populations and that were largely

Table 6. Effects of components included in intervention studies.

Studies	Effects			Effects of components											
	Base I	ES	p-value	Patterns / Portion sizes		Carbohydrates		Fibre		Fruits		Legumes		Balanced meals	
Glucose (mmol/l)				Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No
DH!AAN [9]	5.3	-0.14	0.66		0 X		0 X	0 X		0 X			0 X		0 X
DPM [21]	5.2 ²	0.08	0.05	0 ✓			0 ✓	0 ✓			0 ✓	0 ✓			0 ✓
DPP [22]	5.3 ²	0.05	<0.001	0 ✓			0 ✓	0 ✓			0 ✓	0 ✓			0 ✓
Dutta et al. 2014 [23]	6.1 ²	0.41	0.02		1 ✓		1 ✓		1 ✓		1 ✓		1 ✓		1 ✓
Hegde et al. 2013 [24]	5.3	0.00	0.04		0 ✓		0 ✓		0 ✓		0 ✓		0 ✓		0 ✓
Hurst et al. 2010 [25]	4.7		0.82		X		X		X		X		X		X
IDPP-1 [12]	5.4	0.23	0.03		1 ✓	1 ✓		1 ✓		1 ✓		1 ✓		1 ✓	
IDPP-2 [36]	5.7	0.19	≥0.05		0 X	0 X		0 X		0 X		0 X		0 X	
InnvaDiab-DE-PLAN [30]	5.6	2.78	0.02		3 ✓	3 ✓			3 ✓		3 ✓	3 ✓			3 ✓
Islam et al. 2016 [37]	5.8	1.6	<0.001		3 ✓		3 ✓		3 ✓		3 ✓		3 ✓		3 ✓
PAMH [20]	5.3		0.30		X		X		X		X		X		X
PODOSA [13]	5.8	0.12	0.34	0 X		0 X		0 X		0 X		0 X		0 X	
RICE [26]	6.4 ²	0.70	<0.01	2 ✓		2 ✓			2 ✓	2 ✓			2 ✓	2 ✓	
BMI (kg/m2)															
DH!AAN [9]	28.1	0.08	0.09		0 X		0 X	0 X		0 X			0 X		0 X
DPM [21]	20.6	-0.05	<0.01	0 X			0 X	0 X			0 X	0 X			0 X
DPP [22]	20.7	0.02	<0.01	0 ✓			0 ✓	0 ✓			0 ✓	0 ✓			0 ✓
Dutta et al. 2014 [23]	26.3	0.24	≥0.05		1 X		1 X		1 X		1 X		1 X		1 X
Hegde et al. 2013 [24]	27.2	0.13	0.01		0 ✓		0 ✓		0 ✓		0 ✓		0 ✓		0 ✓
IDPP-1 [12]	25.5	0.06	0.04		0 ✓	0 ✓		0 ✓		0 ✓		0 ✓		0 ✓	
IDPP-2 [36]	26.0	-0.14	≥0.05		0 X	0 X		0 X		0 X		0 X		0 X	
InnvaDiab-DE-PLAN [30]	29.4	0.34	0.18		1 X	1 X			1 X		1 X	1 X			1 X
PAMH [20]	27.1		<0.01		✓		✓		✓		✓		✓		✓
PODOSA [13]	30.6	0.11	0.01	0 ✓		0 ✓		0 ✓		0 ✓		0 ✓		0 ✓	
Ramachandran et al. 2013 [29]	25.8	0.00			0	0		0		0		0		0	
RICE [26]	27.8	0.20	0.08	0 X		0 X			0 X	0 X			0 X	0 X	
WC (cm)															
DH!AAN [9]	94	0.09	0.50		0 X		0 X	0 X		0 X			0 X		0 X
DPM [21]	76 ²	0.43	0.001	1 ✓			1 ✓	1 ✓			1 ✓	1 ✓			1 ✓
DPP [22]	75 ²	0.05	0.01	0 ✓			0 ✓	0 ✓			0 ✓	0 ✓			0 ✓
Hegde et al. 2013 [24]	88.9	-0.02	0.03		0 ✓		0 ✓		0 ✓		0 ✓		0 ✓		0 ✓
IDPP-1 [12]	89.1	0.07	0.43		0 X	0 X		0 X		0 X		0 X		0 X	
IDPP-2 [36]	91.2	0.14	≥0.05		0 X	0 X		0 X		0 X		0 X		0 X	
InnvaDiab-DE-PLAN [30]	95.3	0.62	0.18		1 X	1 X			1 X		1 X	1 X			1 X
PAMH [20]	98		0.01		✓		✓		✓		✓		✓		✓
PODOSA [13]	102.7	0.16	0.01	0 ✓		0 ✓		0 ✓		0 ✓		0 ✓		0 ✓	
Ramachandran et al. 2013 [29]	92.6	-0.01			0	0		0		0		0		0	
RICE [26]	93 ²	0.19	0.39	0 X		0 X			0 X	0 X			0 X	0 X	

1, 2 or 3 are shown to indicate the magnitude of the effect size, which corresponds to none (<0.2), small (>0.2), medium (>0.5) or large (0.8) A ✓ is shown to indicate that the study reported a significant improved p-value and a X is shown to indicate that there was no significant improved p-value. Yes and no were used to indicate whether the component was included in the study. Studies on vitamin D and physical activity were included in the evaluation of effects of components but did not include any of the evaluated components. T2D, type 2 diabetes; 2-h glucose, 2-h post 75g glucose blood glucose; IGT, Impaired Glucose Tolerance; IFG, Impaired Fasting Glucose; BMI, Body Mass Index; WC, Waist Circumference; HC, Hip Circumference; TC, Thigh Circumference; WHR, Waist Hip Ratio; BF, Body Fat; Out, Outcome; I, Intervention group; C, Control group; Base, Baseline; ES, Effect size shown by Cohens d.

²Original units were converted to SI units.

<https://doi.org/10.1371/journal.pone.0200681.t006>

Table 7. Overview additional studies.

Study ID	Ref	End date	Country	Ethnicity	Group	Age (years)	Behaviour change objective	Study type	Study duration (months)
Badaam et al.	[57]	03/2013	India	Indian	Pre-diabetes	25–40	PA; Aerobic vs. Sprint	RCT	3
BanglaDiP	[58]	10/2012	UK	Bangladeshi	At risk T2D	20–70	Diet, PA, Tobacco	RCT	6
DIAbetes PREvention programme	[59]	12/2012	India	Indian	IFG/IGT	18–80	Diet, PA (Yoga)		12
GO for IT	[60]	02/2016	UK	South Asian	At risk T2D	50–75	PA (HIIT)	Crossover trial	7 hours
IPDS	[61]	02/2017	India	Indian	BMI ≥23; IFG/IGT	30–70	PA (Fenugreek/Yoga)	RCT	36
K-DPP	[62, 63]	11/2015	India	Indian		30–60		RCT	24
mDiab	[64]	Ongoing	India	Indian	BMI ≥23	20–65	Diet, PA	RCT	3
PREVENT-WIN	[65]	12/2015	India	Indian	Pre-diabetes	20–60	Diet (Vitamin D)	RCT	24
Ranasinghe et al. 2012	[66, 67]	06/2012	Sri Lanka	Sri Lankan	Pre-diabetes	12–65	Diet (Zinc)	RCT	2
Shekar et al.	[68]	12/2015	India	Indian	IGT; Sedentary lifestyle	24–50	Lifestyle	RCT	2
Thomson et al.	[69]	12/2017	India/UK		Pre-diabetes	18–74		RCT	24
Velho et al.	[70]	06/2016	India	Indian	Pre-diabetes	30–45	Diet	RCT	6
VITALITY	[71]	10/2013	UK	South Asian	Insulin resistance	25–75	Diet (Vitamin D)	RCT	6

IFG, Impaired fasting glucose; T2D, Type 2 diabetes; BMI, Body Mass Index; IGT, Impaired fasting glucose; PA, Physical activity; HIIT, High Intensity Interval Training; RCT, Randomised Controlled Trial.

<https://doi.org/10.1371/journal.pone.0200681.t007>

in line with guidelines not specifically developed for a South Asian origin population, such as the NICE guideline. We could not identify clear patterns in components across subgroups according to study or population characteristics. The included studies and guidelines often did not provide specific evidence that recommended components were effective in reducing T2D risk among South Asian populations or subgroups. Although the overall directionality of the results was towards an improvement of outcome measures, we were not able to assess patterns to determine the contribution of specific components to these effects.

As anticipated, we found a lack of specific adaptations of components and a lack of underpinning of the suitability of the components for the target group in both intervention studies and guidelines. For some components this may be because evidence of potential effectiveness within this population was lacking at the time of development, but the lack of evidence for the effectiveness of components among South Asians was not mentioned in the studies or guidelines. Nor did these studies, in absence of specific data on effectiveness, reference specific observational data supporting the importance of the components among South Asians. However, evidence that was available may also not have been considered because researchers were optimistic concerning the expected effects based on the large reductions in T2D risk observed in efficacy studies [10, 11], including one from India [16]. The assumption that expected effects may be similar across populations may hold for several components, as confirmed by some observational work in South Asian populations. For example, a dietary pattern high in fruits, vegetables, dairy products and monounsaturated fatty acids and low in refined cereals was associated with a lower risk for T2D [72]. Another study showed reduced T2D risk for legume intake [73]. However, evidence for the effectiveness of other specific components is, as yet,

unavailable. Moreover, the specific dose or change that is needed to reduce the risk for T2D in South Asian populations may differ from those in the European population. For instance, the optimal required amount and type of physical activity that is necessary to achieve an effect may differ by ethnic group [19]. We recommend further evaluations of the effectiveness of components within South Asian populations so that, if necessary, targeted recommendations can be developed for these populations. The recommendations should then also take into account the existing dietary patterns and activity levels in South Asian populations. South Asians, for instance, tend to be less physically active, and to more often have a vegetarian diet than populations of European origin [74–76]. In addition, dietary patterns may vary across by South Asian population (e.g. by country of residence) [77].

It is noteworthy that potentially relevant evidence that has only emerged in recent years was lacking from the intervention studies and guidelines for South Asians. These components may also be relevant to the general population, but were shown effective for South Asians. One example related to diet is the infrequent consideration of lacto-, lacto-ovo and semi-vegetarian diets [78]. A study conducted among men and women in India indicated that these types of diets were protective for T2D compared to non-vegetarian diets. An example related to physical activity is the lack of attention to sedentary time, although this might also be relevant for other populations. Emerging evidence among Pakistani immigrant men living in Norway indicates that a reduction of sedentary time may be as important as, or perhaps even more important than, an increase in moderate physical activity to reduce the T2D risk [79]. Secondary analyses of PODOSA reported a cross-sectional relationship between sitting time and 2-h glucose levels in U.K. South Asians [80]. These recent findings may be included in future intervention studies and guidelines, and it might be expected that their inclusion may increase effectiveness. While sedentary behaviour has recently been specified in the health guidelines of many countries they have yet to be included in guidelines / recommendations targeting South Asians.

Information on the extent to which recommendations were truly adopted in daily practice was not available. A change in T2D risk, given theoretical effects of the component, can only be observed when the behaviour of participants is changed towards recommended levels. Ram et al., 2014 showed that among those with greater compliance to the lifestyle modification than among participants that failed to make any changes, incidence of T2D was lower [81]. Unfortunately, such information on implementation and changes in behaviour were not reported for most studies limiting our ability to assess this further.

A general point that follows from our work is that given the high burden of T2D and T2D related complications among South Asians, guidelines to reduce the risk of T2D seem imperative. Yet, only four guidelines aimed at T2D prevention among South Asians were retrieved. This lack of guidelines aiming to reduce the risk of T2D among the high risk South Asian population fits the more general finding that there is a lack of consideration of ethnicity in reviews and health guidelines [82]. On the other hand, dissemination of guidelines for different types of subgroups may be difficult as different health messages for subgroups may lead to confusion across populations. However, identified guidelines were especially developed for South Asians and still did not recommended components especially developed for South Asians. Guidelines were not underpinned by evidence that showed components to be effective for South Asians. At the same time, the identified guidelines may not reflect all recommended components within the South Asian population as we may have missed recommendations that were only spread through other methods of dissemination. For instance, during the searches of the grey literature we identified that strategies and components to prevent T2D in South Asian countries are often communicated via alternative materials such as posters [83–85]. Also, more general (not specifically focused at the prevention of T2D) dietary and physical activity guidelines

are available such as the Indian dietary guidelines, and the guideline for physical activity in Asian Indians [41], [86].

Limitations

Our work has some limitations that merit discussion. Firstly, the search strategy for scientific studies was limited to abstracts written in English. Although we did the utmost to identify further work through reference tracing and expert consultations, this means that we may have missed studies that were not published or referenced in the English language. Similarly, we may also have missed some guidelines as, even though language was not a formal requirement, the search terms were in English.

Secondly, within studies some of the dietary and physical activity components may have been missed which should be considered in the interpretations of the findings. Although study investigators were contacted to retrieve protocols, not all were reached or investigators no longer had protocols available for assessment. In addition, we did not extract components listed in referenced material (e.g. general guidelines reference in the study) as referenced materials may have been used as basis for the trial design, but not fully included in the trial.

Thirdly, we used the Quality Assessment Tool for Quantitative Studies [42] to assess study quality and only included studies in the evaluation of effectiveness if the study obtained at least a moderate rating. However, studies of poor quality were not excluded from the assessment of diet and physical activity components. Although the execution of the study does not directly reflect the intervention development process, it is not unlikely that the design of an intervention from a study with a poor rating is also of poorer quality than that of other studies. Another point is that we observed the Quality Assessment Tool to be vulnerable to minor differences in interpretation, while relatively arbitrary changes in rating of one section can result in a different overall score. Previously, the inter-rater agreement for individual domains was described as fair [87]. Due to the discrepancies in the scoring, it was decided to involve a third reviewer in the scoring of all studies. This also better addressed the point that reviewers were not blinded to the studies. Last, based on the estimated end date of yet unpublished trials, we found evidence to suggest possible publication bias. We found six studies that ended prior to 2014 that have not yet updated their registrations or published their results in the international scientific literature. Although the reason for this underreporting cannot be determined, e.g. unsuccessful inclusion or negative results, it is indicative of a probable publication bias.

Conclusions

In conclusion, our study shows that, as was expected, dietary and physical activity components in intervention studies and guidelines to prevent T2D in South Asians are based upon those developed for the European white population. Intervention studies and guidelines did not reference evidence that shows included components to be effective for South Asian populations in particular. It remains uncertain whether dietary and/or physical activity components in interventions and guidelines should be adapted to fit the metabolic characteristics of South Asians, or whether other aspects must be adapted to increase the effectiveness of interventions. Therefore, we cannot yet formulate recommendations to include or exclude certain components in future intervention studies. Research into the effects of current and emerging components among South Asian populations and subgroups in their specific contexts is needed for further clarification, or for instance by more high quality trials in South Asians that test a specific dietary or physical activity component.

Supporting information

S1 PRISMA Checklist.

(DOC)

S1 Text. Search strategy scientific literature databases.

(DOC)

S2 Text. Search strategy guideline databases.

(DOC)

S1 Table. Effects of studies. ¹ 0, 1, 2 or 3 are shown to indicate the magnitude of the effect size, which corresponds to none (<0.2), small (>0.2), medium (>0.5) or large (0.8) A ✓ is shown to indicate that the study reported a significant improved p-value and a X is shown to indicate that there was no significant improved p-value. Yes and no were used to indicate whether the component was included in the study. Studies on vitamin D and physical activity were included in the evaluation of effects of components but did not include any of the evaluated components. T2D, type 2 diabetes; 2-h glucose, 2-h post 75g glucose blood glucose; IGT, Impaired Glucose Tolerance; IFG, Impaired Fasting Glucose; BMI, Body Mass Index; WC, Waist Circumference; HC, Hip Circumference; TC, Thigh Circumference; WHR, Waist Hip Ratio; BF, Body Fat; Out, Outcome; I, Intervention group; C, Control group; Base, Baseline; ES, Effect size shown by Cohens d *Original units were converted to SI units. Studies shown in cursive obtained a weak score in the quality assessment and were not included in the effects assessment.

(DOC)

S2 Table. Recommended components NICE guideline.

(DOC)

S3 Table. Patterns according to included components. Data are the number of studies reporting the specific component. Components were included in the table, if these were recommended by more than one study.

(DOC)

S4 Table. Dietary components per study compared to the NICE guideline.

(DOC)

Acknowledgments

We thank Faridi S. van Etten-Jamaludin, clinical librarian at the Academic Medical Center, for her help with the development of the search strategy for the different databases. We thank Saadul Islam and our colleague for translations of documents. And we thank study authors of all included studies for additional study data / study clarification.

Author Contributions

Conceptualization: Mirthe Muilwijk, Karien Stronks, Irene G. M. van Valkengoed.

Data curation: Mirthe Muilwijk, Mary Nicolaou, Samera A. Qureshi, Carlos Celis-Morales, Irene G. M. van Valkengoed.

Formal analysis: Mirthe Muilwijk, Irene G. M. van Valkengoed.

Funding acquisition: Karien Stronks, Irene G. M. van Valkengoed.

Investigation: Mirthe Muilwijk.

Methodology: Mirthe Muilwijk, Aziz Sheikh, Anne Karen Jenum, Irene G. M. van Valkengoed.

Supervision: Karien Stronks, Irene G. M. van Valkengoed.

Validation: Mary Nicolaou, Irene G. M. van Valkengoed.

Visualization: Mirthe Muilwijk.

Writing – original draft: Mirthe Muilwijk.

Writing – review & editing: Mirthe Muilwijk, Mary Nicolaou, Samera A. Qureshi, Carlos Celis-Morales, Jason M. R. Gill, Aziz Sheikh, Naveed Sattar, Erik Beune, Anne Karen Jenum, Karien Stronks, Irene G. M. van Valkengoed.

References

- Gujral UP, Pradeepa R, Weber MB, Narayan KM, Mohan V. Type 2 diabetes in South Asians: similarities and differences with white Caucasian and other populations. *Annals of the New York Academy of Sciences*. 2013; 1281:51–63. Epub 2013/01/16. <https://doi.org/10.1111/j.1749-6632.2012.06838.x> PMID: 23317344; PubMed Central PMCID: PMC3715105.
- Meeks KA, Freitas-Da-Silva D, Adeyemo A, Beune EJ, Modesti PA, Stronks K, et al. Disparities in type 2 diabetes prevalence among ethnic minority groups resident in Europe: a systematic review and meta-analysis. *Internal and emergency medicine*. 2015. Epub 2015/09/16. <https://doi.org/10.1007/s11739-015-1302-9> PMID: 26370238.
- Sattar N, Gill JMR. Type 2 diabetes in migrant south Asians: mechanisms, mitigation, and management. *The Lancet Diabetes & Endocrinology*. 3(12):1004–16. [https://doi.org/10.1016/S2213-8587\(15\)00326-5](https://doi.org/10.1016/S2213-8587(15)00326-5)
- Razak F, Anand SS, Shannon H, Vuksan V, Davis B, Jacobs R, et al. Defining obesity cut points in a multiethnic population. *Circulation*. 2007; 115(16):2111–8. Epub 2007/04/11. <https://doi.org/10.1161/CIRCULATIONAHA.106.635011> PMID: 17420343.
- Misra A, Khurana L. Obesity-related non-communicable diseases: South Asians vs White Caucasians. *International journal of obesity (2005)*. 2011; 35(2):167–87. Epub 2010/07/21. <https://doi.org/10.1038/ijo.2010.135> PMID: 20644557.
- Jayawardena R, Ranasinghe P, Byrne NM, Soares MJ, Katulanda P, Hills AP. Prevalence and trends of the diabetes epidemic in South Asia: a systematic review and meta-analysis. *BMC Public Health*. 2012; 12:380–. <https://doi.org/10.1186/1471-2458-12-380> PubMed PMID: PMC3447674. PMID: 22630043
- Misra A, Ramchandran A, Jayawardena R, Shrivastava U, Snehalatha C. Diabetes in South Asians. *Diabetic medicine: a journal of the British Diabetic Association*. 2014; 31(10):1153–62. Epub 2014/07/01. <https://doi.org/10.1111/dme.12540> PMID: 24975549.
- Ramchandran A, Snehalatha C, Samith Shetty A, Nanditha A. Primary prevention of Type 2 diabetes in South Asians—challenges and the way forward. *Diabetic medicine: a journal of the British Diabetic Association*. 2013; 30(1):26–34. Epub 2012/07/26. <https://doi.org/10.1111/j.1464-5491.2012.03753.x> PMID: 22827704.
- Alberti KG, Zimmet P, Shaw J. International Diabetes Federation: a consensus on Type 2 diabetes prevention. *Diabetic medicine: a journal of the British Diabetic Association*. 2007; 24(5):451–63. Epub 2007/05/02. <https://doi.org/10.1111/j.1464-5491.2007.02157.x> PMID: 17470191.
- Lindström J, Louheranta A, Mannelin M, Rastas M, Salminen V, Eriksson J, et al. The Finnish Diabetes Prevention Study (DPS): Lifestyle intervention and 3-year results on diet and physical activity. *Diabetes Care*. 2003; 26(12):3230–6. <https://doi.org/10.2337/diacare.26.12.3230> PMID: 14633807
- The Diabetes Prevention Program Research Group. The Diabetes Prevention Program (DPP): Description of lifestyle intervention. *Diabetes Care*. 2002; 25(12):2165–71. <https://doi.org/10.2337/diacare.25.12.2165> PMID: 12453955
- NICE. Type 2 diabetes prevention: population and community-level interventions. National Institute for Health and Clinical Excellence; 2011.
- Admiraal WM, Vlaar EM, Nierkens V, Holleman F, Middelkoop BJ, Stronks K, et al. Intensive lifestyle intervention in general practice to prevent type 2 diabetes among 18 to 60-year-old South Asians: 1-year effects on the weight status and metabolic profile of participants in a randomized controlled trial. *PloS one*. 2013; 8(7):e68605. Epub 2013/07/31. <https://doi.org/10.1371/journal.pone.0068605> PMID: 23894322; PubMed Central PMCID: PMC3718785.

14. Wallia S, Bhopal RS, Douglas A, Bhopal R, Sharma A, Hutchison A, et al. Culturally adapting the prevention of diabetes and obesity in South Asians (PODOSA) trial. *Health promotion international*. 2014; 29(4):768–79. Epub 2013/04/12. <https://doi.org/10.1093/heapro/dat015> PMID: 23574693.
15. Brown T, Smith S, Bhopal R, Kasim A, Summerbell C. Diet and physical activity interventions to prevent or treat obesity in South Asian children and adults: a systematic review and meta-analysis. *International journal of environmental research and public health*. 2015; 12(1):566–94. Epub 2015/01/15. <https://doi.org/10.3390/ijerph120100566> PMID: 25584423; PubMed Central PMCID: PMC4306880.
16. Ramachandran A, Snehalatha C, Mary S, Mukesh B, Bhaskar AD, Vijay V. The Indian Diabetes Prevention Programme shows that lifestyle modification and metformin prevent type 2 diabetes in Asian Indian subjects with impaired glucose tolerance (IDPP-1). *Diabetologia*. 2006; 49(2):289–97. Epub 2006/01/05. <https://doi.org/10.1007/s00125-005-0097-z> PMID: 16391903.
17. Bhopal RS, Douglas A, Wallia S, Forbes JF, Lean MEJ, Gill JMR, et al. Effect of a lifestyle intervention on weight change in south Asian individuals in the UK at high risk of type 2 diabetes: A family-cluster randomised controlled trial. *The Lancet Diabetes and Endocrinology*. 2014; 2(3):218–27. [https://doi.org/10.1016/S2213-8587\(13\)70204-3](https://doi.org/10.1016/S2213-8587(13)70204-3) PMID: 24622752.
18. Bakker LEH, van Schinkel LD, Guigas B, Streefland TCM, Jonker JT, van Klinken JB, et al. A 5-Day High-Fat, High-Calorie Diet Impairs Insulin Sensitivity in Healthy, Young South Asian Men but Not in Caucasian Men. *Diabetes*. 2014; 63(1):248–58. <https://doi.org/10.2337/db13-0696> PMID: 24357702
19. Celis-Morales CA, Ghouri N, Bailey ME, Sattar N, Gill JM. Should physical activity recommendations be ethnicity-specific? Evidence from a cross-sectional study of South Asian and European men. *PloS one*. 2013; 8(12):e82568. Epub 2013/12/19. <https://doi.org/10.1371/journal.pone.0082568> PMID: 24349313; PubMed Central PMCID: PMC3859604.
20. Iliodromiti S, Ghouri N, Celis-Morales CA, Sattar N, Lumsden MA, Gill JM. Should Physical Activity Recommendations for South Asian Adults Be Ethnicity-Specific? Evidence from a Cross-Sectional Study of South Asian and White European Men and Women. *PloS one*. 2016; 11(8):e0160024. Epub 2016/08/17. <https://doi.org/10.1371/journal.pone.0160024> PMID: 27529339; PubMed Central PMCID: PMC4987009.
21. Chapman J, Qureshi N, Kai J. Effectiveness of physical activity and dietary interventions in South Asian populations: a systematic review. *The British journal of general practice: the journal of the Royal College of General Practitioners*. 2013; 63(607):e104–14. Epub 2013/04/09. <https://doi.org/10.3399/bjgp13X663064> PMID: 23561688; PubMed Central PMCID: PMC3553636.
22. Chowdhury TA, Hitman GA. Type 2 diabetes in people of South Asian origin: potential strategies for prevention. *The British Journal of Diabetes & Vascular Disease*. 2007; 7(6):279–82. <https://doi.org/10.1177/14746514070070060401>
23. Muilwijk M, Stronks K, Qureshi SA, Beune E, Celis-Morales C, Gill J, et al. Dietary and physical activity strategies to prevent type 2 diabetes in South Asian adults: protocol for a systematic review. *BMJ Open*. 2017; 7(6).
24. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *Annals of Internal Medicine*. 2009; 151(4):264–9. <https://doi.org/10.7326/0003-4819-151-4-200908180-00135> PMID: 19622511
25. Ramachandran A, Snehalatha C, Ram J, Selvam S, Simon M, Nanditha A, et al. Effectiveness of mobile phone messaging in prevention of type 2 diabetes by lifestyle modification in men in India: a prospective, parallel-group, randomised controlled trial. *The lancet Diabetes & endocrinology*. 2013; 1(3):191–8. Epub 2014/03/14. [https://doi.org/10.1016/s2213-8587\(13\)70067-6](https://doi.org/10.1016/s2213-8587(13)70067-6) PMID: 24622367.
26. Balagopal P, Kamamma N, Patel TG, Misra R. A community-based diabetes prevention and management education program in a rural village in India. *Diabetes Care*. 2008; 31(6):1097–104. Epub 2008/03/05. <https://doi.org/10.2337/dc07-1680> PMID: 18316397.
27. Balagopal P, Kamamma N, Patel TG, Misra R. A community-based participatory diabetes prevention and management intervention in rural India using community health workers. *The Diabetes educator*. 2012; 38(6):822–34. Epub 2012/10/04. <https://doi.org/10.1177/0145721712459890> PMID: 23033123.
28. Dutta D, Mondal SA, Choudhuri S, Maisnam I, Hasanoor Reza AH, Bhattacharya B, et al. Vitamin-D supplementation in prediabetes reduced progression to type 2 diabetes and was associated with decreased insulin resistance and systemic inflammation: An open label randomized prospective study from Eastern India. *Diabetes Research and Clinical Practice*. 2014; 103(3):e18–e23. <https://doi.org/10.1016/j.diabres.2013.12.044> PMID: 24456991.
29. Hegde SV, Adhikari P, Shetty S, Manjrekar P, D'Souza V. Effect of community-based yoga intervention on oxidative stress and glycemic parameters in prediabetes: A randomized controlled trial. *Complementary therapies in medicine [Internet]*. 2013; 21(6):[571–6 pp.]. Available from: <http://online.library.wiley.com/doi/10.1002/ctm.120913> PMID: 24280463

30. Hurst PR, Stonehouse W, Coad J. Vitamin D supplementation reduces insulin resistance in South Asian women living in New Zealand who are insulin resistant and vitamin D deficient—a randomised, placebo-controlled trial. *British journal of nutrition* [Internet]. 2010; 103(4):[549–55 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1469-7580.2010.02415.x> http://journals.cambridge.org/download.php?file=%2FBJN%2FBJN103_04%2FS0007114509992017a.pdf&code=4665078f8beee0f8b810d4d41529542f. <https://doi.org/10.1017/S0007114509992017> PMID: 19781131
31. Andersen E, Høstmark AT, Holme I, Anderssen SA. Intervention Effects on Physical Activity and Insulin Levels in Men of Pakistani Origin Living in Oslo: A Randomised Controlled Trial. *Journal of Immigrant and Minority Health*. 2013; 15(1):101–10. <https://doi.org/10.1007/s10903-012-9686-3> PMID: 22828963
32. McDermott KA, Rao MR, Nagarathna R, Murphy EJ, Burke A, Nagendra RH, et al. A yoga intervention for type 2 diabetes risk reduction: a pilot randomized controlled trial. *BMC complementary and alternative medicine* [Internet]. 2014; 14:[212 p.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1186/s12900-014-0142-2> <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4096416/pdf/1472-6882-14-212.pdf>.
33. Telle-Hjellset V, Kjollesdal MKR, Bjorge B, Holmboe-Ottesen G, Wandel M, Birkeland KRI, et al. The InnvaDiab-DE-PLAN study: a randomised controlled trial with a culturally adapted education programme improved the risk profile for type 2 diabetes in Pakistani immigrant women. *British Journal of Nutrition*. 2013; 109(3):529–38. <https://doi.org/10.1017/S000711451200133X> PubMed PMID: WOS:000315454700017. PMID: 22717286
34. Islam NS, Zanowiak JM, Wyatt LC, Kavathe R, Singh H, Kwon SC, et al. Diabetes prevention in the New York City Sikh Asian Indian community: a pilot study. *International journal of environmental research and public health*. 2014; 11(5):5462–86. Epub 2014/05/24. <https://doi.org/10.3390/ijerph110505462> PMID: 24852392; PubMed Central PMCID: PMC4053907.
35. Madsen C, Mogensen P, Thomas N, Christensen DL, Bygbjerg IC, Mohan V, et al. Effects of an outdoor bicycle-based intervention in healthy rural Indian men with normal and low birth weight. *Journal of developmental origins of health and disease*. 2015; 6(1):27–37. Epub 2014/12/18. <https://doi.org/10.1017/S2040174414000609> PMID: 25514893.
36. Ramachandran A, Snehalatha C, Mary S, Selvam S, Kumar CKS, Seeli AC, et al. Pioglitazone does not enhance the effectiveness of lifestyle modification in preventing conversion of impaired glucose tolerance to diabetes in Asian Indians: results of the Indian Diabetes Prevention Programme-2 (IDPP-2). *Diabetologia*. 2009; 52(6):1019–26. <https://doi.org/10.1007/s00125-009-1315-x> PMID: 19277602
37. Islam MR, Attia J, Ali L, McEvoy M, Selim S, Sibbritt D, et al. Zinc supplementation for improving glucose handling in pre-diabetes: A double blind randomized placebo controlled pilot study. *Diabetes Res Clin Pract*. 2016; 115:39–46. Epub 2016/06/01. <https://doi.org/10.1016/j.diabres.2016.03.010> PMID: 27242121.
38. Patel RM, Misra R, Raj S, Balasubramanyam A. Effectiveness of a Group-Based Culturally Tailored Lifestyle Intervention Program on Changes in Risk Factors for Type 2 Diabetes among Asian Indians in the United States. *Journal of Diabetes Research*. 2017; 2017:13. <https://doi.org/10.1155/2017/2751980> PMID: 28168201
39. Thirunavukkarasu D, Kirubamani NH, Naidu MB. Effect of soy flour intake on systemic blood pressure and glycemic control in post-menopausal women with pre-diabetes and prehypertension. *Indian Journal of Pharmaceutical Education and Research*. 2017; 51(2):349–54. <https://doi.org/10.5530/ijper.51.2.42>
40. Worldbank. *Migration and Remittances Factbook*. 2011.
41. National Institute of Nutrition (NIN). *Dietary Guidelines for Indians—A Manual*. Indian Council of Medical Research; 2011.
42. Thomas H. *Quality Assessment Tool for Quantitative Studies*.: Effective Public Health Practice Project, McMaster University: Hamilton, ON, Canada; 2003.
43. Brouwers M, Kho ME, Browman GP, Burgers JS, Cluzeau F, Feder G, et al. AGREE II: Advancing guideline development, reporting and evaluation in healthcare. *Can Med Assoc J*. 2010. Epub July 5, 2010. <https://doi.org/10.1503/cmaj.090449> PMID: 20603348
44. Cuijpers P, Turner EH, Koole SL, van Dijke A, Smit F. What is the threshold for a clinically relevant effect? The case of major depressive disorders. *Depression and Anxiety*. 2014; 31(5):374–8. <https://doi.org/10.1002/da.22249> PMID: 24677535
45. Sehat A. *Healthy South Asian meals—The GI way: Apnee Sehat*; 2008 [11-4-2016]. Available from: <http://www.apneesehat.net>.
46. Misra R, Abraham N, Balagopal P, Batheja RS, Bhargava N, Chatterjee S, et al. *Indian Foods: AAPI's Guide to Nutrition, Health, and Diabetes*. American Association of Physicians of Indian Origin.: Sachdev, S.; 2011 [cited 2016 31-5-2016].
47. *The South Asian Diabetes Prevention Care Kit—A cultural and Language Relevant Resource*: Flemingdon Health Center; 2009 [10-03-2016]. Available from: <http://www.fhc-chc.com/diabetes/prevention>.

48. Misra A, Sharma R, Gulati S, Joshi SR, Sharma V, Ghafoorunissa, et al. Consensus dietary guidelines for healthy living and prevention of obesity, the metabolic syndrome, diabetes, and related disorders in Asian Indians. *Diabetes technology & therapeutics*. 2011; 13(6):683–94. Epub 2011/04/15. <https://doi.org/10.1089/dia.2010.0198> PMID: 21488798.
49. Hague HCotNT. Guidelines for a healthy diet: the ecological perspective. 2011.
50. Kemper HCG, Ooijendijk WTM, Stiggelbout M. Concensus over de Nederalndse norm voor gezond bewegen. *Tijdschrift voor Gezondheidswetenschappen (TSG)*. 2000; 78(3):180–3.
51. Mensink M, Corpeleijn E, Feskens EJ, Kruijshoop M, Saris WH, de Bruin TW, et al. Study on lifestyle-intervention and impaired glucose tolerance Maastricht (SLIM): design and screening results. *Diabetes Res Clin Pract*. 2003; 61(1):49–58. Epub 2003/07/10. PMID: 12849923.
52. Cené CW, Haymore LB, Ellis DF, Whitaker S, Henderson S, Lin F-C, et al. Implementation of the Power to Prevent Diabetes Prevention Educational Curriculum Into Rural African American Communities: A Feasibility Study. *The Diabetes educator*. 2013; 39(6):776–85. <https://doi.org/10.1177/0145721713507114> PubMed PMID: PMC4429528. PMID: 24129595
53. Lindström J, Louheranta A, Mannelin M, Rastas M, Salminen V, Eriksson J, et al. The Finnish Diabetes Prevention Study (DPS). Lifestyle intervention and 3-year results on diet and physical activity. 2003; 26(12):3230–6. <https://doi.org/10.2337/diacare.26.12.3230>
54. Counterweight Project Team. The implementation of the Counterweight Programme in Scotland, UK. *Family Practice*. 2012; 29(suppl 1):i139–i44. <https://doi.org/10.1093/fampra/cmr074> PMID: 22399544
55. National Diabetes Education Program. Power to Prevent: A Family Lifestyle Approach to Diabetes Prevention. Available from: <http://ndep.nih.gov/media/power-to-prevent-508.pdf>.
56. Islam NS, Wyatt LC, Patel SD, Shapiro E, Tandon SD, Mukherji BR, et al. Evaluation of a Community Health Worker Pilot Intervention to Improve Diabetes Management in Bangladeshi Immigrants With Type 2 Diabetes in New York City. *The Diabetes educator*. 2013; 39(4):478–93. <https://doi.org/10.1177/0145721713491438> PMID: 23749774
57. Comparative study of traditional aerobic exercise versus sprint interval training in prediabetes: a randomized controlled trial. CTRI/2012/09/002974. [Internet]. CTRI. 2012 [cited 25-5-2016]. Available from: <http://www.ctri.nic.in/Clinicaltrials/pmaindet2.php?trialid=5185>.
58. Diabetes prevention in people from Bangladesh: a pilot trial in East London. ISRCTN37781193 [Internet]. ISRCTN. 2006 [cited 25-5-2016]. Available from: <http://www.isrctn.com/ISRCTN37781193>.
59. DIAbetes PREvention programme [Internet]. 2011 [cited 25-5-216]. Available from: <http://www.ctri.nic.in/Clinicaltrials/pmaindet2.php?trialid=3345>.
60. Ethnic differences in glucose regulation following interval training. ISRCTN12337078 [Internet]. 2015 [cited 12-05-2016]. Available from: <http://www.isrctn.com/ISRCTN12337078>.
61. A Multicentric, Randomized, Controlled Trial of Yoga and Fenugreek in the Prevention of Type 2 Diabetes Mellitus-Indian Prevention of Diabetes Study (IPDS). CTRI/2013/01/03316. [Internet]. 2013 [cited 25/05/2016]. Available from: <http://www.ctri.nic.in/Clinicaltrials/pmaindet2.php?trialid=1017>.
62. Sathish T, Williams ED, Pasricha N, Absetz P, Lorgelly P, Wolfe R, et al. Cluster randomised controlled trial of a peer-led lifestyle intervention program: study protocol for the Kerala diabetes prevention program. *BMC Public Health*. 2013; 13(1):1–11. <https://doi.org/10.1186/1471-2458-13-1035> PMID: 24180316
63. Kerala Diabetes Prevention Program (K-DPP): A cluter RCT trial of its effectiveness and cost-effectiveness. ACTRN12611000262909 [Internet]. ANZCTR. 2011 [cited 12-05-2016]. Available from: <https://www.anzctr.org.au/Trial/Registration/TrialReview.aspx?ACTRN=12611000262909>.
64. Mobile Health Technology for the Prevention of Type 2 Diabetes. CTRI/2015/07/006011 [Internet]. Clinical Trials Registry—India. 2015 [cited 12-05-2016].
65. Prevention of Type 2 Diabetes With Vitamin D Supplementation (PREVENT-WIN). NCT02513888 [Internet]. U.S. National Institutes of Health. 2015 [cited 12-5-2016].
66. Ranasinghe P, Jayawardena R, Pigera AS, Katulanda P, Constantine GR, Galappaththy P. Zinc supplementation in pre-diabetes: study protocol for a randomized controlled trial. *Trials*. 2013; 14:52. Epub 2013/02/21. <https://doi.org/10.1186/1745-6215-14-52> PMID: 23421759; PubMed Central PMCID: PMC3599871.
67. Effects of Zinc supplementation on pre-diabetes: A double-blind, randomized, placebo-controlled trial. SLCTR/2012/010 [Internet]. Sri Lanka Clinical Trials Registry [Internet]. 2012 [cited 12-5-2016].
68. Assessment of lifestyle intervention in men and women (24–50 years) with impaired glucose tolerance (IGT) in Mumbai. CTRI/2015/06/005832 [Internet]. Clinical Trials Registry—India. 2015 [cited 12-05-2016]. Available from: <http://ctri.nic.in/Clinicaltrials/showallp.php?mid1=11141&Enchid=&userName=CTRI/2015/06/005832>.

69. Diabetes Prevention Using SMS Technology. NCT01795833 [Internet]. U.S. National Institutes of Health. 2013 [cited 12-5-2016]. Available from: <https://clinicaltrials.gov/ct2/show/NCT01795833>.
70. A Prospective, Open Label, Parallel Group, Randomized Controlled clinical study to evaluate the effect of flavoured oats and blended oil with antioxidants in pre-diabetes subjects. CTRI/2015/12/006429 [Internet]. National Institute of Medical Statistics. 2015 [cited 14/03/2016]. Available from: <http://ctri.nic.in>.
71. Can vitamin D replacement reduce insulin resistance in South Asians with vitamin D deficiency? ISRCTN18522562. [Internet]. ISRCTN. 2011. Available from: <http://www.isrctn.com/ISRCTN18522562?q=&filters=conditionCategory:Nutritional%5C,%20Metabolic%5C,%20Endocrine,ageRange:Adult,recruitmentCountry:United%20Kingdom&sort=&offset=147&totalResults=389&page=2&pageSize=100&searchType=basic-search>.
72. Anjana RM, Sudha V, Nair DH, Lakshmi priya N, Deepa M, Pradeepa R, et al. Diabetes in Asian Indians-How much is preventable? Ten-year follow-up of the Chennai Urban Rural Epidemiology Study (CURES-142). *Diabetes Res Clin Pract*. 2015; 109(2):253–61. Epub 2015/06/06. <https://doi.org/10.1016/j.diabres.2015.05.039> PMID: 26044609.
73. Agrawal S, Ebrahim S. Association between legume intake and self-reported diabetes among adult men and women in India. *BMC Public Health*. 2013; 13:706. Epub 2013/08/07. <https://doi.org/10.1186/1471-2458-13-706> PMID: 23915141; PubMed Central PMCID: PMC3735389.
74. Ranasinghe CD, Ranasinghe P, Jayawardena R, Misra A. Physical activity patterns among South-Asian adults: a systematic review. *International Journal of Behavioral Nutrition and Physical Activity*. 2013; 10(1):116. <https://doi.org/10.1186/1479-5868-10-116>.
75. Kandola K, Sandhu S, Tang T. Immigration and dietary patterns in South Asian Canadians at risk for diabetes. *Journal of Diabetes and its Complications*. 2016; 30(8):1462–6. <https://doi.org/10.1016/j.jdiacomp.2016.08.003> PMID: 27591030
76. Green R, Milner J, Joy EJ, Agrawal S, Dangour AD. Dietary patterns in India: a systematic review. *The British journal of nutrition*. 2016; 116(1):142–8. Epub 2016/05/06. <https://doi.org/10.1017/S0007114516001598> PMID: 27146890; PubMed Central PMCID: PMC4890343.
77. Holmboe-Ottesen G, Wandel M. Changes in dietary habits after migration and consequences for health: a focus on South Asians in Europe. *Food & Nutrition Research*. 2012; 56: <https://doi.org/10.3402/fnr.v56i0.18891> PubMed PMID: PMC3492807. PMID: 23139649
78. Agrawal S, Millett CJ, Dhillion PK, Subramanian S, Ebrahim S. Type of vegetarian diet, obesity and diabetes in adult Indian population. *Nutrition Journal*. 2014; 13(1):1–18. <https://doi.org/10.1186/1475-2891-13-89> PMID: 25192735
79. Andersen E, Ekelund U, Anderssen SA. Effects of reducing sedentary time on glucose metabolism in immigrant Pakistani men. *Med Sci Sports Exerc*. 2015; 47(4):775–81. Epub 2014/07/25. <https://doi.org/10.1249/MSS.0000000000000460> PMID: 25058329.
80. Gill JM, Bhopal R, Douglas A, Wallia S, Bhopal R, Sheikh A, et al. Sitting time and waist circumference are associated with glycemia in U.K. South Asians: data from 1,228 adults screened for the PODOSA trial. *Diabetes Care*. 2011; 34(5):1214–8. Epub 2011/04/06. <https://doi.org/10.2337/dc10-2313> PMID: 21464463; PubMed Central PMCID: PMC3114490.
81. Ram J, Selvam S, Snehalatha C, Nanditha A, Simon M, Shetty AS, et al. Improvement in diet habits, independent of physical activity helps to reduce incident diabetes among prediabetic Asian Indian men. *Diabetes Res Clin Pract*. 2014; 106(3):491–5. Epub 2014/12/03. <https://doi.org/10.1016/j.diabres.2014.09.043> PMID: 25458326.
82. Davidson EM, Liu JJ, Bhopal RS, White M, Johnson MRD, Netto G, et al. Consideration of ethnicity in guidelines and systematic reviews promoting lifestyle interventions: a thematic analysis. *The European Journal of Public Health*. 2013. <https://doi.org/10.1093/eurpub/ckt093> PMID: 23893129
83. How to prevent diabetes 2016 [25-09-2015]. Available from: <http://www.nepaldiabetesassociation.com/>.
84. Diet chart to prevent and manage diabetes: Ameen Diabetic Association; 2015 [06-10-2015]. Available from: <http://www.diabetes.org.pk>.
85. Choose Healthy Stay Healthy: Diabetes Foundation India; 2016 [31-5-2016]. Available from: <http://www.diabetesfoundationindia.org/about.aspx#>.
86. Misra A, Nigam P, Hills AP, Chadha DS, Sharma V, Deepak KK, et al. Consensus physical activity guidelines for Asian Indians. *Diabetes technology & therapeutics*. 2012; 14(1):83–98. Epub 2011/10/13. <https://doi.org/10.1089/dia.2011.0111> PMID: 21988275.
87. Armijo-Olivo S, Stiles CR, Hagen NA, Biondo PD, Cummings GG. Assessment of study quality for systematic reviews: a comparison of the Cochrane Collaboration Risk of Bias Tool and the Effective Public Health Practice Project Quality Assessment Tool: methodological research. *Journal of evaluation in clinical practice*. 2012; 18(1):12–8. Epub 2010/08/12. <https://doi.org/10.1111/j.1365-2753.2010.01516.x> PMID: 20698919.