



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Imaging radial velocity planets with SPHERE

Citation for published version:

Zurlo, A, Mesa, D, Desidera, S, Messina, S, Gratton, R, Moutou, C, Beuzit, JL, Biller, B, Boccaletti, A, Bonavita, M, Bonnefoy, M, Bhowmik, T, Brandner, W, Buenzli, E, Chauvin, G, Cudel, M, D'Orazi, V, Feldt, M, Hagelberg, J, Janson, M, Lagrange, AM, Langlois, M, Lannier, J, Lavie, B, Lazzoni, C, Maire, AL, Meyer, M, Mouillet, D, Peretti, S, Perrot, C, Potiron, PJ, Salter, G, Schmidt, T, Sissa, E, Vigan, A, Delboulb , A, Petit, C, Ramos, J, Rigal, F & Rochat, S 2018, 'Imaging radial velocity planets with SPHERE' Monthly Notices of the Royal Astronomical Society, vol. 480, no. 1, pp. 35-48. DOI: 10.1093/mnras/sty1809

Digital Object Identifier (DOI):

[10.1093/mnras/sty1809](https://doi.org/10.1093/mnras/sty1809)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Monthly Notices of the Royal Astronomical Society

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Imaging radial velocity planets with SPHERE

A. Zurlo,^{1,2,3*} D. Mesa,^{4,5} S. Desidera,⁵ S. Messina,⁶ R. Gratton,⁵ C. Moutou,³
 J.-L. Beuzit,⁷ B. Biller,⁸ A. Boccaletti,⁹ M. Bonavita,⁸ M. Bonnefoy,⁷ T. Bhowmik,⁹
 W. Brandner,¹⁰ E. Buenzli,¹¹ G. Chauvin,⁷ M. Cudel,⁹ V. D’Orazi,⁵ M. Feldt,¹⁰
 J. Hagelberg,¹² M. Janson,^{10,13} A.-M. Lagrange,⁷ M. Langlois,^{3,14} J. Lannier,⁷
 B. Lavie,¹² C. Lazzoni,⁵ A.-L. Maire,¹⁰ M. Meyer,¹⁵ D. Mouillet,⁷ S. Peretti,¹²
 C. Perrot,⁹ P. J. Potiron,¹⁶ G. Salter,³ T. Schmidt,⁹ E. Sissa,⁵ A. Vigan,³ A. Delboulb e,⁷
 C. Petit,¹⁷ J. Ramos,¹⁰ F. Rigal¹⁸ and S. Rochat⁷

Affiliations are listed at the end of the paper

Accepted 2018 July 3. Received 2018 July 02; in original form 2018 May 14

ABSTRACT

We present observations with the planet finder Spectro-Polarimetric High-contrast Exoplanet REsearch (SPHERE) of a selected sample of the most promising radial velocity (RV) companions for high-contrast imaging. Using a Monte Carlo simulation to explore all the possible inclinations of the orbit of wide RV companions, we identified the systems with companions that could potentially be detected with SPHERE. We found the most favourable RV systems to observe are: HD 142, GJ 676, HD 39091, HIP 70849, and HD 30177 and carried out observations of these systems during SPHERE Guaranteed Time Observing. To reduce the intensity of the starlight and reveal faint companions, we used principal component analysis algorithms alongside angular and spectral differential imaging. We injected synthetic planets with known flux to evaluate the self-subtraction caused by our data reduction and to determine the 5σ contrast in the J band versus separation for our reduced images. We estimated the upper limit on detectable companion mass around the selected stars from the contrast plot obtained from our data reduction. Although our observations enabled contrasts larger than 15 mag at a few tenths of arcsec from the host stars, we detected no planets. However, we were able to set upper mass limits around the stars using AMES-COND evolutionary models. We can exclude the presence of companions more massive than $25\text{--}28 M_{\text{Jup}}$ around these stars, confirming the substellar nature of these RV companions.

Key words: instrumentation: spectrographs – methods: data analysis – techniques: imaging spectroscopy – techniques: radial velocities – stars: individual: HD 142, HIP 70849, GJ 676A, HD 39091 – planetary systems.

1 INTRODUCTION

So far, 751 planets have been discovered with the radial velocity (RV) technique.¹ As stellar activity also produces RV variability and can mimic planet signals, these objects’ hosts are old and non-active stars. This method can detect both close planets and massive long-period objects, but, due to the unknown inclination of the detected companions, only the minimum mass can be determined. The measured parameter is the mass of the companion multiplied

by the sine of the inclination of its orbit, $M \sin i$. As the inclination is unknown from the RV measurements alone, the real mass of the object cannot be directly measured using this technique alone. Combining RV and high-contrast imaging measurements of the same companion allows us to constrain the companion orbit and thus measure its dynamical mass (see e.g. Boden, Torres & Latham 2006), providing a crucial benchmark for evolutionary models of substellar objects (e.g. Baraffe et al. 2015).

In the last years, a number of surveys have been conducted to image previously detected RV companion objects on wide orbits, for instance, direct imaging observations of targets with RV drifts from the High Accuracy Radial velocity Planet Searcher (HARPS) and CORALIE employing Very Large Telescope (VLT)/NACO (Hagel-

* E-mail: alice.zurlo@mail.udp.cl

¹ www.exoplanet.eu

berg 2010), the Targetting Benchmark-objects with Doppler Spectroscopy (TREND) imaging survey of targets with known trends in the RV (see e.g. Crepp et al. 2012b), and the Near-Infrared Coronagraphic Imager (NICI) follow-up survey on long-period RV targets (Salter et al. 2014). So far no planetary companion has both a measurement of dynamical mass and an estimated mass from evolutionary models. The failure of previous attempts is due to the small apparent separation between the planet and the star. The dynamical mass of the brown dwarfs HR 7672 B (Liu et al. 2002) and HD 4747 B (Crepp et al. 2016) has been successfully measured via the coupling of the direct imaging and the RV technique (Crepp et al. 2012a, 2018; Peretti et al. 2018). The combinations of RV and direct imaging have allowed a better constrain on the atmospheric properties of the companion as the constrain on the mass helps to solve degeneracies in the atmospheric models. Since planets with longer and longer periods (and separations) are being discovered by RV surveys extending over longer time spans, repeating this attempt with the latest discoveries offers a higher probability of success relative to previous attempts. Constraining mass, age, and chemical composition in this manner presents a unique opportunity to calibrate theoretical evolutionary models that are fundamental in describing disc instability (DI) companions (Potter et al. 2002).

With this aim, we selected a sample of targets with known companion objects found via the RV method that could potentially be imaged with the Spectro-Polarimetric High-contrast Exoplanet REsearch (SPHERE) instrument (Beuzit et al. 2006) at the VLT on Cerro Paranal. SPHERE is a highly specialized instrument dedicated to high-contrast imaging, built by a wide consortium of European laboratories. It is based on the SAXO, extreme adaptive optics system (Fusco et al. 2006; Petit et al. 2014; Sauvage et al. 2014), with a 41×41 actuators wavefront control, pupil stabilization, differential tip-tilt control, and employs stress polished toric mirrors for beam transportation (Hugot et al. 2012). Several coronagraphic devices for stellar diffraction suppression are provided, including apodized Lyot coronagraphs (Soummer 2005; Guerri et al. 2011) and achromatic four-quadrant phase mask (Boccaletti et al. 2008). The instrument is equipped with three science channels: the Infrared Differential Imaging Camera (IRDIS; Dohlen et al. 2008), an integral field spectrograph (IFS; Claudi et al. 2008), and the Zimpol rapid-switching Imaging Polarimeter (ZIMPOL; Thalmann et al. 2008). The system includes a dedicated data pipeline capable of applying basic reduction steps and higher level differential imaging analysis procedures (Pavlov et al. 2008).

Since old planetary mass objects at large separation from the host star are cool and extremely faint at optical wavelengths, in this paper we utilized IRDIS and IFS, which operate in the infrared (IR). These instruments have been designed to be particularly sensitive to the detection of T dwarf companions, the spectral type that we expect for RV companions. The performance of these two instruments has been widely presented in recent results as e.g. Vigan et al. (2016), Maire et al. (2016a), Zurlo et al. (2016), and Bonnefoy et al. (2016). Based on these results, we expected to reach a contrast of 10^{-6} at a separation of 0.5 arcsec with IFS, which would allow us to detect objects of several Jupiter masses around old > 1 Gyr stars.

We performed a Monte Carlo simulation, presented in Section 2, to select the most promising RV companions observable with SPHERE. We found five companions with a non-null detection probability: HD 142 Ac, GJ 676 b and GJ 676 c, HIP 70849 b, and HD 30177 b. We also added the star HD 39091, which has constraints from astrometry rather than RV but is still an intriguing target. These targets have been observed during the Guaranteed Time Observing (GTO) campaign in 2014–2017.

We present the selection criteria for our RV sample observed with SPHERE in Section 2. Observations and data analysis are described in Section 3; followed by the analysis of each individual target: HD 142 (Section 4), GJ 676 (Section 5), HD 39091 (Section 6), HIP 70849 (Section 7), and HD 30177 (Section 8) and Conclusions (Section 9).

2 SELECTION OF THE RV TARGETS

In 2014, we selected a sample of long-period RV planets to include in SPHERE GTO. The input sample was composed of all RV planets with a projected semimajor axis (as listed in the input catalogues) from the host star larger than 0.1 arcsec and with a declination lower than 40° (to be observable from the VLT). The selected minimum separation was chosen to match the projected radius of the smallest apodized Lyot coronagraph available with SPHERE, which is 145 mas. Only planets with known parameters of the orbit have been included, while objects with RV drifts have been excluded. Using a Monte Carlo simulation we explored the intrinsic magnitude of each companion and its projected separation from its host star as a function of its unknown orbital inclination. The Monte Carlo simulation explores all the possible inclinations and values of the longitude of ascending node, Ω_{node} , while the other orbital parameters are fixed by the orbital solution. The projected separation between each star and its companion at a given epoch is calculated with the Thiele–Innes formalism (e.g. Binnendijk 1960). Two epochs have been explored: the apostron passage, which in the majority of the cases is the moment where the planet is at its farthest position, and the year 2014, when the GTO started.

Several objects within the sample have constraints on the inclination from astrometry (see e.g. Reffert & Quirrenbach 2011) that were included in the simulation. The simulation explores different inclinations of the orbit, providing the value of the true mass of the planet for each inclination. We used the AMES-COND models (Allard et al. 2003) to estimate the intrinsic magnitude of each planet given its true mass.

Because all the objects under consideration are older than ~ 1 Gyr, we do not expect the choice of the model to be crucial, as at ages ≥ 1 Gyr, models with different values of the initial entropy converge (see e.g. Marley et al. 2007). The magnitude strongly depends on the age of the system, thus the error bars are correlated with the uncertainty on the age of the host star. For each target, we calculated the contrast limit reachable with IFS using the official ESO Exposure Time Calculator (ETC), assuming 1 h of exposure time, as requested for the observations. We considered only the IFS curve as it reaches deeper contrasts at small separations compared to IRDIS.

In Fig. 1, we show an example of the output of the Monte Carlo simulation. The green dashed curve represents the expected contrast achievable with IFS in *J* band, for a 1-h observation as calculated using the ETC. The red dots in Fig. 1 represent the calculated contrast in *J* magnitude for different inclinations of the planetary orbit resulting in different planetary masses and different separations at the foreseen passage of the planet at the apostron (in this case at the beginning of 2020). The blue dots represent the same calculation made for the period of the scheduled SPHERE observations. We note that the ETC tends to be conservative – the actual contrast limits we can reach are ~ 2 mag better. The minimum mass listed in the plot is the minimum possible mass of the companion that would be detectable with SPHERE.

As we do not know the true inclination of each system, we calculate a detection probability for each planet by inferring the probability that the mass is larger than the minimum mass for




Figure 1. Contrast in J band of IFS for the planet HD 142 Ac. The blue dots represent the expected contrast of the planet for the observation date (2014 October) calculated with the Monte Carlo simulation with the inclination as a free parameter. The red dots represent the expected contrast of the planet during the apoastron passage. The error bars are related to the uncertainty on the age of the system. The green dashed line shows the predicted detection limit calculated with the ETC for IFS. In the legend of this figure we provide the detection probabilities at both epochs considered. As we will show in the next sections, the expected contrast limits are pessimistic.

detection. To this aim, we used the a posteriori probability distribution of $\sin i$ suggested by Ho & Turner (2011). The details of the detection probability calculation are found in Appendix A. We selected for observation all the objects that have a detection probability higher than 5 per cent. The most promising targets observable with SPHERE had detection probabilities spanning from ~ 6 to ~ 20 per cent: HD 142, GJ 676, HD 39091, HIP 70849, and HD 30177. We have continued to update all the RV information regarding the orbits of these objects after the first selection made in 2014.

3 OBSERVATIONS OF THE TARGETS

As part of the SPHERE GTO, we observed five promising targets with detection probabilities >5 per cent: HD 142, GJ 676, HD 39091, HIP 70849, and HD 30177. We adopted the same observing strategy for all objects: IRDIFS observations, with IRDIS in $H2H3$ bands ($H2$ centred at $1.587 \mu\text{m}$ and $H3$ centred at $1.667 \mu\text{m}$; Vigan et al. 2010) and IFS in YJ mode ($0.95\text{--}1.35 \mu\text{m}$; Zurlo et al. 2014), in pupil stabilized mode during meridian passage, to take advantage of angular differential imaging (ADI; Marois et al. 2006). The stars HD 142, HIP 70849, HD 39091, and HD 30177 have been observed once, while GJ 676 was observed twice in order to obtain astrometric follow-up for several candidate companions identified during the initial observing epoch (all candidate companions were background objects, see Appendix B). More details on each observation can be found in Table 1.

For all the data sets, data from each of the IRDIS dual-band filters images, $H2$ and $H3$, were reduced separately. After background subtraction and flat-fielding, we suppressed the residual speckle noise using ADI with the Karhunen–Loève Image Processing (KLIP) algorithm (Soumerai, Pueyo & Larkin 2012). The procedure is the same used to reduce the data for HR 8799, and we refer to the related paper Zurlo et al. (2016) for the full description.

Table 1. Observation details for the RV special GTO targets.

Target	Epoch	Seeing (arcsec)	ΔPA ($^\circ$)
HD 142	2014-10-13	0.9	35
GJ 676	2015-05-06	0.7	34
	2016-05-30	1.0	38
HD 39091	2015-12-20	0.6	24
HIP 70849	2015-05-05	0.9	34
HD 30177	2017-11-03	0.5	35




Figure 2. ADI reduction of SPHERE/IRDIS $H2H3$ channels for the object HD 142 A (the mean images of the two filters are shown). In the image, no candidate(s) with signal-to-noise ratio (S/N) above 5σ are detected. The stellar companion is not visible as only the inner part of the field of view (FOV) is shown. The scale indicates the contrast.

As an example of the typical results from this procedure, the reduced IRDIS image of HD 142 is shown in Fig. 2. We also performed ADI+spectral differential imaging (SDI; Lafrenière et al. 2007), obtaining similar results. The contrast reached is not as deep as that reached by IFS at close-in separations, thus we adopt the IFS results for the following analysis.

The IFS data were reduced with the data reduction and handling (DRH) recipes (Pavlov et al. 2008). The DRH produces a calibrated data cube for each input raw frame. The speckle subtraction on these data cubes was performed with KLIP and with a second principal component analysis (PCA) method utilizing a single value decomposition (SVD) algorithm adapted to the case of the SPHERE IFS (see Mesa et al. 2015). The median collapsed final result is displayed in Fig. 3, left-hand side, using 100 principal components. To select the appropriate number of principal components to apply, we performed PCA using various numbers of components and evaluated the final contrast after taking into account self-subtraction. In this manner we verified that 100 principal components maximize the achieved contrast in these data relative to more or fewer principal components.

The contrast limits for both instruments have been calculated using the same method as in Mesa et al. (2015) and are shown in Fig. 4. Self-subtraction is the most important issue to consider when calculating the final contrast obtained with high-contrast imaging methods. To properly account for it, we performed a number of




Figure 3. Left: result of the PCA SVD-based procedure for the IFS data for HD 142 A, after median combination along IFS’ 39 spectral channels and using 100 principal components in the data reduction. Right: final result of the PCA procedure for the IFS data for HD 142 A with simulated planets inserted at separations ranging between 0.2 and 0.8 arcsec and with $\log(\text{contrast})$ ranging from -6.1 to -5.6 with a step of 0.1 (increasing counter-clockwise). Again, 100 principal components were used in the data reduction.

different tests with synthetic planets. First, we inserted synthetic planets [created by scaling the off-axis point spread function (PSF)] at a variety of separations from the central star with star–planet contrasts very close to the detection limit estimated without accounting for self-subtraction. In this way we were able to evaluate the self-subtraction caused by our method and subsequently to correct the contrast previously calculated. Second, we introduced simulated planets with contrast just above and below the contrast limit calculated with the previous step. Checking that the simulated planets inserted above the detection limit were indeed detected allows us to confirm the contrast curve obtained with the method described above. An example of the results of this test is provided in Fig. 3, right-hand panel. To define contrast curves for companions with both L and T spectral types, we injected simulated planets both with L0- (Testi et al. 2001) and T5-type spectrum (Burgasser et al. 2004). We note that we perform these detailed planet simulation and retrieval tests only for the IFS data, as the IFS data provide deeper contrasts at the separations of interest compared to the IRDIS data. The transmission of the coronagraph at close separations has been taken into account.

4 HD 142

The first target observed as part of this study, HD 142, is in fact a double-star system. The primary, HD 142 A, is a F7V star with a mass of $1.25 \pm 0.01 M_{\odot}$ (Bonfanti, Ortolani & Nascimbeni 2016) and a distance to the Earth of 26.2 ± 0.1 pc (Gaia Collaboration et al. 2018). The secondary, HD 142 B, is K8.5-M1.5 companion, with a smaller mass of $0.59 \pm 0.02 M_{\odot}$, at a projected separation of ~ 4 arcsec from the primary.

The primary, HD 142 A, has two RV-detected companion objects: HD 142 Ab has a $M \sin i$ of $1.03 M_{\text{Jup}}$ and a semimajor axis of 1.0 au, corresponding to ~ 0.04 arcsec (Tinney et al. 2002), while HD 142 Ac has a $M \sin i$ of $5.3 M_{\text{Jup}}$ and a semimajor axis of 6.8 au,

corresponding to ~ 0.26 arcsec, with an eccentricity of 0.21 and a period of 6005 d (Wittenmyer et al. 2012). For HD 142 Ac, our Monte Carlo simulations predicted a probability of detection of 11.30 per cent for the epoch of 2014 October. The orbital parameters used for the simulation are listed in Table 2. However, we stress that, given the very low minimum mass obtained from the RV measurement, we could realistically expect to be able to image HD 142 Ac only if the inclination of its orbit is quite far from being edge-on.

4.1 System age

As the age of the system strongly impacts the interpretation of the observational data, we reconsidered the age estimate for the HD 142 system here. Activity and rotation age indicators provide contradictory age estimates for this system: the low chromospheric activity (Jenkins et al. 2006; Ramírez, Meléndez & Asplund 2014) and non-detection in X-ray (Poppenhaeger, Robrade & Schmitt 2010) point to an age older than the Sun, when using Mamajek & Hillenbrand (2008) calibrations, however, the primary is a moderately fast rotator, with a rotational velocity of $v \sin i = 10.4 \pm 0.4 \text{ km s}^{-1}$ (Fischer & Valenti 2005; Butler et al. 2006), implying a younger age. Combining the projected rotational velocity and the stellar radius ($R = 1.44 \pm 0.14 R_{\odot}$), we derive an upper limit for the stellar rotation period $P_{\text{rot}} < 7 \pm 1$ d. This corresponds to an age of $< 1.3 \pm 0.3$ Gyr, using the gyro-chronology relations of Mamajek & Hillenbrand (2008).

On the basis of our derived upper limit on its rotation period and its F7V spectral type, we expect that HD 142 A may exhibit magnetic activity and, therefore, photometric variability arising from activity centres on its photosphere. The light rotational modulation induced by activity centres can be used to derive the rotation period using Fourier analysis. While this star has been included in a few photometric surveys, given its brightness the only unsaturated

Table 2. Orbital parameters for the planets of the systems presented in this analysis.

Planet	Period (d)	T_0 (JD - 240 0000)	e	ω ($^\circ$)	$M\sin(i)$ (M_{Jup})	Ref.
HD 142 c	6005 ± 477	$55\,954 \pm 223$	0.21 ± 0.07	250 ± 20	5.3 ± 0.7	Wittenmyer et al. (2012)
GJ 676 c	$7462.9^{+105.4}_{-101.4}$	$405.4^{+63.5}_{-65.6}$	0	–	6.9 ± 0.1	Sahlmann et al. (2016)
HIP 70849 b	$17\,349.4 \pm 600$	–	0.715 ± 0.245	–	9 ± 6	Ségransan et al. (2011)
HD 39091 b	2049 ± 150	$50\,073 \pm 150$	0.61 ± 0.03	330 ± 20	10.3	Jones et al. (2002)
HD 30771 c	$11\,613 \pm 1837$	$51\,660 \pm 573$	0.22 ± 0.14	11 ± 13	7.6 ± 3.1	Wittenmyer et al. (2017)

photometry available is from *Hipparcos*, which obtained 214 photometric measurements with a photometric precision $\sigma_V = 0.006$ mag between 1989 December 14 and 1993 March 17. We used the Lomb–Scargle (LS; Lomb 1976; Scargle 1982) and the CLEAN periodogram analysis techniques to search for significant periodicities (due to the stellar rotation period) in the magnitude time series. The false alarm probability (FAP = 1 – confidence level) associated with our detected period, which is the probability that a peak of given height in the periodogram is caused simply by statistical variations, i.e. Gaussian noise, was computed through Monte Carlo simulations, i.e. by generating 1000 artificial light curves obtained from the real light curve, keeping the date but permuting the magnitude values (see e.g. Herbst et al. 2002). We followed the method used by Lamm et al. (2004) to compute the errors associated with the period determination. No highly significant periodicities were detected for HD 142 A.

The discrepancy between age estimates based on rotation and activity is likely due to the fact that the temperature/spectral type of this star is close to the blue edge of the range of valid spectral types/colours used in the Mamajek & Hillenbrand (2008) age–activity calibration (valid for F7–K2 dwarfs, with $0.5 < B - V < 0.9$ mag and that its evolutionary phase is somewhat off of the main sequence; see below). Indeed, the observed rotational velocity is qualitatively consistent with rotation evolution model from van Saders & Pinsonneault (2013), for a star matching the spectroscopic parameters of HD 142 A from Ramírez et al. (2014). Lithium absorption features in the spectrum of this star do not provide a useful age constraint, as lithium is not particularly sensitive to stellar age for stars with temperatures similar to HD 142 A. However, a young stellar age (≤ 1 Gyr) can be ruled out by the kinematic parameters for this star (Nordström et al. 2004), well outside the space populated by young stars (Montes et al. 2001).

The most reliable age estimate can be obtained through isochrone fitting. From such an isochronal age analysis, Ramírez et al. (2014) recently derived an age of $2.6^{+0.2}_{-0.3}$ Gyr. This estimate is compatible with the value of 2.8 ± 0.5 Gyr found by Bonfanti et al. (2016). These estimates suggest an intermediate age for this system. As these isochronal age estimates are more accurate than the estimate based on other methods, we adopt an age of 2.6 Gyr for HD 142 in the following analysis.

4.2 Results

From our data reduction, we do not detect any companions either at the expected separation of 0.3 arcsec for planet c (blue curve in Fig. 1), or at any other separations. This lack of detections can be seen in the ADI image in Fig. 2 for IRDIS and in the PCA image in Fig. 3 for IFS.

Contrast limits in flux and J -band magnitude are shown in Fig. 4, after correcting for self-subtraction via the procedure described in Section 3. Adopting a stellar age of 2.6 Gyr, as justified in Section 4.1, we calculated the mass limits displayed in Fig. 6, and




Figure 4. 5σ residual noise levels as a function of separation for IRDIS observations of HD 142 A. The mean azimuthal profile of the off-axis PSF (black) and the coronagraphic profile (red) are shown for the two IRDIS channels $H2$ (continuous line) and $H3$ (dashed line). The IRDIS contrast plot after the implementation of ADI using the KLP algorithm is shown in green both for the $H2$ (solid line) and the $H3$ filters (dashed line). The light blue lines give the contrast obtained for IFS using the PCA algorithm. The solid line gives the contrast obtained injecting a T5-type spectrum planet, while the dashed line gives the contrast obtained injecting a L0-type spectrum planet.




Figure 5. 5σ residual noise levels as a function of separation for the IFS data only. The curves refer to the observations of GJ 676, HIP 70849, HD 39091, and HD 30177. Contrasts presented here were obtained with the PCA algorithm.




Figure 6. Mass limit versus the separation in au from the central star as obtained from converting the contrast curves displayed in Figs 4 and 5 to minimum detectable companion mass using the AMES-COND models.

Table 3. Summary table with the upper limit of the mass of planets around each of the five systems of this analysis at different separations from the star. The maximum possible inclination of each RV planet is also shown in the last column.

System	5 au	10 au	15 au	max(<i>i</i>)
HD 142 A	28 M_{Jup}	26 M_{Jup}	26 M_{Jup}	11°
GJ 676	25 M_{Jup}	25 M_{Jup}	25 M_{Jup}	16°
HIP 70849	28 M_{Jup}	26 M_{Jup}	26 M_{Jup}	20°
HD 39091	26 M_{Jup}	26 M_{Jup}	26 M_{Jup}	23°
HD 30177	30 M_{Jup}	29 M_{Jup}	28 M_{Jup}	15°

listed in Table 3, by converting our contrast limits to minimum detectable companion mass using the AMES-COND models (Allard et al. 2000). As previously mentioned, we used for this work only the contrast limits obtained with IFS, since the contrast we obtain with this subsystem is deeper than that gathered with IRDIS. At the expected separation for HD 142 Ac of 0.3 arcsec, we obtained an upper mass limit of $22.6^{+0.9}_{-1.7} M_{\text{Jup}}$ for a T-type spectrum and $23.6^{+0.9}_{-1.7} M_{\text{Jup}}$ for an L-type spectrum. Using the AMES-COND models we can infer an upper limit for T_{eff} of ~ 600 K for HD 142 Ac from the mass limit calculated in this work, while using the minimum mass obtained through the RV we can infer a minimum value for the T_{eff} of ~ 270 K.

Our upper mass limit for HD 142 Ac constrains the possible inclination of its orbit, excluding all inclinations smaller than $13.6^{+1.1}_{-0.5}$ from a pole-on orbit. Given the minimum mass of $5.3 M_{\text{Jup}}$ for this planet obtained through the RV technique and the upper mass limit obtained with SPHERE, the orbit of HD 142 Ac would have had to have been very close to the pole-on case in order for SPHERE to have been able to image it.

The Multi-purpose Exoplanet Simulation System (MESS; see Bonavita et al. 2012; Bonavita, de Mooij & Jayawardhana 2013) code was used to evaluate the probability of detection of companions around HD 142 A. The code is a recently developed tool with a well-demonstrated utility for analysis of exoplanet data (see e.g. Rameau et al. 2013; Chauvin et al. 2015).

4.3 HD 142 B

HD 142 A has a stellar companion, a K8.5-M1.5 star with a mass of $0.59 \pm 0.02 M_{\odot}$ (Eggenberger et al. 2007). Combining our SPHERE astrometry with astrometry from the literature, we can place new constraints on the orbital elements for this stellar companion. Since HD 142 B is strongly saturated in our coronagraphic images, we measured its position and contrast using non-coronagraphic IRDIS images that are normally taken for flux calibration purposes. For these flux calibration images, a neutral density filter is used, preventing saturation for both stars in this binary system. For astrometric calibrations (true north, plate-scale, distortion), we refer the reader to Maire et al. (2016b). The astrometric and photometric results for this object are shown in Table 4, together with the values available in the literature. We retrieved all the relative astrometric measurements available in the Washington Double Star Catalog (WDS; Mason et al. 2001), starting from 1894. To these data we added the results of Eggenberger et al. (2007) and Wittenmyer et al. (2012), as well as our own measurement, which are listed in Table 4. All these astrometric positions are displayed in Fig. 7 together with the error bars that we adopted to use with our Monte Carlo simulation of potential orbital parameters. As no error bar was given for the WDS point, we assumed a reasonable value for the given instrumentation of that epoch.

Using all the available astrometric points, we performed a Monte Carlo simulation of potential orbits, following the model by Zurlo et al. (2013) and Desidera et al. (2011). From the results of this simulation, we excluded the orbits that may cause instability in the system, following equation (1) of Holman & Wiegert (1999). Orbits where the critical semimajor axis is greater than the periastron of planet c were thus excluded from the results. Although equation (1) in Holman & Wiegert (1999) refers to circular and coplanar orbits, it is a good approximation to exclude solutions where the eccentricity is too high to assure the stability of the system.

From this Monte Carlo simulation, we constrain the orbital parameters of the stellar companion. The histograms of the resulting orbital parameter distributions are plotted in Fig. 8. The orbit of HD 142 B is most likely nearly edge-on (peak at 96°) and the semimajor axis is most likely around $a = 150$ au. A caveat to mention is that as we used a uniform linear distribution to generate the semimajor axis values, we might be biased towards long-period orbits. Considering the limits placed on system inclination from the non-detection of HD 142 Ac and from the measured orbit of the stellar companion HD 142 B, the orbits of HD 142 Ac and HD 142 B are compatible with coplanarity but substantial misalignments are also possible. If the coplanarity of HD 142 B and HD 142 Ac were confirmed, the mass of the latter would be very near to its minimum mass of $\sim 5 M_{\text{Jup}}$, beyond the limits of detection with current direct imaging instrumentation.

5 GJ 676

GJ 676 Ab (Forveille et al. 2011) is a substellar companion orbiting the binary star Gliese 676 A. The system is composed of two M stars with an angular separation of about 50 arcsec at a distance of 16.0 ± 0.1 pc (Gaia Collaboration et al. 2018); GJ 676 A has been classified as a MOV star (Koen et al. 2010).

As with HD 142, we reconsider the age estimate for GJ 676 A as well. Spectroscopic time series data collected by Suárez Mascareño et al. (2015) for this star revealed a rotation period $P = 41.2 \pm 3.8$ d. Suárez Mascareño et al. (2015) measured magnetic

Table 4. Astrometric positions of the stellar companion HD 142 B.

Epoch	PA ($^{\circ}$)	σ_{PA} ($^{\circ}$)	ρ (arcsec)	σ_{ρ} (arcsec)	Δmag	Ref.
1900.01	177.1	–	5.18	–	–	1
1900.86	175.9	–	5.44	–	–	1
1901.90	177.4	–	5.68	–	–	1
1914.77	174.6	–	5.49	–	–	1
1928.85	176.5	–	5.38	–	–	1
2004.4762	184.16	0.18	4.10	0.02	$\Delta H = 3.09 \pm 0.05$	2
2004.4788	184.16	0.18	4.11	0.02	$\Delta H = 3.52 \pm 0.07$	2
2004.8505	184.18	0.18	4.10	0.02	$\Delta H = 3.09 \pm 0.04$	2
2005.9373	184.13	0.29	4.08	0.02	$\Delta H = 2.89 \pm 0.15$	2
2010.5377	184.47	0.26	3.965	0.013	–	3
2014.7847	185.462	0.04	3.908	0.002	$\Delta H2 = 3.39 \pm 0.04$ $\Delta H3 = 3.08 \pm 0.08$	4

Note. References: 1 – Mason et al. (2001, WDS); 2 – Eggenberger et al. (2007); 3 – Wittenmyer et al. (2012); 4 – this work.

activity level for GJ 676 A via the calcium R'_{HK} index, obtaining a value of $\log R'_{\text{HK}} = -4.96 \pm 0.04$. The presence of activity cycles was reported by Suárez Mascareño, Rebolo & González Hernández (2016) with a period $P = 7.5 \pm 0.6$ yr and an amplitude of 0.08 mag from photometric time series. Gomes da Silva et al. (2012) also report activity cycles with a $P = 3.35$ yr based on Na I EW time series.

Using the Mamajek & Hillenbrand (2008) gyro–age relation, we inferred an age of 3.6 ± 0.5 Gyr, based on the known rotation period and the stellar colour of $B - V = 1.44$ mag (Koen et al. 2010). We derived a similarly old age 4.6 ± 0.4 Gyr from the gyro–age relation of Engle & Guinan (2011), which is valid for M0V stars. Based on the known level of chromospheric activity, we can also infer a chromospheric age of about 5.8 Gyr using the activity–age relation by Mamajek & Hillenbrand (2008). Despite the divergence of age estimates drawn from different methods, it is clear that GJ 676 has an age equal to or older than about 3.6 Gyr.

GJ 676 Ab has a minimum mass of $4.9 M_{\text{Jup}}$ and a period of 1056.8 d. The planet is on a low-eccentricity orbit, with no constraints on inclination from astrometry. Based on the Monte Carlo simulations described in Section 2, the minimum companion mass detectable by SPHERE is $41 M_{\text{Jup}}$ and the probability of detecting GJ 676 Ab is 10 per cent. Anglada-Escudé & Tuomi (2012) red-derived Doppler measurements in the HARPS-South data base for this system and discovered a number of additional candidate companions; GJ 676 A may hosts up to four substellar companions. Of these companions, GJ 676 Ac, which shows a trend in the RV, may also be observable with SPHERE, with a minimum mass of $19.3 M_{\text{Jup}}$. Forveille et al. (2011) provide orbital parameters for GJ 676 Ab, but the uncertainties on these values are quite high. Recently, Sahlmann et al. (2016) constrained the mass of planet b using astrometry. These authors observed the primary star with Focal Reducer and low dispersion Spectrograph 2 (FOR2) at VLT to obtain precise astrometric positions over a period of 2 yr starting in 2010 April and determined an orbit for planet b of 1052 d based on the motion of the star around the centre of mass of the system. From this value they derived a mass for this planet of $6.7^{+1.8}_{-1.5} M_{\text{Jup}}$. This value is compatible with our upper mass limit of $27 M_{\text{Jup}}$ for the planet based on our SPHERE observations. For planet c, we find an upper mass limit of $25 M_{\text{Jup}}$, but since the two objects are likely coplanar the expected mass for planet c would be close to its minimum mass of $6.8 M_{\text{Jup}}$ – thus, it is too small and faint to be imaged with current instruments, as the contrast needed would be of the order of almost 40 mag.




Figure 7. Astrometric positions of HD 142 B with respect to HD 142 A obtained from measurements starting from 1894. The error bars for each measurement are shown. More details are found in Table 4. The black star at the centre of the figure is the position of HD 142 A.

6 HD 39091

HD 39091 b (Jones et al. 2002) is a massive object with a minimum mass of $10.09 M_{\text{Jup}}$, long period (2151 d), and very eccentric (0.641) orbit. It orbits a G1 star at a distance of 18.28 ± 0.2 pc from the Earth (Gaia Collaboration et al. 2018). Reffert & Quirrenbach (2011) provide constraints on the astrometry of this system that limit the inclination of this object to 20° – 150° . Unfortunately, with this range of inclinations, this object is undetectable with SPHERE. The existing astrometric constraints are directly taken into account in the simulation, when generating all the possible random inclinations. For this target we reached a contrast of 10^{-6} at 0.3–0.8 arcsec (see Fig. 5). From this contrast limit, the companion must be more massive than $26 M_{\text{Jup}}$ at separations wider than 5 au as shown in Fig. 6 and listed in Table 3.

Bonfanti et al. (2016) estimate an age of 2.8 ± 0.8 Gyr for HD 39091 using the isochronal fitting method. Pace (2013) estimate a chromospheric age of 4.03 ± 1.33 Gyr, while Saffe, Gómez & Chavero (2005) derive chromospheric ages of 3.83 and 1.83 Gyr,




Figure 8. Histograms with the distribution of the orbital parameters of HD 142 B from our Monte Carlo simulation. The orbital parameters do not follow uniform distributions, but do present clear maximum probability values. Orbits that produce unstable systems have been excluded as explained in the text.

respectively, depending on the adopted calibration (either Donahue 1993 or Rocha-Pinto & Maciel 1998).

We used archival time series photometry to estimate a gyrochronological age for this system. As this star is relatively bright, the only time series photometry of HD 39091 available is from the *Hipparcos* archive (ESA 1997). We retrieved a total of 127 measurements collected between 1989 November and 1993 March, with an average photometric precision $\sigma_V = 0.006$ mag. We inferred a stellar radius $R = 1.14 R_\odot$ from the visual magnitude $V = 5.67$ mag (SIMBAD data base) for this star. Combining this stellar radius with the distance $d = 18.28$ pc [the bolometric correction $BC_V = -0.06$ mag (Pecaut & Mamajek 2013) appropriate for its G0V spectral type, and the average projected rotational velocity $v \sin i = 2.96$ km s $^{-1}$ (Delgado Mena et al. 2015)], we expect a rotation period $P \lesssim 20$ d. However, for the possible rotational light modulation to be detectable in the *Hipparcos* time series, the inclination of the stellar rotation axis must be $i \gtrsim 20^\circ$ – 30° . If this was the case, we would expect a much shorter rotation period $P \geq 5$ d. Thus, we carried out our LS and CLEAN periodogram analyses in the period range from 5 to 20 d and detected a period of $P = 18.3 \pm 1.0$ d with a 99 per cent confidence level, which can be interpreted as the stellar rotation period. We fit a sinusoid to the time series photometry phased by the rotation period, finding a peak-to-peak amplitude for the best-fitting sinusoid of $\Delta V = 0.008$ mag. Using the Mamajek &

Hillenbrand (2008) gyro-age relation, we inferred from the known rotation period and the stellar colour $B - V = 0.58$ mag (SIMBAD data base) an age of 3.4 ± 0.4 Gyr. Combining results from these three different age determination methods, we estimate the age of HD 39091 to be ~ 3.5 Gyr.

7 HIP 70849

HIP 70849 is an old (1–5 Gyr) K7V star, with a mass of $0.63 \pm 0.03 M_\odot$ (Ségransan et al. 2011) and a distance of 24.1 ± 0.1 pc (Gaia Collaboration et al. 2018) from the Earth. Ségransan et al. (2011) detected HIP 70849 b, a long-period (5–90 yr), high-eccentricity planet orbiting this star, with a mass between 3 and $15 M_{\text{Jup}}$. Lodieu et al. (2014) identified as well an extremely wide T4.5 dwarf companion at a distance of 6.3 arcmin, or ~ 9100 au, well outside of the field of view (FOV) of the IRDIS detector.

HIP 70849 exhibits clear evidence of magnetic activity from both photometric and chromospheric proxies. HIP 70849 was observed by SuperWASP from 2006 May to 2008 April. A total of 10 160 measurements were collected. The V -band magnitude time series exhibits a decreasing linear trend that was removed before our period search. After outlier removal and data binning (24-h bin width), we ran the LS and CLEAN periodograms on a total of 123 mean




Figure 9. Top panel: (left) ASAS V magnitude time series for HD 39091 versus Julian Day; (centre) Lomb-Scargle periodogram (black) with the window function overplotted (dotted red) and the power level corresponding to a 99 per cent confidence level. The vertical line marks the power peak corresponding to the stellar rotation period; (right) CLEAN periodogram. Bottom panel: ASAS V magnitude plotted versus rotation phase, computed using the period $P = 18.3$ d. The red solid line is a fit to the data with a sinusoidal function.




Figure 10. As in Fig. 9, but for the target HD 30177.

magnitudes and detected a significant (FAP < 0.01) power peak at $P = 41.2 \pm 0.4$ d in both periodograms. The sinusoidal fit to the light curve has an amplitude of $\Delta V = 0.007$ mag. The 41.2-d periodic light modulation is likely the stellar rotation period and arises from the presence of surface brightness inhomogeneities that are carried in and out of view by the stellar rotation. This star also shows a starspot cycle of $P_{\text{cyc}} = 10.1 \pm 1.4$ yr with an amplitude of $A_{\text{cyc}} = 6.9 \pm 0.9$ mmag (Suárez Mascareño et al. 2016).

To derive the basic physical parameters of this star we analysed the spectral energy distribution (SED). We used the *vosa* tool (Bayo et al. 2008) to build the observed SED, which was best fitted by a model spectrum from the BT-NextGen-GNS93 suite of models (Allard, Homeier & Freytag 2012) with $T_{\text{eff}} = 4000 \pm 50$ K, gravity $\log g = 4.5$, and metallicity $[\text{Fe}/\text{H}] = 0.3$. We additionally inferred a bolometric luminosity $L_{\text{bol}} = 0.0892 \pm 0.0005 L_{\odot}$ and a stellar radius $R = 0.62 \pm 0.02 R_{\odot}$. Rotation period, stellar radius, and the projected rotational velocity $v \sin i = 1.93 \text{ km s}^{-1}$ (Ségransan et al. 2011) can provide an estimate of the rotation axis inclination. However, we inferred an inconsistently large $\sin i = 2.7$, likely arising from an overestimated $v \sin i$ value. In fact, the combined effect on spectral lines by the macro- and microturbulence in main-sequence K7-type stars (Gray 1984; Husser et al. 2013), although relatively small ($< 1 \text{ km s}^{-1}$), is enough to explain the overestimated rotational velocity. Taking this into account, we infer an inclination $i \sim 90^{\circ}$.

Suárez Mascareño et al. (2016) used All Sky Automated Survey (ASAS) photometry to investigate the rotational and magnetic activity properties of this target. They detect a starspot cycle of $P_{\text{cyc}} = 10.1 \pm 1.4$ yr with an amplitude of $A_{\text{cyc}} = 6.9 \pm 0.9$ mmag. However, as this period is longer than the 8.8-yr time span of the ASAS time series data, it should be regarded as tentative. They also marginally detect periodic modulation of the light curve with an initial period estimate of $P_{\text{rot}} = 133.55 \pm 0.75$ d. This star has chromospheric emission with an index $R'_{\text{HK}} = -4.74 \pm 0.05$ and a projected rotational velocity $v \sin i = 1.93 \text{ km s}^{-1}$ measured from the HARPS cross-correlation function.

Using the age–rotation relationship from Mamajek & Hillenbrand (2008), we derive a gyrochronological age of 3.6 ± 0.15 Gyr from the $P = 41.2$ d rotation period. Using the age–activity relationship from Mamajek & Hillenbrand (2008), we derive a chromospheric age of 2.43 Gyr from the activity index $R'_{\text{HK}} = -4.74 \pm 0.05$. Finally, the relatively long starspot cycle collocates this star in the active branch of the rotation period–Rossby number relation, which is populated mainly by old field stars (Saar & Brandenburg 1999).

We found a detection probability of ~ 20 per cent for HIP 70849 b using our Monte Carlo simulation, however, no companion was detected with IFS or IRDIS in our SPHERE observations. Our observations obtained a mean contrast of 2×10^{-6} at 0.3–0.8 arcsec (see Fig. 5). Adopting a relatively old age for the system (as discussed above), the companion must be more massive than $26 M_{\text{Jup}}$ at separations wider than 7 au as shown in Fig. 6 and summarized in Table 3.

8 HD 30177

Wittenmyer et al. (2017) detected a candidate massive Saturn-analog orbiting the solar-type star HD 30177. The system was previously known to host an interior planet with a minimum mass of $9.7 M_{\text{Jup}}$; the outer planet has $m \sin i = 7.6 \pm 3.1 M_{\text{Jup}}$ and $a = 9.9 \pm 1.0$ au.

Isochronal and chromospheric age estimates of HD 30177 suggest this star is rather old. Bonfanti et al. (2016) and Ramírez et al.

(2012) estimated ages of 5.9 ± 1.1 and 6.18 ± 2 Gyr, respectively, via isochronal fitting, while Saffe et al. (2005) found a chromospheric age for the system of 8.30 or 1.50 Gyr, depending on the adopted calibration (Donahue 1993 or Rocha-Pinto & Maciel 1998, respectively). We used archival time series photometry to estimate a gyrochronological age as well for this system from the rotation period of the star. The stellar rotation period in late-type stars, as HD 30177, can be inferred from the period of the light rotational modulation arising from the presence of surface temperature inhomogeneities related to the magnetic activity. HD 30177 was observed by the ASAS (Pojmanski 1997) from 2000 to 2009. After removing outliers with a 3σ moving boxcar filter and discarding a few inaccurate ($\sigma_V \geq 0.05$ mag) data points, a time series of 510 measurements remained, characterized by an average photometric precision $\sigma_V = 0.03$ mag. This time series was analysed with the LS (Scargle 1982; Horne & Baliunas 1986) and the CLEAN (Roberts, Lehar & Dreher 1987) periodogram methods to search for significant periodicities.

HD 30177 has an estimated radius $R = 1.16 \pm 0.31 R_{\odot}$. This value is derived from the visual magnitude $V = 8.37$ mag (Høg et al. 2000), the distance $d = 55.7 \pm 0.1$ pc (Gaia Collaboration et al. 2018), and the bolometric correction $BC_V = -0.14$ (Pecaut & Mamajek 2013) corresponding to an average effective temperature $T_{\text{eff}} = 5596$ K (see table 1 in Wittenmyer et al. 2017). Combining the estimated radius and the projected rotational velocity $v \sin i = 2.96 \pm 0.5 \text{ km s}^{-1}$ (Butler et al. 2006), the stellar rotation period is expected to be shorter than about 30 d. On the other hand, for the possible light rotational modulation induced by surface inhomogeneities to be detectable, the stellar rotation axis of HD 30177 must have an inclination sufficiently far from a pole-on configuration (i.e. $i \geq 20^{\circ}$ – 30°). These circumstances made reasonable a search from periodicities in the photometric time series of HD 30177, covering a range of possible rotation periods from 10 to 30 d.

The LS periodogram analysis detected a period $P = 24.9 \pm 1$ d, which can be likely interpreted as the stellar rotation period, although only with a confidence level ≤ 90 per cent. However, the same period is also found by the CLEAN algorithm, rendering this period more believable.

The peak-to-peak amplitude of the sinusoid function used to fit the magnitudes phased with the rotation period is $\Delta V = 0.007$ mag. We note, for instance, that the very low amplitude of the photometric variability is consistent with the slow rotation of HD 30177, whose level of activity is expected consequently low. Combining the stellar radius, the rotation period, and the projected rotational velocity we derive an inclination of the stellar rotation axis close to $i \simeq 90^{\circ}$. This circumstance likely allowed us to detect the rotational light variation, despite the very low level of magnetic activity and the relatively small ratio of the light-curve amplitude over the photometric precision.

Using the Mamajek & Hillenbrand (2008) gyro–age relation, we inferred an age of 2.8 ± 0.3 Gyr from the known rotation period and the stellar colour $B - V = 0.81$ mag (Høg et al. 2000). This result is in qualitative agreement with the earlier results from the literature and points toward an old age for HD 30177.

From the contrast limit that we obtained with SPHERE, the companion must be more massive than $30 M_{\text{Jup}}$ at separations wider than 5 au as shown in Fig. 6 and listed in Table 3.

9 CONCLUSIONS

We present in this paper the analysis of a sample of RV systems imaged with SPHERE. We selected a sample of five promising

RV-detected, massive, wide orbit companions using a Monte Carlo simulation that explored all the possible inclinations for these objects. These five sample stars, specifically, HD 142 Ac, GJ 676 b and GJ 676 c, HD 39091 b, HIP 70849 b, and HD 30177 c, were observed during GTO using the NIR arm of SPHERE, reaching contrasts of $1\text{--}3 \times 10^{-6}$ at separations larger than 0.3 arcsec. We detected none of the known RV companions for these stars, however, for the system HD 142 AB, the known stellar companion was detected in the IRDIS FOV.

We thus obtained an additional high precision astrometric point for HD 142 B, which, when combined with literature astrometry, provides a baseline of 130 yr of observations. We used a Monte Carlo simulation to investigate the probability distribution of the parameters of its orbit, finding that the stellar companion most likely has an edge-on orbit with a semimajor axis of about 150 au. The orbit of HD 142 Ac might be coplanar with that of HD 142 B. If this is the case, the mass of HD 142 Ac may be very near to the minimum mass obtained through the RV measurements for this planet. For this reason, and given the mass limit found in this work, we can conclude that HD 142 Ac is considerably too low mass and too faint to be imaged with SPHERE.

From the contrast limits reached by SPHERE for the four other systems observed, we similarly conclude that the inclinations of these orbits are quite close to edge-on and the companion masses are close to the minimum mass given by RV measurements.

In the next few years, *Gaia* will provide astrometric measurements with precisions down to a few tens of microarcseconds. *Gaia* astrometry will strongly constrain the inclination of the orbits of these and other similar planets and will also yield numerous new planet discoveries. Given the performance of the current suite of planet imagers, detecting cool and old RV companions is highly challenging, however, with future 30-m class telescopes, these objects will be easily detected and characterized. This will eventually yield vital constraints on evolutionary models for planetary mass companions.

ACKNOWLEDGEMENTS

We want to thank the anonymous referee for comments that improved the quality of this paper. We are grateful to the SPHERE team and all the people at Paranal for the great effort during SPHERE early-GTO run. AZ acknowledges support from the CONICYT+PAI/Convocatoria nacional subvención a la instalación en la academia, convocatoria 2017+Folio PAI77170087. SD and RG acknowledge support from the ‘Progetti Premiali’ funding scheme of the Italian Ministry of Education, University, and Research. We acknowledge support from the French National Research Agency (ANR) through the GUEPARD project grant ANR10-BLANC0504-01.

SPHERE was funded by ESO, with additional contributions from CNRS (France), MPIA (Germany), INAF (Italy), FINES (Switzerland), and NOVA (Netherlands). SPHERE also received funding from the European Commission Sixth and Seventh Framework Programmes as part of the Optical Infrared Coordination Network for Astronomy (OPTICON) under grant number RII3-Ct-2004-001566 for FP6 (2004–2008), grant number 226604 for FP7 (2009–2012), and grant number 312430 for FP7 (2013–2016).

This research has made use of the Washington Double Star Catalog maintained at the US Naval Observatory. We thank Dr B. Mason for making the individual measurements of HD 142 available to us.

REFERENCES

- Allard F., Hauschildt P. H., Alexander D. R., Ferguson J. W., Tamanai A., 2000, in Griffith C. A., Marley M. S., eds, ASP Conf. Ser. Vol. 212, From Giant Planets to Cool Stars. Astron. Soc. Pac., San Francisco, p. 127
- Allard F., Guillot T., Ludwig H.-G., Hauschildt P. H., Schweitzer A., Alexander D. R., Ferguson J. W., 2003, in Martín E., ed., Proc. IAU Symp. 211, Brown Dwarfs. Astron. Soc. Pac., San Francisco, p. 325
- Allard F., Homeier D., Freytag B., 2012, *Philos. Trans. R. Soc. Lond. Ser. A*, 370, 2765
- Anglada-Escudé G., Tuomi M., 2012, *A&A*, 548, A58
- Baraffe I., Homeier D., Allard F., Chabrier G., 2015, *A&A*, 577, A42
- Bayo A., Rodrigo C., Barrado Y Navascués D., Solano E., Gutiérrez R., Morales-Calderón M., Allard F., 2008, *A&A*, 492, 277
- Beuzit J.-L. et al., 2006, *The Messenger*, 125, 29
- Binnendijk L., 1960, *Properties of Double Stars: A Survey of Parallaxes and Orbits*. Univ. Pennsylvania Press, Philadelphia, PA
- Boccaletti A. et al., 2008, *Proc. SPIE*, 7015, 70151B
- Boden A. F., Torres G., Latham D. W., 2006, *ApJ*, 644, 1193
- Bonavita M., Chauvin G., Desidera S., Gratton R., Janson M., Beuzit J. L., Kasper M., Mordasini C., 2012, *A&A*, 537, A67
- Bonavita M., de Mooij E. J. W., Jayawardhana R., 2013, *PASP*, 125, 849
- Bonfanti A., Ortolani S., Nascimbeni V., 2016, *A&A*, 585
- Bonnefoy M. et al., 2016, *A&A*, 587, A58
- Burgasser A. J., McElwain M. W., Kirkpatrick J. D., Cruz K. L., Tinney C. G., Reid I. N., 2004, *AJ*, 127, 2856
- Butler R. P. et al., 2006, *ApJ*, 646, 505
- Chauvin G. et al., 2015, *A&A*, 573, A127
- Claudi R. U. et al., 2008, *Proc. SPIE*, 7014, 70143E
- Crepp J. R. et al., 2012a, *ApJ*, 751, 97
- Crepp J. R. et al., 2012b, *ApJ*, 761, 39
- Crepp J. R., Gonzales E. J., Bechter E. B., Montet B. T., Johnson J. A., Piskorz D., Howard A. W., Isaacson H., 2016, *ApJ*, 831, 136
- Crepp J. R. et al., 2018, *ApJ*, 853, 192
- Delgado Mena E. et al., 2015, *A&A*, 576, A69
- Desidera S. et al., 2011, *A&A*, 533, A90
- Dohlen K. et al., 2008, *Proc. SPIE*, 7014, 70143L
- Donahue R. A., 1993, PhD thesis, New Mexico State University, University Park
- Eggenberger A., Udry S., Chauvin G., Beuzit J.-L., Lagrange A.-M., Ségransan D., Mayor M., 2007, *A&A*, 474, 273
- Engle S. G., Guinan E. F., 2011, in Qain S., Leung K., Zhu L., Kwok S., eds, ASP Conf. Ser. Vol. 451, 9th Pacific Rim Conference on Stellar Astrophysics. Astron. Soc. Pac., San Francisco, p. 285
- Fischer D. A., Valenti J., 2005, *ApJ*, 622, 1102
- Forveille T. et al., 2011, *A&A*, 526, A141
- Fusco T. et al., 2006, *Opt. Express*, 14, 7515
- Gaia Collaboration, Brown A. G. A., Vallenari A., Prusti T., de Bruijne J. H. J., Babusiaux C., Bailer-Jones C. A. L., 2018, *A&A*, in press ([arXiv:1804.09365](https://arxiv.org/abs/1804.09365))
- Gomes da Silva J., Santos N. C., Bonfils X., Delfosse X., Forveille T., Udry S., Dumusque X., Lovis C., 2012, *A&A*, 541, A9
- Gray D. F., 1984, *ApJ*, 281, 719
- Grether D., Lineweaver C. H., 2006, *ApJ*, 640, 1051
- Guéri G. et al., 2011, *Exp. Astron.*, 30, 59
- Hagelberg J., 2010, in Boccaletti A., ed., Proc. Conf. In the Spirit of Lyot 2010: Direct Detection of Exoplanets and Circumstellar Disks. University of Paris Diderot, Paris, p. 22
- Heinze A. N., Hinz P. M., Kenworthy M., Meyer M., Sivanandam S., Miller D., 2010, *ApJ*, 714, 1570
- Herbst W., Bailer-Jones C. A. L., Mundt R., Meisenheimer K., Wackermann R., 2002, *A&A*, 396, 513
- Ho S., Turner E. L., 2011, *ApJ*, 739, 26
- Høg E. et al., 2000, *A&A*, 355, L27
- Holman M. J., Wiegert P. A., 1999, *AJ*, 117, 621
- Horne J. H., Baliunas S. L., 1986, *ApJ*, 302, 757

Hugot E., Ferrari M., El Hadi K., Costille A., Dohlen K., Rabou P., Puget P., Beuzit J. L., 2012, *A&A*, 538, A139

Husser T.-O., Wende-von Berg S., Dreizler S., Homeier D., Reiners A., Barman T., Hauschildt P. H., 2013, *A&A*, 553, A6

Jenkins J. S. et al., 2006, *MNRAS*, 372, 163

Jones H. R. A., Paul Butler R., Tinney C. G., Marcy G. W., Penny A. J., McCarthy C., Carter B. D., Pourbaix D., 2002, *MNRAS*, 333, 871

Koen C., Kilkenny D., van Wyk F., Marang F., 2010, *MNRAS*, 403, 1949

Lafrenière D. et al., 2007, *ApJ*, 670, 1367

Lamm M. H., Bailer-Jones C. A. L., Mundt R., Herbst W., Scholz A., 2004, *A&A*, 417, 557

Liu M. C., Fischer D. A., Graham J. R., Lloyd J. P., Marcy G. W., Butler R. P., 2002, *ApJ*, 571, 519

Lodieu N., Pérez-Garrido A., Béjar V. J. S., Gauza B., Ruiz M. T., Rebolo R., Pinfield D. J., Martín E. L., 2014, *A&A*, 569, A120

Lomb N. R., 1976, *Ap&SS*, 39, 447

Maire A.-L. et al., 2016a, *A&A*, 587, A56

Maire A.-L. et al., 2016b, *Proc. SPIE*, 9908, 990834

Mamajek E. E., Hillenbrand L. A., 2008, *ApJ*, 687, 1264

Marley M. S., Fortney J. J., Hubickyj O., Bodenheimer P., Lissauer J. J., 2007, *ApJ*, 655, 541

Marois C., Lafrenière D., Doyon R., Macintosh B., Nadeau D., 2006, *ApJ*, 641, 556

Mason B. D., Wycoff G. L., Hartkopf W. I., Douglass G. G., Worley C. E., 2001, *AJ*, 122, 3466

Mesa D. et al., 2015, *A&A*, 576, A121

Metchev S. A., Hillenbrand L. A., 2009, *ApJS*, 181, 62

Montes D., López-Santiago J., Gálvez M. C., Fernández-Figueroa M. J., De Castro E., Cornide M., 2001, *MNRAS*, 328, 45

Nordström B. et al., 2004, *A&A*, 418, 989

Pace G., 2013, *A&A*, 551, L8

Pavlov A., Möller-Nilsson O., Feldt M., Henning T., Beuzit J.-L., Mouillet D., 2008, *Proc. SPIE*, 7019, 701939

Pecaut M. J., Mamajek E. E., 2013, *ApJS*, 208, 9

Peretti S. et al., 2018, preprint (arXiv:1805.05645)

Petit C. et al., 2014, *Proc. SPIE*, 9148, 914800

Pojmanski G., 1997, *Acta Astron.*, 47, 467

Poppenhaeger K., Robrade J., Schmitt J. H. M. M., 2010, *A&A*, 515, A98

Potter D., Martín E. L., Cushing M. C., Baudoz P., Brandner W., Guyon O., Neuhauser R., 2002, *ApJ*, 567, L133

Rameau J. et al., 2013, *A&A*, 553, A60

Ramírez I., Fish J. R., Lambert D. L., Allende Prieto C., 2012, *ApJ*, 756, 46

Ramírez I., Meléndez J., Asplund M., 2014, *A&A*, 561, A7

Reffert S., Quirrenbach A., 2011, *A&A*, 527, A140

Roberts D. H., Lehar J., Dreher J. W., 1987, *AJ*, 93, 968

Rocha-Pinto H. J., Maciel W. J., 1998, *MNRAS*, 298, 332

Saar S. H., Brandenburg A., 1999, *ApJ*, 524, 295

Saffé C., Gómez M., Chavero C., 2005, *A&A*, 443, 609

Sahlmann J. et al., 2016, *A&A*, 595, A77

Salter G. S., Tinney C. G., Wittenmyer R. A., Jenkins J. S., Jones H. R. A., O'Toole S. J., 2014, in Booth M., Matthews B. C., Graham J. R., eds, *Proc. IAU Symp. Vol. 299, Exploring the Formation and Evolution of Planetary Systems*. Cambridge Univ. Press, Cambridge, p. 66

Sauvage J.-F. et al., 2014, *Proc. SPIE*, 9148, 914847

Scargle J. D., 1982, *ApJ*, 263, 835

Ségransan D. et al., 2011, *A&A*, 535, A54

Soumer R., 2005, *ApJ*, 618, L161

Soumer R., Pueyo L., Larkin J., 2012, *ApJ*, 755, L28

Suárez Mascareño A., Rebolo R., González Hernández J. I., Esposito M., 2015, *MNRAS*, 452, 2745

Suárez Mascareño A., Rebolo R., González Hernández J. I., 2016, *A&A*, 595, A12

Testi L. et al., 2001, *ApJ*, 552, L147

Thalman C. et al., 2008, *Proc. SPIE*, 7014, 70143F

Tinney C. G., Butler R. P., Marcy G. W., Jones H. R. A., Penny A. J., McCarthy C., Carter B. D., 2002, *ApJ*, 571, 528

van Saders J. L., Pinsonneault M. H., 2013, *ApJ*, 776, 67

Vigan A., Moutou C., Langlois M., Allard F., Boccaletti A., Carillet M., Mouillet D., Smith I., 2010, *MNRAS*, 407, 71

Vigan A. et al., 2016, *A&A*, 587, A55

Wittenmyer R. A. et al., 2012, *ApJ*, 753, 169

Wittenmyer R. A. et al., 2017, *AJ*, 153

Zurlo A. et al., 2013, *A&A*, 554, A21

Zurlo A. et al., 2014, *A&A*, 572, A85

Zurlo A. et al., 2016, *A&A*, 587, A57

APPENDIX A: PROBABILITY OF DETECTION

Concerning the probability of detection, it is clear that massive objects, for a given age, have a higher chance of being detected. We do not know the true inclination for a given planet, but we can calculate the probability that the planet mass is greater than the minimum detectable mass by integrating over a reasonable distribution of potential inclination values. For this aim, we used the a posteriori probability distribution of $\sin(i)$ suggested by Ho & Turner (2011). This inclination distribution a priori is isotropic with the constraints given by Reffert & Quirrenbach (2011).

The a priori probability distribution function of the observed mass M_0 given the real mass M_T is

$$P(M_0 | M_T) = \frac{M_0/M_T^2}{\sqrt{1 - \left(\frac{M_0}{M_T}\right)^2}}. \quad (\text{A1})$$

If we want the posterior probability, using Bayes' theorem,

$$P(A | B) = \frac{P(B | A)P(A)}{P(B)}, \quad (\text{A2})$$

and assuming that

$$P(M_0) = \int P(M_0 | M_T)P(M_T)dM_T, \quad (\text{A3})$$

we finally obtain

$$P(M_T | M_0) = \frac{P(M_0 | M_T)P(M_T)}{\int P(M_0 | M_T)P(M_T)dM_T}, \quad (\text{A4})$$

that is the posterior distribution of the probability. We can write it as

$$P(M_T | M_0) = \frac{\frac{M_0/M_T^2}{\sqrt{1 - \left(\frac{M_0}{M_T}\right)^2}} P(M_T)}{\int \frac{M_0/M_T^2}{\sqrt{1 - \left(\frac{M_0}{M_T}\right)^2}} P(M_T)dM_T}. \quad (\text{A5})$$

If we want the probability of finding a mass greater than a value X :

$$P(M_T > X | M_0) = \frac{\int_{M_0}^X \frac{M_0/M_T^2}{\sqrt{1 - \left(\frac{M_0}{M_T}\right)^2}} P(M_T)dM_T}{\int_{M_0}^{M_{\max}} \frac{M_0/M_T^2}{\sqrt{1 - \left(\frac{M_0}{M_T}\right)^2}} P(M_T)dM_T}. \quad (\text{A6})$$

Finally, to calculate the probability we need the distribution of the true mass $P(M_T)$. In this work we assumed that the distribution in the planetary regime is a power-law function with index -1 (Heinze et al. 2010) and for the brown dwarf regime a power law with index approximately 0 (Metchev & Hillenbrand 2009).

So, the function $P(M_T)$ is $\propto M^{-1}$ from M_0 to the upper limit of the brown dwarf desert and constant for greater values of the mass. The final function is the sum of the two different curves connected in the point where the mass is equal to the value of the upper limit of the brown dwarf desert. The boundaries of the brown dwarf desert are assumed to be $20\text{--}57 M_{\text{Jup}}$ (Grether & Lineweaver 2006).

If we want to know how many object we can detect with SPHERE in the substellar regime, the upper limit of the mass is $80 M_{\text{Jup}}$. If the physical maximum mass of the target is in the planetary regime, we changed the upper limit of the integral.

In general we have to solve the integral

$$y = \int_{M_0}^X \frac{M_0/M_T^2}{\sqrt{1 - \left(\frac{M_0}{M_T}\right)^2}} A M_T^\alpha dM_T \quad (\text{A7})$$

for $\alpha = -1$ and $\alpha = 0$:

$$y(\alpha = -1) = A \int_{M_0}^X \frac{M_0/M_T^3}{\sqrt{1 - \left(\frac{M_0}{M_T}\right)^2}} dM_T = A \frac{\sqrt{X^2 - M_0^2}}{M_0 X}, \quad (\text{A8})$$

$$y(\alpha = 0) = A' \int_{M_0}^X \frac{M_0/M_T^2}{\sqrt{1 - \left(\frac{M_0}{M_T}\right)^2}} dM_T = A' \arccos\left(\frac{M_0}{X}\right). \quad (\text{A9})$$

We can split the two zones below and above the upper limit of the brown dwarf desert, $M_{\text{des}} = 57 M_{\text{Jup}}$, and the total area for the normalization is

$$\Phi = A \int_{M_0}^{M_{\text{des}}} \frac{M_0/M_T^3}{\sqrt{1 - \left(\frac{M_0}{M_T}\right)^2}} dM_T + A' \int_{M_{\text{des}}}^{80 M_{\text{J}}} \frac{M_0/M_T^2}{\sqrt{1 - \left(\frac{M_0}{M_T}\right)^2}} dM_T, \quad (\text{A10})$$

where the constant $A' = A/M_{\text{des}}$ because we want the two curves to be connected in the value of M_{des} .

If the mass of detection is $M_0 \leq X \leq M_{\text{des}}$, then the probability is calculated through

$$P(M_T > X | M_0) = 1 - \frac{A \int_{M_0}^X \frac{M_0/M_T^3}{\sqrt{1 - \left(\frac{M_0}{M_T}\right)^2}} dM_T}{A \int_{M_0}^{M_{\text{des}}} \frac{M_0/M_T^3}{\sqrt{1 - \left(\frac{M_0}{M_T}\right)^2}} dM_T + \frac{A}{M_{\text{des}}} \int_{M_{\text{des}}}^{80 M_{\text{J}}} \frac{M_0/M_T^2}{\sqrt{1 - \left(\frac{M_0}{M_T}\right)^2}} dM_T} \quad (\text{A11})$$

that gives

$$P(M_T > X | M_0) = 1 - \frac{\frac{\sqrt{X^2 - M_0^2}}{M_0 X}}{\frac{\sqrt{M_{\text{des}}^2 - M_0^2}}{M_0 M_{\text{des}}} + \frac{1}{M_{\text{des}}} \arccos\left(\frac{M_0}{80 M_{\text{J}}}\right)}. \quad (\text{A12})$$

For $M_{\text{des}} < X \leq 80 M_{\text{J}}$ in the same way we obtain

$$P(M_T > X | M_0) = 1 - \frac{\frac{\sqrt{M_{\text{des}}^2 - M_0^2}}{M_0 M_{\text{des}}} + \frac{1}{M_{\text{des}}} \arccos\left(\frac{M_0}{X}\right)}{\frac{\sqrt{M_{\text{des}}^2 - M_0^2}}{M_0 M_{\text{des}}} + \frac{1}{M_{\text{des}}} \arccos\left(\frac{M_0}{80 M_{\text{J}}}\right)}. \quad (\text{A13})$$

In this way we obtained the probability that an object is more massive than the minimum mass value for the detection and that it is not a stellar-type companion.

APPENDIX B: GJ 676 CANDIDATE MONITORING

During the first epoch of observation of the star GJ 676, 14 companions candidates were identified with an S/N above 5. Two other candidates have been identified with lower S/N and all 16 candidates are shown in Fig. B1. The second epoch, taken after 1 yr, confirmed that all of them are background sources. Table B1 presents the list of the astrometric relative positions of each star with respect to GJ 676.




Figure B1. Background stars found in the IRDIS FOV of GJ 676. The candidates are in the red circles.

Table B1. Astrometric positions of the background stars around GJ 676. The sky coordinates of the star for the epoch of 2016 May are RA = 17.50298568 and Dec. = -51.63782484 .

Star ID	ΔRA (arcsec)	$\Delta\text{Dec.}$ (arcsec)
1	3.95	-2.28
2	3.90	-3.84
3	3.71	-1.18
4	1.98	0.12
5	1.79	0.48
6	1.61	0.70
7	-1.18	-4.53
8	-1.66	-3.22
9	-1.81	-3.84
10	-3.86	-1.91
11	-4.82	0.04
12	-5.07	-1.22
13	-5.42	0.80
14	-5.85	0.69

¹Núcleo de Astronomía, Facultad de Ingeniería y Ciencias, Universidad Diego Portales, Av. Ejercito 441, Santiago, Chile

²Escuela de Ingeniería Industrial, Facultad de Ingeniería y Ciencias, Universidad Diego Portales, Av. Ejercito 441, Santiago, Chile

³Aix Marseille Université, CNRS, LAM – Laboratoire d’Astrophysique de Marseille, UMR 7326, F-13388 Marseille, France

⁴INCT, Universidad De Atacama, calle Copayapu 485, Copiapó, Atacama, Chile

⁵INAF – Osservatorio Astronomico di Padova, Vicolo dell’Osservatorio 5, I-35122 Padova, Italy

⁶INAF – Osservatorio Astrofisico di Catania, Via S. Sofia 78, I-95123 Catania, Italy

⁷Université Grenoble Alpes, IPAG, F-38000 Grenoble, France

⁸Institute for Astronomy, The University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, UK

⁹LESIA, Observatoire de Paris, Université PSL, CNRS, Sorbonne Université, Univ. Paris Diderot, Sorbonne Paris Cité, 5 place Jules Janssen, F-92195 Meudon, France

¹⁰*Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany*

¹¹*ETH Zurich, Institute for Particle Physics and Astrophysics, Wolfgang-Pauli-Strasse 27, CH-8093 Zurich, Switzerland*

¹²*Observatoire de Genève, University of Geneva, 51 Chemin des Maillettes, CH-1290 Versoix, Switzerland*

¹³*Department of Astronomy, Stockholm University, AlbaNova University Center, SE-106 91 Stockholm, Sweden*

¹⁴*CRAL, UMR 5574, CNRS, Université Lyon 1, 9 avenue Charles André, F-69561 Saint Genis Laval Cedex, France*

¹⁵*Department of Astronomy, University of Michigan, 1085 S. University Ave, Ann Arbor, MI 48109-1107, USA*

¹⁶*Université Nice-Sophia Antipolis, CNRS, Observatoire de la Côte d'Azur, Laboratoire J.-L. Lagrange, CS 34229, F-06304 Nice cedex 4, France*

¹⁷*DOTA, ONERA, Université Paris Saclay, F-91123 Palaiseau, France*

¹⁸*Anton Pannekoek Institute for Astronomy, Science Park 904, NL-1098 XH Amsterdam, the Netherlands*

This paper has been typeset from a \TeX/L\TeX file prepared by the author.