

1 Introduction

2 **The ecology and evolution of microbial CRISPR-Cas adaptive**
3 **immune systems**

4

5 Edze R. Westra¹, Stineke van Houte¹, Sylvain Gandon², Rachel Whitaker³

6

7 ¹ESI and CEC, Biosciences, University of Exeter, Cornwall Campus, Penryn TR10 9EZ, UK

8 ²CEFE UMR 5175, CNRS Université de Montpellier Université Paul-Valéry Montpellier EPHE,
9 34293 Montpellier Cedex 5, France

10 ³ Department of Microbiology, University of Illinois, Urbana-Champaign, 601 S. Goodwin Ave,
11 Urbana, IL 61801

12

13 **Authors for correspondence:**

14 Edze R. Westra - E.R.Westra@exeter.ac.uk,

15 Stineke van Houte – C.van-Houte@exeter.ac.uk,

16 Sylvain Gandon – Sylvain.GANDON@cefe.cnrs.fr,

17 Rachel Whitaker - rwhitaker@life.illinois.edu

18

19

20 **Keywords:** CRISPR, ecology, evolution, adaptive immunity, phage, plasmids, bacteria, archaea

21

22

23

24

1 **Introduction**

2 Over the past decade, the field of CRISPR-Cas research has received a lot of attention from the
3 scientific community. While initially this mostly concerned microbiologists who were fascinated
4 by the discovery that some bacteria encode RNA-guided adaptive immune systems, this rapidly
5 spread to other scientific disciplines following the development of groundbreaking molecular
6 biology tools [1], and more recently to the public domain where the societal and ethical
7 implications and legislation surrounding CRISPR applications are being heavily debated. Some
8 of the potential CRISPR applications that are currently being explored in the lab would involve
9 the release of CRISPR genes into confined or open environments – for example, when CRISPR
10 would be used to protect focal bacterial species against phage infections, when it is applied to
11 suppress the spread of antimicrobial resistance or to control vectors of disease [2-4]. One
12 component of the societal impacts of these applications entails an assessment of the potential
13 risks associated with these strategies (e.g. [5-7]), which requires an understanding how these
14 CRISPR-Cas behaves in an ecological context. In this special issue we explore this question, by
15 examining the evolutionary history of CRISPR-Cas immune systems, where they occur
16 naturally, when they evolve and how this impacts the spread and evolution of other DNA
17 elements. Finally we return to the question how CRISPR-Cas may be exploited in an ecological
18 context for the benefit of human health, and the ethical challenges that are associated with this.

19
20 ***CRISPR-Cas adaptive immune systems – a brief overview***

21 CRISPR-Cas adaptive immune systems were discovered around 15 years ago [8-11], and are
22 estimated to exist in approximately 50% of all bacterial genomes and roughly 90% of all
23 archaeal genomes [12]. A CRISPR immunity phenotype is genetically encoded by a so-called
24 CRISPR locus (Clustered Regularly Interspaced Short Palindromic Repeats) – an array of
25 repetitive and unique sequences (repeats and spacers, respectively), both of which are typically
26 around 30 nt in length. Spacers are derived from (foreign) genetic elements, such as plasmids
27 and viruses, and provide immunity to re-infection based on recognition of the cognate sequence
28 (known as “protospacer”) [13-15]. Bacteria or archaea may carry a single linked array of spacers
29 interspersed with repeats (one CRISPR locus) or multiple loci. CRISPR loci can evolve very
30 rapidly due to insertion of new spacers and the occasional loss of spacers or deletion of the
31 CRISPR locus itself, which can cause very closely related strains to carry unique combinations

1 of spacer sequences, known as a CRISPR allele. The overall length of CRISPR loci will increase
2 and decrease with the acquisition and loss of spacers, and can vary from as little as a single
3 spacer flanked by two repeats to hundreds of spacers and repeats [16]. Since new spacers are
4 added at the so-called leader-end of the locus, which is the sequence that contains the CRISPR
5 promoter, CRISPR loci form an inverse chronological record of previous infections from the
6 leader to the trailer end of the locus [17]. The extent to which different strains share the same
7 spacer sequences in the same order (usually at the trailer end of the CRISPR locus), is commonly
8 used to define related allele groups (RAGs) as a measure of their evolutionary relatedness.

9 Apart from the genetic CRISPR memory, a functional CRISPR-Cas immune system also
10 requires a set of CRISPR-associated genes (*cas* genes), which encode the protein machinery
11 required for carrying out the immune response [18]. Cas operons vary in their *cas* gene
12 composition and gene synteny, resulting in a classification of CRISPR-Cas systems into 2
13 classes, 6 Types and 33 subtypes [19-22]. These diverse CRISPR-Cas variants differ in many of
14 their mechanistic details, which have been discussed elsewhere [23, 24], yet also have
15 commonalities in the basic steps of the immune pathway. For example, two Cas proteins – Cas1
16 and Cas2 – are almost invariably part of CRISPR-Cas immune systems and are responsible for
17 inserting new spacer sequences into CRISPR arrays, sometimes assisted by other Cas proteins
18 (reviewed in [15, 25]). CRISPR transcripts are processed by either Cas proteins or housekeeping
19 RNases [26], and the resulting processed CRISPR RNAs (crRNAs) are bound by Cas proteins to
20 form a ribonucleoprotein complex that serves to detect and cleave complementary nucleic acid
21 sequences [23].

22 23 ***Ecology and diversity of CRISPR-Cas immunity***

24
25 CRISPR-Cas immune systems are unevenly distributed across taxa and environments. For
26 example, only less than half of mesophilic bacteria encode CRISPR-Cas immune systems,
27 compared to over 90% of bacterial thermophiles and archaea (both mesophilic and thermophilic)
28 [11, 16, 27-29]. Moreover, some uncultured bacterial lineages are virtually devoid of CRISPR-
29 Cas immune systems [30]. A recent computer learning approach suggested that abiotic factors
30 such as oxygen levels and temperature are important predictors of whether microorganisms
31 encode CRISPR-Cas immune systems [31]. However, the ecological drivers of CRISPR

1 distribution remain unclear. In this context, the key question is when CRISPR-Cas is favoured
2 over alternative defense strategies [32]. Experimental, theoretical and correlational studies have
3 suggested a role for viral abundance and diversity [27, 28, 33, 34], direct and indirect fitness
4 costs of CRISPR-Cas immune systems (e.g. autoimmunity, reduced horizontal gene transfer,
5 induced fitness costs, see below) [35-38], and epistasis with other host genes [39, 40].

6 In natural environments, whether that be the human lung, a hot spring or a fermenter, the
7 ecological and evolutionary impact of CRISPR depends on the population level diversity of
8 immune alleles. The diversity of alleles in the real world may reflect the previous history of
9 interactions and can be used to predict whether infection epidemics will occur in the future. For
10 example, the spread of a virulent virus can eliminate immune diversity from a population by
11 eliminating all susceptible (non-immune) cells, resulting in a selective sweep of an individual
12 strain with a matching CRISPR allele. In the future, this immunodominance across the CRISPR
13 allele might make a population susceptible to the subsequent epidemic spread of a new virus or
14 other mobile element in the population. Starting with a single strain of host or virus, diversity has
15 been shown to evolve in experimental studies [33, 41] and to protect bacteria from virus invasion
16 [42]. Data suggest that immune diversity shapes the evolution of viral pathogens by selecting for
17 recombinant genotypes that are more likely to escape immunity [43] and, recombinant microbial
18 CRISPR alleles that are more likely to increase immune profiles of a single strain [44, 45]. This
19 inevitable interaction between viruses and mobile elements with immune diversity will thereby
20 broadly impact dynamics of multiple microbial pathogens and link their dynamics through
21 CRISPR.

22 Interestingly, studies of CRISPR diversity have observed differences in different
23 microbial species from different environments. *Leptospirillum* species from acid mine drainage
24 biofilms [43], and *Sulfolobus islandicus* strains from a single hot spring [46], Halophilic archaea
25 from a saline pond [47], *Helicobacter cinaedi* [48] and *Pseudomonas aeruginosa* from a single
26 hospital [49] show diversity of CRISPR spacers co-exist. In contrast, *Pseudomonas aeruginosa*
27 within a human lung, or *Prevotella* strains in a single gut sample are clonal with complete
28 immunodominance at one time and place [49, 50]. These differences in diversity among CRISPR
29 populations may result from difference in CRISPR dynamics or demographics (e.g. colonization
30 bottle-necks [51]) and biological constraints (e.g. an inactive spacer acquisition machinery [52]).
31 As CRISPR studies have mainly focused on mechanistic details, our understanding of microbial

1 population dynamics in real populations is surprisingly limited. Additional studies from a diverse
2 range of CRISPR-containing organisms and their viruses, across a range of environments are
3 needed to identify the basis for the differences in the genetic structure of CRISPR and predict
4 how they impact epidemics and evolutionary dynamics in local populations.

6 *Co-evolution between CRISPR hosts and viruses*

7 Given the detailed molecular understanding of how CRISPR-Cas systems work, one can often
8 predict resistance phenotypes of an individual on the basis of its genotype and that of its phage
9 (or other foreign DNA element). Specifically, while bacteria gain resistance through the
10 acquisition of spacers, phages and other mobile genetic elements can overcome immunity
11 through the acquisition of “escape mutations” in (or sometimes near) the protospacer (e.g. [53,
12 54]). The fascinating details of the biology of CRISPR immunity and phage evasion challenge
13 the classical theoretical framework used to understand host-parasite interactions. In particular,
14 the Lamarckian ability of CRISPR to acquire new spacers allows the bacteria to accumulate a
15 diverse range of resistance alleles against a focal phage [28, 55, 56]. This multiplicity of
16 resistance within individual bacteria and/or within bacterial populations can overwhelm the
17 evolutionary potential of the phage and drive it to extinction [42, 56-58]. Alternative
18 mechanisms of escape are also possible and, recently, a number of different anti-CRISPR (Acr)
19 systems have been described in different phages [59]. These Acr proteins have the ability to
20 down-regulate the immunity of CRISPR-Cas systems and allow the phage to exploit CRISPR
21 resistant bacteria. Interestingly, this immunosuppression has been shown to require the
22 cooperation between multiple phages, where the first phages reduce the efficacy of CRISPR-Cas
23 and allow subsequent infections to exploit the host [60, 61].

24 However, even if phage lack *acr* genes, CRISPR immunity is far from being the silver
25 bullet against phages because it carries several different fitness costs. Some fitness costs are
26 likely to result from the production of an effective interference against invading pathogens and
27 mobile genetic elements [33, 62]. Some other costs are associated with self-targeting and auto-
28 immunity [35, 36]. Also, when CRISPR targets mobile genetic elements or prophages that carry
29 some adaptive mutation for the bacteria, CRISPR immunity may also be viewed as a form of
30 self-targeting, since immunity directly harms the bacterial host. For instance, when CRISPR
31 targets beneficial plasmids carrying antibiotic resistance, this immunity can be counter selected

1 and lead to the loss of CRISPR loci [37, 63-65], potentially explaining why many bacteria lack
2 CRISPR-Cas immune systems. Note, however, that apart from a few studies [66], the
3 experimental study of the coevolution between CRISPR immunity and phage evasion is often
4 limited to a few generations. The monitoring of longer coevolution experiments in the laboratory
5 or in natural environments are very welcome to better understand the coevolutionary dynamics
6 driven by CRISPR immunity (see several papers in this special issue).

7

8 *Application of CRISPR in an ecological context*

9 While CRISPR-mediated targeting of antibiotics resistance plasmids may be maladaptive for
10 individual bacteria (see above), it was soon realized that this feature of CRISPR systems may be
11 exploited to limit the spread of antibiotics resistance [67-69], and more generally to engineer
12 microbial communities. As the majority of bacteria do not encode end-joining mechanisms to
13 repair double-strand DNA breaks, CRISPR-based technologies are particularly useful for either
14 sequence-specific killing of pathogenic bacteria or removal of accessory genes (e.g.
15 antimicrobial resistance, virulence, etc.). Synthetic CRISPR systems can be delivered to a target
16 bacterium using so-called phagemids, which are replication-deficient bacteriophage particles,
17 [67, 69], or through conjugative delivery of CRISPR systems [67, 70], but research is still in its
18 infancy and considerable challenges are associated with these approaches [3].

19 CRISPR-mediated ecological engineering has also been explored in higher organisms,
20 where it has raised major interest given its potential use in pest and invasive species control and
21 reducing vector-borne diseases. Ecological engineering can be achieved by CRISPR-based
22 genome engineering of wild animals followed by introducing these engineered animals in a local
23 wild population. However, this method is likely to be effective only if the target population is
24 relatively small (see Buchthal *et al* in this issue). A more powerful, but also riskier approach is to
25 apply synthetic CRISPR-based gene drives [4, 71, 72]. These elements spread with super-
26 Mendelian inheritance through a population and can disrupt genetic loci or facilitate the spread
27 of genetically linked genes. CRISPR-based gene drives have been successfully used in confined
28 laboratory settings in a range of organisms, including yeast, fruit flies, mosquitoes and, very
29 recently, mice [73-76], and some first insights into the evolutionary dynamics of these drives,
30 including evolution of resistance, are emerging [77-80].

1 For all these ecological engineering approaches, there is a clear need to better understand
2 the potential risks of these gene drives and whether there is any appetite in society for these types
3 of interventions. For example, what is the durability of these interventions, what are their long-
4 term ecological consequences for (microbial) communities, and what countermeasures can be
5 taken to limit or reverse gene drive spread? Being able to predict risks and benefits will be one
6 important factor to get support from local communities, and for ethical and legislative approval.
7 Recognizing the urgent need for new guidelines, scientists from a range of disciplines, including
8 ethicists, social scientists and biologists, are working together to develop improved safety
9 recommendations for CRISPR-based ecological engineering technologies [6, 81, 82], and ethical
10 guidelines and the involvement of the public at an early stage in the development of these
11 applications [81, 83, 84].

12

13 **Aims of the theme issue**

14 This theme issue aims to examine when and where CRISPR-Cas immune systems are important,
15 how this impacts the coexistence and coevolution between bacteria and phages, and how this ties
16 in with more practical and ethical considerations concerning the application of CRISPR-based
17 technologies in an ecological context. The studies presented in this issue use a wide range of
18 different approaches, from bioinformatics and metagenomics correlational studies to
19 experimental and theoretical analyses, to answer an equally wide range of questions all somehow
20 linked to CRISPR ecology and evolution: from understanding the basic principles of CRISPR
21 evolution in bacteria to community engagement studies to examine if CRISPR-based
22 applications can be applied in the real world to limit the spread of Lyme disease. The theme issue
23 highlights the need to not only understand how CRISPR works in an ecological context, but also
24 to engage with important practical issues when translating this knowledge to real-world
25 applications to improve human health.

26

27 **Overview of the papers**

28 This theme issue is broadly divided into four areas that deal respectively with (i) the evolutionary
29 history and relative importance of CRISPR-Cas immune systems, (ii) the role of CRISPR-Cas
30 immune systems during bacteria-phage or Archaea-virus coevolution in real environments, (iii)
31 understanding the drivers and consequences of coevolution in controlled lab environments, and

1 (iv) the way CRISPR may be harnessed to remove infectious diseases in real environments, and
2 the associated considerations that need to be taken into account.

3 4 **A) Evolutionary history and relative importance of CRISPR-Cas immune systems**

5 One of the key questions in the field of CRISPR-Cas is how these sophisticated immune systems
6 evolved in the first place, and how the enormous diversity of CRISPR Classes, Types and
7 subtypes emerged in this process. Koonin & Makarova (ref) use comparative genomics to bring
8 the evolutionary history of this system in focus. They suggest a key role for a Cas1-related
9 transposase in the early evolution of the “Adaptation” module of the system, followed by
10 evolution of increased complexity through a series of gene duplication and displacement events,
11 recruitment of genes with nuclease activities and in some cases signal transduction genes to form
12 the “Interference” module of the system. While a cloud of uncertainty inevitably surrounds such
13 analyses, their work provides a plausible explanation for the way in which a highly complex
14 adaptive immune system may have evolved. A next obvious question is then, why do not all
15 bacteria and Archaea carry this adaptive immune system. This is a recurrent question, and one
16 factor that clearly matters is how well CRISPR-Cas immune systems performs (in terms of the
17 benefits it provides) relative to alternative immune strategies. For example, many bacteria and
18 archaea encode restriction-modification systems or they can mutate receptors that phages use to
19 attach to the cell surface. So why put up with CRISPR immunity, is the question that is raised by
20 Levin and co-workers (ref). Using mathematical modeling they demonstrate that in theory the
21 conditions where CRISPR is favoured over these alternative defenses is restricted. In other
22 words, microbes have many options available to defend themselves against phages, and CRISPR
23 may simply not always be the best one. However, the benefits of CRISPR are not only
24 determined by the presence or absence of phage, but also by the genetic context of these immune
25 systems, as Aude Bernheim and her colleagues show using bioinformatics approaches (ref), and
26 as Anne Chevallereau and co-workers show using experimental manipulations (ref). Specifically,
27 Bernheim et al find evidence for both positive and negative epistatic interactions between
28 CRISPR-Cas subtypes and double stranded DNA repair pathways in bacteria. This therefore
29 suggests that the distribution of CRISPR-Cas subtypes across bacterial species is shaped by the
30 genetic context, but the underlying mechanism for this association is unclear. Earlier work from
31 the same team demonstrated that one *cas* gene can inhibit the NHEJ pathway, which conceivably

1 can be maladaptive, explaining why these systems hardly ever co-occur in the same genome
2 [39]. Chevallereau et al experimentally examined the impact of *mutS*, which is part of the DNA
3 mismatch repair system, on the evolution of CRISPR resistance by *Pseudomonas aeruginosa*
4 against its phage DMS3vir. They found that due to the resulting increase in mutation supply,
5 bacteria were much more likely to evolve resistance through mutation of the phage receptor,
6 suggesting no selective benefit of carrying CRISPR-Cas immune systems in this host genetic
7 background. Together, these studies show how CRISPR-Cas immune systems may have evolved
8 in the first place, and how natural selection for these systems not only depends on the presence or
9 absence of phage, but also on the presence of alternative defense mechanisms as well as the
10 genetic context of the CRISPR-Cas system.

11

12 **B) Patterns of spacer acquisition in an ecological context**

13 One of the key questions in the field of CRISPR ecology and evolution is what drives the
14 generation and maintenance of CRISPR diversity. Using mathematical modeling, Bradde et al
15 show that spacer acquisition rates can be constrained by a cost of autoimmunity (ref). Their
16 model assumes that higher spacer acquisition rates not only increase the benefits of phage
17 resistance, but also the costs of autoimmunity due to spacer sampling from the bacterial genome.
18 This results in selection for an optimum spacer acquisition rate that maximizes bacterial survival
19 (a balance of phage resistance and autoimmunity), which in turn depends on factors that
20 influence the infection risk, such as bacterial and phage population sizes. In addition to this
21 theoretical approach, several papers in this issue apply comparative population studies to
22 describe spacer diversity in natural environments, and to identify the factors that drive this
23 diversity. Lopatina et al. (ref) examine the CRISPR loci of populations of the bacterium *Thermus*
24 in different geographical locations in Chile, Italy and Russia. They demonstrate that within a
25 single population of the bacterium *Thermus* more spacers are shared among strains than among
26 distant populations. The authors infer from the pattern that local dynamics between hosts and
27 viruses define the local diversity although the shared spacers among populations indicate some
28 evidence for gene flow. Similarly, Pauly et al. (ref) show patterns of persistent local diversity
29 within a single hot spring of *Sulfolobus islandicus*. Pauly also shows that these patterns vary for
30 different viruses indicating that virus lifestyle impacts immune diversity and explore different
31 mechanisms of CRISPR escape from different viruses through mutation or virus genome

1 replacement. Finally, Hoikkala et al. review the importance of CRISPR-Cas immune systems in
2 aquaculture, and the observed dynamics of coevolution of a bacterial fish pathogen and its
3 phage. They discuss how CRISPR diversity of bacterial strains in these environments can be
4 exploited for tracking the epidemiology of bacterial pathogens, and for phage therapy
5 interventions (ref). Their review highlights how a better understanding of CRISPR-diversity may
6 be used to stabilize and shape healthy microbial populations and prevent invasions of bacterial
7 pathogens into aquaculture environments. Collectively, these studies provide novel insights into
8 the generation, maintenance and potential application of spacer diversity in natural
9 environments.

11 C) Coevolution between CRISPR-Cas immune systems and phages in the lab

12 Experiments carried out in controlled environments in the lab allow to explore the details of the
13 coevolutionary processes between CRISPR systems and phages. Common et al (ref) carried out a
14 coevolution experiment between *Streptococcus thermophilus* and a lytic phage over 30 days. The
15 monitoring of phenotypic and genotypic changes in both the bacteria and the phage across time
16 confirmed the emergence of an antagonist arms race that often results in the extinction of the
17 phage populations. These extinctions suggest that this coevolution is asymmetric because the
18 phage cannot cope with the accumulation of escape mutations. The existence of a fitness cost
19 associated with most escape mutations was confirmed by a separate study by Chabas et al (ref)
20 on the same biological model. In addition, Chabas et al. found a dramatic variation in the speed
21 at which the phage can escape different CRISPR mediated resistance alleles. The resistance
22 induced by some spacers can be easily bypassed by the rapid emergence of escape mutations,
23 while the resistance induced by other spacers is much more durable. The measurement of the
24 phage mutation rate using Luria-Delbruck fluctuation tests allowed Chabas et al to suggest that
25 this variation in resistance durability is likely to be driven by heterogeneity in the mutation rate
26 across different protospacers. The evolution of escape mutations was also studied by Watson et
27 al (ref) in two lytic phages infecting CRISPR resistant *Pectobacterium atrosepticum*. This study
28 contrasted the evolution of the phage against single or multiple spacers. When the resistance was
29 mediated by a single spacer, most escape mutations were due to a single point mutation in the
30 PAM or the seed sequence. When the resistance is due to multiple spacers, most phages escaped
31 with deletions in genes encoding structural proteins. These mutations were viable but affect the

1 morphology and the fitness of the virus, which further support the existence of a cost of escape
2 and an asymmetry in the coevolutionary arms race. The details of phage adaptation was also
3 studied by McKitterick et al (ref) in a fascinating system where the phage ICP1 uses a fully
4 functional CRISPR-Cas system to down-regulate bacterial immunity of *Vibrio cholerae*
5 mediated by a phage inducible chromosomal island-like element (PLE). High-throughput
6 sequencing allowed the authors to monitor the acquisition of new spacers against PLE in the
7 CRISPR of ICP1, which provide quantitative resistance against PLE that depends on the number
8 and sequence of spacers. Furthermore, even spacers not targeting the PLE but instead targeting
9 the small chromosome of the bacterium could still reduce bacterial immunity if the chromosomal
10 targeting occurred close to the PLE integration site. Collectively, these different experiments in
11 the lab shed a new light on the interference mediated by CRISPR and by the costs associated
12 with different mutations allowing phages to escape bacterial immunity.

13

14 **D) Implications of CRISPR-Cas for human health and wellbeing**

15 Natural CRISPR-Cas systems can have an important impact on human health by for example
16 altering the spread of antimicrobial resistance and virulence genes [85]. However, even though
17 experimental studies show that CRISPR-Cas forms an important barrier for horizontal gene
18 transfer, correlational studies have shown mixed results [63, 86]. Here, Shereen et al (ref)
19 examine if the presence of anti-CRISPR genes may impact the correlation between CRISPR-Cas
20 immune systems and antibiotic resistance genes (ARGs) genes. They find a very high degree of
21 variation between bacterial species in the abundance of CRISPR-Cas systems and no correlation
22 with ARG for most species. However, for *Pseudomonas aeruginosa* they found a positive
23 correlation between anti-CRISPR and ARG genes, indicating that anti-CRISPRs may facilitate
24 the spread of clinically relevant genes such as those encoding antibiotic resistance or virulence
25 factors in the face of CRISPR-Cas immune systems. These genes may be removed using
26 CRISPR-based ecological engineering technologies. Although this is still a very young field the
27 developments have moved fast and scientists now have a variety of CRISPR-based tools at hand
28 to alter microbiome composition and function, and use them as next-generation antimicrobials.
29 Ramachandran & Bikard (ref) provide a thorough overview of the these various CRISPR-based
30 applications to alter the microbiome, including its application in editing bacteria and phage
31 genomes, controlling their gene expression using CRISPRi, killing pathogenic bacteria using

1 CRISPR-based antimicrobials and removing antibiotic resistance genes or other virulence
2 determinants.

3 While these CRISPR-based ecological engineering approaches have great potential to
4 solve ecological problems, there are many technological, regulatory, societal and ethical
5 challenges. In this issue, De Graeff et al (ref) provide a thorough overview of the arguments
6 reported in the scientific literature for and against genome editing in animals, including gene
7 drives, which relate to human health, efficiency, risks and uncertainty, animal welfare, animal
8 dignity, environment and public acceptability. The authors also pointed out that the ethical
9 debate on genome editing in animals is predominantly shaped by biomedical and veterinary
10 scientists, and less so by ethicists and social scientists. They argue that involvement of ethicists
11 and social scientists from the very early research stages of technology development may help
12 facilitate responsible governance of animal genome editing applications [87]. Furthermore, there
13 is a need for engagement with the public to address amongst others concerns around equity of
14 access (who will benefit from the new technology?) and the commercialization of the technology
15 (will businesses prioritize profit-making over providing a safe public good?). Buchthal et al (ref)
16 provide an excellent example of involving the public in decision-making on CRISPR-based
17 ecological engineering. ‘Mice against Ticks’ is an exciting new community-guided ecological
18 engineering project aimed at reducing the incidence of tick-borne diseases vectored by mice on
19 the islands of Nantucket and Martha’s Vineyard off the US east coast. Their idea is to introduce
20 CRISPR-edited mice that are heritably resistant to ticks and/or tick-borne disease in the islands’
21 local mouse population, which is hypothesized to disrupt the disease transmission cycle. From
22 the stages of conception of the project the local communities of both islands have been actively
23 involved in all decision-making. Although at a relatively early stage still, successes and
24 challenges from this exciting pilot project will provide future ecological engineering projects
25 with highly valuable insights on how to set up community-driven projects.

26

27 **Conclusion and outlook**

28 While our mechanistic understanding of CRISPR-Cas immune systems has raced ahead, our
29 understanding of the evolutionary ecology of these fascinating immune systems is still relatively
30 limited. As outlined above, this issue fills some important gaps in our knowledge of the
31 distribution and impact of CRISPR immune systems, but there are also many outstanding

1 questions that need to be answered. For example, why are there so many different CRISPR-Cas
2 variants, what are their costs and benefits, and how does this depend on the environment? Why
3 do so many bacteria not encode CRISPR-Cas immune systems, and what drives their loss? Why
4 is it that in the lab bacteria that encode CRISPR-Cas systems often evolve receptor-based
5 resistance against their phages? Is this because of a lack of ecological realism, widespread anti-
6 CRISPR strategies of phages, or are as yet unknown phage life history traits important for the
7 evolution of CRISPR resistance? As argued in this introduction, a better understanding of the
8 principles that govern evolution of CRISPR-Cas immune systems and their coevolution with
9 mobile genetic elements can help to inform applications of CRISPR-based technologies in real
10 environments. Apart from this quest for a better understanding of CRISPR ecology and
11 evolution, contributions in this issue also highlight the need for community engagement by
12 CRISPR biologists and participation in debates surrounding the ethics and legislative aspects of
13 these technologies, and provide some excellent examples of how this can be done.

14

15 **Competing interests**

16 The authors declare no competing interests.

17

18 **Funding**

19 ERW is supported by a NERC Independent Research Fellowship (NE/M018350/1), and received
20 funding in the area of CRISPR ecology and evolution from the ERC (ERC-STG-2016-714478 –
21 EVOIMMECH, and EPICUT) BBSRC (BB/N017412/1), and NERC (NE/S001921/1). SVH is
22 supported by a BBSRC Future Leader Fellowship (BB/R010781/1). RJW is supported for this
23 work by the Cystic Fibrosis Foundation WHITAK16PO, NSF Dimensions of Biodiversity
24 #1342876 and an Allen Distinguished Investigator Award.

25

26 **Acknowledgements**

27 The authors thank the Royal Society and all participants of the Hooke meeting on CRISPR
28 ecology and evolution in London, February 2019, and all contributors of the special issue on
29 CRISPR ecology and evolution.

30

31 **References**

- 1 1. Adli M. 2018 The CRISPR tool kit for genome editing and beyond. *Nat Commun* **9**(1), 1911.
2 (doi:10.1038/s41467-018-04252-2).
- 3 2. Selle K., Barrangou R. 2015 CRISPR-Based Technologies and the Future of Food Science. *J*
4 *Food Sci* **80**(11), R2367-2372. (doi:10.1111/1750-3841.13094).
- 5 3. Pursey E., Sunderhauf D., Gaze W.H., Westra E.R., van Houte S. 2018 CRISPR-Cas
6 antimicrobials: Challenges and future prospects. *PLoS Pathog* **14**(6), e1006990.
7 (doi:10.1371/journal.ppat.1006990).
- 8 4. Noble C., Adlam B., Church G.M., Esvelt K.M., Nowak M.A. 2018 Current CRISPR gene drive
9 systems are likely to be highly invasive in wild populations. *Elife* **7**. (doi:10.7554/eLife.33423).
- 10 5. Webber B.L., Raghu S., Edwards O.R. 2015 Opinion: Is CRISPR-based gene drive a biocontrol
11 silver bullet or global conservation threat? *Proc Natl Acad Sci U S A* **112**(34), 10565-10567.
12 (doi:10.1073/pnas.1514258112).
- 13 6. Akbari O.S., Bellen H.J., Bier E., Bullock S.L., Burt A., Church G.M., Cook K.R., Duchek P.,
14 Edwards O.R., Esvelt K.M., et al. 2015 BIOSAFETY. Safeguarding gene drive experiments in
15 the laboratory. *Science* **349**(6251), 927-929. (doi:10.1126/science.aac7932).
- 16 7. Brokowski C., Adli M. 2019 CRISPR Ethics: Moral Considerations for Applications of a
17 Powerful Tool. *J Mol Biol* **431**(1), 88-101. (doi:10.1016/j.jmb.2018.05.044).
- 18 8. Mojica F.J., Diez-Villasenor C., Garcia-Martinez J., Soria E. 2005 Intervening sequences of
19 regularly spaced prokaryotic repeats derive from foreign genetic elements. *J Mol Evol* **60**(2), 174-
20 182. (doi:10.1007/s00239-004-0046-3).
- 21 9. Bolotin A., Quinquis B., Sorokin A., Ehrlich S.D. 2005 Clustered regularly interspaced short
22 palindrome repeats (CRISPRs) have spacers of extrachromosomal origin. *Microbiology* **151**(Pt
23 8), 2551-2561. (doi:10.1099/mic.0.28048-0).
- 24 10. Pourcel C., Salvignol G., Vergnaud G. 2005 CRISPR elements in *Yersinia pestis* acquire new
25 repeats by preferential uptake of bacteriophage DNA, and provide additional tools for
26 evolutionary studies. *Microbiology* **151**(Pt 3), 653-663. (doi:10.1099/mic.0.27437-0).
- 27 11. Makarova K.S., Grishin N.V., Shabalina S.A., Wolf Y.I., Koonin E.V. 2006 A putative RNA-
28 interference-based immune system in prokaryotes: computational analysis of the predicted
29 enzymatic machinery, functional analogies with eukaryotic RNAi, and hypothetical mechanisms
30 of action. *Biol Direct* **1**, 7. (doi:10.1186/1745-6150-1-7).
- 31 12. Grissa I., Vergnaud G., Pourcel C. 2007 The CRISPRdb database and tools to display CRISPRs
32 and to generate dictionaries of spacers and repeats. *BMC Bioinformatics* **8**, 172.
33 (doi:10.1186/1471-2105-8-172).
- 34 13. Barrangou R., Fremaux C., Deveau H., Richards M., Boyaval P., Moineau S., Romero D.A.,
35 Horvath P. 2007 CRISPR provides acquired resistance against viruses in prokaryotes. *Science*
36 **315**(5819), 1709-1712. (doi:10.1126/science.1138140).
- 37 14. Marraffini L.A., Sontheimer E.J. 2008 CRISPR interference limits horizontal gene transfer in
38 staphylococci by targeting DNA. *Science* **322**(5909), 1843-1845. (doi:10.1126/science.1165771).
- 39 15. Amitai G., Sorek R. 2016 CRISPR-Cas adaptation: insights into the mechanism of action. *Nat*
40 *Rev Microbiol* **14**(2), 67-76. (doi:10.1038/nrmicro.2015.14).
- 41 16. Westra E.R., Dowling A.J., Broniewski J.M., van Houte S. 2016 Evolution and Ecology of
42 CRISPR. *Annual Review of Ecology, Evolution, and Systematics, Vol 47* **47**, 307-331.
43 (doi:10.1146/annurev-ecolsys-121415-032428).
- 44 17. Vale P.F., Little T.J. 2010 CRISPR-mediated phage resistance and the ghost of coevolution past.
45 *Proc Biol Sci* **277**(1691), 2097-2103. (doi:10.1098/rspb.2010.0055).
- 46 18. Jansen R., Embden J.D., Gastra W., Schouls L.M. 2002 Identification of genes that are
47 associated with DNA repeats in prokaryotes. *Mol Microbiol* **43**(6), 1565-1575.
- 48 19. Shmakov S.A., Makarova K.S., Wolf Y.I., Severinov K.V., Koonin E.V. 2018 Systematic
49 prediction of genes functionally linked to CRISPR-Cas systems by gene neighborhood analysis.
50 *Proc Natl Acad Sci U S A* **115**(23), E5307-E5316. (doi:10.1073/pnas.1803440115).

- 1 20. Shmakov S., Smargon A., Scott D., Cox D., Pyzocha N., Yan W., Abudayyeh O.O., Gootenberg
2 J.S., Makarova K.S., Wolf Y.I., et al. 2017 Diversity and evolution of class 2 CRISPR-Cas
3 systems. *Nat Rev Microbiol* **15**(3), 169-182. (doi:10.1038/nrmicro.2016.184).
- 4 21. Koonin E.V., Makarova K.S., Zhang F. 2017 Diversity, classification and evolution of CRISPR-
5 Cas systems. *Curr Opin Microbiol* **37**, 67-78. (doi:10.1016/j.mib.2017.05.008).
- 6 22. Makarova K.S., Wolf Y.I., Alkhnbashi O.S., Costa F., Shah S.A., Saunders S.J., Barrangou R.,
7 Brouns S.J., Charpentier E., Haft D.H., et al. 2015 An updated evolutionary classification of
8 CRISPR-Cas systems. *Nat Rev Microbiol* **13**(11), 722-736. (doi:10.1038/nrmicro3569).
- 9 23. van der Oost J., Westra E.R., Jackson R.N., Wiedenheft B. 2014 Unravelling the structural and
10 mechanistic basis of CRISPR-Cas systems. *Nat Rev Microbiol* **12**(7), 479-492.
11 (doi:10.1038/nrmicro3279).
- 12 24. Marraffini L.A. 2015 CRISPR-Cas immunity in prokaryotes. *Nature* **526**(7571), 55-61.
13 (doi:10.1038/nature15386).
- 14 25. Jackson S.A., McKenzie R.E., Fagerlund R.D., Kieper S.N., Fineran P.C., Brouns S.J. 2017
15 CRISPR-Cas: Adapting to change. *Science* **356**(6333). (doi:10.1126/science.aal5056).
- 16 26. Charpentier E., Richter H., van der Oost J., White M.F. 2015 Biogenesis pathways of RNA
17 guides in archaeal and bacterial CRISPR-Cas adaptive immunity. *FEMS Microbiol Rev* **39**(3),
18 428-441. (doi:10.1093/femsre/fuv023).
- 19 27. Weinberger A.D., Wolf Y.I., Lobkovsky A.E., Gilmore M.S., Koonin E.V. 2012 Viral diversity
20 threshold for adaptive immunity in prokaryotes. *MBio* **3**(6), e00456-00412.
21 (doi:10.1128/mBio.00456-12).
- 22 28. Iranzo J., Lobkovsky A.E., Wolf Y.I., Koonin E.V. 2013 Evolutionary dynamics of the
23 prokaryotic adaptive immunity system CRISPR-Cas in an explicit ecological context. *J Bacteriol*
24 **195**(17), 3834-3844. (doi:10.1128/JB.00412-13).
- 25 29. Anderson R.E., Brazelton W.J., Baross J.A. 2011 Using CRISPRs as a metagenomic tool to
26 identify microbial hosts of a diffuse flow hydrothermal vent viral assemblage. *FEMS Microbiol*
27 *Ecol* **77**(1), 120-133. (doi:10.1111/j.1574-6941.2011.01090.x).
- 28 30. Burstein D., Sun C.L., Brown C.T., Sharon I., Anantharaman K., Probst A.J., Thomas B.C.,
29 Banfield J.F. 2016 Major bacterial lineages are essentially devoid of CRISPR-Cas viral defence
30 systems. *Nat Commun* **7**, 10613. (doi:10.1038/ncomms10613).
- 31 31. Weissman J., Laljani R., Fagan W., Johnson P. 2018 Ecology shapes microbial immune strategy:
32 Temperature and oxygen as determinants of the incidence of CRISPR adaptive immunity. *bioRxiv*.
- 33 32. van Houte S., Buckling A., Westra E.R. 2016 Evolutionary Ecology of Prokaryotic Immune
34 Mechanisms. *Microbiol Mol Biol Rev* **80**(3), 745-763. (doi:10.1128/MMBR.00011-16).
- 35 33. Westra E.R., van Houte S., Oyesiku-Blakemore S., Makin B., Broniewski J.M., Best A., Bondy-
36 Denomy J., Davidson A., Boots M., Buckling A. 2015 Parasite exposure drives selective
37 evolution of constitutive versus inducible defense. *Curr Biol* **25**(8), 1043-1049.
38 (doi:10.1016/j.cub.2015.01.065).
- 39 34. Chabas H., van Houte S., Hoyland-Kroghsbo N.M., Buckling A., Westra E.R. 2016 Immigration
40 of susceptible hosts triggers the evolution of alternative parasite defence strategies. *Proc Biol Sci*
41 **283**(1837). (doi:10.1098/rspb.2016.0721).
- 42 35. Vercoe R.B., Chang J.T., Dy R.L., Taylor C., Gristwood T., Clulow J.S., Richter C., Przybilski
43 R., Pitman A.R., Fineran P.C. 2013 Cytotoxic chromosomal targeting by CRISPR/Cas systems
44 can reshape bacterial genomes and expel or remodel pathogenicity islands. *PLoS Genet* **9**(4),
45 e1003454. (doi:10.1371/journal.pgen.1003454).
- 46 36. Stern A., Keren L., Wurtzel O., Amitai G., Sorek R. 2010 Self-targeting by CRISPR: gene
47 regulation or autoimmunity? *Trends Genet* **26**(8), 335-340. (doi:10.1016/j.tig.2010.05.008).
- 48 37. Bikard D., Hatoum-Aslan A., Mucida D., Marraffini L.A. 2012 CRISPR interference can prevent
49 natural transformation and virulence acquisition during in vivo bacterial infection. *Cell Host*
50 *Microbe* **12**(2), 177-186. (doi:10.1016/j.chom.2012.06.003).

- 1 38. van Sluijs L., van Houte S., van der Oost J., Brouns S.J., Buckling A., Westra E.R. 2019
2 Addiction systems antagonize bacterial adaptive immunity. *FEMS Microbiology Letters* **in press**.
- 3 39. Bernheim A., Calvo-Villamanan A., Basier C., Cui L., Rocha E.P.C., Touchon M., Bikard D.
4 2017 Inhibition of NHEJ repair by type II-A CRISPR-Cas systems in bacteria. *Nat Commun* **8**(1),
5 2094. (doi:10.1038/s41467-017-02350-1).
- 6 40. Levy A., Goren M.G., Yosef I., Auster O., Manor M., Amitai G., Edgar R., Qimron U., Sorek R.
7 2015 CRISPR adaptation biases explain preference for acquisition of foreign DNA. *Nature*
8 **520**(7548), 505-510. (doi:10.1038/nature14302).
- 9 41. Paez-Espino D., Sharon I., Morovic W., Stahl B., Thomas B.C., Barrangou R., Banfield J.F. 2015
10 CRISPR immunity drives rapid phage genome evolution in *Streptococcus thermophilus*. *MBio*
11 **6**(2). (doi:10.1128/mBio.00262-15).
- 12 42. van Houte S., Ekroth A.K., Broniewski J.M., Chabas H., Ashby B., Bondy-Denomy J., Gandon
13 S., Boots M., Paterson S., Buckling A., et al. 2016 The diversity-generating benefits of a
14 prokaryotic adaptive immune system. *Nature* **532**(7599), 385-388. (doi:10.1038/nature17436).
- 15 43. Andersson A.F., Banfield J.F. 2008 Virus population dynamics and acquired virus resistance in
16 natural microbial communities. *Science* **320**(5879), 1047-1050. (doi:10.1126/science.1157358).
- 17 44. Held N.L., Herrera A., Whitaker R.J. 2013 Reassortment of CRISPR repeat-spacer loci in
18 *Sulfolobus islandicus*. *Environ Microbiol.* (doi:10.1111/1462-2920.12146).
- 19 45. Tyson G.W., Banfield J.F. 2008 Rapidly evolving CRISPRs implicated in acquired resistance of
20 microorganisms to viruses. *Environ Microbiol* **10**(1), 200-207. (doi:10.1111/j.1462-
21 2920.2007.01444.x).
- 22 46. Held N.L., Herrera A., Cadillo-Quiroz H., Whitaker R.J. 2010 CRISPR associated diversity
23 within a population of *Sulfolobus islandicus*. *PLoS One* **5**(9).
24 (doi:10.1371/journal.pone.0012988).
- 25 47. Emerson J.B., Andrade K., Thomas B.C., Norman A., Allen E.E., Heidelberg K.B., Banfield J.F.
26 2013 Virus-host and CRISPR dynamics in Archaea-dominated hypersaline Lake Tyrrell, Victoria,
27 Australia. *Archaea* **2013**, 370871. (doi:10.1155/2013/370871).
- 28 48. Tomida J., Morita Y., Shibayama K., Kikuchi K., Sawa T., Akaike T., Kawamura Y. 2017
29 Diversity and microevolution of CRISPR loci in *Helicobacter cinaedi*. *PLoS One* **12**(10),
30 e0186241. (doi:10.1371/journal.pone.0186241).
- 31 49. England W.E., Kim T., Whitaker R.J. 2018 Metapopulation Structure of CRISPR-Cas Immunity
32 in *Pseudomonas aeruginosa* and Its Viruses. *mSystems* **3**(5). (doi:10.1128/mSystems.00075-18).
- 33 50. Devoto A.E., Santini J.M., Olm M.R., Anantharaman K., Munk P., Tung J., Archie E.A.,
34 Turnbaugh P.J., Seed K.D., Blekhman R., et al. 2019 Megaphages infect *Prevotella* and variants
35 are widespread in gut microbiomes. *Nat Microbiol.* (doi:10.1038/s41564-018-0338-9).
- 36 51. Common J., Westra E.R. 2019 CRISPR evolution and bacteriophage persistence in the context of
37 population bottlenecks. *RNA Biol.* (doi:10.1080/15476286.2019.1578608).
- 38 52. Touchon M., Rocha E.P. 2010 The small, slow and specialized CRISPR and anti-CRISPR of
39 *Escherichia* and *Salmonella*. *PLoS One* **5**(6), e11126. (doi:10.1371/journal.pone.0011126).
- 40 53. Semenova E., Jore M.M., Datsenko K.A., Semenova A., Westra E.R., Wanner B., van der Oost J.,
41 Brouns S.J., Severinov K. 2011 Interference by clustered regularly interspaced short palindromic
42 repeat (CRISPR) RNA is governed by a seed sequence. *Proc Natl Acad Sci U S A* **108**(25),
43 10098-10103. (doi:10.1073/pnas.1104144108).
- 44 54. Deveau H., Barrangou R., Garneau J.E., Labonte J., Fremaux C., Boyaval P., Romero D.A.,
45 Horvath P., Moineau S. 2008 Phage response to CRISPR-encoded resistance in *Streptococcus*
46 *thermophilus*. *J Bacteriol* **190**(4), 1390-1400. (doi:10.1128/JB.01412-07).
- 47 55. Levin B.R., Moineau S., Bushman M., Barrangou R. 2013 The population and evolutionary
48 dynamics of phage and bacteria with CRISPR-mediated immunity. *PLoS Genet* **9**(3), e1003312.
49 (doi:10.1371/journal.pgen.1003312).

- 1 56. Childs L.M., England W.E., Young M.J., Weitz J.S., Whitaker R.J. 2014 CRISPR-induced
2 distributed immunity in microbial populations. *PLoS One* **9**(7), e101710.
3 (doi:10.1371/journal.pone.0101710).
- 4 57. Chabas H., Lion S., Nicot A., Meaden S., van Houte S., Moineau S., Wahl L.M., Westra E.R.,
5 Gandon S. 2018 Evolutionary emergence of infectious diseases in heterogeneous host
6 populations. *PLoS Biol* **16**(9), e2006738. (doi:10.1371/journal.pbio.2006738).
- 7 58. Westra E.R., Sunderhauf D., Landsberger M., Buckling A. 2017 Mechanisms and consequences
8 of diversity-generating immune strategies. *Nat Rev Immunol* **17**(11), 719-728.
9 (doi:10.1038/nri.2017.78).
- 10 59. Bondy-Denomy J., Pawluk A., Maxwell K.L., Davidson A.R. 2013 Bacteriophage genes that
11 inactivate the CRISPR/Cas bacterial immune system. *Nature* **493**(7432), 429-432.
12 (doi:10.1038/nature11723).
- 13 60. Borges A.L., Zhang J.Y., Rollins M.F., Osuna B.A., Wiedenheft B., Bondy-Denomy J. 2018
14 Bacteriophage Cooperation Suppresses CRISPR-Cas3 and Cas9 Immunity. *Cell* **174**(4), 917-925
15 e910. (doi:10.1016/j.cell.2018.06.013).
- 16 61. Landsberger M., Gandon S., Meaden S., Rollie C., Chevallereau A., Chabas H., Buckling A.,
17 Westra E.R., van Houte S. 2018 Anti-CRISPR Phages Cooperate to Overcome CRISPR-Cas
18 Immunity. *Cell* **174**(4), 908-916 e912. (doi:10.1016/j.cell.2018.05.058).
- 19 62. Vale P.F., Lafforgue G., Gatchitch F., Gardan R., Moineau S., Gandon S. 2015 Costs of CRISPR-
20 Cas-mediated resistance in *Streptococcus thermophilus*. *Proc Biol Sci* **282**(1812), 20151270.
21 (doi:10.1098/rspb.2015.1270).
- 22 63. Palmer K.L., Gilmore M.S. 2010 Multidrug-resistant enterococci lack CRISPR-cas. *MBio* **1**(4).
23 (doi:10.1128/mBio.00227-10).
- 24 64. Jiang W., Maniv I., Arain F., Wang Y., Levin B.R., Marraffini L.A. 2013 Dealing with the
25 evolutionary downside of CRISPR immunity: bacteria and beneficial plasmids. *PLoS Genet* **9**(9),
26 e1003844. (doi:10.1371/journal.pgen.1003844).
- 27 65. Gandon S., Vale P.F. 2014 The evolution of resistance against good and bad infections. *J Evol*
28 *Biol* **27**(2), 303-312. (doi:10.1111/jeb.12291).
- 29 66. Paez-Espino D., Morovic W., Sun C.L., Thomas B.C., Ueda K., Stahl B., Barrangou R., Banfield
30 J.F. 2013 Strong bias in the bacterial CRISPR elements that confer immunity to phage. *Nat*
31 *Commun* **4**, 1430. (doi:10.1038/ncomms2440).
- 32 67. Citorik R.J., Mimee M., Lu T.K. 2014 Sequence-specific antimicrobials using efficiently
33 delivered RNA-guided nucleases. *Nat Biotechnol* **32**(11), 1141-1145. (doi:10.1038/nbt.3011).
- 34 68. Gomaa A.A., Klumpe H.E., Luo M.L., Selle K., Barrangou R., Beisel C.L. 2013 Programmable
35 removal of bacterial strains by use of genome-targeting CRISPR-Cas systems. *MBio* **5**(1),
36 e00928-00913. (doi:10.1128/mBio.00928-13).
- 37 69. Bikard D., Euler C.W., Jiang W., Nussenzweig P.M., Goldberg G.W., Duportet X., Fischetti
38 V.A., Marraffini L.A. 2014 Exploiting CRISPR-Cas nucleases to produce sequence-specific
39 antimicrobials. *Nat Biotechnol* **32**(11), 1146-1150. (doi:10.1038/nbt.3043).
- 40 70. Dong H., Xiang H., Mu D., Wang D., Wang T. 2019 Exploiting a conjugative CRISPR/Cas9
41 system to eliminate plasmid harbouring the *mcr-1* gene from *Escherichia coli*. *Int J Antimicrob*
42 *Agents* **53**(1), 1-8. (doi:10.1016/j.ijantimicag.2018.09.017).
- 43 71. Esvelt K.M., Smidler A.L., Catteruccia F., Church G.M. 2014 Concerning RNA-guided gene
44 drives for the alteration of wild populations. *Elife*, e03401. (doi:10.7554/eLife.03401).
- 45 72. Godfray H.C.J., North A., Burt A. 2017 How driving endonuclease genes can be used to combat
46 pests and disease vectors. *BMC Biol* **15**(1), 81. (doi:10.1186/s12915-017-0420-4).
- 47 73. DiCarlo J.E., Chavez A., Dietz S.L., Esvelt K.M., Church G.M. 2015 Safeguarding CRISPR-Cas9
48 gene drives in yeast. *Nat Biotechnol* **33**(12), 1250-1255. (doi:10.1038/nbt.3412).
- 49 74. Gantz V.M., Jasinskiene N., Tatarenkova O., Fazekas A., Macias V.M., Bier E., James A.A. 2015
50 Highly efficient Cas9-mediated gene drive for population modification of the malaria vector

- 1 mosquito *Anopheles stephensi*. *Proc Natl Acad Sci U S A* **112**(49), E6736-6743.
2 (doi:10.1073/pnas.1521077112).
- 3 75. Gantz V.M., Bier E. 2015 Genome editing. The mutagenic chain reaction: a method for
4 converting heterozygous to homozygous mutations. *Science* **348**(6233), 442-444.
5 (doi:10.1126/science.aaa5945).
- 6 76. Grunwald H.A., Gantz V.M., Poplawski G., Xu X.S., Bier E., Cooper K.L. 2019 Super-
7 Mendelian inheritance mediated by CRISPR-Cas9 in the female mouse germline. *Nature*
8 **566**(7742), 105-109. (doi:10.1038/s41586-019-0875-2).
- 9 77. Champer J., Liu J., Oh S.Y., Reeves R., Luthra A., Oakes N., Clark A.G., Messer P.W. 2018
10 Reducing resistance allele formation in CRISPR gene drive. *Proc Natl Acad Sci U S A* **115**(21),
11 5522-5527. (doi:10.1073/pnas.1720354115).
- 12 78. Unckless R.L., Clark A.G., Messer P.W. 2017 Evolution of Resistance Against CRISPR/Cas9
13 Gene Drive. *Genetics* **205**(2), 827-841. (doi:10.1534/genetics.116.197285).
- 14 79. Noble C., Olejarz J., Esvelt K., Church G., Nowak M. 2017 Evolutionary dynamics of CRISPR
15 gene drives. *Bioarchiv*.
- 16 80. Kyrou K., Hammond A.M., Galizi R., Kranjc N., Burt A., Beaghton A.K., Nolan T., Crisanti A.
17 2018 A CRISPR-Cas9 gene drive targeting doublesex causes complete population suppression in
18 caged *Anopheles gambiae* mosquitoes. *Nat Biotechnol* **36**(11), 1062-1066.
19 (doi:10.1038/nbt.4245).
- 20 81. James S., Collins F.H., Welkhoff P.A., Emerson C., Godfray H.C.J., Gottlieb M., Greenwood B.,
21 Lindsay S.W., Mbogo C.M., Okumu F.O., et al. 2018 Pathway to Deployment of Gene Drive
22 Mosquitoes as a Potential Biocontrol Tool for Elimination of Malaria in Sub-Saharan Africa:
23 Recommendations of a Scientific Working Group(dagger). *Am J Trop Med Hyg* **98**(6_Suppl), 1-
24 49. (doi:10.4269/ajtmh.18-0083).
- 25 82. Min J., Smidler A.L., Najjar D., Esvelt K.M. 2018 Harnessing gene drive. *Journal of Responsible*
26 *Innovation* **5** (sup1), S40-S65.
- 27 83. National Academies of Sciences E., and Medicine. 2016 Gene Drives on the Horizon: Advancing
28 Science, Navigating Uncertainty, and Aligning Research with Public Values. (Washington, DC,
29 The National Academies Press.
- 30 84. Benedict M., Bonsall M., James A.A., James S., Lavery J., Mumford J., Quemada H., Rose R.,
31 Thompson P., Toure Y., et al. 2014 Guidance framework for testing of genetically modified
32 mosquitoes. (FNIH, WHO, TDR.
- 33 85. Hatoum-Aslan A., Marraffini L.A. 2014 Impact of CRISPR immunity on the emergence and
34 virulence of bacterial pathogens. *Curr Opin Microbiol* **17C**, 82-90.
35 (doi:10.1016/j.mib.2013.12.001).
- 36 86. Gophna U., Kristensen D.M., Wolf Y.I., Popa O., Drevet C., Koonin E.V. 2015 No evidence of
37 inhibition of horizontal gene transfer by CRISPR-Cas on evolutionary timescales. *ISME J* **9**(9),
38 2021-2027. (doi:10.1038/ismej.2015.20).
- 39 87. Hartley S., Gillund F., van Hove L., Wickson F. 2016 Essential Features of Responsible
40 Governance of Agricultural Biotechnology. *PLoS Biol* **14**(5), e1002453.
41 (doi:10.1371/journal.pbio.1002453).
- 42
43
44