

Research Article

Andrew J. Corbett*

A proof of the refined Gan–Gross–Prasad conjecture for non-endoscopic Yoshida lifts

DOI: 10.1515/forum-2015-0164

Received August 31, 2015; revised January 29, 2016

Abstract: We prove a precise formula relating the Bessel period of certain automorphic forms on $\mathrm{GSp}_4(\mathbb{A}_F)$ to a central L -value. This is a special case of the refined Gan–Gross–Prasad conjecture for the groups $(\mathrm{SO}_5, \mathrm{SO}_2)$ as set out by Ichino–Ikeda [12] and Liu [14]. This conjecture is deep and hard to prove in full generality; in this paper we succeed in proving the conjecture for forms lifted, via automorphic induction, from $\mathrm{GL}_2(\mathbb{A}_E)$ where E is a quadratic extension of F . The case where $E = F \times F$ has been previously dealt with by Liu [14].

Keywords: Automorphic period integrals, special values of L -functions, the refined Gan–Gross–Prasad conjecture

MSC 2010: 11F67, 11F70

Communicated by: Freydoon Shahidi

1 Introduction

The aim of this paper is to prove a special case of a deep conjectural relation between periods of automorphic forms and central values of L -functions. An early prototype of such a result is due to Waldspurger [21], who computed toric integrals of automorphic forms on GL_2 to be an ‘Euler-product’ of local integrals scaled by a global constant of certain L -values. Soon after, Gross–Prasad [8] made a series of fascinating conjectures relating periods of $\mathrm{SO}_{n+1} \times \mathrm{SO}_n$ -forms along SO_n (embedded diagonally) to central L -values – the case $n = 2$ is implied by the work of Waldspurger. These conjectures were extended to include all classical groups by Gan–Gross–Prasad [6].

In their original form, the Gross–Prasad conjectures omit a precise description of the factorisation of the global automorphic period. However, a recent work of Liu [14], extending that of Ichino–Ikeda [12], offers a refined conjecture by giving a precise conjectural formula for the Bessel period of a wide family of automorphic forms in terms of the central values of certain L -functions. In its full generality, Liu’s conjecture appears out of reach of our current methods, even for specific groups. Nevertheless, one can try to prove special cases of it; Liu himself proved his conjecture in the case of *endoscopic* automorphic forms on GSp_4 [14], motivated by Prasad–Takloo-Bighash [15]. These endoscopic forms are classically known as Yoshida lifts and essentially correspond to lifts from $\mathrm{GL}_2 \times \mathrm{GL}_2$.

In this paper we prove such a formula for the *non-endoscopic Yoshida lifts*: the automorphic forms on GSp_4 lifted from the *non-split* orthogonal group GO_4 (that is, the underlying quadratic space defining GO_4 has non-square discriminant). Making use of exceptional isomorphisms, we see that such forms are obtained by automorphic induction from $\mathrm{GL}_2(E)$ where E is a quadratic extension of the base field F . (Liu’s result covers the *split* case where $E = F \times F$.) For our proof we require both a much finer analysis of the four-dimensional quadratic spaces governing GO_4 (of non-square discriminant) and a more detailed construction of the

*Corresponding author: Andrew J. Corbett: School of Mathematics, University of Bristol, Bristol, BS8 1TW, United Kingdom, e-mail: andrew.corbett@bristol.ac.uk

automorphic representations of this group than that found in [14]. This analysis provides a notable diversion from Liu’s method, especially in the final deduction of our explicit formula Section 7.

Before describing our results in more detail we also remark on a conjecture of Böcherer [2] (see also [19]). In this work Böcherer formulates an equality between sums of Fourier coefficients (indexed by ideal classes of a fixed quadratic field K) of Siegel modular forms and certain L -values. The present paper considers the Bessel period of an automorphic form on $\mathrm{GSp}_4(\mathbb{A})$; if the form in question is the adèlisation of a Siegel modular form, then (by [4] for example) one computes the Bessel period to be precisely the Fourier coefficients that Böcherer considered. Thus our result provides a proof of (a refinement of) Böcherer’s conjecture for non-endoscopic Yoshida lifts.

1.1 The Bessel period

Let F be a (totally real) number field with adèle ring $\mathbb{A} = \mathbb{A}_F$. We consider the refined Gan–Gross–Prasad conjecture for the groups $(\mathrm{SO}_5, \mathrm{SO}_2)$. In this case we extend SO_2 to the *Bessel subgroup* $R = U \rtimes \mathrm{SO}_2$, with $R \hookrightarrow \mathrm{SO}_5$, where U is a certain unipotent subgroup of SO_5 . The conjecture describes the explicit form of a period integral of automorphic forms on $\mathrm{SO}_5 \times R$ along the (diagonally embedded) subgroup R . Our approach to the problem makes use of the exceptional isomorphisms

$$\mathrm{SO}_5 \cong \mathrm{PGSp}_4 \quad \text{and} \quad \mathrm{SO}_2 \cong \mathrm{Res}_{K/F} K^\times / F^\times$$

where K is a quadratic field extension of F .

More specifically, let χ be a unitary Hecke character of \mathbb{A}_K^\times , simultaneously thought of as a character of $\mathrm{SO}_2(F) \backslash \mathrm{SO}_2(\mathbb{A})$, and let π be an irreducible, cuspidal automorphic representation of $\mathrm{GSp}_4(\mathbb{A})$ in the space of cusp forms \mathcal{V}_π . Impose the central character condition that $\pi \otimes \chi|_{\mathbb{A}^\times} = 1$. Additionally, make a (standard and inconsequential) choice of automorphic character ψ of U so that $\psi \boxtimes \chi$ is an automorphic character of R . We then define the χ -Bessel period of $\varphi \in \mathcal{V}_\pi$ to be the absolutely convergent integral

$$\mathcal{P}(\varphi, \chi) = \int_{\mathbb{A}^\times R(F) \backslash R(\mathbb{A})} \varphi(g) (\psi \boxtimes \chi)(g) dg. \quad (1.1)$$

This integral defines an element of $\mathrm{Hom}_{R(\mathbb{A})}(\pi \otimes (\psi \boxtimes \chi), \mathbb{C})$. The unrefined conjecture claims that there exists some vector φ^* in (the Vogan L -packet of) π such that

$$\mathcal{P}(\varphi^*, \chi) \neq 0 \iff L(1/2, \pi \boxtimes \chi) \neq 0$$

where $\mathcal{P}(\varphi^*, \chi)$ may be defined for more general elements φ^* of the Vogan L -packet in a similar way to (1.1). It is this unrefined dependence which we make explicit.

To discuss the local side, assume the factorisations $\pi = \otimes_v \pi_v$; $\chi = \otimes_v \chi_v$; $\psi = \otimes_v \psi_v$ and suppose that $\varphi = \otimes_v \varphi_v$. Associated to this data, we follow Liu in defining

$$\alpha^\natural(\varphi_v, \chi_v) \in \mathrm{Hom}_{R(F_v)}(\pi_v \otimes (\psi_v \boxtimes \chi_v), \mathbb{C})$$

at each place v to be an integral over local matrix coefficients (see Section 6). Roughly speaking – up to a normalisation constant (see (6.1)) – the integral defining $\alpha^\natural(\varphi_v, \chi_v)$ is equal to

$$\int_{F_v^\times \backslash R(F_v)} \mathcal{B}_{\pi_v}(\pi_v(g_v) \varphi_v, \bar{\varphi}_v) (\chi_v \boxtimes \psi_v)(g_v) dg_v$$

where \mathcal{B}_{π_v} is a local unitary pairing for π_v . The foundation on which Liu is able to generalise the refined conjecture is the regularisation of these integrals. They are shown to converge absolutely and a natural normalisation is found such that $\alpha^\natural(\varphi_v, \chi_v) = 1$ for almost all places v (see [14, Theorem 2.1 and 2.2]). We may thus make sense of the infinite product $\prod_v \alpha^\natural(\varphi_v, \chi_v)$. The refined Gan–Gross–Prasad conjecture then asks for the constant of proportionality between this product of local factors and the square of the absolute value of the Bessel period.

1.2 Lifted representations

The representations of $\mathrm{SO}_5(\mathbb{A}) \cong \mathrm{PGSp}_4(\mathbb{A})$ are precisely those representations of $\mathrm{GSp}_4(\mathbb{A})$ with trivial central character. We consider a class of representations of $\mathrm{PGSp}_4(\mathbb{A})$ which are lifted from representations of the group $\mathrm{GO}_4(\mathbb{A})$, when GO_4 is non-split, via the theta correspondence for $(\mathrm{GO}_4, \mathrm{GSp}_4)$ – we call such lifted representations the *non-endoscopic Yoshida lifts*. The domain of this lift comprises of the representations of $\mathrm{GO}_4(\mathbb{A})$ (of trivial central character); these are uniquely determined by representations of $D^\times(\mathbb{A}_E)$ for a canonical choice of quadratic extension E/F and quaternion algebra D over F . Thus, via Jacquet–Langlands transfer, one may view a non-endoscopic Yoshida lift π as being of the form $\pi = \mathrm{AI}(\pi')$: the automorphic induction, to $\mathrm{GSp}_4(\mathbb{A})$, of a representation π' of $\mathrm{GL}_2(\mathbb{A}_E)$.

1.3 Main result

We refer the reader to Theorem 7.5 for a more precise statement of our result. To simplify notation here assume the following decompositions for both the Petersson inner product \mathcal{B}_π on π and the Tamagawa measure dg on $\mathbb{A}^\times \backslash R(\mathbb{A})$:

$$\mathcal{B}_\pi = \prod_v \mathcal{B}_{\pi_v}, \quad dg = \prod_v dg_v \quad (1.2)$$

where \mathcal{B}_{π_v} and dg_v are the local factors used to define $\alpha^{\natural}(\varphi_v, \chi_v)$.

Theorem. *Let $\pi = \mathrm{AI}(\pi')$ be a non-endoscopic Yoshida lift to $\mathrm{PGSp}_4(\mathbb{A})$, as per Section 1.2, where π' is an irreducible, cuspidal automorphic representation of $\mathrm{GL}_2(\mathbb{A}_E)$ with trivial central character. Let K be a quadratic field extension of F such that $\mathrm{SO}_2 \cong K^\times/F^\times$. Let χ be a unitary Hecke character of \mathbb{A}_K^\times with $\chi|_{\mathbb{A}^\times} = 1$. Then χ is simultaneously an automorphic character of $\mathrm{SO}_2(\mathbb{A})$. Denote by $\chi_{K/F}$ the quadratic character associated to K by class field theory. Assume the choices of (1.2) and that the local integrals $\alpha^{\natural}(\varphi_v, \chi_v)$ are properly normalised (as in Definition 7.1). Then for a cusp form $\varphi = \otimes_v \varphi_v$ in the space associated to π we have*

$$|\mathcal{P}(\varphi, \chi)|^2 = \frac{1}{4} \frac{\zeta_F(2) \zeta_F(4) L(1/2, \pi \boxtimes \chi)}{L(1, \pi, \mathrm{Ad}) L(1, \chi_{K/F})} \prod_v \alpha^{\natural}(\varphi_v, \chi_v).$$

1.4 Remarks

The case where $E = F \times F$ is dealt with by Liu [14, Section 4]. Liu’s theorem determines the Bessel period attached to an automorphic form on GSp_4 which is a lift from $\mathrm{GL}_2 \times \mathrm{GL}_2$. These lifts are precisely the *endoscopic* representations of GSp_4 . Moreover, Qui has proved a formula for $|\mathcal{P}(\varphi, \chi)|^2$ when π is in the *nontempered* cuspidal spectrum of SO_5 (see [16]). This is achieved by considering the so-called Saito–Kurukawa and Soudry lifts.

Following these two works, this paper uses the functorial lift from $\mathrm{GL}_2(E)$ to give a wide class of nonendoscopic, tempered, cuspidal automorphic representations of PGSp_4 that conform to the refined Gan–Gross–Prasad conjecture. Further works on attempting to prove such a formula in general have been approached by using tools such as relative trace formulae (see [5] for example).

The assumption that F is a totally real number field is needed only to permit the application of a result of [7] on the Petersson inner product of a theta lift; they, in turn, only require this assumption to use the Siegel–Weil formula in their calculation.

Finally, we would like to highlight the occurrence of the constant $1/4$ in our formula, to be compared with the constant $1/8$ appearing in [14]. This falls in line with the general conjecture of Liu [14] in that it relates precisely to the (conjectural) Arthur parameters of π and χ (as first pointed out by Ichino–Ikeda [12, Section 2] and then by Gan–Ichino [7, Remark 1.2]). Specifically, the constant should be $\frac{1}{|\mathcal{S}_\pi||\mathcal{S}_\chi|}$ where \mathcal{S}_π (resp. \mathcal{S}_χ) is the centraliser of the image of the Arthur parameter of π (resp. χ); note that in our case we trivially have $|\mathcal{S}_\chi| = 2$.

The discrepancy of $1/2$ between our result and that of [14] is supported by the observation that

$$|\mathcal{S}_\pi| = \begin{cases} 4 & \text{if } E = F \times F, \\ 2 & \text{if } E = F(\sqrt{e}) \text{ for some } e \notin (F^\times)^2. \end{cases}$$

It is interesting to see this factor arise naturally due to the structure of the representations of $\mathrm{GO}_4(\mathbb{A})$: in [14] the Bessel period boils down to twice the period considered by Waldspurger [21] in contrast to the single occurrence that we observe in our computation.

This paper is set out as follows: after some preliminary definitions regarding the Bessel period (Section 2) we review the theta correspondence for $(\mathrm{GO}_4, \mathrm{GSp}_4)$ (Section 3) and discuss the representation theory of GO_4 (Section 4), explaining the lift we use and its domain. We then analyse the global (Section 5) and local (Section 6) periods before uniting these quantities (Section 7) via a theorem of Waldspurger and proving the result at hand.

2 Preliminary discussion

2.1 Some conventions

We work over a fixed number field F which we assume to be totally real. Put \mathcal{O} for the ring of integers of F and \mathbb{A} for the ring of F -adèles. Given an extension $L \supset F$ let $\mathbb{A}_L = \mathbb{A} \otimes_F L$.

If G is a linear algebraic group defined over F and R is an F -algebra, write $G(R)$ for the R -points of G . At a place v of F simplify the notation $G(F_v)$ to G_v . Given a function f on G , denote left and right translation by elements $g \in G$ by

$$L(g)f(x) = f(g^{-1}x) \quad \text{and} \quad R(g)f(x) = f(xg).$$

If S is a finite set of places of F , then introduce the following notation: $F_S = \prod_{v \in S} F_v$ and $\mathbb{A}^S = \prod'_{v \notin S} F_v$. Note the compatibility of the products $G(F_S) = \prod_{v \in S} G(F_v)$ and $G(\mathbb{A}^S) = \prod'_{v \notin S} G(F_v)$ meaning that we can formally identify $G(\mathbb{A}) = G(F_S)G(\mathbb{A}^S)$.

2.1.1 Measures

For an algebraic group G we fix a Haar measure on $G(\mathbb{A})$ by taking the Tamagawa measure dg (as originally defined in [22]). Let dg_v be a specified choice of local Haar measures on G_v for each v such that $\prod_v dg_v$ is a well-defined measure on $G(\mathbb{A})$. By the uniqueness of Haar measures there exists a constant of proportionality $C \in \mathbb{C}$ such that $dg = C \prod_v dg_v$. We call such a C *Haar measure constant*, as in [12].

2.1.2 Automorphic representations and pairings

The space of automorphic (resp. cusp) forms on $G(\mathbb{A})$ shall be denoted $\mathcal{A}(G)$ (resp. $\mathcal{A}_0(G)$). For an irreducible, cuspidal automorphic representation π of $G(\mathbb{A})$ we denote by \mathcal{V}_π the realisation of π in $\mathcal{A}_0(G)$ and put ω_π for its central character. One has $\pi \cong \otimes_v \pi_v$ (and $\mathcal{V}_\pi \cong \otimes'_v \mathcal{V}_{\pi_v}$) where at each place v of F , π_v is an irreducible, admissible, unitary representation of G_v on \mathcal{V}_{π_v} . Let $\bar{\pi}$ denote the *conjugate* representation of π realised on the space

$$\mathcal{V}_{\bar{\pi}} = \{\bar{f} : f \in \mathcal{V}_\pi\}.$$

There is a canonical bilinear pairing $\mathcal{B}_\pi : \mathcal{V}_\pi \otimes \mathcal{V}_{\bar{\pi}} \rightarrow \mathbb{C}$ given by the Petersson inner product

$$\mathcal{B}_\pi(f, \bar{f}) = \int_{Z_G(\mathbb{A})G(F)\backslash G(\mathbb{A})} f(g)\bar{f}(g) dg$$

where Z_G is the maximal split torus in the centre of G and dg is the Tamagawa measure on $(Z_G \backslash G)(\mathbb{A})$ as always. In particular, since \mathcal{V}_π is a complex Hilbert space and π is unitary, one can show that $\bar{\pi}$ is isomorphic to π^\vee , the *contragredient representation* of π realised on the space of smooth vectors in the dual space \mathcal{V}_π^\vee of \mathcal{V}_π . Moreover, any pairing on a unitary Hilbert space representation is unique up to a scalar factor. Both of these facts are corollaries to the Riesz representation theorem. Throughout, any local, irreducible, admissible representation of G_v is always considered to be unitary.

2.1.3 L-functions

Given a representation r of the Langlands dual group and an automorphic representation π of G we have the Langlands L -function $L(s, \pi, r)$. When r is the standard representation of the dual group, which we assume is a subgroup of $\mathrm{GL}_n(\mathbb{C})$, we write $L(s, \pi)$ for $L(s, \pi, r)$. The notation $\pi_1 \boxtimes \pi_2$ denotes the (external tensor product) representation of the direct product $G_1 \times G_2$, where π_i are representations of the groups G_i for $i = 1, 2$, respectively.

The most interesting L -function for us is given as follows. Let π be an automorphic representation of $\mathrm{PGSp}_4(\mathbb{A}) \cong \mathrm{SO}_5$ and let χ be a character of $\mathrm{SO}_2(F) \backslash \mathrm{SO}_2(\mathbb{A})$ corresponding to a Hecke character of \mathbb{A}_K^\times as in the introduction. Then we consider the $\mathrm{SO}_5 \times \mathrm{SO}_2$ L -function $L(s, \pi \boxtimes \chi)$. However, other authors interpret this L -function as:

- the $\mathrm{GSp}_4 \times \mathrm{GL}_2$ L -function $L(s, \pi \boxtimes \mathrm{AI}(\chi))$, where $\mathrm{AI}(\chi)$ is the automorphic induction of χ from \mathbb{A}_K^\times to $\mathrm{GL}_2(\mathbb{A})$, or
 - the $\mathrm{GSp}_4(K)$ L -function $L(s, \mathrm{BC}(\pi) \otimes \chi)$, where $\mathrm{BC}(\pi)$ is the base change of π from $\mathrm{GSp}_4(\mathbb{A}_F)$ to $\mathrm{GSp}_4(\mathbb{A}_K)$.
- Each of these representations arises due to a functorial transfer from the original representation $\pi \boxtimes \chi$. The characteristic property of such a transfer implies that these L -functions are indeed all equal.

Other notation includes: ζ_F , the Dedekind zeta function for a number field F , and $\chi_{K/F}$ which always denotes the quadratic character of K^\times given by class field theory. Note that for any Hecke character χ of \mathbb{A}_K^\times , the adjoint L -function is trivially $L(s, \chi, \mathrm{Ad}) = L(s, \chi_{K/F})$.

2.1.4 Quadratic spaces

Let (V, q) be a quadratic space over F of even dimension $2m$ (we always assume such a V is non-degenerate). The quadratic form q corresponds to a symmetric matrix $S_q \in M_{2m}^{\mathrm{sym}}(F)$ such that $q(v) = {}^t v S_q v$ for $v \in V$. One defines the *discriminant* of V to be $\mathrm{disc} V = (-1)^m \det S_q$ and the associated *discriminant algebra* as

$$K_V = \begin{cases} F(\sqrt{\mathrm{disc} V}) & \text{if } \mathrm{disc} V \notin (F^\times)^2, \\ F \times F & \text{if } \mathrm{disc} V \in (F^\times)^2. \end{cases} \tag{2.1}$$

We intend to study the *orthogonal similitude group* of V :

$$\mathrm{GO}(V) = \{g \in \mathrm{GL}(V) : q(gv) = \lambda(g)q(v) \text{ for all } v \in V\} = \{g \in \mathrm{GL}_{2m}(F) : {}^t g S_q g = \lambda(g) S_q\}$$

where $\lambda : \mathrm{GO}(V) \rightarrow F^\times$ is the *similitude character*. One observes that $(\det g)^2 = \lambda(g)^{2m}$, so there is a natural *sign character* on $\mathrm{GO}(V)$:

$$\mathrm{sgn} : g \mapsto \det g / \lambda(g)^m \in \mu_2$$

where $\mu_2 = \mu_2(F)$. We define the connected component of $\mathrm{GO}(V)$ to be the normal subgroup $\mathrm{GSO}(V) = \ker(\mathrm{sgn})$ which sits in the exact sequence

$$1 \longrightarrow \mathrm{GSO}(V) \longrightarrow \mathrm{GO}(V) \xrightarrow{\mathrm{sgn}} \mu_2 \longrightarrow 1.$$

Similarly, if one defines the classical *orthogonal group* $\mathrm{O}(V) = \ker(\lambda)$, then the *special orthogonal group* $\mathrm{SO}(V)$ is found in the exact sequence

$$1 \longrightarrow \mathrm{SO}(V) \longrightarrow \mathrm{O}(V) \xrightarrow{\det} \mu_2 \longrightarrow 1$$

where $\det = \text{sgn}$ here. When $\dim V = 4$, we see later in Section 4.1 that the sign character is surjective and we exhibit a natural choice of representatives for $GO(V)/GSO(V)$. In essence, there is a unique element $\iota \in GO(V)$ with

$$\lambda(\iota) = 1, \quad \iota^2 = 1, \quad \det \iota = -1. \quad (2.2)$$

We are then able to fix a splitting such that μ_2 is identified with the subgroup of $GO(V)$ generated by ι . In particular, we arrive at the decomposition $GO(V) = GSO(V) \rtimes \mu_2$.

Remark 2.1. For an F -algebra A , the above comments apply more generally to the exact sequence

$$1 \longrightarrow GSO(V)(A) \longrightarrow GO(V)(A) \xrightarrow{\text{sgn}} \mu_2(A) \longrightarrow 1$$

where the A points of $GSO(V)$ coincide with the kernel of the sign function on $GO(V)(A)$. In particular, we have a well-defined notion of $\mu_2(\mathbb{A})$, $GSO(V)(\mathbb{A})$, $GSO(V)_V$ and so on.

2.2 The Bessel period and definitions

2.2.1 $GSp_4(F)$ in coordinates

Let $W = F^4$ and endow W with an antisymmetric bilinear form $(\cdot, \cdot)_W$ so that W becomes a four-dimensional symplectic vector space over F . In the coordinates of F^4 one may choose

$$(u, v)_W = {}^t u \begin{pmatrix} 0 & 1_2 \\ -1_2 & 0 \end{pmatrix} v$$

where 1_2 is the 2×2 identity matrix. Setting $W_1 = F^2$ then $W = W_1 \oplus W_1^\vee$ gives a complete polarisation of W such that W_1^\vee is identified with the dual space of W_1 under the form $(\cdot, \cdot)_W$. Recall the definition for the *symplectic similitude group*:

$$GSp_4(F) = GSp(W) = \{g \in GL(W) : (gu, gv)_W = \lambda(g)(u, v)_W \text{ for all } u, v \in W\}$$

where $\lambda(g) \in F^\times$. We use λ for the similitude character of any similitude group.

2.2.2 The torus

Fix a choice of anisotropic, symmetric matrix

$$S = \begin{pmatrix} a & b/2 \\ b/2 & c \end{pmatrix} \in M_2^{\text{sym}}(F)$$

to represent the quadratic form $q_S(v) = {}^t v S v$ for $v \in W_1$. Then (W_1, q_S) is a two-dimensional quadratic space over F of (scaled) discriminant

$$d = -4 \det S = b^2 - 4ac.$$

By the anisotropy of S (that $q_S(v) = 0 \Rightarrow v = 0$) it is clear that d is not a square in F . Hence the discriminant algebra $K_{W_1} = F(\sqrt{d})$ is a quadratic field extension of F . Fix the notation $K = K_{W_1}$. We consider a maximal, non-split torus in $GL_2(F)$ given by the orthogonal group

$$T = T_S = \{g \in GL_2(F) : {}^t g S g = (\det g) S\} = GSO(W_1).$$

One has the isomorphism $T \cong \text{Res}_{K/F} K^\times$ of algebraic groups over F . Specifically, one shows that

$$T(F) = \left\{ x + y \begin{pmatrix} b/2 & c \\ -a & -b/2 \end{pmatrix} : x, y \in F \right\}^\times$$

and defines an isomorphism $T(F) \rightarrow K^\times = F(\sqrt{d})^\times$ by

$$x + y \begin{pmatrix} b/2 & c \\ -a & -b/2 \end{pmatrix} \mapsto x + y \frac{\sqrt{d}}{2}.$$

2.2.3 The Bessel subgroup

Consider the following subgroups of $\mathrm{GSp}_4(F)$.

- Let U be the *unipotent radical* stabilising the flag $\{0\} \subset W_1 \subset W$; explicitly we have

$$U = \left\{ u(A) = \begin{pmatrix} 1_2 & A \\ 0 & 1_2 \end{pmatrix} : A \in M_2^{\mathrm{sym}}(F) \right\}.$$

All elements of U have similitude $\lambda(u(A)) = 1$. We also identify U with the space of symmetric F -linear maps $W_1^\vee \rightarrow W_1$. Taking the standard additive character

$$\psi : F \backslash \mathbb{A} \rightarrow \mathbb{C}^\times, \tag{2.3}$$

we define a character ψ_M of $U(F) \backslash U(\mathbb{A})$, for a matrix $M \in M_2^{\mathrm{sym}}(F)$, by

$$\psi_M(u(A)) = \psi(\mathrm{Tr}(MA)). \tag{2.4}$$

All characters of U arise in this way for some M .

- One has an embedding $T \hookrightarrow \mathrm{GSp}(W)$ by mapping $g \in T$ to

$$\hat{g} = \begin{pmatrix} g & \\ & (\det g)^t g^{-1} \end{pmatrix} \in \mathrm{GSp}(W).$$

This element has similitude factor $\lambda(\hat{g}) = \det g$. Moreover, if $u \in U$, then $ug = gu$.

- The *Bessel subgroup* of $\mathrm{GSp}_4(F)$ is then the semidirect product

$$R = U \rtimes T.$$

2.2.4 The Bessel period

Let π be an automorphic representation of $\mathrm{GSp}_4(\mathbb{A})$. All automorphic representations of the abelian group $T(\mathbb{A})$ are given by characters

$$\chi : T(F) \backslash T(\mathbb{A}) \rightarrow \mathbb{C}^\times,$$

of which we now fix a χ such that $\omega_\pi \cdot \chi|_{\mathbb{A}^\times} = 1$. We shall simultaneously think of χ as a character of $K^\times \backslash \mathbb{A}_K^\times$. For $\varphi_\pi \in \mathcal{V}_\pi$, the *Bessel period* of φ_π (with respect to χ) is defined by the period integral

$$\mathcal{P}(\varphi_\pi, \chi) = \int_{\mathbb{A}^\times T(F) \backslash T(\mathbb{A})} \int_{U(F) \backslash U(\mathbb{A})} \varphi_\pi(u\hat{g})\chi(g)\psi_S^{-1}(u) du dg \tag{2.5}$$

where du and dt are the Tamagawa measures on $U(\mathbb{A})$ and $\mathbb{A}^\times \backslash T(\mathbb{A})$ respectively. We realise \mathbb{A}^\times as the scalar matrices in the domain of integration $\mathbb{A}^\times R(F) \backslash R(\mathbb{A})$.

2.3 Notation for groups

For a fixed four-dimensional quadratic space V over F and the four-dimensional symplectic vector space $W = F^4$ (from Section 2.2.1) assign the notation

$$\begin{aligned} G &= \mathrm{GSp}(W), & H &= \mathrm{GO}(V), & H^0 &= \mathrm{GSO}(V), \\ G_1 &= \mathrm{Sp}(W), & H_1 &= \mathrm{O}(V), & H_1^0 &= \mathrm{SO}(V) \end{aligned}$$

which will be used freely throughout. Also define the groups

$$Y = \mathrm{G}(\mathrm{Sp}(W) \times \mathrm{O}(V)) = \{ (g, h) \in \mathrm{GSp}(W) \times \mathrm{GO}(V) : \lambda(g) = \lambda(h) \}$$

and

$$G^+ = \{ g \in G : \lambda(g) = \lambda(h) \text{ for some } h \in H \}.$$

3 The theta correspondence for $(\mathrm{GO}_4, \mathrm{GSp}_4)$

This section is devoted to constructing certain representations of GSp_4 from representations of GO_4 both locally and globally.

3.1 The local theta correspondence

Let v be a place of F and omit the subscript v from the notation in this section ($F = F_v$, $G = G(F_v)$, $W = W \otimes_F F_v$ and so on). Define the space $\mathbb{W} = W \otimes V$ which is given the symplectic form $(\cdot, \cdot)_{\mathbb{W}} = (\cdot, \cdot)_W \otimes (\cdot, \cdot)_V$. Then groups G_1 and H_1 form a *reductive dual pair* as subgroups of $\mathrm{Sp}(\mathbb{W})$. The polarisation of $W = W_1 \oplus W_1^\vee$ induces a polarisation

$$\mathbb{W} = (W_1 \otimes V) \oplus (W_1^\vee \otimes V)$$

on which we make some remarks:

- Having chosen the natural basis for W we may identify $W_1^\vee \otimes V \cong V^2$.
- There is an isomorphism $W_1^\vee \otimes V \cong \mathrm{Hom}_F(W_1, V)$.

(These comments are also relevant in the global setting, considering the adèlic points of the above spaces.)

Choose a non-trivial additive character ψ of F by taking it to be a local component of the standard (additive) adèlic character (2.3). Let $\omega = \omega_\psi$ be the *Weil representation* of $G_1 \times H_1$, with respect to ψ , which may be extended to a representation of Y as in [9, p. 82]. We realise ω in the space of Schwartz functions $\mathcal{V}_\omega = \mathcal{S}(V^2)$ where Y acts as follows. For $(g, h) \in G_1 \times H_1$ and $\phi \in \mathcal{S}(V^2)$:

$$\begin{aligned} \omega(1, h)\phi(x) &= \phi(h^{-1}x), \\ \omega(J_2, 1)\phi(x) &= \gamma_4 \hat{\phi}(x), \\ \omega(u(A), 1)\phi(x) &= \psi(\mathrm{Tr}(M_x A))\phi(x), \\ \omega(m(B), 1)\phi(x) &= \chi_V(\det B) |\det B|_F^2 \phi(xB) \end{aligned} \tag{3.1}$$

where the elements

$$J_2 = \begin{pmatrix} 0 & 1_2 \\ -1_2 & 0 \end{pmatrix}, \quad u(A) = \begin{pmatrix} 1_2 & A \\ 0 & 1_2 \end{pmatrix}, \quad m(B) = \begin{pmatrix} B & 0 \\ 0 & {}^t B^{-1} \end{pmatrix}$$

generate $G_1 = \mathrm{Sp}(W)$ where $A \in \mathrm{Hom}(W_1^\vee, W_1)$ and $B \in \mathrm{GL}(W_1)$. The character $\chi_V(\det B)$ is the quadratic character of F^\times ; it is defined using the Hilbert symbol. The action of the unipotent group U is dependent on the *Gram matrix* of $x = {}^t(x_1, x_2) \in V^2$ defined to be

$$M_x = ((x_i, x_j)_V)_{i,j}.$$

We define the character $\psi_S(u(A)) = \psi(\mathrm{Tr}(SA))$. We also have that $\gamma_4 \in \mu_4$ is a certain fourth root of unity and $\hat{\phi}$ is the Fourier transform of the Schwartz function ϕ (see [18, Section 1] for more details on this action). As in [9], the extended action of ω to Y is obtained by taking $(g, h) \in Y$, $\phi(x) \in \mathcal{S}(V^2)$ and setting

$$\omega(g, h)\phi(x) = |\lambda(h)|_F^{-2} \omega(g_1, 1)\phi(h^{-1}x) \tag{3.2}$$

where

$$g_1 = g \begin{pmatrix} 1_2 & 0 \\ 0 & \lambda(g)^{-1} 1_2 \end{pmatrix} \in G_1.$$

We now closely follow [7, Section 5]. Define the induced Weil representation by compact induction:

$$\Omega = \mathrm{Ind}_R^{H \times G^+}(\omega).$$

If σ is an irreducible, unitary, admissible representation of H and $\bar{\sigma}$ is the conjugate representation of σ , then the maximal $\bar{\sigma}$ -isotypic quotient of Ω is given by $\Omega / \bigcap \ker(\Psi)$ where Ψ runs over $\mathrm{Hom}_H(\Omega, \bar{\sigma})$. This is

a $\bar{\sigma}$ -isotypic direct sum as an H -representation. Since G^+ naturally commutes with H in $G^+ \times H$, the space of $\Omega / \cap \ker(\Psi)$ inherits an action of G^+ and as a representation of $G^+ \times H$ thus we may write

$$\Omega / \bigcap \ker(\Psi) = \bar{\sigma} \boxtimes \Theta^+(\sigma)$$

where $\Theta^+(\sigma)$ is a smooth representation of G^+ . We call $\Theta^+(\sigma)$ the *big theta lift* of σ to G^+ . Whilst $\Theta^+(\sigma)$ may be zero, it is known that if this is not the case then $\Theta^+(\sigma)$ is of finite length, and hence is admissible, and has a unique, maximal, irreducible quotient [7, Theorem A.1] which we denote $\theta^+(\sigma)$. This allow us to finally define the (local) *theta lift* of σ to G as

$$\theta(\sigma) = \text{Ind}_{G^+}^G(\theta^+(\sigma)).$$

By [7, Lemma 5.2], if σ is non-zero and unitary,¹ then $\theta(\sigma)$ is an irreducible representation of G . We obtain a unique (up to scalar) Y -equivariant, surjective map

$$\theta : \mathcal{V}_\sigma \otimes \mathcal{V}_\omega \rightarrow \mathcal{V}_{\theta(\sigma)}. \tag{3.3}$$

Remark 3.1. That $\theta^+(\sigma)$ exists as a unique, maximal, irreducible representation is in fact the statement of the local Howe conjectures.

3.2 The global theta correspondence

In this section we return to our original notation where F is a number field. The following construction follows [7, Section 7.2].

We have the fixed, non-trivial, additive character $\psi = \otimes_v \psi_v$ of \mathbb{A}/F (chosen in (2.3)). For each place v of F we let $\omega_v = \omega_{\psi_v}$ be the Weil representation of $Y(F_v)$, with respect to ψ_v , realised in the Schwartz space $\mathcal{V}_{\omega_v} = \mathcal{S}(V^2(F_v))$. Let $\mathcal{B}_{\omega_v} : \mathcal{V}_{\omega_v} \otimes \mathcal{V}_{\bar{\omega}_v} \rightarrow \mathbb{C}$ be the canonical pairing defined by

$$\mathcal{B}_{\omega_v}(\phi, \tilde{\phi}) = \int_{V^2(F_v)} \phi(x) \tilde{\phi}(x) dx.$$

The Weil representation of $Y(\mathbb{A})$ is given by $\omega = \otimes_v \omega_v$, and comes equipped with the decomposable unitary pairing $\mathcal{B}_\omega = \prod_v \mathcal{B}_{\omega_v}$. The action of ω in $\mathcal{V}_\omega = \otimes_v \mathcal{S}(V^2(F_v))$ is applied place-by-place using the local action in (3.1) and (3.2).

The global theta correspondence, in our setting, provides a cuspidal automorphic form on $G(\mathbb{A})$ from one on $H(\mathbb{A})$. We define this cusp form now. For a Schwartz function $\phi \in \mathcal{V}_\omega$ we note that the series

$$\sum_{x \in V^2(F)} \omega(g, h)\phi(x)$$

is a smooth function on $(g, h) \in Y(F) \backslash Y(\mathbb{A})$ of moderate growth.

Definition 3.1. Let σ be an irreducible, cuspidal automorphic representation of $H(\mathbb{A})$ and let $\phi \in \mathcal{V}_\omega$. Then for any $f \in \mathcal{V}_\sigma \subset \mathcal{A}_0(H)$ we define the *theta integral*

$$\theta(f, \phi; g) = \int_{H_1(F) \backslash H_1(\mathbb{A})} \sum_{x \in V^2(F)} \omega(g, hh_g)\phi(x) f(hh_g) dh \tag{3.4}$$

where h_g is any element in $H(\mathbb{A})$ such that $\lambda(h_g) = \lambda(g)$.

This integral is absolutely convergent and independent of the choice h_g since all such elements are of the form $h_g h_0$ for $h_0 \in H_1(\mathbb{A})$. One computes the central character of $\theta(f, \phi)$ to be equal to ω_σ , the central character of f (since $\dim V = 4$ is even).

By construction, $\theta(f, \phi)$ is a function on $G^+(F) \backslash G^+(\mathbb{A})$. By the natural inclusion of $G^+ \hookrightarrow G$ we extended $\theta(f, \phi)$ to a function on $G(F) \backslash G(\mathbb{A})$ by letting it take the value zero outside $G^+(\mathbb{A})$. This extension is unique.

¹ This is indeed the case when σ is a local component of an irreducible, unitary, cuspidal automorphic representation of $H(\mathbb{A})$ with a non-zero, cuspidal *global* theta lift to $G(\mathbb{A})$.

Remark 3.2. For any $h \in H = H^0 \rtimes \mu_2$ there is an $h_0 \in H^0$ with $\lambda(h) = \lambda(h_0)$ since $h = h_0 \varepsilon$ for $\varepsilon \in \mu_2 \cong \langle \iota \rangle$ where $\iota \in H$ is the element defined in (2.2) with $\lambda(\iota) = 1$. Thus we may interchange H with H^0 in the definition of G^+ .

Definition 3.2. Let $\theta(\sigma)$ be the automorphic representation of $G(\mathbb{A})$ realised in the space

$$\mathcal{V}_{\theta(\sigma)} = \{R(g)\theta(f, \phi) : f \in \mathcal{V}_\sigma, \phi \in \mathcal{S}(V^2(\mathbb{A})), g \in G(\mathbb{A})\}.$$

We call $\theta(\sigma)$ the (*global*) *theta lift* of σ to $G(\mathbb{A})$.

We shall fix assumptions on σ (see Assumption 4.1) under which $\theta(\sigma)$ is *cuspidal*. Under these conditions [7, Lemma 7.12] applies so that $\mathcal{V}_{\theta(\sigma)} \neq 0$. We then obtain a $Y(\mathbb{A})$ -equivariant, surjective map

$$\theta : \mathcal{V}_\sigma \otimes \mathcal{V}_\omega \rightarrow \mathcal{V}_{\theta(\sigma)}. \quad (3.5)$$

We may restrict θ to $\mathcal{V}_{\omega_v} \otimes \mathcal{V}_{\sigma_v}$ at each place v and conclude that, by the uniqueness of the local maps (3.3), for $\sigma = \otimes_v \sigma_v$,

$$\theta(\sigma) \cong \otimes_v \theta(\sigma_v)$$

and is irreducible [7, Lemma 7.2]. In particular, the local factors $\theta(\sigma_v)$ are unitary and non-zero at each v .

3.3 Automorphic induction

An alternative description of the theta lift is that it arises due to a functorial transfer of representations from $H'(\mathbb{A})$ to $\mathrm{GSp}_4(\mathbb{A})$ where $H' = \mathrm{Res}_{E/F}(\mathrm{GL}_2)$ is the Weil restriction of scalars (meaning that H' is unique in that $H'(F) = \mathrm{GL}_2(E)$ as algebraic groups) and E is a quadratic extension of F . For simplicity let us consider the trivial central character interpretation: the automorphic induction transfer between automorphic representations of the groups $H'_1 = \mathrm{Res}_{E/F} \mathrm{SL}_2$ and $G_1 = \mathrm{Sp}_4$. On the one hand, the L -group of G_1 is ${}^L G_1 = \mathrm{SO}_5(\mathbb{C}) \times \mathcal{G}_F$ where \mathcal{G}_F is the absolute Galois group of F . On the other hand, the L -group of H' is

$${}^L H'_1 \cong \prod_{\mathcal{G}_E \backslash \mathcal{G}_F} \mathrm{SL}_2(\mathbb{C}) \rtimes \mathcal{G}_F$$

noting $\mathcal{G}_E \backslash \mathcal{G}_F \cong \mathrm{Gal}(E/F)$ acts on the first factor in the product via permutations of the index set. Once again make note of the isomorphism $\mathrm{SO}_5(\mathbb{C}) \cong \mathrm{PGSp}_4(\mathbb{C})$ which gives rise to an embedding

$$\mathrm{SL}_2(\mathbb{C}) \times \mathrm{SL}_2(\mathbb{C}) \rightarrow \mathrm{Sp}_4(\mathbb{C}), \quad \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix} \right) \mapsto \begin{pmatrix} a & & b & \\ & a' & & b' \\ c & & d & \\ & c' & & d' \end{pmatrix}$$

which in turn induces an L -homomorphism

$$u : {}^L H'_1 \rightarrow {}^L G_1.$$

On composing u with a representation r of the Weil–Deligne group W'_E of E into ${}^L H'_1$ we obtain a representation $u \circ r$ that lands in ${}^L G_1$. Noting $W'_E \subset W'_F$, this acquired representation is precisely the induced representation

$$u \circ r = \mathrm{Ind}_{W'_E}^{W'_F} r$$

(on the Galois side). Whilst on the automorphic-side we have an irreducible, cuspidal automorphic representation $\mathrm{AI}(\pi')$ of $G_1(\mathbb{A})$ for each π' on $H'_1(\mathbb{A}) = \mathrm{SL}_2(\mathbb{A}_E)$. A more general review in support of this exposition is given in [3].

A characteristic property of such a lift is that the L -function of the representations ($\mathrm{AI}(\pi')$ and π') are equal, thus uniquely characterising the target L -packet. By the work of Roberts [18, Section 8] we find that this is also the case for the theta lift discussed in the previous two sections. Then, due to an exceptional isomorphism (see the next section, Section 4.1), we may realise the group GO_2 as $\mathrm{Res}_{E/F}(\mathrm{GL}_2)$ and hence any representation given by the above theta lift is functorial in this sense.

4 Automorphic Representations of GO_4

To classify the image of the theta correspondence for (GO_4, GSp_4) we provide a thorough review concerning the domain of the lift: we determine the structure of all four-dimensional quadratic spaces V , giving rise to $GO(V) \cong GO_4$, and with this analysis we examine the irreducible, cuspidal automorphic representations of $GO(V)(\mathbb{A})$. The review in this section is largely expository, however it includes new notation and crucial results which are used freely later on.

4.1 Four-dimensional quadratic spaces and their similitude groups

Any four-dimensional quadratic space is isomorphic to a member of a family of spaces whose structure is explicit and indexed by two invariants: a quaternion algebra and a square-free integer (corresponding to the discriminant). For more details we refer to the exposition given in [18, Section 2].

Consider a four-dimensional quadratic space V over F with $\text{disc}(V) = e$. Let $E = K_V$ be the discriminant algebra of V (defined in (2.1)) and put $\text{Gal}(E/F) = \{1, \kappa\}$, using both $\kappa(z)$ and z^κ to denote the image of $z \in E$ under κ . The usual norm and trace of E/F are given by

$$N_{E/F}(z) = zz^\kappa \quad \text{and} \quad \text{Tr}_{E/F}(z) = z + z^\kappa.$$

Definition 4.1. Let B be an arbitrary F -algebra whose centre is E with an involution $x \mapsto x^*$ that fixes E . Call B a *quadratic-quaternion algebra* over F if there is a quaternion algebra D , over F , contained in B such that the natural map $D \otimes_F E \rightarrow B$, given by $x \otimes z \mapsto xz$, is an isomorphism of E -algebras and the canonical involution on D is given by $x \mapsto x^*$. Choosing a D , there is no loss in generality in considering $B = D(E)$, the E -points of the F -algebra D . The *norm* and *trace* on B are defined respectively as

$$N_B(x) = xx^* \quad \text{and} \quad \text{Tr}_B(x) = x + x^*.$$

When restricted to D these are the usual reduced norm N_D and trace Tr_D . Endow B with the unique Galois action (with respect to D) by linearly extending the automorphism κ of E to B , that is $\kappa(xz) = x\kappa(z)$ for $z \in E$, $x \in D$. Denote this Galois action by κ as well. Finally, define a second four-dimensional quadratic space (over F) by

$$X = X_{D,e} = \{x \in D(E) : \kappa(x) = x^*\},$$

whose quadratic form, denoted N_X , is given by the restriction of N_B to X . We find that this new space has $\text{disc } X_{D,e} = \det N_X = e$ upon computing the determinant of N_X .

Remark 4.1. A *Galois action* on B is an F -automorphism $a : B \rightarrow B$ such that $a^2 = 1$ and $a(xz) = a(x)\kappa(z)$ for $z \in E$, $x \in B$. There is a bijection between Galois actions on B and quaternion F -algebras contained in B .

By [18, Proposition 2.7] we have the exact sequence

$$1 \rightarrow E^\times \xrightarrow{\Delta} F^\times \times B^\times \xrightarrow{\rho} \text{GSO}(X) \rightarrow 1 \quad (4.1)$$

where the injection $\Delta : E^\times \rightarrow F^\times \times B^\times$ is given by $\Delta(z) = (N_{E/F}(z), z)$ and the action of $F^\times \times B^\times$ on X is given by

$$\rho(s, a)x = s^{-1}ax\iota(a)^*.$$

In particular, writing ΔE^\times for $\text{Im}(\Delta)$, we have

$$F^\times \times B^\times / \Delta E^\times \cong \text{GSO}(X). \quad (4.2)$$

The similitude factor of an element $\rho(s, a) \in \text{GSO}(X)$ is given by

$$\lambda(\rho(s, a)) = s^{-2} N_{E/F}(N_B(a)).$$

We denote by ι the restriction of the Galois action κ to the subspace $X \subset B$ (again writing $\iota(x)$ and x' for the image of x under ι). The notation ι rightfully coincides with that already introduced in Section 2.1.4 since the map ι is precisely the unique element of $\text{GO}(X)$ satisfying the properties $\iota \in \text{O}(X)$, $\iota^2 = 1$ and $\det \iota = -1$ by [18, Proposition 2.5 and 2.7]. We choose this element to fix, once and for all, the splitting

$$\mu_2(F) \cong \langle \iota \rangle \quad \text{and} \quad \text{GO}(X) \cong \text{GSO}(X) \rtimes \langle \iota \rangle.$$

Conjugating an element $\rho(s, a) \in \text{GSO}(X)$ by ι gives the relation $\iota\rho(s, a)\iota = \rho(s, a')$; we denote this *adjoint* of ι action by

$$\text{Ad}(\iota) : \rho(s, a) \mapsto \rho(s, a'). \quad (4.3)$$

Proposition 4.2. *Let V be an arbitrary four-dimensional quadratic space over F of discriminant e . Then there exists a quaternion algebra D over F and an isomorphism $\gamma : V \rightarrow X_{D,e}$ such that the map*

$$c_\gamma : \text{GSO}(V) \rightarrow \text{GSO}(X_{D,e}),$$

given by $c_\gamma(g) = \gamma \circ g \circ \gamma^{-1}$, is an isomorphism of similitude groups. There is therefore no loss in generality in considering the space $\text{GSO}(X_{D,e})$ in place of $\text{GSO}(V)$

Proof. See [18, Proposition 2.8]. □

From here on in, fix a quaternion algebra D over F and a square free integer e . We shall work with the four-dimensional quadratic space $X = X_{D,e}$. Fix notation for: the quadratic extension $E = F(\sqrt{e})$ and the quadratic quaternion algebra $B = D(E)$. We assume the application of $V = X$ to the notations $H = \text{GO}(V)$ etc. of Section 2.3.

4.2 Local representation theory for $H(F_\nu)$

In this section let ν be a place of F and suppress the subscript ν from the notation (for example, F now denotes a local field). We shall systematically discuss the local (and later global) representation theory of H in terms of that of H^0 . We use this section to fix notation; this material has been previously considered in the expositions [10, Section 1], [18, Section 2–4] and [7, Section A] – we advise the reader to look there for details and proof. In [20], all restrictions in [18] are removed, in particular the quadratic space X may be of any signature.

4.2.1 Admissible representations of H^0

In light of the isomorphism in (4.2),

$$\rho : F^\times \times B^\times / \Delta E^\times \xrightarrow{\sim} H^0,$$

let (τ, \mathcal{V}_τ) be an irreducible, admissible, unitary representation of $B^\times = B^\times(F)$ with central character ω_τ (noting $Z_{B^\times} = E^\times$). Further assume that ω_τ is $\text{Gal}(E/F)$ -invariant; thus we let ν be the unitary character of F^\times such that

$$\omega_\tau = \nu^{-1} \circ \text{N}_{E/F}. \quad (4.4)$$

Every irreducible, admissible, unitary representation of H^0 may then be written in the form $\sigma_0 = \sigma_0(\nu, \tau)$, for such a ν and τ , by defining

$$\sigma_0(\rho(s, a)) = \nu(s)\tau(a).$$

Both σ_0 and τ are realised in the same space $\mathcal{V}_{\sigma_0} = \mathcal{V}_\tau$. The requirement on ν (4.4) ensures that $\sigma_0(\nu, \tau)$ is indeed trivial on ΔE^\times . We identify the centre $Z_{H^0} \cong F^\times$, through ρ , as the set

$$\{(x^{-1}, 1) : x \in F^\times\} \subset F^\times \times B^\times / \Delta E^\times,$$

from which we note that σ_0 has central character

$$\omega_{\sigma_0} = \nu^{-1}.$$

Definition 4.2. Suppose that v is not split in E (so that $E = E(F_v)$ is a field). In this case, we call an irreducible admissible representation σ_0 of H^0 *distinguished* if

$$\sigma_0 = \sigma_0(\omega_\rho^{-1}, \varrho_E^D)$$

for some irreducible admissible representation ϱ of $\mathrm{GL}_2(F)$; denoting by ϱ_E the base-change lift of ϱ from $\mathrm{GL}_2(F)$ to $\mathrm{GL}_2(E)$, and appending the superscript D to mean that ϱ_E^D is the Jacquet–Langlands transfer of ϱ_E from $\mathrm{GL}(E)$ to $D^\times(E) = B^\times$.

The central character of such a distinguished $\sigma_0(\omega_\rho^{-1}, \varrho_E^D)$ is ω_ρ , the central character of ϱ . This follows from properties of the base-change lift (that $\omega_{\varrho_E} = \omega_\rho \circ N_{E/F}$). Distinguished representations are invariant under the adjoint action of ι on H^0 (4.3). Hence a distinguished representation has the property that $\sigma_0 \cong \sigma_0 \circ \mathrm{Ad}(\iota)$ since we have $\varrho_E \cong \varrho_E \circ \iota$ (see [1, Section 3]).

4.2.2 Admissible representations of H

To describe the irreducible, admissible representations of H it suffices² to consider the induction of some σ_0 as σ_0 varies over the irreducible, admissible representations of H^0 . To make this explicit, put $\sigma_0^t = \sigma_0 \circ \mathrm{Ad}(\iota)$ and consider a second representation of H^0 in \mathcal{V}_{σ_0} given by

$$\sigma_0^t(h)v = \sigma_0(\iota h \iota)v \quad \text{for } v \in \mathcal{V}_{\sigma_0}.$$

Now define the representation $(\hat{\sigma}, \mathcal{V}_{\hat{\sigma}})$ of H by setting $\mathcal{V}_{\hat{\sigma}} = \mathcal{V}_{\sigma_0} \oplus \mathcal{V}_{\sigma_0^t}$ and letting H act on $u \oplus v$ by

$$\begin{cases} \hat{\sigma}(h_0)u \oplus v = \sigma_0(h_0)u \oplus \sigma_0^t(h_0)v, \\ \hat{\sigma}(\iota)u \oplus v = v \oplus u \end{cases}$$

noting that any $h \in H$ may be written uniquely as $h = h_0 \varepsilon$ for some $h_0 \in H^0$ and $\varepsilon \in \mu_2$.

On the other hand, recall that $\mathrm{Ind}_{H^0}^H(\sigma_0)$ is given by right translation in the space

$$\{f : H \rightarrow \mathcal{V}_{\sigma_0} \mid f(h_0 h) = \sigma_0(h_0)f(h) \text{ for } h_0 \in H^0\}.$$

One may check that there is an H -module isomorphism between the representations $\hat{\sigma} \cong \mathrm{Ind}_{H^0}^H(\sigma_0)$. We will use $\hat{\sigma}$ as a model for $\mathrm{Ind}_{H^0}^H(\sigma_0)$ from now on and proceed by dividing our analysis into two cases.

Definition 4.3. Let σ_0 be an irreducible, admissible representation of H^0 .

- We say σ_0 is *regular* if $\hat{\sigma} \cong \mathrm{Ind}_{H^0}^H(\sigma_0)$ is irreducible. We find $\hat{\sigma} \cong \hat{\sigma} \otimes \mathrm{sgn}$ and, as H^0 -representations, $\sigma_0 \neq \sigma_0^t$. In this case denote $\sigma_0^+ = \mathrm{Ind}_{H^0}^H(\sigma_0)$.
- We say σ_0 is *invariant* if $\hat{\sigma} \cong \mathrm{Ind}_{H^0}^H(\sigma_0)$ is reducible. We find $\hat{\sigma} \cong \hat{\sigma} \otimes \mathrm{sgn}$ and the adjoint action of ι in \mathcal{V}_{σ_0} is trivial, that is, $\sigma_0 \cong \sigma_0^t$. In this case

$$\mathrm{Ind}_{H^0}^H(\sigma_0) \cong \sigma_0^+ \oplus \sigma_0^-$$

where σ_0^\pm are two non-isomorphic irreducible representations of H .

Remark 4.3. If σ_0 is distinguished, then we have already noted that σ_0 is invariant. In this instance exactly one of σ_0^\pm occurs in the theta correspondence with GSp_4 (see [18, Theorem 3.4]), denoting this representation by σ_0^+ . Then for an irreducible, admissible representation σ of H we have that $\theta(\sigma) \neq 0$ if and only if $\sigma \neq \sigma_0^-$ for some distinguished σ_0 of H^0 .

² Let σ be an irreducible, admissible representation of H . Then either $\mathrm{Res}_{H^0}^H(\sigma)$ is irreducible, in which case σ is an irreducible constituent of $\mathrm{Ind}_{H^0}^H(\mathrm{Res}_{H^0}^H(\sigma))$ and we are in the ‘invariant’ case, or

$$\mathrm{Res}_{H^0}^H(\sigma) = \sigma_{0,1} \oplus \sigma_{0,2},$$

in which case $\sigma \cong \mathrm{Ind}_{H^0}^H(\sigma_{0,i})$ for either $i = 1, 2$; this is the ‘regular’ case. Definition 4.3 provides a full explanation of the invariant and regular cases.

4.3 Global representation theory and automorphic forms for $H(\mathbb{A})$

In this section we reinstate F as a number field. Our purpose is now to review the theory of automorphic forms on $H(\mathbb{A})$. The following sources should be referred to for more detail: [10, Section 1], [18, Sections 5–7] and [7, Section 2].

4.3.1 Automorphic representations of $H^0(\mathbb{A})$

The exactness of the sequence in (4.1) (taking $F = F_\nu$ for each place ν) implies that

$$1 \rightarrow \mathbb{A}_E^\times \xrightarrow{\Delta} \mathbb{A}^\times \times B^\times(\mathbb{A}) \xrightarrow{\rho} H^0(\mathbb{A}) \rightarrow 1$$

is also exact, where ρ and Δ operate as in the exact sequence (4.1) at each place. We identify $E(\mathbb{A})$ with \mathbb{A}_E and note $B^\times(\mathbb{A}) = D^\times(\mathbb{A}_E)$. As subspaces of $L^2(\mathbb{A}^\times \times B^\times(\mathbb{A}))$, the tensor product of the spaces of cusp forms $\mathcal{A}_0(F^\times) \otimes \mathcal{A}_0(B^\times)$ is dense in $\mathcal{A}_0(F^\times \times B^\times)$ and since these are spaces of smooth functions they are isomorphic. Any function on $\mathbb{A}^\times \times B^\times(\mathbb{A})/\Delta\mathbb{A}_E^\times$ is a function on $\mathbb{A}^\times \times B^\times(\mathbb{A})$ subject to the constraint that it is constant on equivalence classes modulo $\Delta\mathbb{A}_E^\times = \text{Im}(\Delta)$. In particular, if $\nu : F^\times \backslash \mathbb{A}^\times \rightarrow \mathbb{C}^\times$ is a unitary Hecke character and (τ, \mathcal{V}_τ) an irreducible, cuspidal automorphic representation of $B^\times(\mathbb{A})$ then, given some $\eta \in \mathcal{V}_\tau$, we have that $\nu \otimes \eta \in \mathcal{A}_0(F^\times \times B^\times/\Delta E^\times)$ if and only if

$$\omega_\tau(z) = \nu^{-1} \circ N_{E/F}(z) \quad \text{for all } z \in \mathbb{A}_E^\times$$

where $\omega_\tau : E^\times \backslash \mathbb{A}_E^\times \rightarrow \mathbb{C}^\times$ is the central character of τ . Hence any irreducible, cuspidal automorphic representation of $H^0(\mathbb{A})$ is of the form $\sigma_0 = \sigma_0(\nu, \tau)$, for such a ν and τ , where σ_0 is realised in the space of cusp forms $\mathcal{V}_{\sigma_0} = \{\nu \otimes \eta : \eta \in \mathcal{V}_\tau\}$ by $\sigma_0(\rho(s, a))\nu \otimes \eta = \nu(s)\nu \otimes \tau(a)\eta$. Once again, the central character of σ_0 is $\omega_{\sigma_0} = \nu^{-1}$.

4.3.2 Factorising automorphic representations of $B^\times(\mathbb{A}_E)$ and $H^0(\mathbb{A})$

Consider the isomorphism

$$E \otimes_F F_\nu \cong \prod_{w|\nu} E_w \tag{4.5}$$

where the product is over all places w of E above ν [17, Proposition 4-40]. One deduces

$$B^\times(F_\nu) \cong \prod_{w|\nu} B^\times(E_w).$$

Thus smooth representations of $B^\times(F_\nu)$ are of the form $\tau_\nu = \otimes_{w|\nu} \tau_w$ where the τ_w are smooth representations of $B^\times(E_w)$ for $w|\nu$. If $\sigma_0 = \sigma_0(\nu, \tau)$ is an irreducible, cuspidal automorphic representation of $H^0(\mathbb{A})$, as in Section 4.3.1, then by the tensor product theorem we may assume $\sigma_0 \cong \otimes_\nu \sigma_{0,\nu}$ and $\nu = \otimes_\nu \nu_\nu$, over places ν of F , and $\tau \cong \otimes_w \tau_w$ over places w of E . Then, by the previous remark, these local factors are related by $\sigma_{0,\nu} = \sigma_{0,\nu}(\nu_\nu, \tau_\nu)$ where $\tau_\nu = \otimes_{w|\nu} \tau_w$ and the space $\mathcal{V}_{\sigma_{0,\nu}} = \mathcal{V}_{\tau_\nu} = \otimes_{w|\nu} \mathcal{V}_{\tau_w}$ (as per Section 4.2.1).

4.3.3 Automorphic representations of $H(\mathbb{A})$

Assumption 4.1. Let $\sigma \cong \otimes_\nu \sigma_\nu$ be an irreducible, cuspidal automorphic representation of $H(\mathbb{A})$ realised on the space $\mathcal{V}_\sigma \subset \mathcal{A}_0(H)$. For the remainder of this paper we shall assume the following for such a representation σ .

- (1) The Jacquet–Langlands transfer of $\sigma|_{B^\times(\mathbb{A}_E)}$ to $\text{GL}_2(\mathbb{A}_E)$ is cuspidal.
- (2) There is at least one place ν for which $\sigma_\nu \cong \sigma_\nu \otimes \text{sgn}$.
- (3) If $\sigma_\nu \neq \sigma_\nu \otimes \text{sgn}$, then $\sigma_\nu \neq \sigma_{0,\nu}^-$ for any distinguished (and invariant) admissible representation $\sigma_{0,\nu}$ of $H^0(F_\nu)$.

These conditions are imposed in [7], thus ensuring that $\theta(\sigma)$ is both cuspidal (1) and non-zero (3). Condition (2) is necessary to compute the Petersson inner product of the theta lift $\theta(\sigma)$ in (4.9).

We now determine all such σ by considering their restriction to $H^0(\mathbb{A})$. (This top-down approach contrasts with the bottom-up analysis used in the local setting.) To this end, define a (possibly infinite) subset of the places of F by

$$\mathfrak{S} = \{v : \sigma_v \cong \sigma_v \otimes \text{sgn}\}.$$

Assumption 4.1 (2) implies $\mathfrak{S} \neq \emptyset$. By the tensor product theorem, fix an isomorphism of $H(\mathbb{A})$ -representations

$$\mathcal{V}_\sigma \cong \bigotimes'_v \mathcal{V}_{\sigma_v} = \varinjlim_S \left(\bigotimes_{v \in S} \mathcal{V}_{\sigma_v} \right) \otimes \left(\bigotimes_{v \notin S} f_v^\circ \right)$$

where \mathcal{V}_{σ_v} is the space of σ_v and, for a sufficiently large set of places S outside which σ_v is unramified, $f_v^\circ \in \mathcal{V}_{\sigma_v}$ is an $H(\mathcal{O}_v)$ -invariant (spherical) vector for $v \notin S$. By analogy with our local discussion Section 4.2.2, the restriction of σ_v to H_v^0 gives rise to two cases.

- If $v \in \mathfrak{S}$, then $\sigma_v|_{H_v^0} \cong \sigma_{0,v} \oplus \sigma_{0,v}^t$ where $\sigma_{0,v}$ is an irreducible representation of H_v^0 with $\sigma_{0,v} \neq \sigma_{0,v}^t$. Earlier, we called such a $\sigma_{0,v}$ *regular* and noted that its induction, $\hat{\sigma}_v$, was irreducible. The space of σ_v decomposes as $\mathcal{V}_{\sigma_v} = \mathcal{V}_{\sigma_{0,v}} \oplus \mathcal{V}_{\sigma_{0,v}^t}$, realising the space $\mathcal{V}_{\sigma_{0,v}^t} \cong \sigma_v(t)\mathcal{V}_{\sigma_{0,v}}$. For almost all $v \in \mathfrak{S}$, the spherical vector $f_v^\circ = \mathfrak{F}_v^\circ + \sigma_v(t)\mathfrak{F}_v^\circ \in \mathcal{V}_{\sigma_{0,v}} \oplus \mathcal{V}_{\sigma_{0,v}^t}$ where \mathfrak{F}_v° is an $H^0(\mathcal{O}_v)$ -invariant vector.
- If $v \notin \mathfrak{S}$, then $\sigma_v|_{H_v^0}$ is irreducible and *invariant*; we have $\mathcal{V}_{\sigma_v} = \mathcal{V}_{\sigma_{0,v}}$ and the spherical vector $f_v^\circ = \mathfrak{F}_v^\circ$ is $H^0(\mathcal{O}_v)$ -invariant. Write $\sigma_{0,v} = \sigma_v|_{H_v^0}$ in this case.

Let S be a sufficiently large set of places of F and put $S' = S \setminus (S \cap \mathfrak{S})$. For each $\varepsilon = (\varepsilon_v) \in \mu_2(F_{S \cap \mathfrak{S}})$ define $\mathcal{V}_{\sigma,S}^\varepsilon \subset \mathcal{V}_\sigma$ by

$$\mathcal{V}_{\sigma,S}^\varepsilon \cong \left(\bigotimes_{v \in S \cap \mathfrak{S}} \sigma_v(\varepsilon_v) \mathcal{V}_{\sigma_{0,v}} \right) \otimes \left(\bigotimes_{v \in S'} \mathcal{V}_{\sigma_{0,v}} \right) \otimes \left(\bigotimes_{v \notin S} f_v^\circ \right).$$

Viewing σ from a different perspective, consider the space of restricted functions

$$\mathcal{V}_\sigma|_{H^0(\mathbb{A})} = \{f|_{H^0(\mathbb{A})} : f \in \mathcal{V}_\sigma\}.$$

By [10, Lemma 2] there exists an irreducible, cuspidal automorphic representation σ_0 of $H^0(\mathbb{A})$ realised in a space of cusp forms \mathcal{V}_{σ_0} such that

$$\mathcal{V}_\sigma|_{H^0(\mathbb{A})} = \mathcal{V}_{\sigma_0} \oplus \mathcal{V}_{\sigma_0}^t, \quad (4.6)$$

defining $\mathcal{V}_{\sigma_0}^t = \{f^t = f \circ \text{Ad}(t) : f \in \mathcal{V}_{\sigma_0}\}$, and such that $\sigma_0 \neq \sigma_0^t$. In this circumstance we shall say σ *lies above* σ_0 .

Applying the tensor product theorem and comparing the local components of σ and σ_0 with those $\sigma_{0,v}$ already defined, we may assume that $\sigma_0 \cong \otimes_v \sigma_{0,v}$. Moreover, choosing $\varepsilon = 1$, the restriction of the space of functions $\mathcal{V}_{\sigma,S}^1 = \mathcal{V}_{\sigma_0}$.

As a final remark, (4.6) shows that $\mathcal{V}_{\sigma,S}^\varepsilon|_{H^0(\mathbb{A})} = \{0\}$ unless $\varepsilon \in \mu_2(F)$ (else contradicting that $\sigma_0 \neq \sigma_0^t$). In particular, consider evaluating a function $f \in \mathcal{V}_{\sigma,S}^1$ on

$$H(\mathbb{A}) = \bigcup_{\varepsilon \in \mu_2(F_{S \cap \mathfrak{S}})} H^0(\mathbb{A}) \mu_2(\mathbb{A}^{S \cap \mathfrak{S}}) \varepsilon.$$

For $\varepsilon \in \mu_2(F_{S \cap \mathfrak{S}})$ we have $\mathcal{V}_{\sigma,S}^\varepsilon = \sigma(\varepsilon) \mathcal{V}_{\sigma,S}^1$ and hence $\sigma(\varepsilon)f = 0$ unless $\varepsilon \in \mu_2(F)$. We then obtain [7, Lemma 2.2]

$$\text{supp}(f) \subset H^0(\mathbb{A}) \mu_2(\mathbb{A}^{S \cap \mathfrak{S}}) \cup H^0(\mathbb{A}) \mu_2(\mathbb{A}^{S \cap \mathfrak{S}}) t. \quad (4.7)$$

4.4 Explicit unitary pairings and the Petersson inner product

The unique (up to scalar) unitary pairings $\mathcal{B}_{\sigma_{0,v}} : \mathcal{V}_{\sigma_{0,v}} \otimes \mathcal{V}_{\bar{\sigma}_{0,v}} \rightarrow \mathbb{C}$ associated to the local components $\sigma_{0,v} = \sigma_{0,v}(\nu_v, \tau_v)$ of $\sigma_0(\nu, \tau)$, as in Section 4.3.2, are precisely the pairings on $\mathcal{V}_{\tau_v} \otimes \mathcal{V}_{\bar{\tau}_v}$ since $\mathcal{V}_{\sigma_{0,v}} = \mathcal{V}_{\tau_v}$ and ν_v is unitary.

We therefrom assume that, whenever \mathcal{B}_{τ_v} is specified, by $\mathcal{B}_{\sigma_{0,v}}$ we always mean the pairing $\mathcal{B}_{\sigma_{0,v}} = \mathcal{B}_{\tau_v}$. The possible splitting of ν in E must also be accounted for in our choice of pairing: we make the convention

that if \mathcal{B}_{τ_w} is a specified pairing on \mathcal{V}_{τ_w} (for each place w of E lying above v), then

$$\mathcal{B}_{\tau_v} = \bigotimes_{w|v} \mathcal{B}_{\tau_w}$$

is the fixed pairing on $(\otimes_{w|v} \mathcal{V}_{\tau_w}) \otimes (\otimes_{w|v} \mathcal{V}_{\bar{\tau}_w})$ and hence also on $\mathcal{V}_{\sigma_{0,v}} \otimes \mathcal{V}_{\bar{\sigma}_{0,v}}$.

If $\mathcal{V}_{\sigma_{0,v}}$ carries a pairing $\mathcal{B}_{\sigma_{0,v}}$ and σ_v is an irreducible, admissible representation above $\sigma_{0,v}$, then we choose to consider a specific pairing on \mathcal{V}_{σ_v} :

- If $v \in \mathfrak{S}$, then $\mathcal{V}_{\sigma_v}|_{H_v^0} = \mathcal{V}_{\sigma_{0,v}} \oplus \mathcal{V}_{\sigma'_{0,v}}$ is irreducible; take the pairing

$$\mathcal{B}_{\sigma_v} : (\mathcal{V}_{\sigma_{0,v}} \oplus \mathcal{V}_{\sigma'_{0,v}}) \otimes (\mathcal{V}_{\bar{\sigma}_{0,v}} \oplus \mathcal{V}_{\bar{\sigma}'_{0,v}}) \rightarrow \mathbb{C}$$

given by $\mathcal{B}_{\sigma_v}((x + \sigma_v(t)y), (\tilde{x} + \bar{\sigma}_v(t)\tilde{y})) = \frac{1}{2}(\mathcal{B}_{\sigma_{0,v}}(x, \tilde{x}) + \mathcal{B}_{\sigma_{0,v}}(y, \tilde{y}))$.

- If $v \notin \mathfrak{S}$, then $\mathcal{V}_{\sigma_v}|_{H_v^0} = \mathcal{V}_{\sigma_{0,v}}$ is irreducible; take $\mathcal{B}_{\sigma_v} = \mathcal{B}_{\sigma_{0,v}}$.

This pairing is chosen carefully so that we may factorise the Petersson inner products \mathcal{B}_{σ} and $\mathcal{B}_{\bar{\sigma}}$ when $\sigma = \otimes_v \sigma_v$ is an automorphic representation of $H(\mathbb{A})$ that lies above $\sigma_0 = \otimes_v \sigma_{0,v}$. As before, fix an isomorphism for the conjugate representation $\bar{\sigma} \cong \otimes_v \bar{\sigma}_v$,

$$\mathcal{V}_{\bar{\sigma}} \cong \bigotimes_v \mathcal{V}_{\bar{\sigma}_v} = \varinjlim_S \left(\bigotimes_{v \in S} \mathcal{V}_{\bar{\sigma}_v} \right) \otimes \left(\bigotimes_{v \notin S} \tilde{f}_v^\circ \right)$$

where, for a sufficiently large set of places S outside which $\bar{\sigma}_v$ is unramified, $\tilde{f}_v^\circ \in \mathcal{V}_{\bar{\sigma}_v}$ is an $H(\mathcal{O}_v)$ -invariant (spherical) vector for $v \notin S$. If $v \in \mathfrak{S}$, then $\tilde{f}_v^\circ = \tilde{\mathfrak{F}}_v^\circ + \sigma_v(t)\tilde{\mathfrak{F}}_v^\circ$ where $\tilde{\mathfrak{F}}_v^\circ$ is an $H^0(\mathcal{O}_v)$ -invariant vector and if $v \notin \mathfrak{S}$ then $\tilde{f}_v^\circ = \tilde{\mathfrak{F}}_v^\circ$ is $H^0(\mathcal{O}_v)$ -invariant.

Lemma 4.4. *For almost all v suppose that $\mathcal{B}_{\sigma_{0,v}}$ is normalised by $\mathcal{B}_{\sigma_{0,v}}(\tilde{\mathfrak{F}}_v^\circ, \tilde{\mathfrak{F}}_v^\circ) = 1$. Then, if the pairings $\mathcal{B}_{\sigma_{0,v}}$ are normalised so that the Petersson inner product may be factorised as $\mathcal{B}_{\sigma_0} = \prod_v \mathcal{B}_{\sigma_{0,v}}$, we additionally have the following decomposition:*

$$\mathcal{B}_{\sigma} = \prod_v \mathcal{B}_{\sigma_v}.$$

Proof. See [7, Lemma 2.3]. □

The Petersson inner products for both the automorphic representations $\sigma_0 = \sigma_0(v, \tau)$ and τ agree: if we have $(\eta, \tilde{\eta}) \in \mathcal{V}_{\tau} \otimes \mathcal{V}_{\bar{\tau}}$ and $f_0 = v \otimes \eta, \tilde{f}_0 = \bar{v} \otimes \tilde{\eta}$, then

$$\mathcal{B}_{\sigma_0}(f_0, \tilde{f}_0) = \mathcal{B}_{\tau}(\eta, \tilde{\eta}).$$

The Petersson inner product associated to the unitary Hecke character χ of \mathbb{A}_K^\times (trivially) coincides with the Tamagawa number of $F^\times \backslash K^\times$, given by $\text{Vol}(\mathbb{A}^\times K^\times \backslash \mathbb{A}_K^\times) = 2$ (see [14, p. 44]). Underlying our calculations we choose local pairings $\mathcal{B}_{\chi_v} = 1$ at all v .

4.5 The Petersson inner product for theta lifts

Gan–Ichino prove a decomposition of the Petersson inner product for the theta lift $\theta(\sigma)$ with respect to some specified pairings for the local factors $\theta(\sigma_v)$. This result assumes that F is a totally real number field and that $\sigma = \otimes_v \sigma_v$ is an irreducible, cuspidal automorphic representation of $H(\mathbb{A})$ satisfying Assumption 4.1. In particular, in this assumption, conditions (2) and (3) are used explicitly in the proof of this formula whereas the totally real assumption is required for an application of the Siegel–Weil formula.

Fix a choice of local pairings $\mathcal{B}_{\sigma_{0,v}}$ such that $\mathcal{B}_{\sigma_0} = \prod_v \mathcal{B}_{\sigma_{0,v}}$ and consider the pairings \mathcal{B}_{σ_v} , defined in Section 4.4. For $(f, \tilde{f}) \in \mathcal{V}_{\sigma} \otimes \mathcal{V}_{\bar{\sigma}}$ and Schwartz functions $(\phi_v, \tilde{\phi}_v) \in \mathcal{V}_{\omega_v} \otimes \mathcal{V}_{\bar{\omega}_v}$ define

$$\mathcal{B}_{\theta(\sigma_v)}(\theta(f_v, \phi_v), \theta(\tilde{f}_v, \tilde{\phi}_v)) = \frac{\zeta_{F_v}(2)\zeta_{F_v}(4)}{L(\text{std}, \sigma_v, 1)} \int_{H_1(F_v)} \mathcal{B}_{\omega_v}(\omega_v(h_v)\phi_v, \tilde{\phi}_v) \mathcal{B}_{\sigma_v}(\sigma_v(h_v)f_v, \tilde{f}_v) dh_v \quad (4.8)$$

where the Haar measures dh_v on $H_{1,v}$ are those determined by a differential form (of top degree) on H_1 and the self-dual Haar measure on F_v^\times (with respect to ψ_v) – these in fact give the Tamagawa measure $dh = \prod_v dh_v$ of $H_1(\mathbb{A})$ (as constructed in [22]).

Gan–Ichino take care in deriving the constant of proportionality between the Petersson inner product for $\theta(\sigma)$ and $\prod_v \mathcal{B}_{\theta(\sigma_v)}$. With Assumption 4.1 we have [7, Proposition 7.13]

$$\mathcal{B}_{\theta(\sigma)} = \frac{L(\text{std}, \sigma, 1)}{\zeta_F(2)\zeta_F(4)} \prod_v \mathcal{B}_{\theta(\sigma_v)}. \tag{4.9}$$

5 Global calculation: The Bessel period

Preliminary remarks aside, we use this section to determine the form of the Bessel period (2.5) for the theta integral (3.4). First of all we explicitly highlight any running assumptions and notations (in addition to those in Assumption 4.1).

5.1 Hypotheses and variables

We have fixed the (base) number field F to be totally real. This assumption permits the use of the Siegel–Weil formula (or rather its corollary; the Rallis inner product formula) in a calculation made in [7] whereby the Petersson inner product for a theta lift is computed in terms of local pairings (see Proposition 4.9).

In Section 4.1 we acquired the following notation and assumptions: V is a four-dimensional quadratic space (over F) of discriminant $\text{disc } V = e$; we assume that e is not a square in F^\times (since the case when e is a square has been settled by Liu); Proposition 4.2 implies that it suffices to fix such an $e \in F^\times$ and a (possibly split) quaternion algebra D over F and consider instead the space $X = X_{D,e}$ – we do this and apply $V = X$ to the notations $H = \text{GO}(V)$ etc. of Section 2.3; fix once and for all $E = F(\sqrt{e})$ and $B = D(E) \cong D \otimes_F E$.

Our result is concerned with irreducible, cuspidal automorphic representations of $\text{GSp}_4(\mathbb{A})$ lifted from $\text{GO}(V)(\mathbb{A})$ by the theta correspondence (Section 3).

Assumption 5.1. We only consider representations of $\text{PGSp}_4(\mathbb{A}) \cong \text{SO}_5(\mathbb{A})$; these are precisely the representations of $\text{GSp}_4(\mathbb{A})$ with *trivial central character*.

Note that the theta lift $\theta(\sigma)$ has central character $\omega_{\theta(\sigma)} = \omega_\sigma$ so we assume $\omega_\sigma = 1$. If σ lies above $\sigma_0 = \sigma_0(\nu, \tau)$, as in (4.6), then $\nu = \omega_\sigma^{-1} = 1$. For the remainder of this paper, we keep in mind a fixed irreducible, cuspidal automorphic representation $\sigma \cong \otimes_v \sigma_v$ of $H(\mathbb{A})$ (in the space \mathcal{V}_σ) lying above $\sigma_0 = \sigma_0(1, \tau)$ where $\tau \cong \otimes_w \tau_w$ is an irreducible, cuspidal automorphic representation of $B^\times(\mathbb{A})$ whose central character $\omega_\tau = 1$. Also fix a factorisation for the conjugate representation $\bar{\sigma} \cong \otimes_v \bar{\sigma}_v$. There exists a set of places $\mathfrak{S} = \{v : \sigma_v \cong \sigma_v \otimes \text{sgn}\}$ which determine σ uniquely given σ_0 (see Section 4.3.3).

Let $f = \otimes_v f_v \in \mathcal{V}_\sigma$ be a pure tensor, fixing this choice throughout the remainder of this paper. We identify a factorisation for the conjugate of f by

$$\bar{f} = \otimes_v \bar{f}_v \tag{5.1}$$

so that it makes sense to talk about a specific \bar{f}_v corresponding to a local factor f_v of f . Similarly, we fix factorisations for the Schwartz functions $\phi = \otimes_v \phi_v \in \mathcal{V}_\omega$ and $\bar{\phi} = \otimes_v \bar{\phi}_v \in \mathcal{V}_{\bar{\omega}}$.

Choose a series of local unitary pairings \mathcal{B}_{τ_w} on $\mathcal{V}_{\tau_w} \otimes \mathcal{V}_{\bar{\tau}_w}$, for each place w of E , such that the Petersson pairing has the factorisation $\mathcal{B}_\tau = \prod_w \mathcal{B}_{\tau_w}$. Due to the choices of Section 4.4, we then automatically obtain the pairings $\mathcal{B}_{\sigma_{0,v}}$ and \mathcal{B}_{σ_v} for $\sigma_{0,v}$ and σ_v , respectively. Note that these depend on the place v of F . The Petersson pairings will satisfy a similar factorisation

$$\mathcal{B}_{\sigma_0} = \prod_v \mathcal{B}_{\sigma_{0,v}} \quad \text{and} \quad \mathcal{B}_\sigma = \prod_v \mathcal{B}_{\sigma_v}. \tag{5.2}$$

Fix another non-square element $d \in F^\times$. Let $K = F(\sqrt{d})$ and define a Hecke character $\chi : K^\times \backslash \mathbb{A}_K^\times \rightarrow \mathbb{C}^\times$. Then K and χ index a unique Bessel period (see Section 2.2.4). We impose the following assumption, which is essentially the trivial central character assumption when considering χ as a representation of $\text{GSO}(X)$.

Assumption 5.2. Suppose that χ is unitary and satisfies $\chi|_{\mathbb{A}_K^\times} = 1$.

5.2 Explicit vectors

We shall consider vectors $\varphi = \theta(f, \phi)$ for $f \in \mathcal{V}_\sigma$ and $\phi \in \mathcal{V}_\omega$ such that $\varphi = \otimes_v \varphi_v$ is a pure tensor. The global map θ of (3.5) is linear in each variable and hence φ is a pure tensor when both $f = \otimes_v f_v$ and $\phi = \otimes_v \phi_v$ are pure tensors (as we have assumed). We fix the notation $\varphi_v = \theta(f_v, \phi_v)$ for the local components in the factorisation of $\theta(f, \phi)$ (noting that this is necessary as each local map θ (3.3) is only unique up to a scalar constant).

Our choice of local vectors $\bar{f}_v \in \mathcal{V}_{\bar{\sigma}_v}$ (see (5.1)) and $\bar{\phi}_v \in \mathcal{V}_{\bar{\omega}_v}$ give rise to the factors in $\bar{\varphi} = \otimes_v \bar{\varphi}_v$ in the sense that

$$\bar{\varphi}_v = \overline{\theta(f_v, \phi_v)} = \theta(\bar{f}_v, \bar{\phi}_v) \quad (5.3)$$

by the uniqueness of (3.3) and (3.5), the choice of vectors $\varphi_v = \theta(f_v, \phi_v)$ and then applying [7, Proposition 5.5].

Lemma 5.1. *Define $f^!(h) = f(h\iota)$. One has $\theta(f, \phi) = \theta(f^!, \omega(\iota)\phi)$.*

Proof. We compute

$$\begin{aligned} \theta(f^!, \omega(\iota)\phi; g) &= \int_{H_1(F) \backslash H_1(\mathbb{A})} \sum_{x \in X^2(F)} \omega(g, hh_g \iota) \phi(x) f(\iota h h_g \iota) dh \\ &= \int_{H_1(F) \backslash H_1(\mathbb{A})} \sum_{\iota x \in X^2(F)} \omega(g, \iota h \iota h'_g) \phi(\iota x) f(\iota h \iota h'_g) dh \\ &= \theta(f, \phi; g) \end{aligned}$$

where $h'_g = \iota h_g \iota$ has $\lambda(h'_g) = \lambda(g)$. Here we use the automorphy of f under $\iota \in \mu_2(F)$ and rearrange the summation by $x \mapsto \iota x$. The Tamagawa measure dh is invariant under the transformation $h \mapsto \iota h \iota$. \square

Since an arbitrary element f of \mathcal{V}_σ is of the form $f = f_1 + f_2^!$ for some $f_1, f_2 \in \mathcal{V}_{\sigma, S}^1$ (by (4.6)), Lemma 5.1 implies

$$\theta(f_1 + f_2^!, \phi) = \theta(f_1, \phi) + \theta(f_2, \omega(\iota)\phi).$$

There is then no loss in generality in restricting our choice of $f \in \mathcal{V}_\sigma$ to the following.

Assumption 5.3. For a fixed, finite set S , assume that $f = \otimes_v f_v \in \mathcal{V}_{\sigma, S}^1$ is a pure tensor. Such an f satisfies the property that $f|_{H^0(\mathbb{A})} \in \mathcal{V}_{\sigma_0, v}$.

Recalling that τ is the automorphic representation of $B^\times(\mathbb{A})$ such that $\sigma_0 = \sigma_0(1, \tau)$, we denote by

$$\eta = \otimes_w \eta_w \in \mathcal{V}_\tau$$

(decomposed over places w of E) the function such that

$$f(\rho(s, a)) = \eta(a).$$

The local factors of these functions are identified by $f_v = \otimes_w \eta_w$ (see Section 4.3.2). Note that $f^! = \sigma(\iota)f$, and since $\theta(f, \phi) = \otimes_v \theta(f_v, \phi_v)$, Lemma 5.1 implies that for each v

$$\theta(f_v, \phi_v) = \theta(\sigma_v(\iota)f_v, \omega_v(\iota)\phi_v).$$

5.3 A calculation in terms of the variant theta integral

To simplify matters (overall) we introduce the *variant* theta integral (to be compared with (3.4)):

$$\theta^0(f, \phi; g) = \int_{H_1^0(F) \backslash H_1^0(\mathbb{A})} \sum_{x \in X^2(F)} \omega(g, h_0 h_g) \phi(x) f(h_0 h_g) dh_0$$

where the domain is defined in terms of the connected, index-two subgroup H_1^0 of H_1 . For this function we also have

$$\theta^0(f, \phi) = \theta^0(f^!, \omega(\iota)\phi) \quad (5.4)$$

by a computation identical to Lemma 5.1. Observe how $\theta^0(f, \phi)$ is related to $\theta(f, \phi)$.

Lemma 5.2. *For any integrable function Φ on $H_1(F)\backslash H_1(\mathbb{A})$ we have*

$$\int_{H_1(F)\backslash H_1(\mathbb{A})} \Phi(h) dh = \int_{\mu_2(F)\backslash\mu_2(\mathbb{A})} \int_{H_1^0(F)\backslash H_1^0(\mathbb{A})} \Phi(h_0\varepsilon) dh_0 d\varepsilon$$

where $d\varepsilon$ is the Tamagawa measure on $\mu_2(\mathbb{A})$.

Since $\theta^0(f, \phi; g)$ is independent of a particular choice of h_g , we may apply Lemma 5.2 and substitute $h_g \mapsto \varepsilon h_g \varepsilon$ (as $\lambda(\varepsilon) = 1$) to find

$$\theta(f, \phi; g) = \int_{\mu_2(F)\backslash\mu_2(\mathbb{A})} \theta^0(\sigma(\varepsilon)f, \omega(\varepsilon)\phi; g) d\varepsilon. \quad (5.5)$$

This relation permits one to consider the refined quantity $\mathcal{P}(\theta^0(f, \phi), \chi)$.

5.4 Unfolding the Weil representation

By definition (see (2.5)) we have

$$\mathcal{P}(\theta^0(f, \phi), \chi) = \int_{\mathbb{A} \times T(F)\backslash T(\mathbb{A})} \int_{U(F)\backslash U(\mathbb{A})} \theta^0(f, \phi; u\hat{g})\chi(g)\psi_S^{-1}(u) du dg$$

so we start out by computing

$$\theta^0(f, \phi; u\hat{g}) = \int_{H_1^0(F)\backslash H_1^0(\mathbb{A})} \sum_{x \in X^2(F)} \omega(u\hat{g}, h_0 h_g)\phi(x)f(h_0 h_g) dh_0.$$

Applying the action of ω to $\phi = \otimes_v \phi_v$ (place-by-place) we find that

$$\omega(u\hat{g}, h_0 h_g)\phi(x) = \left(\prod_v \chi_{V,v}(\det(g_v)) |\lambda(g_v)|_v^{-2} |\det g_v|_v^2 \right) \psi_{M_x}(u)\phi(h_g^{-1}h_0^{-1}xg) = \psi_{M_x}(u)\phi(h_g^{-1}h_0^{-1}xg),$$

recalling $\prod_v \chi_{V,v}(\det(g_v)) = 1$ (by quadratic reciprocity) and ψ_{M_x} is the character of U defined in (2.4). On removing the factor containing the integral over $U(F)\backslash U(\mathbb{A})$ we obtain

$$\mathcal{P}(\theta^0(f, \phi), \chi) = \int_{\mathbb{A} \times T(F)\backslash T(\mathbb{A})} \int_{H_1^0(F)\backslash H_1^0(\mathbb{A})} \sum_{x \in X^2(F)} \phi(h_g^{-1}h_0^{-1}xg)f(h_0 h_g)\Phi(x) dh_0 dg$$

where we have introduced the notation

$$\Phi(x) = \int_{U(F)\backslash U(\mathbb{A})} \psi_{M_x}(u)\psi_S^{-1}(u) du.$$

This integral of orthogonal characters simply boils down to

$$\Phi(x) = \begin{cases} \text{Vol}(U(F)\backslash U(\mathbb{A})), & \psi_{M_x} = \psi_S, \\ 0, & \text{otherwise.} \end{cases} \quad (5.6)$$

The group U is abelian (and hence unimodular) so the Tamagawa number $\text{Vol}(U(F)\backslash U(\mathbb{A})) = 1$ (see [22]). Writing $u = u(A)$ for $A \in M_2^{\text{sym}}(\mathbb{A})$ we then have

$$\Phi(x) = 1 \iff \psi(\text{Tr}(SA - M_x A)) = 1 \iff M_x = S.$$

Thus $\Phi(x)$ is an indicator function allowing only those $x \in X^2(F)$ with $M_x = S$ to contribute non-zero terms to the summation in $\mathcal{P}(\theta^0(f, \phi), \chi)$. Define

$$X_S^2 = \{x \in X^2 : M_x = S\}$$

so that

$$\mathcal{P}(\theta^0(f, \phi), \chi) = \int_{\mathbb{A} \times T(F)\backslash T(\mathbb{A})} \int_{H_1^0(F)\backslash H_1^0(\mathbb{A})} \sum_{x \in X_S^2(F)} \phi(h_g^{-1}h_0^{-1}xg)f(h_0 h_g) dh dg. \quad (5.7)$$

We are interested in decomposing the algebra $B \cong D \otimes E$ into its subalgebras, in particular the role played by the field $L \cong K \otimes E$. Hence we make the following observation.

Proposition 5.3. *If L does not embed into B as a subalgebra, then $X_S^2(F) = \emptyset$ and consequently*

$$\mathcal{P}(\theta^0(f, \phi), \chi) = 0.$$

Proof. Suppose $L \not\hookrightarrow B$ and assume the contrary: there exists $\xi \in X_S^2(F)$ with $\xi \neq 0$. Then ξ gives a realisation of W_1 as a quadratic subspace of X and we have $X = W_1 \oplus W_1^\perp$ as before. Since $E \cap X = F$, we have that $X \otimes E \cong B$, so we may decompose B as

$$B = (W_1 \otimes E) \oplus (W_1^\perp \otimes E).$$

But Lemma 5.4 gives us that $W_1 = Kw$ for any $w \in W_1$. Noting that $1 \in X$ we proceed by checking two cases: Firstly, if $1 \in W_1$, we may take $w = 1$ so that $W_1 = K$. Then $W_1 \otimes E = L$ and $B = L \oplus L^\perp$. Thus $L \hookrightarrow B$ as a quadratic subalgebra (over E), a contradiction. Secondly, if $1 \notin W_1$, then $J = W_1^\perp \otimes E$ is a field and subalgebra of B . In fact, this field has to be L : for any $j \in J^\perp = W_1 \otimes E$ we may write $J^\perp = Jj$ but $W_1 = Kw$ implies $Jj = Lw$ for any $w \in W_1 \subset J^\perp$. Taking $j = w$ gives $J = L$ and thus, once again, we have the contradiction $L \hookrightarrow B$. \square

Assumption 5.4. Without loss in generality we assume that $X_S^2(F) \neq \emptyset$.

Indeed it is clear from (5.6) that $X_S^2(F) = \emptyset$ implies $\mathcal{P}(\theta^0(f, \phi), \chi) = 0$. Under Assumption 5.4 we may conclude that, by Proposition 5.3, one has an algebra-embedding $L \hookrightarrow B$ and subsequently that $K \hookrightarrow D$ as a subalgebra too. Note that this assumption is truly on the choice of K (or equivalently d) since E has been fixed in advance.

We continue by expressing X_S^2 in terms of the group $\mathrm{SO}(X)$ acting on it, reconsidering points of X_S^2 via the isomorphism $X^2(F) \cong \mathrm{Hom}_F(W_1, X)$. Fix a base point $\xi \in X_S^2(F)$, to be considered as an F -homomorphism $\xi : W_1 \rightarrow X$ satisfying the properties:

- (1) ξ is injective (since the Gram matrix $M_\xi = S$ is invertible).
- (2) ξ is an isometry onto its image in X .

We briefly justify (2). Recall that (Section 2.2.2) W_1 is endowed with the quadratic form q_S ; a simple calculation shows that for $w \in W_1$ we have $q_S(w) = q_{M_\xi}(w) = N_X(\xi(w))$. Thus W_1 is identified with a quadratic subspace of X via ξ . (We abuse notation and call this subspace W_1 too.) Consider the orthogonal decomposition

$$X = W_1 \oplus W_1^\perp.$$

Lemma 5.4. *The image of W_1 in X is a one-dimensional K -vector space: for any $w \in W_1$ we have $W_1 = Kw$. In particular, there is an F -vector space isomorphism $W_1 \cong K$.*

Proof. Recall $M_\xi = S$ and fix

$$S = \begin{pmatrix} a & b/2 \\ b/2 & c \end{pmatrix} \in M_2^{\mathrm{sym}}(F)$$

so that $d = -4 \det S = b^2 - 4ac$. Fix a basis $\{e_1, e_2\}$ of W_1 and let $\xi_i = \xi(e_i)$ for $i = 1, 2$. We show that any two vectors in W_1 are linearly dependent over K . Note that the polynomial $p(X) = X^2 - bX + ac$ has the root $\xi_2 \xi_1^* = \frac{1}{2}(b - \sqrt{d})$. Multiplying each side by ξ_1 , and noting $a = N_X(\xi_1)$ by assumption, we see that

$$\xi_2 = \frac{1}{2a}(b - \sqrt{d})\xi_1 \in K\xi_1.$$

Since ξ is injective, ξ_1 and ξ_2 constitute a basis for $W_1 \subset X$ over F . Hence the K -span of any vector $w \in W_1$ is equal to W_1 as F -vector spaces. \square

We proceed by continuing to exploit the base point ξ . The group $\mathrm{SO}(X)$ acts transitively on $X_S^2(F)$ in which the stabiliser of ξ is $\mathrm{SO}(W_1^\perp)$ by construction. Then after some calculation the isomorphism

$$X_S^2(F) \cong \mathrm{SO}(W_1^\perp) \backslash \mathrm{SO}(X)$$

permits the following reformulation of (5.7):

$$\mathcal{P}(\theta^0(f, \phi), \chi) = \int_{\mathrm{SO}(W_1^\perp)(\mathbb{A}) \backslash \mathrm{SO}(X)(\mathbb{A})} \phi(h_0^{-1}\xi) \Lambda_\xi(R(h_0)f, \chi) dh_0,$$

by defining

$$\Lambda_\xi(f, \chi) = \int_{\mathbb{A} \times \mathrm{G}(\mathrm{SO}(W_1^\perp) \times \mathrm{SO}(W_1)) \backslash \mathrm{G}(\mathrm{SO}(W_1^\perp) \times \mathrm{SO}(W_1))(\mathbb{A})} \chi(g) f(y_g h_g) dy_g dg$$

where $h_g \in H^0(\mathbb{A})$ is any element such that

$$\lambda(h_g) = \lambda(g)$$

with the additional constraints that $h_g(\xi(v)) = \xi(g(v))$ for $v \in W_1$ and $h_g(w) = w$ when $w \in W_1^\perp$. The variable of integration (y_g, g) is an element of $\mathrm{G}(\mathrm{SO}(W_1^\perp) \times \mathrm{SO}(W_1))(\mathbb{A})$ whence $\lambda(y_g) = \lambda(g)$.

5.5 Exploiting exceptional isomorphisms

In this section we analyse the domain of $\Lambda_\xi(f, \chi)$ and apply the representation theory of H to rewrite this integral as a period of automorphic forms on $B^\times(\mathbb{A})$. By the decomposition $X = W_1 \oplus W_1^\perp$, we look to reinterpret the subgroup $\mathrm{G}(\mathrm{SO}(W_1^\perp) \times \mathrm{SO}(W_1)) \leq \mathrm{GSO}(X)$ (featured in $\Lambda_\xi(f, \chi)$) as a subgroup of $F^\times \times B^\times / \Delta E^\times$ via the isomorphism ρ of (4.2).

5.5.1 Structural decomposition of quadratic spaces

Since $K \hookrightarrow D$, the standard involution $*$ on D restricts to the non-trivial Galois automorphism of K . We may write

$$D = K \oplus Kj$$

for any $j \in K^\perp$ since for such a j we have $K^\perp = Kj$. Extending this decomposition to $B \cong D \otimes E$ (where $*$ extends to a Galois action on B , trivial on E , as in Section 4.1) define

$$L = K(E) \cong K \otimes_F E.$$

Then $L = E(\sqrt{d})$ is a quadratic extension of E such that we have an embedding $L \hookrightarrow B$. The standard involution on B (given by $x \mapsto x^*$) restricts to the non-trivial Galois involution on L . Then, for the same $j \in K^\perp$ as before, we have $B = L \oplus Lj$.

Focusing now on the subspace $X \subset B$ define

$$X_L = \{x \in L : \iota(x) = x^*\}.$$

Both $X_L \subset X$ and $1 \in X_L$. Moreover, we may realise X_L as a quadratic extension of F . Under the quadratic form N_X we have the orthogonal decomposition $X = X_L \oplus X_L^\perp$ which is described by the following lemma.

Lemma 5.5. *For any $z_0 \in E$ with $\mathrm{Tr}_{E/F}(z_0) = 0$ we have the orthogonal decomposition*

$$X = X_L \oplus z_0 Kj.$$

Proof. The orthogonal complement X_L^\perp is given by $X \cap Lj$ (otherwise $X_L \cap X_L^\perp \neq 0$). Hence X_L^\perp contains elements xj where $x \in L$ such that $\iota(xj) = (xj)^*$; these are the elements $x \in L$ such that $x + \iota(x) = 0$ since j and x are orthogonal under N_X . Fix some $z_0 \in E$ with $\mathrm{Tr}_{E/F}(z_0) = 0$ then for any $k \in K$ we have $\iota(z_0 k) = -z_0 k$. Hence $z_0 Kj \subseteq X_L^\perp$, and since both are two-dimensional F -vector spaces we have equality. \square

Lemma 5.4 gave us an interpretation of $W_1 \subset X$ as the space $W_1 \cong K$. Combining this with Lemma 5.5 allows one to deduce the following (F -vector space) isomorphisms:

$$W_1 \cong X_L^\perp \quad \text{and} \quad W_1^\perp \cong X_L.$$

Consequently, we have the reinterpretation of the orthogonal groups

$$\mathrm{GSO}(X_L^\perp) \cong \mathrm{GSO}(W_1) \cong K^\times \quad \text{and} \quad \mathrm{GSO}(X_L) \cong \mathrm{GSO}(W_1^\perp),$$

justifying our conclusion that

$$G(\mathrm{SO}(X_L) \times \mathrm{SO}(X_L^\perp)) \cong G(\mathrm{SO}(W_1^\perp) \times \mathrm{SO}(W_1)). \quad (5.8)$$

Proposition 5.6. *There is an F -isomorphism of algebraic groups*

$$\Phi : F^\times \times L^\times / \Delta E^\times \xrightarrow{\sim} G(\mathrm{SO}(W_1^\perp) \times \mathrm{SO}(W_1))$$

where the projection onto the second component is given by $(s, k) \mapsto s^{-1}kk^t \in K^\times$ (whereby K^\times acts on $W_1 \cong K$ by left multiplication).

Proof. By the isomorphism (5.8), it suffices to find an isomorphism Φ such that the following diagram commutes:

$$\begin{array}{ccc} F^\times \times B^\times / \Delta E^\times & \xrightarrow{\rho} & \mathrm{GSO}(X) = \mathrm{GSO}(X_L \oplus X_L^\perp) \\ \cup & & \cup \\ F^\times \times L^\times / \Delta E^\times & \xrightarrow{\Phi} & G(\mathrm{SO}(X_L) \times \mathrm{SO}(X_L^\perp)). \end{array} \quad (5.9)$$

We consider the surjective map

$$\Phi : F^\times \times L^\times \rightarrow \{(s^{-1}k\iota(k)^*, s^{-1}k\iota(k)) : s \in F^\times, k \in L^\times\}.$$

One can check that the projections of $\mathrm{Im}(\Phi)$, onto the first and second components, act on X_L and X_L^\perp , respectively, by left multiplication. Noting that the similitude factors of each component in the image are equal, hence we may extend Φ to a mapping into $G(\mathrm{SO}(X_L) \times \mathrm{SO}(X_L^\perp))$. Since the kernel of Φ is ΔE^\times , we have an injection

$$\Phi : F^\times \times L^\times / \Delta E^\times \hookrightarrow G(\mathrm{SO}(X_L) \times \mathrm{SO}(X_L^\perp)).$$

To demonstrate the surjectivity of Φ we need only check that diagram (5.9) commutes. Observe that, for $(s, k) \in F^\times \times L^\times$,

$$\rho(s, k)(X_L \oplus X_L^\perp) = s^{-1}k(X_L \oplus X_L^\perp)\iota(k)^* = s^{-1}k\iota(k)^*X_L \oplus s^{-1}k\iota(k)X_L^\perp = \Phi(s, k)(X_L \oplus X_L^\perp).$$

Thus meaning that, up to an automorphism of $X_L \oplus X_L^\perp$, $\rho|_{F^\times \times L^\times} = \Phi$. Since ρ is one-to-one then Φ must also be surjective. \square

5.5.2 Interpretation of the integral $\Lambda_\xi(f, \chi)$

Considering the domain of $\Lambda_\xi(f, \chi)$, one uses Proposition 5.6 to deduce the isomorphism

$$\mathbb{A}^\times G(\mathrm{SO}(W_1^\perp) \times \mathrm{SO}(W_1)) \backslash G(\mathrm{SO}(W_1^\perp) \times \mathrm{SO}(W_1))(\mathbb{A}) \cong \mathbb{A}_E^\times L^\times \backslash \mathbb{A}_L^\times.$$

The application of this isomorphism to $\Lambda_\xi(f, \chi)$ requires a change of integration variable. This is accomplished by substituting $(y_g, g) \mapsto \rho(1, k)$ where $k \in \mathbb{A}_E^\times L^\times \backslash \mathbb{A}_L^\times$. For this we note that the original variables $h_g \in \mathrm{GSO}(W_1)(\mathbb{A})$ and $(y_g, g) \in G(\mathrm{SO}(W_1^\perp) \times \mathrm{SO}(W_1))(\mathbb{A})$ satisfy:

- $h_g \in \mathrm{GSO}(W_1)(\mathbb{A})$ fixes $W_1^\perp(\mathbb{A})$ and acts as g on $W_1(\mathbb{A})$,
- $y_g \in \mathrm{GSO}(W_1^\perp)(\mathbb{A})$ fixes $W_1(\mathbb{A})$ and acts as y_g on $W_1^\perp(\mathbb{A})$,
- $\lambda(h_g) = \lambda(y_g)$.

Hence the product $y_g h_g$, corresponding to (y_g, g) , is substituted with $\rho(1, k)$ and element $g \in \mathrm{GSO}(W_1)(\mathbb{A})$, the projection of (y_g, g) onto its second factor, is substituted with kk^t (as in Proposition 5.6). This substitution returns

$$\Lambda_\xi(f, \chi) = \int_{\mathbb{A}_E^\times L^\times \backslash \mathbb{A}_L^\times} \chi(kk^t) f(\rho(1, k)) dk.$$

For any $k \in \mathbb{A}_L^\times$ we have $kk^t \in \mathbb{A}_K^\times$ so we have a character $\Omega : L^\times \backslash \mathbb{A}_L^\times \rightarrow \mathbb{C}^\times$ by defining

$$\Omega(k) = \chi(kk^t) \quad (5.10)$$

such that upon restricting Ω to \mathbb{A}_E^\times we have $\Omega|_{\mathbb{A}_E^\times} = \chi \circ N_{E/F}$. Since we have chosen $f \in \mathcal{V}_{\sigma,S}^1$ to correspond to some $\eta \in \mathcal{V}_\tau$ such that $f|_{H^0(\mathbb{A})}(\rho(s, a)) = \eta(a)$, the integral above becomes

$$\Lambda_\xi(f, \chi) = \int_{\mathbb{A}_E^\times L^\times \backslash \mathbb{A}_L^\times} \Omega(k) \eta(k) dk. \quad (5.11)$$

6 Local calculation: Integrals over matrix coefficients

We will ultimately show that $|\mathcal{P}(\theta(f, \phi), \chi)|^2$ factorises into a product of special L -values and a finite number of local integrals. In this section we follow [14] in defining these local integrals and make use of the excellent results proved by Liu to rearrange them for our purposes. Throughout this section we work locally at a place v of F suppressing the subscript v from the notation (so that $F = F_v$, σ denotes one local component in the tensor product $\otimes_v \sigma_v$ and so on).

6.1 Local integrals

To provide a complete picture, we define the local integrals in full generality for any (local) irreducible, admissible representation π of G . The definition is divided into a non-archimedean and an archimedean case; this is due to the nature of the analysis in [14, Section 3] in ‘regularising’ these integrals. Immediately after this definition we specialise to choosing $\pi = \theta(\sigma)$, the (local) theta lift of σ , and unify the integrals from each case since they have the same form in this specialisation. We point out that such a $\pi = \theta(\sigma)$ is always tempered and thus the regularisation results of [14] apply.

6.1.1 The non-archimedean case

Suppose that F is a non-archimedean local field. We consider the notion of a stable integral as defined in [13]. We refer the reader to there for more information since it is not of central importance to our discussion.

Definition 6.1 (The non-archimedean local factors). Given $\varphi \in \mathcal{V}_\pi$, $\tilde{\varphi} \in \mathcal{V}_{\tilde{\pi}}$ and a unitary paring

$$\mathcal{B}_\pi : \mathcal{V}_\pi \otimes \mathcal{V}_{\tilde{\pi}} \rightarrow \mathbb{C}$$

we define

$$\alpha(\varphi, \tilde{\varphi}; \chi) = \int_{F^\times \backslash T} \int_U^{\text{st}} \mathcal{B}_\pi(\pi(ug)\varphi, \tilde{\varphi}) \chi(g) \psi_S^{-1}(u) du dg$$

where the integral over U is called a *stable integral* (see [13, Definition 2.1]) and is evaluated on a certain compact open subgroup $N \subset U$. This N is chosen to be ‘maximally’ in the sense that if N' is another compact open subgroup with $N \subset N' \subset U$, then the integral over N' equals the integral over N . The product of Haar measures $du dg$ is again a Haar measure on the Bessel subgroup $F^\times \backslash R$.

Indeed it is not obvious that the integrals of Definition 6.1 converge, nor should such an N exist, but Liu proves these facts in [14, Theorem 2.1] and [14, Lemma 3.2], respectively.

6.1.2 The archimedean case

Let F be an archimedean local field. The method of regularisation here is to consider the Fourier transform of certain matrix coefficients in a so-called *regular* subset of U .

Recall that the abelian unipotent group $U \cong M_2^{\text{sym}}(F)$ is self-dual and all its characters are given by ψ_M , for some $M \in M_2^{\text{sym}}(F)$, as in (2.4). We denote by $M_2^{\text{sym}}(F)^{\text{reg}}$ the open and dense subset of non-singular symmetric matrices in $M_2^{\text{sym}}(F)$ and define its image in U as

$$U^{\text{reg}} \cong M_2^{\text{sym}}(F)^{\text{reg}}.$$

Definition 6.2 (The archimedean local factors). Given $\varphi \in \mathcal{V}_\pi$, $\tilde{\varphi} \in \mathcal{V}_{\bar{\pi}}$ and a unitary paring

$$\mathcal{B}_\pi : \mathcal{V}_\pi \otimes \mathcal{V}_{\bar{\pi}} \rightarrow \mathbb{C}$$

we define

$$\alpha(\varphi, \tilde{\varphi}; \chi) = \int_{F^\times \setminus T} \int_{U^{\text{reg}}} \mathcal{B}_\pi(\pi(ug)\varphi, \tilde{\varphi})\chi(g)\psi_S^{-1}(u) du dg.$$

Here, for a fixed $g \in T$, the map

$$\psi_S \mapsto \int_{U^{\text{reg}}} \mathcal{B}_\pi(\pi(ug)\varphi, \tilde{\varphi})\psi_S^{-1}(u) du$$

is the Fourier transform (in U^{reg}) of the function $u \mapsto \mathcal{B}_\pi(\pi(ug)\theta(f, \phi), \theta(\tilde{f}, \tilde{\phi}))$.

Once again, Liu proves that this integral converges absolutely in [14, Theorem 2.1].

6.1.3 Normalisation of local integrals

In his paper [14], Liu goes on to show that there exists a specified set of *good* places, which exclude a *finite* number of places of the base number field (including the archimedean ones), for which the local integrals may be computed as follows (see [14, p. 7] for details).

Proposition 6.1. *If v is a good place of the base number field, then for the local vectors $\varphi \in \mathcal{V}_\pi$, $\tilde{\varphi} \in \mathcal{V}_{\bar{\pi}}$ one has*

$$\alpha(\varphi, \tilde{\varphi}; \chi) = \frac{\zeta_F(2)\zeta_F(4)L(1/2, \pi \boxtimes \chi)}{L(1, \pi, \text{Ad})L(1, \chi_{K/F})}.$$

Hence we normalise the local factors by setting

$$\alpha^{\natural}(\varphi, \tilde{\varphi}; \chi) = \frac{L(1, \pi, \text{Ad})L(1, \chi_{K/F})}{\zeta_F(2)\zeta_F(4)L(1/2, \pi \boxtimes \chi)} \alpha(\varphi, \tilde{\varphi}; \chi) \quad (6.1)$$

so that $\alpha^{\natural}(\varphi, \tilde{\varphi}; \chi) = 1$ for almost all v .

Given *any* place v , if, instead of considering an arbitrary vector $\tilde{\varphi} \in \mathcal{V}_{\bar{\pi}}$, we take the local vector $\tilde{\varphi} = \bar{\varphi}$ – in the context of being local factors of functions on adèle groups as in (5.3) – then we define the notation

$$\alpha(\varphi, \chi) = \alpha(\varphi, \bar{\varphi}; \chi) \quad \text{and} \quad \alpha^{\natural}(\varphi, \chi) = \alpha^{\natural}(\varphi, \bar{\varphi}; \chi). \quad (6.2)$$

As well as absolute convergence, [14, Theorem 2.1] states that whenever such a π is tempered, we have the positivity result

$$\alpha(\varphi, \chi) \geq 0.$$

Remark 6.2. The integrals defining $\alpha(\varphi, \tilde{\varphi}; \chi)$ have a unipotent part (over U) which is given by either a stable integral (over a compact open $N \subset U$) or a Fourier transform (with respect to $U^{\text{reg}} \subset U$) when v is non-archimedean or archimedean, respectively. We consider these integrals for π tempered. The choices of regularisation for these integrals are justified by noting that when π is *square integrable* we may take the entire space U in each definition. That is, for *any* v , when π is square integrable we have

$$\alpha(\varphi, \tilde{\varphi}; \chi) = \int_{F^\times \setminus T} \int_U \mathcal{B}_\pi(\pi(ug)\varphi, \tilde{\varphi})\chi(g)\psi_S^{-1}(u) du dg,$$

by [14, Propositions 3.5 and 3.15].

6.1.4 A unified result for theta lifts

Let us specialise now by assuming $\pi = \theta(\sigma)$ is the theta lift of σ , a local factor of the fixed representation in Section 5.1. We select the pairing \mathcal{B}_π to be defined as in (4.8); this depends on a choice of \mathcal{B}_σ which we made in (5.2). Retaining some generality in what follows, we note that by [7, Proposition 5.5] the conjugate representation $\tilde{\pi}$ is generated by elements $\theta(\tilde{f}, \tilde{\phi})$ for $\tilde{f} \in \mathcal{V}_{\tilde{\sigma}}$ and $\tilde{\phi} \in \mathcal{V}_{\tilde{\omega}}$.

Proposition 6.3. *In either the non-archimedean or archimedean cases, if $\theta(f, \phi) \in \mathcal{V}_\pi$ and $\theta(\tilde{f}, \tilde{\phi}) \in \mathcal{V}_{\tilde{\pi}}$ then the local integrals become*

$$\begin{aligned} & \alpha(\theta(f, \phi), \theta(\tilde{f}, \tilde{\phi}); \chi) \\ &= \frac{\zeta_F(2)\zeta_F(4)}{L(1, \sigma, \text{std})} \int_{F^\times \backslash \text{GSO}(W_1)} \int_{\text{O}(X)} \int_{\text{SO}(W_1^+) \backslash \text{SO}(X)} \phi(h_g^{-1}h^{-1}h_1^{-1}\xi g)\tilde{\phi}(h_1^{-1}\xi)\mathcal{B}_\sigma(\sigma(hh_g)f, \tilde{f}) dh_1 dh dg \end{aligned}$$

where $h_g \in H^0(\mathbb{A})$ is any element such that $\lambda(h_g) = \lambda(g)$ with the additional constraints that $h_g(\xi(v)) = \xi(g(v))$ for $v \in W_1$ and $h_g(w) = w$ when $w \in W_1^\perp$ (for comparison see Section 5.4); the element $\xi \in X_S^2$ is the base point chosen in Section 5.4; dh is the Haar measure for $\text{O}(X)$ fixed in the definition for $\mathcal{B}_{\theta(\sigma)}$, see (4.8); and finally dh_1 is the Siegel–Weil measure on $\text{SO}(W_1^+) \backslash \text{SO}(X)$.

Proof. This follows immediately from [14, Lemma 4.2]. □

Remark 6.4. The product of local Siegel–Weil measures is precisely the Tamagawa measure on the adèlic points of the group in question (see [14, Remark 3.18]).

6.2 Explicit local factors for theta lifts

We analyse the terms $\alpha^{\natural}(\theta(f, \phi), \theta(\tilde{f}, \tilde{\phi}); \chi)$ where $\theta(f, \phi) \in \mathcal{V}_\pi$ and $\theta(\tilde{f}, \tilde{\phi}) \in \mathcal{V}_{\tilde{\pi}}$ are as before. We point out again that, even though the subscripts are removed, everything is local here. We will determine the quantity

$$\begin{aligned} & \left(\frac{\zeta_F(2)\zeta_F(4)}{L(1, \sigma, \text{std})} \right)^{-1} \alpha(\theta(f, \phi), \theta(\tilde{f}, \tilde{\phi}); \chi) \\ &= \int_{F^\times \backslash \text{GSO}(W_1)} \int_{\text{O}(X)} \int_{\text{SO}(W_1^+) \backslash \text{SO}(X)} (\omega(h)\phi)(\xi)(\tilde{\omega}(h_1)\tilde{\phi})(\xi)\mathcal{B}_\sigma(\sigma(h_1^{-1}h_g h)f, \tilde{f})\chi(g) dh_1 dh dg \end{aligned}$$

after making the substitution $h \mapsto h_1^{-1}h_g h h_g^{-1}$ and recalling that $\xi g = h_g \xi$, by definition. We decompose the integral over $\text{O}(X)$ in terms of its connected component $\text{SO}(X)$ and replace the measure dh with

$$dh_2 = 2dh|_{\text{SO}(X)}$$

so that the volumes

$$\text{Vol}(\text{O}(X), dh) = \text{Vol}(\text{SO}(X), dh_2).$$

Then we find that the above quantity is equal to

$$\frac{1}{2} \sum_{\varepsilon \in \mu_2(F)} \int_{F^\times \backslash \text{GSO}(W_1)} \int_{\text{SO}(X)} \int_{\text{SO}(W_1^+) \backslash \text{SO}(X)} (\omega(h_2\varepsilon)\phi)(\xi)(\tilde{\omega}(h_1)\tilde{\phi})(\xi)\mathcal{B}_\sigma(\sigma(h_1^{-1}h_g h_2\varepsilon)f, \tilde{f})\chi(g) dh_1 dh_2 dg.$$

To simplify further, note that

$$\text{SO}(X) \cong (\text{SO}(W_1^+) \backslash \text{SO}(X)) \times \text{SO}(W_1^\perp)$$

where we substitute $h_2 \mapsto (h_2, y)$, with measure $dh_2 \mapsto dh_2 dy$, so that

$$\begin{aligned} & \left(\frac{\zeta_F(2)\zeta_F(4)}{L(1, \sigma, \text{std})} \right)^{-1} \alpha(\theta(f, \phi), \theta(\tilde{f}, \tilde{\phi}); \chi) \\ &= \frac{1}{2} \sum_{\varepsilon \in \mu_2(F)} \int_{F^\times \backslash \text{GSO}(W_1)} \int_{\text{SO}(W_1^+) \backslash \text{SO}(X)} \int_{(\text{SO}(W_1^+) \backslash \text{SO}(X))^2} (\omega(h_2\varepsilon)\phi)(\xi)(\tilde{\omega}(h_1)\tilde{\phi})(\xi)\mathcal{B}_\sigma(\sigma(h_1^{-1}yh_g h_2\varepsilon)f, \tilde{f})\chi(g) dh_1 dh_2 dy dg, \end{aligned}$$

recalling that $y \in \mathrm{SO}(W_1^\perp)$ stabilises ξ and commutes with h_g . Using that σ is unitary under \mathcal{B}_σ we finally obtain

$$\begin{aligned} & \alpha(\theta(f, \phi), \theta(\tilde{f}, \tilde{\phi}); \chi) \\ &= \frac{1}{2} \frac{\zeta_F(2)\zeta_F(4)}{L(1, \sigma, \mathrm{std})} \sum_{\varepsilon \in \mu_2(F)} \int_{(\mathrm{SO}(W_1^\perp) \backslash \mathrm{SO}(X))^2} (\omega(h_2\varepsilon)\phi)(\xi)(\tilde{\omega}(h_1)\tilde{\phi})(\xi) \Gamma_{\xi, \nu}(\sigma(h_2\varepsilon)f, \bar{\sigma}(h_1)\tilde{f}; \chi) dh_1 dh_2 \end{aligned}$$

by defining

$$\Gamma_{\xi, \nu}(f, \tilde{f}; \chi) = \int_{F^\times \backslash \mathrm{GSO}(W_1)} \int_{\mathrm{SO}(W_1^\perp)} \mathcal{B}_\sigma(\sigma(yh_g)f, \tilde{f})\chi(g) dy dg.$$

7 The result: Local and global assembly

This section concludes with the unification of the global period in Section 5 and the rearranged local integrals in Section 6. The connection is facilitated by the work of Waldspurger [21] who, in 1985, gave the pioneering example of refined Gan–Gross–Prasad conjecture: a proof for the pair $(\mathrm{SO}_3, \mathrm{SO}_2)$. We apply his formula to our calculation.

7.1 A theorem of Waldspurger

Let B be a (possibly split) quaternion algebra over E . Let L be a quadratic extension of a number field E such that there exists an embedding $L \hookrightarrow B$ and let Ω be a Hecke character of \mathbb{A}_L^\times . Let $\tau = \otimes_w \tau_w$ be an irreducible, cuspidal automorphic representation of $B^\times(\mathbb{A}_E)$, realised in \mathcal{V}_τ , such that $\omega_\tau \cdot \Omega|_{\mathbb{A}_E^\times} = 1$. For $\eta \in \mathcal{V}_\tau$ define the global period integral

$$\mathcal{Q}(\eta, \Omega) = \int_{\mathbb{A}_E^\times \backslash L^\times \backslash \mathbb{A}_L^\times} \Omega(k)\eta(k) dk.$$

For each place w of E let \mathcal{B}_{τ_w} be a unitary pairing on $\mathcal{V}_{\tau_w} \otimes \mathcal{V}_{\bar{\tau}_w}$. For each $\eta_w \in \mathcal{V}_{\tau_w}$ and $\tilde{\eta}_w \in \mathcal{V}_{\bar{\tau}_w}$ define the local integrals

$$\beta_w(\eta_w, \tilde{\eta}_w; \Omega_w) = \int_{E_w^\times \backslash L_w^\times} \mathcal{B}_{\tau_w}(\tau_w(k_w)\eta_w, \tilde{\eta}_w)\Omega_w(k_w) dk_w$$

and their natural normalisation,

$$\beta_w^h(\eta_w, \tilde{\eta}_w; \Omega_w) = \frac{L(1, \tau_w, \mathrm{Ad})L(1, \chi_{L_w/E_w})}{\zeta_{E_w}(2)L(1/2, \tau_{L,w} \otimes \Omega_w)} \beta_w(\eta_w, \tilde{\eta}_w; \Omega_w)$$

where $\tau_{L,w}$ is the base change lift of τ_w to $B^\times(L_w)$.

The following theorem was originally given in [21, Section III.3] (and then stated in terms of the refined Gan–Gross–Prasad conjecture in [12, Section 6]). Fix a choice of Haar measures dk_w such that the Tamagawa measure on $(E^\times \backslash L^\times)(\mathbb{A})$ decomposes as $dk = \prod_w dk_w$, and a choice of local pairings \mathcal{B}_{τ_w} such that the Petersson inner product decomposes as $\mathcal{B}_\tau = \prod_w \mathcal{B}_{\tau_w}$.

Theorem 7.1 (Waldspurger). *The integrals $\beta_w(\eta_w, \tilde{\eta}_w; \Omega_w)$ are absolutely convergent and*

$$\beta_w^h(\eta_w, \tilde{\eta}_w; \Omega_w) = 1$$

for almost all places w of E . If, in addition, τ has trivial central character, $\omega_\tau = 1$, and Ω is unitary, then

$$\mathcal{Q}(\eta, \Omega)\mathcal{Q}(\tilde{\eta}, \bar{\Omega}) = \frac{1}{2} \frac{\zeta_E(2)L(1/2, \tau'_L \otimes \Omega)}{L(1, \tau, \mathrm{Ad})L(1, \chi_{L/E})} \prod_w \beta_w^h(\eta_w, \tilde{\eta}_w; \Omega_w)$$

where τ_L denotes the base change lift of τ to $B^\times(\mathbb{A}_L)$ and τ'_L is the Jacquet–Langlands transfer of τ_L to $\mathrm{GL}_2(\mathbb{A}_L)$.

We remark that the L -function $L(1/2, \tau'_L \otimes \Omega)$ may be interpreted in various ways due to the low-dimensional isomorphisms that occur (see Section 2.1.3).

7.2 Application of Waldspurger

Let the arbitrary notation introduced in Section 7.1 now assume the running meanings that we assigned in Section 5.1 (for the representation $\tau = \otimes_w \tau_w$ and the pairings \mathcal{B}_{τ_w}) and Section 5.5 (for the algebras $B \cong D \otimes E$, $L \cong K \otimes E$). We draw special attention to the assumption that $f \in \mathcal{V}_{\sigma, S}^1$ with $f|_{H^0(\mathbb{A})} \circ \rho = \eta$. The set \mathfrak{S} contains those places of F such that $\sigma_v \cong \sigma_v \otimes \text{sgn}$ and S is the fixed, finite set of places of F outside which $f_v = f_v^\circ$ is $H(\mathcal{O}_v)$ -invariant (see Section 4.3.3). We choose $\eta = \otimes_w \eta_w$, implying $f = \otimes_v f_v$ with $f_v = \otimes_{w|v} \eta_w$ as in Section 4.3.2. The pairings \mathcal{B}_{τ_w} , for $w|v$, determine the pairings $\mathcal{B}_{\sigma_{0,v}}$ and \mathcal{B}_{σ_v} (as in Section 4.4) which are used to define the local integrals (Section 6).

Lemma 7.2. *The global period integral in Waldspurger’s formula satisfies*

$$\Lambda_\xi(f, \chi) = \Omega(\eta, \Omega).$$

Proof. We only need to remark that $\Omega|_{\mathbb{A}_E^\times} = \chi \circ N_{E/F}$ implying the condition $\Omega|_{\mathbb{A}_E^\times} = 1$ is satisfied since $\chi|_{\mathbb{A}^\times} = 1$ (Assumption 5.2). Moreover, Ω is unitary because χ is assumed so. We then have that the form of $\Lambda_\xi(f, \chi)$ in (5.11) is given precisely by $\Omega(\eta, \Omega)$. \square

In a similar manner, we identify the local period integrals in Waldspurger’s formula with our own terms $\Gamma_{\xi, v}(f_v, \tilde{f}_v; \chi_v)$. The following lemma is a local analogue of the analysis of $\Lambda_\xi(f, \chi)$ in Section 5.5.

Lemma 7.3. *Let v be a place of F . Then, for $f_v \in \mathcal{V}_{\sigma_v}$ and $\tilde{f}_v \in \mathcal{V}_{\tilde{\sigma}_v}$ as above,*

$$\Gamma_{\xi, v}(f_v, \tilde{f}_v; \chi_v) = \frac{1}{2^{c_v}} \prod_{w|v} \beta_w(\eta_w, \tilde{\eta}_w; \Omega_w)$$

where

$$c_v = \begin{cases} 1 & \text{if } v \in \mathfrak{S} \cap S, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Analogous to the global setting (discussed in Section 5.5.2) we have

$$F_v^\times \backslash \text{GSO}(W_1)_v \times \text{SO}(W_1^\perp)_v \cong F_v^\times \backslash \text{G}(\text{SO}(W_1^\perp) \times \text{SO}(W_1))_v$$

so that

$$\Gamma_{\xi, v}(f_v, \tilde{f}_v; \chi_v) = \int_{F_v^\times \backslash \text{G}(\text{SO}(W_1^\perp) \times \text{SO}(W_1))_v} \mathcal{B}_{\sigma_v}(\sigma_v(y h_g) f_v, \tilde{f}_v) \chi_v(g) dy_g dg \quad (7.1)$$

where $h_g \in H_v^0$ is any element such that

$$\lambda(h_g) = \lambda(g)$$

with the additional constraints that $h_g(\xi(v)) = \xi(g(v))$ for $v \in W_{1,v}$ and $h_g(w) = w$ when $w \in W_{1,v}^\perp$. The variable of integration (y_g, g) is an element of $\text{G}(\text{SO}(W_1^\perp) \times \text{SO}(W_1))_v$ whence $\lambda(y_g) = \lambda(g)$. By Proposition 5.6 there is an F_v -isomorphism

$$F_v^\times \backslash \text{G}(\text{SO}(W_1^\perp) \times \text{SO}(W_1))_v \cong (E^\times \backslash L^\times)(F_v).$$

Applying this isomorphism to (7.1) (checking Section 5.5.2 for comparison), we substitute the element $y_g h_g$, which corresponds to (y_g, g) by definition, with $\rho(1, k)$ where $k \in (E^\times \backslash L^\times)(F_v)$. The element $g \in \text{GSO}(W_1)_v$ is the projection of (y_g, g) onto its second factor; as in Proposition 5.6, this projection corresponds to $\rho(1, k) \mapsto k k^t$. This substitution returns

$$\Gamma_{\xi, v}(f_v, \tilde{f}_v; \chi_v) = \int_{(E^\times \backslash L^\times)(F_v)} \mathcal{B}_{\sigma_v}(\sigma_v(\rho(1, k_v)) f_v, \tilde{f}_v) \chi_v(k_v k_v^t) dk_v.$$

The automorphic character $\Omega = \otimes_w \Omega_w$ of (5.10), factorised over places of E , may be divided into factors corresponding to each place v of F by $\Omega_v = \otimes_{w|v} \Omega_w$. These factors coincide with the factorisation of $\chi = \otimes_v \chi_v$ in that $\Omega_v : k_v \mapsto \chi_v(k_v k_v^t)$.

The measures dk_v are chosen so that the Tamagawa measure dk on $(E^\times \backslash L^\times)(\mathbb{A}_E)$ factorises as

$$dk = \prod_w dk_w,$$

over places of E , with $dk_v = \prod_{w|v} dk_w$. The dk_v are precisely the measures $dh_{1,v}$ of $H_{1,v}$ in (4.8) (defining $\mathcal{B}_{\theta(\sigma_v)}$). We now express the domain in terms of places w of E . By (4.5) we have

$$(E^\times \backslash L^\times)(F_v) \cong \prod_{w|v} E_w^\times \backslash L_w^\times.$$

Our calculation now depends on whether or not $v \in \mathfrak{S}$. With the vectors $f_v = \otimes_{w|v} \eta_w$ and $\tilde{f}_v = \otimes_{w|v} \tilde{\eta}_w$ we have

$$\mathcal{B}_{\sigma_v}(f_v, \tilde{f}_v) = \frac{1}{2^{c_v}} \mathcal{B}_{\sigma_{0,v}}(f_v, \tilde{f}_v) = \frac{1}{2^{c_v}} \prod_{w|v} \mathcal{B}_{\tau_w}(\eta_w, \tilde{\eta}_w).$$

This is clear from the definition of the pairing \mathcal{B}_{σ_v} in Section 4.4 if $v \notin S$ or $v \in \mathfrak{S}$. If $v \in S \cap \mathfrak{S}$, then

$$f_v = f_v + 0 \in \mathcal{V}_{\sigma_{0,v}} \oplus \mathcal{V}_{\sigma'_{0,v}}$$

so we pick up the factor of $1/2^{c_v} = 1/2$.

At last we obtain

$$\begin{aligned} \Gamma_{\xi,v}(f_v, \tilde{f}_v; \chi_v) &= \int_{\prod_{w|v} E_w^\times \backslash L_w^\times} \frac{1}{2^{c_v}} \prod_{w|v} \mathcal{B}_{\tau_w}(\tau_w(k_w)\eta_w, \tilde{\eta}_w)\Omega_w(k_w) dk_w \\ &= \frac{1}{2^{c_v}} \prod_{w|v} \beta_w(\eta_w, \tilde{\eta}_w; \Omega_w). \end{aligned} \quad \square$$

Combining the previous two lemmas allows Waldspurger’s formula to be rewritten in terms of the integrals defining Λ_ξ and Γ_ξ . Recall the notation $S' = S \setminus (S \cap \mathfrak{S})$ and introduce

$$s = |S \cap \mathfrak{S}| \quad \text{and} \quad s' = |S'|.$$

Proposition 7.4. *For all pure tensors $f = \otimes_v f_v \in \mathcal{V}_{\sigma,S}^1$ and $\tilde{f} = \otimes_v \tilde{f}_v \in \mathcal{V}_{\bar{\sigma},S}^1$ we have*

$$\Lambda_\xi(f, \chi)\Lambda_\xi(\tilde{f}, \bar{\chi}) = 2^{s-1} \prod_v \Gamma_{\xi,v}(f_v, \tilde{f}_v; \chi_v).$$

7.3 The explicit formula

Applying the definition of the variant theta integral (5.5) we begin computing the Bessel period’s square:

$$|\mathcal{P}(\theta(f, \phi), \chi)|^2 = \int_{\mu_2(F) \backslash \mu_2(\mathbb{A})} \int_{\mu_2(F) \backslash \mu_2(\mathbb{A})} \mathcal{P}(\theta^0(\sigma(\delta)f, \boldsymbol{\omega}(\delta)\phi), \chi) \overline{\mathcal{P}(\theta^0(\sigma(\varepsilon)f, \boldsymbol{\omega}(\varepsilon)\phi), \chi)} d\delta d\varepsilon.$$

As $\mu_2(F)$ is of index two in $\mu_2(\mathbb{A})$, we rearrange so that the above integral equals

$$\begin{aligned} &\frac{1}{4} \int_{\mu_2(\mathbb{A})} \int_{\mu_2(\mathbb{A})} \mathcal{P}(\theta^0(\sigma(\delta)f, \boldsymbol{\omega}(\delta)\phi), \chi) \overline{\mathcal{P}(\theta^0(\sigma(\varepsilon)f, \boldsymbol{\omega}(\varepsilon)\phi), \chi)} d\delta d\varepsilon \\ &= \frac{1}{4^{1+s+s'}} \sum_{\mu_2(F_S)} \sum_{\mu_2(F_S)} \mathcal{P}(\theta^0(\sigma(\delta)f, \boldsymbol{\omega}(\delta)\phi), \chi) \overline{\mathcal{P}(\theta^0(\sigma(\varepsilon)f, \boldsymbol{\omega}(\varepsilon)\phi), \chi)}. \end{aligned} \quad (7.2)$$

This equality follows since, as $\varepsilon_v \in H(\mathcal{O}_v)$, the integrals for $v \notin S$ fix the integrand and elsewhere we have the (normalised) counting Haar measure. We further reduce the sum by noting that, for $h_0 \in H^0(\mathbb{A})$,

$$\sigma(\varepsilon)f(h_0) = f(h_0\varepsilon) = 0$$

unless $\varepsilon \in \mu_2(\mathbb{A}^{S \cap \mathfrak{S}})\{1, \iota\}$ (by (4.7) or [7, Lemma 2.2]). Hence (7.2) equals

$$\frac{1}{4^{1+s+s'}} \sum_{\mu_2(F_{S'})} \sum_{\mu_2(F_{S'})} \left(\mathcal{P}(\theta^0(\sigma(\delta)f, \boldsymbol{\omega}(\delta)\phi), \chi) + \mathcal{P}(\theta^0(\sigma(\delta\iota)f, \boldsymbol{\omega}(\delta\iota)\phi), \chi) \right) \\ \times \left(\overline{\mathcal{P}(\theta^0(\sigma(\varepsilon)f, \boldsymbol{\omega}(\varepsilon)\phi), \chi)} + \overline{\mathcal{P}(\theta^0(\sigma(\varepsilon\iota)f, \boldsymbol{\omega}(\varepsilon\iota)\phi), \chi)} \right).$$

The invariance under ι , noted in (5.4), implies we have the equality

$$|\mathcal{P}(\theta(f, \phi), \chi)|^2 = \frac{1}{4^{s+s'}} \sum_{\mu_2(F_{S'})} \sum_{\mu_2(F_{S'})} \mathcal{P}(\theta^0(\sigma(\delta)f, \boldsymbol{\omega}(\delta)\phi), \chi) \overline{\mathcal{P}(\theta^0(\sigma(\varepsilon)f, \boldsymbol{\omega}(\varepsilon)\phi), \chi)}.$$

Hence it suffices to proceed by considering the summands

$$\mathcal{P}(\theta^0(\sigma(\delta)f, \boldsymbol{\omega}(\delta)\phi), \chi) \overline{\mathcal{P}(\theta^0(\sigma(\varepsilon)f, \boldsymbol{\omega}(\varepsilon)\phi), \chi)} \\ = \int_{((\mathrm{SO}(W_1^\perp) \backslash \mathrm{SO}(X))(\mathbb{A}))^2} (\boldsymbol{\omega}(h_2\delta)\phi)(\xi) \overline{(\boldsymbol{\omega}(h_1\varepsilon)\phi)(\xi)} \Lambda_\xi(\sigma(h_2\delta)f, \chi) \overline{\Lambda_\xi(\sigma(h_1\varepsilon)f, \chi)} dh_1 dh_2.$$

We have $\overline{\Lambda_\xi(\sigma(h_1\varepsilon)f, \chi)} = \Lambda_\xi(\bar{\sigma}(h_1\varepsilon)\bar{f}, \bar{\chi})$ where $\bar{f} = \otimes_v \bar{f}_v \in \mathcal{V}_{\bar{\sigma}, S}^1$ and the vectors

$$\sigma(h_1\varepsilon)f = \otimes_v \sigma_v(h_{1,v}\varepsilon_v)f_v \in \mathcal{V}_{\sigma, S}^1 \quad \text{and} \quad \bar{\sigma}(h_1\varepsilon)\bar{f} = \otimes_v \bar{\sigma}_v(h_{1,v}\varepsilon_v)\bar{f}_v \in \mathcal{V}_{\bar{\sigma}, S}^1$$

are pure tensors. Thus the hypotheses of Proposition 7.4 are satisfied; we have

$$\Lambda_\xi(\sigma(h_2\delta)f, \chi) \overline{\Lambda_\xi(\sigma(h_1\varepsilon)f, \chi)} = 2^{s-1} \prod_v \Gamma_{\xi, v}(\sigma_v(h_{2,v}\delta_v)f_v, \bar{\sigma}_v(h_{1,v}\varepsilon_v)\bar{f}_v; \chi_v).$$

Subsequently,

$$\mathcal{P}(\theta^0(\sigma(\delta)f, \boldsymbol{\omega}(\delta)\phi), \chi) \overline{\mathcal{P}(\theta^0(\sigma(\varepsilon)f, \boldsymbol{\omega}(\varepsilon)\phi), \chi)} \\ = 2^{s-1} \prod_v \int_{(\mathrm{SO}(W_1^\perp)_v \backslash \mathrm{SO}(X)_v)^2} (\boldsymbol{\omega}_v(h_{2,v}\delta_v)\phi_v)(\xi) \overline{(\boldsymbol{\omega}_v(h_{1,v}\varepsilon_v)\phi_v)(\xi)} \\ \times \Gamma_{\xi, v}(\sigma_v(h_{2,v}\delta_v)f_v, \bar{\sigma}_v(h_{1,v}\varepsilon_v)\bar{f}_v; \chi_v) dh_{1,v} dh_{2,v}.$$

In summary, we have the following formula:

$$|\mathcal{P}(\theta(f, \phi), \chi)|^2 = \frac{1}{4^{s+s'}} 2^{s-1} \sum_{\delta \in \mu_2(F_{S'})} \sum_{\varepsilon \in \mu_2(F_{S'})} \prod_v \mathcal{J}_v(\delta_v, \varepsilon_v) \quad (7.3)$$

for which we have introduced the place-holder notation

$$\mathcal{J}_v(\delta_v, \varepsilon_v) = \int_{(\mathrm{SO}(W_1^\perp)_v \backslash \mathrm{SO}(X)_v)^2} (\boldsymbol{\omega}_v(h_{2,v}\delta_v)\phi_v)(\xi) \overline{(\boldsymbol{\omega}_v(h_{1,v}\varepsilon_v)\phi_v)(\xi)} \\ \times \Gamma_{\xi, v}(\sigma_v(h_{2,v}\delta_v)f_v, \bar{\sigma}_v(h_{1,v}\varepsilon_v)\bar{f}_v; \chi_v) dh_{1,v} dh_{2,v}.$$

The $\mathcal{J}_v(\delta_v, \varepsilon_v)$ are connected to the local integrals of Section 6.2 by

$$\alpha(\theta(f_v, \phi_v), \chi_v) = \frac{1}{2} \frac{\zeta_{F_v}(2)\zeta_{F_v}(4)}{L(1, \sigma_v, \mathrm{std})} \sum_{\varrho_v \in \mu_2(F_v)} \mathcal{J}_v(\varrho_v, 1),$$

recalling $\alpha(\theta(f_v, \phi_v), \chi_v) = \alpha(\theta(f_v, \phi_v), \theta(\bar{f}_v, \bar{\phi}_v); \chi_v)$. We now separate the sum in (7.3) according to the representation σ_v at v . The index set for the double summation runs over $\delta, \varepsilon \in \mu_2(F_{S'})$, with $\delta = (\delta_v)$ and $\varepsilon = (\varepsilon_v)$, where $\delta_v = \varepsilon_v = 1$ if $v \in \mathfrak{S}$ or $v \notin S$.

• If $v \notin S$, then, since $\varrho_v \in H(\mathcal{O}_v)$, $\mathcal{J}_v(\varrho_v, 1) = \mathcal{J}_v(1, 1)$ meaning

$$\mathcal{J}_v(1, 1) = \frac{1}{2} \sum_{\varrho_v \in \mu_2(F_v)} \mathcal{J}_v(\varrho_v, 1) = \frac{L(1, \sigma_v, \mathrm{std})}{\zeta_{F_v}(2)\zeta_{F_v}(4)} \alpha(\theta(f_v, \phi_v), \chi_v).$$

- If $v \in S \cap \mathfrak{S}$, then $J_v(t, 1) = 0$. Indeed, for $f = f + 0 \in \mathcal{V}_{\sigma_{0,v}} \oplus \mathcal{V}_{\sigma'_{0,v}}$ we have

$$\mathcal{B}_{\sigma_v}(\sigma_v(t)f_v, \bar{f}_v) = \frac{1}{2}(\mathcal{B}_{\sigma_{0,v}}(0, \bar{f}_v) + \mathcal{B}_{\sigma_{0,v}}(f_v, 0)) = 0 + 0.$$

The remaining term is

$$J_v(1, 1) = 2 \left(\frac{L(1, \sigma_v, \text{std})}{\zeta_{F_v}(2)\zeta_{F_v}(4)} \right) \alpha(\theta(f_v, \phi_v), \chi_v).$$

- If $v \in S'$, we have a four-term summation. Using that $J_v(t, t) = J_v(1, 1)$ we find

$$\sum_{\delta_v, \varepsilon_v \in \mu_2(F_v)} J_v(\delta_v, \varepsilon_v) = 2 \sum_{\varrho_v \in \mu_2(F_v)} J_v(\varrho_v, 1) = 4 \left(\frac{L(1, \sigma_v, \text{std})}{\zeta_{F_v}(2)\zeta_{F_v}(4)} \right) \alpha(\theta(f_v, \phi_v), \chi_v).$$

Together, these three points prove that (7.3) becomes

$$\begin{aligned} |\mathcal{P}(\theta(f, \phi), \chi)|^2 &= \frac{1}{4^{s+s'}} 2^{s-1} 2^s 4^{s'} \left(\frac{L(1, \sigma, \text{std})}{\zeta_F(2)\zeta_F(4)} \right) \prod_v \alpha(\theta(f_v, \phi_v), \chi_v) \\ &= \frac{1}{2} \left(\frac{L(1, \sigma, \text{std})}{\zeta_F(2)\zeta_F(4)} \right) \frac{\zeta_F(2)\zeta_F(4)L(\pi, \chi, 1/2)}{L(\text{Ad}, \pi, 1)L(\chi_{K/F}, 1)} \prod_v \alpha^\natural(\theta(f_v, \phi_v), \chi_v). \end{aligned}$$

Finally, for our formula to be independent of choice of local pairings (see Remark 7.7) we normalise the Bessel period and instead calculate

$$\frac{|\mathcal{P}(\varphi, \chi)|^2}{\mathcal{B}_{\theta(\sigma)}(\varphi, \bar{\varphi})\mathcal{B}_\chi(\chi, \bar{\chi})} \tag{7.4}$$

for $\varphi \in \mathcal{V}_{\theta(\sigma)}$. The Petersson pairing for the one-dimensional representation χ is trivially constant in this case and is easily seen to equal the Tamagawa number

$$\mathcal{B}_\chi(\chi, \bar{\chi}) = \mathcal{B}_\chi(1, 1) = \text{Vol}(\mathbb{A}^\times K^\times \backslash \mathbb{A}_K^\times) = 2.$$

The Petersson pairing for the theta lift $\theta(\sigma)$ is dealt with by the formula of Gan–Ichino (4.9) which states that the Petersson inner product for $\theta(\sigma)$ equals

$$\mathcal{B}_{\theta(\sigma)} = \frac{L(1, \sigma, \text{std})}{\zeta_F(2)\zeta_F(4)} \prod_v \mathcal{B}_{\theta(\sigma_v)}.$$

Combining these final comments gives the main result.

Theorem 7.5. *Let (π, \mathcal{V}_π) be an irreducible, cuspidal automorphic representation of $\text{PGSp}_4(\mathbb{A})$ lifted, via the theta correspondence in Section 3, from (the Jacquet–Langlands transfer of) a cuspidal automorphic representation of $\text{GL}_2(\mathbb{A}_E)$ with trivial central character. Let K be a quadratic field extension of F such that $\text{SO}_2 \cong K^\times/F^\times$. Let χ be a unitary Hecke character of \mathbb{A}_K^\times such that $\chi|_{\mathbb{A}^\times} = 1$; such a χ may also be viewed as an automorphic representation of $\text{SO}_2(\mathbb{A})$. For the cusp forms $\varphi = \otimes_v \varphi_v \in \mathcal{V}_\pi$ and $\bar{\varphi} = \otimes_v \bar{\varphi}_v \in \mathcal{V}_{\bar{\pi}}$ define the local integrals $\alpha^\natural(\varphi_v, \chi_v)$ as in Section 6: we have $\alpha^\natural(\varphi_v, \chi_v) = 1$ for almost all v . For any choice of local Haar measures defining $\alpha^\natural(\varphi_v, \chi_v)$ let $C \in \mathbb{C}$ be the Haar measure constant (the constant of proportionality given by the ratio of the Tamagawa measure divided by the product of local measures). For each v , let \mathcal{B}_{π_v} be any choice of local unitary pairing. We have proved that*

$$\frac{|\mathcal{P}(\varphi, \chi)|^2}{\mathcal{B}_\pi(\varphi, \bar{\varphi})\mathcal{B}_\chi(\chi, \bar{\chi})} = \frac{C}{4} \frac{\zeta_F(2)\zeta_F(4)L(1/2, \pi \boxtimes \chi)}{L(1, \pi, \text{Ad})L(1, \chi_{K/F})} \prod_v \frac{\alpha^\natural(\varphi_v, \chi_v)}{\mathcal{B}_{\pi_v}(\varphi_v, \bar{\varphi}_v)}.$$

Definition 7.1. We define the local integrals to be properly normalised in the following way: choose local unitary pairings \mathcal{B}_{χ_v} on each one-dimensional space $\mathcal{V}_{\chi_v} \otimes \mathcal{V}_{\bar{\chi}_v}$ such that the Petersson pairing decomposes as $\mathcal{B}_\chi = \prod_v \mathcal{B}_{\chi_v}$. We then take the normalised quantity

$$\mathcal{B}_{\chi_v}(\chi_v, \bar{\chi}_v) \alpha^\natural(\varphi_v, \chi_v)$$

in place of the local integrals in the formula of Theorem 7.5. Note that in the original definition of the local integrals (Section 6.1) we implicitly take $\mathcal{B}_{\chi_v} = 1$ for each v , as per Section 4.4, and we found the decomposition $\mathcal{B}_\chi = 2 \prod_v \mathcal{B}_{\chi_v}$.

Corollary 7.6. Assuming $C = 1$, $\mathcal{B}_\pi = \prod_v \mathcal{B}_{\pi_v}$, and that the local integrals $\alpha^{\natural}(\varphi_v, \chi_v)$ are properly normalised (as in Definition 7.1), Theorem 7.5 becomes

$$|\mathcal{P}(\varphi, \chi)|^2 = \frac{1}{4} \frac{\zeta_F(2)\zeta_F(4)L(1/2, \pi \boxtimes \chi)}{L(1, \pi, \text{Ad})L(1, \chi_{K/F})} \prod_v \alpha^{\natural}(\varphi_v, \chi_v).$$

Remark 7.7. In a more general setting, the representation χ need not be one-dimensional (when considering other groups). Normalising the left-hand-side of the equation in Theorem 7.5 by the Petersson pairings for π and χ , and including the Haar measure constant, ensures that the local choices of pairings and measures are independent of the global setting. These objects may be chosen and may be chosen arbitrarily without affecting the formula and, in particular, the local integrals are independent of such choices (see [11, Remark 1.3]).

Our normalisations may seem ad hoc at first, due to the trivial pairings on χ , however we state our theorem in this way so that it sits in the more general framework of Liu’s conjecture. In Liu’s work one sees that the issue of normalisation appears in a natural setting and we invite the reader to check [14, Conjecture 2.5] for consolidation.

Acknowledgment: The author would like to offer sincere thanks to both Yifeng Liu, for his helpful comments and discussions, and Abhishek Saha, for his valuable guidance. Thanks are also due to Katharine Thornton for her many insightful suggestions.

References

- [1] J. Arthur and L. Clozel, *Simple Algebras, Base Change, and the Advanced Theory of the Trace Formula*, Ann. of Math. Stud. 120, Princeton University Press, Princeton, 1989.
- [2] S. Böcherer, Bemerkungen über die Dirichletreihen von Koecher und Maass, *Math. Gottingensis, Schriftenr. Sonderforschungsbereichs Geom. Anal.* **68** (1986).
- [3] J. Cogdell, Dual groups and Langlands functoriality, in: *An Introduction to the Langlands Program*, Birkhäuser, Boston (2004), 251–268.
- [4] M. Furusawa, On L -functions for $\text{GSp}(4) \times \text{GL}(2)$ and their special values, *J. Reine Angew. Math.* **438** (1993), no. 4, 187–218.
- [5] M. Furusawa, K. Martin and J. Shalika, On central critical values of the degree four L -functions for $\text{GSp}(4)$: The fundamental lemma. III, *Mem. Amer. Math. Soc.* **225** (2013), no. 1057.
- [6] W. T. Gan, B. Gross and D. Prasad, Symplectic local root numbers, central critical L values, and restriction problems in the representation theory of classical groups, in: *Sur les conjectures de Gross et Prasad. I*, Astérisque 346, Société Mathématique de France, Paris (2012), 1–109.
- [7] W. T. Gan and A. Ichino, On endoscopy and the refined Gross–Prasad conjecture for $(\text{SO}_5, \text{SO}_4)$, *J. Inst. Math. Jussieu* **10** (2011), no. 2, 235–324.
- [8] B. Gross and D. Prasad, On the decomposition of a representation of SO_n when restricted to SO_{n-1} , *Canad. J. Math.* **44** (1992), no. 5, 974–1002.
- [9] M. Harris and S. Kudla, Arithmetic automorphic forms for the nonholomorphic discrete series of $\text{GSp}(2)$, *Duke Math. J.* **66** (1992), no. 1, 59–121.
- [10] M. Harris, D. Soudry and R. Taylor, l -adic representations associated to modular forms over imaginary quadratic fields. I: Lifting to $\text{GSp}_4(\mathbb{Q})$, *Invent. Math.* **112** (1993), no. 2, 377–411.
- [11] A. Ichino, Trilinear forms and the central values of triple product L -functions, *Duke Math. J.* **145** (2008), no. 2, 281–307.
- [12] A. Ichino and T. Ikeda, On the periods of automorphic forms on special orthogonal groups and the Gross–Prasad conjecture, *Geom. Funct. Anal.* **19** (2010), no. 5, 1378–1425.
- [13] E. Lapid and Z. Mao, A conjecture on Whittaker–Fourier coefficients of cusp forms, *J. Number Theory* **146** (2015), 448–505.
- [14] Y. Liu, Refined global Gan–Gross–Prasad conjecture for Bessel periods, *J. Reine Angew. Math.* (2014), DOI 10.1515/crelle-2014-0016.
- [15] D. Prasad and R. Takloo-Bighash, Bessel models for $\text{GSp}(4)$, *J. Reine Angew. Math.* **655** (2011), 189–243.
- [16] Y. Qiu, The Bessel period functional on $\text{SO}(5)$: The nontempered case, preprint (2013), <http://arxiv.org/abs/1312.5793>.
- [17] D. Ramakrishnan and R. Valenza, *Fourier Analysis on Number Fields*, Grad. Texts in Math. 186, Springer, New York, 1999.
- [18] B. Roberts, Global L -packets for $\text{GSp}(2)$ and theta lifts, *Doc. Math.* **6** (2001), 247–314.
- [19] A. Saha, A relation between multiplicity one and Böcherer’s conjecture, *Ramanujan J.* **33** (2014), 263–268.

- [20] S. Takeda, Some local-global non-vanishing results for theta lifts from orthogonal groups, *Trans. Amer. Math. Soc.* **361** (2009), no. 10, 5575–5599.
- [21] J.-L. Waldspurger, Sur les valeurs de certaines fonctions L automorphes en leur centre de symétrie, *Compos. Math.* **54** (1985), no. 2, 173–242.
- [22] A. Weil, *Adeles and Algebraic Groups*, Progr. Math. 23, Birkhäuser, Boston, 1982.