



Baritussio, A., Ghosh Dastidar, A., Frontera, A., Ahmed, N., De Garate, E., Harries, I., ... Bucciarelli-Ducci, C. (2017). Diagnostic yield of cardiovascular magnetic resonance in young-middle aged patients with high-grade atrio-ventricular block. *International Journal of Cardiology*, 244, 335-339. <https://doi.org/10.1016/j.ijcard.2017.06.080>

Peer reviewed version

License (if available):  
CC BY-NC-ND

Link to published version (if available):  
[10.1016/j.ijcard.2017.06.080](https://doi.org/10.1016/j.ijcard.2017.06.080)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via Elsevier at <https://doi.org/10.1016/j.ijcard.2017.06.080> . Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/pure/about/ebr-terms>

**Diagnostic Yield of Cardiovascular Magnetic Resonance in Young-Middle Aged Patients with High-Grade Atrio-Ventricular Block**

A. Baritussio, A. Ghosh Dastidar, A. Frontera, N. Ahmed, E. De Garate, I. Harries, I. Diab, E. Duncan, G. Thomas, A. Nisbet, C. Bucciarelli-Ducci

Bristol NIHR Cardiovascular Biomedical Research Unit, Bristol Heart Institute, Bristol, United Kingdom

Authors take responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation

Corresponding author:

Dr Chiara Bucciarelli-Ducci

NIHR Bristol Cardiovascular Biomedical Research Unit,

CMR Unit, Bristol Heart Institute

Upper Maudlin Street

Bristol, BS2 8HW, United Kingdom

Email: [c.bucciarelli-ducci@bristol.ac.uk](mailto:c.bucciarelli-ducci@bristol.ac.uk)

Telephone: +44 117 342 5888

## **Conflict of interest**

C. Bucciarelli-Ducci is Consultant for Circle Cardiovascular Imaging.

## **Keywords**

High grade atrio-ventricular block; cardiovascular magnetic resonance; myocardial tissue characterization

## **List of abbreviations**

AVB Atrio-ventricular block

CMR Cardiovascular magnetic resonance

ECG Electrocardiogram

IHD Ischemic heart disease

LGE Late gadolinium enhancement

LMNA Lamin A/C

LV Left ventricle

NIHD Non-ischemic heart disease

PM Pace-maker

RV Right ventricle

TAVI Trans-catheter aortic valve implantation

TTE Transthoracic echocardiogram

## **Abstract**

**Background:** Atrio-ventricular block (AVB) is a rare finding in young or middle-aged adults, often leading to pacemaker implantation (PM) without further investigation. We sought to assess the diagnostic role of cardiovascular magnetic resonance (CMR) in young and middle-aged adults with high-grade AVB.

**Methods:** We consecutively enrolled young-middle aged (18-65yrs) patients with high grade AVB referred to CMR after standard clinical assessment (history, electrocardiogram and cardiac rhythm monitoring) prior to PM implantation. Cine and post-contrast imaging were performed in a 1.5T scanner.

**Results:** 34 patients (59% male, mean age  $42 \pm 12$  years) with high grade AVB were referred to CMR for suspected ischemic heart disease (IHD)(n=4) and non-ischemic heart disease (NIHD)(n=20); no clear cause was found in 9 patients prior to CMR and 1 patient had suspected lung disease. A pathologic substrate was found on CMR in 15 patients (44%), while a structurally normal heart was reported in 18 (53%). Non-specific findings were reported in 1 patient (3%). There was a fair agreement between CMR and echocardiographic findings (Cohen's kappa 0.243), and CMR provided an entirely new diagnosis in 34% of patients. As compared to the standard clinical assessment, CMR had an additional role in 65% of patients and guided further testing (genetic testing, extra-cardiac imaging) in 9%.

**Conclusions:** CMR found a pathologic substrate in 44% of patients, mainly NIHD (32%). Half of patients (53%) had a structurally normal heart. When added to the

standard clinical assessment, CMR had an incremental diagnostic role in two thirds of patients.

## **Introduction**

Atrio-ventricular block (AVB) typically shows a bimodal distribution, with a major peak in incidence in the elderly, commonly due to degeneration of the conduction system, and the other in infancy and early childhood, when it is mainly congenital. While the lower-grade AVB is considered benign as it can be observed in up to 1–2% of healthy young people, especially during sleep [1], the higher-grade AVB, such as second degree Mobitz II or complete AVB, is expression of the severity of the underlying cause and more often requires a pacemaker (PM) implantation [2]. Higher grade AVB is uncommon in young or middle-aged adults, but when identified it poses a dilemma, both in terms of diagnosis and clinical management. According to international guidelines [2], patients are frequently offered permanent PM implantation without further investigation, and up to 3-5% of all the patients undergoing PM implantation for AVB are aged 18-55 years [3,4]. PM implantation in young adults is generally recognized as a safe procedure, but it has been associated with peri- and post-procedural complications [5,6]. Ischemic, autoimmune and infiltrative diseases are the recognized causes of complete AVB in young or middle-aged adults [5,6]. Correct identification of the underlying pathophysiologic mechanism has a great impact both on therapeutic strategies and on the prognosis of AVB [4,7]. International guidelines recommend the clinical assessment of patients, comprehensive of clinical history, ECG and cardiac rhythm monitoring, but do not provide indications with regards to the need for cardiac imaging prior to treatment [2]. Cardiac magnetic resonance (CMR) is a well-established diagnostic imaging technique with increasing applications in clinical practice: CMR represents the gold standard for the assessment of biventricular volumes and function [8] and offers a uniquely advanced,

non-invasive, myocardial tissue characterization, providing an accurate assessment of myocardial fibrosis by late gadolinium enhancement, which has been validated against histology [9,10] . We sought to assess the diagnostic role of CMR in young and middle-aged adults with high-grade AVB.

## **Materials and Methods**

We retrospectively enrolled consecutive young and middle-age patients (aged 18-65 years) referred for CMR prior to PM implantation because of higher-grade AVB. Higher-grade AVB was defined as advanced II degree AVB (Mobitz II AVB, 2:1 AVB) and complete or III degree AVB. All patients underwent a 1.5T CMR (Avanto, Siemens Healthcare, Germany), with a comprehensive protocol that included long and short axis cine and late gadolinium enhancement (LGE) images. Steady state free precession sequences were performed to acquire the long and short axis cine images; typical parameters were TR 38 ms, TE 1.07 ms, flip angle 80°, slice-thickness 8 mm, inter-slice gap 0 mm, bandwidth 930 Hz/Px, voxel size 2.0x2.0x8.0 mm and temporal resolution  $\leq 45$  ms between phases. For LGE imaging, a standard inversion recovery gradient-echo sequence was adopted. The LGE images were acquired 10-15 minutes after intravenous injection of 0.1 mmol/Kg of body weight of gadolinium-chelate contrast agent (Gadovist 1.0 mmol/ml, Bayer-Schering, Berlin, Germany) in identical short-axis planes to cine images, using an inversion recovery prepared breath-hold gradient-echo technique. Typical image parameters were TR 700 ms, TE 3.15 ms; flip angle 25°; slice thickness 8.0 mm, no interslice gap, bandwidth 140 Hz/Px and voxel size  $2.0 \times 1.5 \times 8.0$  mm. The inversion time was progressively optimized to null normal myocardium (typical values,



250–350 ms). Each slice was obtained during a breath-hold of 10–15 s depending on the patient's heart rate.

Ventricular function was assessed with dedicated software (Circle Cardiovascular Imaging, Calgary, Canada), by tracing endo- and epicardial borders on each short axis cine slice in end-diastole and end-systole. All volumes measurements were indexed to body surface area. The presence/absence, localization, and distribution pattern of LGE were assessed visually by a SCMR/ESC level 3 certified individual with > 10 years of experience. Pattern and extent of LGE were assessed by using short- and long-axis views and were defined as present only if they were detectable in two orthogonal planes. The pattern of LGE distribution was defined as subendocardial or transmural, if involving <50% or  $\geq$ 50% of wall thickness, respectively, and as mid-wall/epicardial if patchy/spotty intra-mural or sub-epicardial enhancement was detected. The presence of LGE at the right ventricle/left ventricle insertion points, in the absence of other LGE distribution patterns, was defined as non-specific findings, as its diagnostic and prognostic meaning is still unclear. All the analysis was carried out in accordance with the recommendation of the Society for Cardiovascular Magnetic Resonance [11]. Based on a standard clinical assessment, including clinical history, ECG and cardiac rhythm monitoring, all patients were referred with a pre-CMR working diagnosis, classified as ischemic heart disease (IHD), non-ischemic heart disease (NIHD), normal heart and unknown cause. The study was reviewed by the local Institutional Research and Innovation Department and in view of the retrospective design of the study, formal ethical approval was waived off. All patients gave written informed consent.

### **Statistical analysis**

Continuous and categorical variables are expressed as mean±SD or median (IQR), and n (%), respectively. Categorical variables were compared by using the chi-square test or Fisher exact test, as appropriate. Inter-rater agreement for categorical variables was assessed by Cohen's kappa coefficient. A p-value of <0.05 was considered statistically significant. Data were analysed with SPSS® version 23 (IBM®).

## **Results**

We consecutively enrolled thirty-four patients (59% male, mean age 42 ±12 years, minimum-maximum 21-61 years) referred to CMR prior to PM implantation secondary to higher grade AVB. Twenty-two patients (65%) had evidence of complete AVB and 12 patients (35%) had second degree Mobitz II AVB. Twenty-three patients (68%) showed intermittent high grade AVB on 24 hours Holter monitoring or on implantable loop recorder monitoring. Eleven patients (32%), all but one presenting with III degree AVB, were admitted to hospital following syncope; they were all haemodynamically stable and underwent CMR as inpatients. Based on the standard clinical assessment, 4 patients were referred for suspected ischemic heart disease (IHD) and 20 for suspected NIHD; no clear cause was identified in 9 patients prior to CMR and one patient had suspected interstitial lung disease. Median LVEF was 64% (IQR 58-68), median indexed left ventricular end-diastolic volume (LVEDV) was 88 ml/m<sup>2</sup> (IQR 77-109) and median indexed left ventricular end-systolic volume (LVESV) was 31 ml/m<sup>2</sup> (IQR 27-54). On tissue characterization, LGE was found in twelve patients (35%), mainly with a mid-wall/epicardial pattern (7/12 patients, 58%); LGE had a septal distribution in 5 patients (42%). Based on CMR findings, a pathologic substrate was found in 15 patients (44%): 4

patients (12%) were diagnosed with IHD and 11 (32%) with NIHD. A structurally normal heart was found in 18 patients (53%) and non-specific findings were reported in 1 patient (3%). Among patients presenting findings consistent with NIHD, myocarditis was the most common (5 patients, of which 2 were diagnosed with cardiac sarcoidosis, Figure 1A and B), followed by athlete's heart and valvular heart disease (moderate aortic regurgitation and severe pulmonary regurgitation) which were each reported in 2 patients, respectively; dilated cardiomyopathy (Figure 1C and D) and left ventricular non-compaction were each found in 1 patient. CMR findings are listed in Table 1.

### **Standard clinical assessment vs CMR findings**

Based on clinical and family history and on ECG findings, IHD was suspected prior to cardiac imaging in 4 patients and NIHD in 20 patients; no clear cause could be identified in 9 patients and 1 patient was diagnosed with suspected lung disease. CMR had an additional role in re-defining the final findings in 22/34 (65%) patients, mainly by re-defining the final diagnosis in 4 patients and by ruling out an underlying structural heart disease in 18 patients (Table 2). Among the 20 patients with a pre-imaging diagnosis of suspected NIHD, CMR diagnosed IHD in 2 patients and found a structurally normal heart in 9 patients. Among the 9 patients with no clear cause for AVB prior to imaging, CMR found a structurally normal heart in 7 patients and NIHD in one.

### **Diagnostic performance of CMR and TTE**

Trans-thoracic echocardiogram (TTE) data were available in 29/34 patients (85%). A pathologic substrate was found on TTE in 3/29 patients (10%): one patient (3%) was

diagnosed with IHD and 2 (7%) with NIHD. There was no significant difference between the ability of CMR and TTE to identify a pathologic substrate, but there was a trend in favour of CMR (15/34 vs 3/29,  $p=0.539$ ). A structurally normal heart was found on TTE in 21 patients (73%) and non-specific findings were reported in 5 patients (17%). CMR and TTE provided the same diagnosis in 17/29 patients (59%) with an overall fair agreement (Cohen's kappa 0.243,  $p=0.019$ ); CMR had mainly a role in re-defining patients diagnosed with normal heart and those with non-specific findings on TTE, overall providing a new diagnosis in 10/29 patients (34%). Out of 21 patients diagnosed with normal heart on TTE, 5 (24%) received a different diagnosis on CMR: one patient had findings of IHD and 4 of NIHD. CMR was diagnostic in all 5 patients with non-specific findings on TTE: three patients had findings consistent with NIHD, one with IHD and one with structurally normal heart (Table 3).

## **Discussion**

The main findings of our study were: 1) CMR identified an underlying pathologic substrate in nearly half of patients, 2) the most prevalent underlying conditions was NIHD (32%); 3) CMR demonstrated a significant incremental diagnostic value (in 65% of patients) when added to standard clinical work-up of patients with AVB; 4) when added to transthoracic echocardiography CMR provided an entirely new diagnosis in 34% of patients.

Higher grade AVB is rare in young or middle-aged adults and the underlying etiologic mechanisms are multiple and different from the degenerative process mainly responsible for AVB in the elderly [12]. To the best of our knowledge this is the largest study

investigating the role of CMR in patients with high-grade AVB, and its added value to the current standard clinical practice. In our cohort, CMR could identify an underlying pathologic substrate in nearly half of patients (44%). We found a 12% incidence of IHD, which is in keeping with previous studies on middle-aged (45-65 years) patients presenting with AVB [13] and on patients with severe LV systolic dysfunction after acute myocardial infarction [14]. More than a third (32%) of our patients had evidence of NIHD on CMR; as confirmed by endomyocardial biopsy (EMB) studies in patients with high grade AVB [15-17], we have also found that myocarditis is the most common non-ischemic cardiomyopathy. It has been shown that Lamin A/C gene mutation in dilated cardiomyopathy accounts for 33% of AVB [18] cases, but similar scenarios are also seen in cases of LV non-compaction and hypertrophic cardiomyopathy, as also confirmed in our cohort. Infiltrative diseases are frequently associated with AVB, which often represents the first and most common clinical manifestation, as in the case of cardiac sarcoidosis [19]. Although cardiac sarcoidosis commonly results in AVB, it is a rare disease, and its prevalence in the spectrum of AVB is not well known [20]. Based on CMR findings we diagnosed cardiac sarcoidosis in 2 patients (6%, 18% of the NIHD). Other causes of high grade AVB in young-middle aged patients include cardiac amyloidosis [21,22] and can be rarely encountered among trained athlete's [23,24], where it is usually considered pathological.

### **CMR vs standard of care clinical assessment**

In patients presenting with AVB, international guidelines recommend a clinical assessment, comprehensive of clinical history, ECG and cardiac rhythm monitoring, but

there is no indication with regards to the need for cardiac imaging prior to treatment [2]. Patients are thus often offered a PM implantation without further investigation, although the underlying pathology has important implications on prognosis. CMR showed an additional role in 65% of patients when added to the standard clinical assessment, mainly by re-defining the final diagnosis in 4 patients and by ruling out an underlying structural heart disease in 18 patients (Table 2). European guidelines on cardiac pacing [2] do not recommend TTE in the work up of high-grade AVB patients. Nevertheless, nearly all of our patients (85%) underwent a TTE in addition to the standard clinical assessment. TTE could identify a pathologic substrate in 10% of patients, which was lower compared to CMR (44%) although it did not reach statistical significance, most likely because of the small sample size. The agreement between TTE and CMR findings was fair, and CMR provided an entirely new diagnosis in 10/29 (34%) patients, mainly by re-defining diagnosis among patients with a structurally normal heart (5/21, 24%) and being diagnostic in all 5 patients with non-specific findings on TTE.

### **Clinical Implications**

Although CMR did not change the final treatment strategy with regards to AVB, as all patients received a PM, information provided by CMR, as compared to both the standard clinical assessment and TTE, had a clinical impact in patients' management with direct implications on their treatment strategies.

The superior diagnostic capability of CMR in our cohort was mainly related to the analysis of LGE sequences: LGE was found in 35% of patients, mainly with a non-ischemic distribution pattern (58%) and with a septal location in 42% of patients. It is

well established that LGE distribution pattern allows non-invasive identification of the structural aetiology of cardiomyopathies [25-27], with a diagnostic accuracy in some cases exceeding that of trans-thoracic echocardiogram and even of EMB [28]. The advanced tissue characterization provided by CMR has also been shown to allow the identification of genetic variants of common cardiomyopathies [29,30]: a recent study by Holmstrom et al. demonstrated that 88% of asymptomatic or mildly symptomatic carriers of Lamin A/C (LMNA) mutations causing dilated cardiomyopathy had typical myocardial fibrosis, involving the mid-myocardium of the basal septum (Figure 1D, white arrow) [29]. The fibrosis was observed in all individuals with an AV conduction defect and LGE-pattern was typically linear with involvement of less than 50% of the area of the segment. In our series, a patient showed mildly dilated LV with the typical mid-wall fibrosis of the basal antero-septum on CMR (Figure 1D, white arrow): late genetic testing was positive for LMNA mutation. Moreover, in our patients cohort, CMR diagnosis of cardiac sarcoidosis triggered further testing, leading to the finding of extra-cardiac manifestations of the disease in both patients.

Finally, it has been shown that CMR is able to detect myocardial infarction in a non-negligible proportion of patients (approximately one third) in which the diagnosis was missed on non-CMR imaging [31]: although this was not the case in our patients' cohort, finding evidence of unknown IHD on CMR in patients with high-grade AVB might lead to diagnostic invasive coronary angiogram, as underlying coronary artery disease can explain the conduction disorder [13].

### **Study Limitations**

The main limitation of the present study is the limited number of patients; nevertheless, AVB in this age group (18-65 years) is a very rare condition and to our knowledge, this is the first and largest series of young and middle-aged patients with high-grade AVB referred for CMR. Half of our patients were found to have structurally normal hearts. It is acknowledged that transient tissue damage, in the form of reversible myocardial oedema, as in the few Tako-Tsubo cases reported in literature [32,33], may induce atrio-ventricular conduction abnormalities; T2-weighted sequences for myocardial oedema were not performed in our cohort, so transient acute tissue damage could not be assessed. We might speculate that this aspect might explain the high prevalence of structurally normal hearts in our cohort. A structurally normal heart on CMR reflects the absence of gross ischemic or non-ischemic abnormalities, but cannot exclude the presence of microscopic, ultra-structural abnormalities, which are below the resolution of the technique. EMB is the gold standard to assess ultra-structural abnormalities, but carries non-negligible complications, and is not routinely performed in clinical practice, and we had no histologic data in our patients' cohort. Finally, LGE imaging accurately allows the identification and characterization of focal fibrosis; however, exploring the presence of diffuse fibrosis on T1 mapping technique [34] might provide a new insight in the aetiology of high-grade AVB. Further studies, to assess tissue characterization with more advanced sequences, such as the T1 and T2 mapping, could provide additional insight on interstitial changes in this patients' cohort.

## **Conclusions**

In young patients presenting with high grade AVB, CMR demonstrated incremental



diagnostic yield by identifying a pathologic substrate in 44% of patients. There was a fair agreement between CMR and TTE final diagnosis, and CMR provided an entirely new diagnosis in 34% of patients. When added to the standard clinical assessment (history, ECG and cardiac rhythm monitoring) CMR had an additional diagnostic role in two thirds of patients. Our results show that CMR could be a valuable test to be included in the clinical work-up of patients with AVB.

**Funding**

This study was funded by the NIHR Biomedical Research Unit in Cardiovascular Disease at the University Hospitals Bristol NHS Foundation Trust and the University of Bristol.

The views expressed are those of the authors and not necessarily those of the UK National Health Service, National Institute for Health Research, or Department of Health.

**Acknowledgements**

We are thankful to Mr Christopher Benjamin Lawton for his skilled help in image acquisition.

## References

1. Barra SNC, Providência R, Paiva L, Nascimento J, Marques AL. A review on advanced atrioventricular block in young or middle-aged adults. *Pacing Clin Electrophysiol.* 2012;35(11):1395-1405. doi:10.1111/j.1540-8159.2012.03489.
2. Brignole M, Auricchio A, Baron-Esquivias G, et al. 2013 ESC Guidelines on cardiac pacing and cardiac resynchronization therapy: the Task Force on cardiac pacing and resynchronization therapy of the European Society of Cardiology (ESC). Developed in collaboration with the European Heart Rhythm Association . *Eur Heart J.* 2013;34(29):2281-2329. doi:10.1093/eurheartj/eh150.
3. Kandolin R, Lehtonen J, Kupari M. Cardiac sarcoidosis and giant cell myocarditis as causes of atrioventricular block in young and middle-aged adults. *Circ Arrhythm Electrophysiol.* 2011;4(3):303-309. doi:10.1161/CIRCEP.110.959254.
4. Pyatt J, Somauroo JD, Jackson M, et al. Long-term survival after permanent pacemaker implantation: analysis of predictors for increased mortality. *Europace.* 2002;4(2):113-119. doi:10.1053/eupc.2002.0233.

5. Harcombe AA, Newell SA, Ludman PF, et al. Late complications following permanent pacemaker implantation or elective unit replacement. *Heart*. 1998;80(3):240-244. doi:10.1136/hrt.80.3.240.
6. Poole JE, Gleva MJ, Mela T, et al. Complication rates associated with pacemaker or implantable cardioverter-defibrillator generator replacements and upgrade procedures: results from the REPLACE registry. *Circulation*. 2010;122(16):1553-1561. doi:10.1161/CIRCULATIONAHA.110.976076.
7. Zanini R, Facchinetti A, Gallo G, et al. Survival Rates after Pacemaker Implantation: A Study of Patients Paced for Sick Sinus Syndrome and Atrioventricular Block. *Pacing Clin Electrophysiol*. 1989;12(7):1065-1069. doi:10.1111/j.1540-8159.1989.tb01927.
8. Pennell DJ. Ventricular volumes and mass by CMR. *J Cardiovasc Magn Reson* 2002; 4: 507-13.
9. Mahrholdt H, Wagner A, Judd RM, Seechurn U, Kim RJ. Delayed enhancement cardiovascular magnetic resonance assessment of non-ischaemic cardiomyopathies. *Eur H J* 2005; 26: 1461-74.

10. Kim RJ, Fieno DS, Parrish TB, et al. Relationship of MRI delayed contrast enhancement to irreversible injury, infarct age, and contractile function. *Circulation* 1999; 100: 1992-2002.
11. Schulz-Menger J, Bluemke DA, Bremerich J, et al. Standardized image interpretation and post processing in cardiovascular magnetic resonance: Society for Cardiovascular Magnetic Resonance (SCMR) Board of Trustees Task Force on standardized post processing. *J Cardiovasc Magn Reson* 2013; 15: 35.
12. Chow GV, Marine JE, Fleg JL. Epidemiology of arrhythmias and conduction disorders in older adults. *Clin Geriatr Med* 2012; 28: 539-53.
13. Ginks W, Sutton R, Siddons H, Leatham A. Unsuspected coronary artery disease as cause of chronic atrioventricular block in middle age. *Br Heart J*. 1980;44(6):699-702.
14. Bloch TPE, Jons C, Raatikainen MJP, et al. Long-term recording of cardiac arrhythmias with an implantable cardiac monitor in patients with reduced ejection fraction after acute myocardial infarction: the Cardiac Arrhythmias and Risk

Stratification After Acute Myocardial Infarction (CARISMA) study. *Circulation*. 2010;122(13):1258-1264. doi:10.1161/CIRCULATIONAHA.109.902148.

15. Hufnagel G, Pankuweit S, Richter A, Schönian U, Maisch B. The European Study of Epidemiology and Treatment of Cardiac Inflammatory Diseases (ESETCID). First epidemiological results. *Herz*. 2000;25(3):279-285.
16. Uemura A, Morimoto S, Hiramitsu S, Hishida H. Endomyocardial biopsy findings in 50 patients with idiopathic atrioventricular block: presence of myocarditis. *Jpn Heart J*. 2001;42(6):691-700.
17. Davidoff R, Palacios I, Southern J, Fallon JT, Newell J, Dec GW. Giant cell versus lymphocytic myocarditis. A comparison of their clinical features and long-term outcomes. *Circulation*. 1991;83(3):953-961.
18. Arbustini E, Pilotto A, Repetto A, et al. Autosomal dominant dilated cardiomyopathy with atrioventricular block: a lamin A/C defect-related disease. *J Am Coll Cardiol*. 2002;39(6):981-990. doi:10.1016/S0735-1097(02)01724-2.
19. Banba K, Kusano KF, Nakamura K, et al. Relationship between arrhythmogenesis

and disease activity in cardiac sarcoidosis. *Heart Rhythm*. 2007;4(10):1292-1299.  
doi:10.1016/j.hrthm.2007.06.006.

20. Yoshida Y, Morimoto S, Hiramitsu S, Tsuboi N, Hirayama H, Itoh T. Incidence of cardiac sarcoidosis in Japanese patients with high-degree atrioventricular block. *Am Heart J*. 1997;134(3):382-386.
21. Cheng Z, Zhu K, Tian Z, Zhao D, Cui Q, Fang Q. The findings of electrocardiography in patients with cardiac amyloidosis. *Ann Noninvasive Electrocardiol*. 2013;18(2):157-162. doi:10.1111/anec.12018.
22. Reisinger J, Dubrey SW, Lavalley M, Skinner M, Falk RH. Electrophysiologic abnormalities in AL (primary) amyloidosis with cardiac involvement. *J Am Coll Cardiol*. 1997;30(4):1046-1051.
23. Ector H, Bourgois J, Verlinden M, et al. Bradycardia, ventricular pauses, syncope, and sports. *Lancet*. 1984;2(8403):591-594.
24. Stein R, Medeiros CM, Rosito GA, Zimmerman LI, Ribeiro JP. Intrinsic sinus and atrioventricular node electrophysiologic adaptations in endurance athletes. *J Am*

*Coll Cardiol.* 2002;39(6):1033-1038.

25. Cheong BYC, Muthupillai R, Nemeth M, et al. The utility of delayed-enhancement magnetic resonance imaging for identifying nonischemic myocardial fibrosis in asymptomatic patients with biopsy-proven systemic sarcoidosis. *Sarcoidosis Vasc Diffuse Lung Dis.* 2009;26(1):39-46.
26. Tadamura E, Yamamuro M, Kubo S, et al. Effectiveness of delayed enhanced MRI for identification of cardiac sarcoidosis: comparison with radionuclide imaging. *AJR Am J Roentgenol.* 2005;185(1):110-115. doi:10.2214/ajr.185.1.01850110.
27. Selvanayagam JB, Leong DP. MR Imaging and Cardiac Amyloidosis Where to Go From Here? *JACC Cardiovasc Imaging.* 2010;3(2):165-167.  
doi:10.1016/j.jcmg.2009.10.013.
28. Lurz P, Eitel I, Adam J, et al. Diagnostic performance of CMR imaging compared with EMB in patients with suspected myocarditis. *JACC Cardiovasc Imaging.* 2012;5(5):513-524. doi:10.1016/j.jcmg.2011.11.022.
29. Holmström M, Kivistö S, Heliö T, et al. Late gadolinium enhanced cardiovascular



magnetic resonance of lamin A/C gene mutation related dilated cardiomyopathy. *J Cardiovasc Magn Reson*. 2011;13:30. doi:10.1186/1532-429X-13-30.

30. Raman SV, Sparks EA, Baker PM, McCarthy B, Wooley CF. Mid-myocardial fibrosis by cardiac magnetic resonance in patients with lamin A/C cardiomyopathy: possible substrate for diastolic dysfunction. *J Cardiovasc Magn Reson*. 2007;9(6):907-913. doi:10.1080/10976640701693733.
31. White JA, Fine NM, Gula L, Yee R, Skanes A, Klein G, et al. Utility of cardiovascular magnetic resonance in identifying substrate for malignant ventricular arrhythmias. *Circ Cardiovasc Imaging* 2012; 5: 12-20.
32. Inoue M, Kanaya H, Matsubara T, Uno Y, Yasuda T, Miwa K. Complete atrioventricular block associated with takotsubo cardiomyopathy. *Circ J*. 2009;73(3):589-592. <http://www.ncbi.nlm.nih.gov/pubmed/19075523>.
33. Shanmugasundaram R, Tamilarasu K, Rajendiran G, Murali A. An uncommon presentation of a rare disease – High-degree AV block with takotsubo cardiomyopathy. *Indian Heart J*. 2012;64(5):511-514.  
doi:10.1016/j.ihj.2012.07.013.

34. Bulluck H, Maestrini V, Rosmini S, et al. Myocardial T1 Mapping. *Circ J*. 2015;79(3):487-494. doi:10.1253/circj.CJ-15-0054.

## Tables

**Table 1. CMR Findings**

CMR Findings	n (%)
LVEF, %, median (IQR)	64 (58-68)
LVEDV, ml/m2, median (IQR)	88 (77-109)
LVESV, ml/m2, median (IQR)	31 (27-54)
LGE, n (%)	12 (35)
Ischemic Heart Disease, n (%)	4 (12)
Non-ischemic Heart Disease, n (%)	11 (32)
- Myocarditis, n	5
- Dilated cardiomyopathy, n	1
- Athlete's heart, n	2
- Aortic/pulmonary valve regurgitation, n	2
- Left ventricular non compaction, n	1
Structurally Normal Heart, n (%)	18 (53)

LVEF, left ventricular ejection fraction; LVEDV, left ventricular end-diastolic volume;

LVESV, left ventricular end-systolic volume; LGE, late gadolinium enhancement.

**Table 2. CMR findings vs standard clinical assessment**

		CMR				TOTAL
		IHD	NIHD	Non-specific Findings	Structurally Normal Heart	
Standard Clinical Assessment	IHD	<b>2</b>	1	0	1	4
	NIHD	2	<b>9</b>	0	9	20
	Non-specific Findings	0	1	<b>1</b>	7	9
	Other	0	0	0	1	1
	TOTAL	4	11	1	18	34

Comparison between pre-CMR working diagnosis based on the standard clinical assessment (history, ECG and cardiac rhythm monitoring) and CMR findings. Boxes in bold show the number of patients that received the same diagnosis before and after CMR.

CMR, cardiovascular magnetic resonance; IHD, ischemic heart disease; NIHD, non-ischemic heart disease.

**Table 3. Diagnostic performance of CMR and trans-thoracic echocardiography**

		CMR				
		IHD	NIHD	Non-specific Findings	Structurally Normal Heart	TOTAL
TTE	IHD	1	0	0	0	1
	NIHD	1	1	0	0	2
	Non-specific Findings	1	3	0	1	5
	Structurally Normal Heart	1	4	1	15	21
	TOTAL	4	8	1	16	29

Comparison between CMR and TTE findings. Boxes in bold show the number of patients that received the same diagnosis on both imaging modalities. CMR, cardiovascular magnetic resonance; TTE, trans-thoracic echocardiogram; IHD, ischemic heart disease; NIHD, non-ischemic heart disease.

## **Figure Legend**

### **Figure 1.**

Complete atrio-ventricular block (AVB) (A) in a patient with prominent mid-wall/epicardial late gadolinium enhancement in the basal inferoseptum (B, black arrow); a suspicion of cardiac sarcoidosis was raised and confirmed on laboratory testing.

Complete AVB (C) in a patient with lamin A/C dilated cardiomyopathy; on late gadolinium enhancement sequences, septal enhancement was noted (D, white arrow).