



Murphy, B., Roche-Newton, O., & Shkredov, I. (2017). Variations on the Sum-Product Problem II. *SIAM Journal on Discrete Mathematics*, 31(3), 1878-1894. <https://doi.org/10.1137/17M112316X>

Publisher's PDF, also known as Version of record

License (if available):  
Other

Link to published version (if available):  
[10.1137/17M112316X](https://doi.org/10.1137/17M112316X)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the final published version of the article (version of record). It first appeared online via Siam at <https://doi.org/10.1137/17M112316X> . Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/pure/about/ebr-terms>

## VARIATIONS ON THE SUM-PRODUCT PROBLEM II\*

BRENDAN MURPHY<sup>†</sup>, OLIVER ROCHE-NEWTON<sup>‡</sup>, AND ILYA D. SHKREDOV<sup>§</sup>

**Abstract.** This paper is a sequel to a paper entitled *Variations on the sum-product problem* by the same authors [*SIAM J. Discrete Math.*, 29 (2015), pp. 514–540]. In this sequel, we quantitatively improve several of the main results of the first paper as well as generalize a method from it to give a near-optimal bound for a new expander. The main new results are the following bounds, which hold for any finite set  $A \subset \mathbb{R}$ :  $\exists a \in A$  such that  $|A(A + a)| \gtrsim |A|^{\frac{3}{2} + \frac{1}{186}}$ ,  $|A(A - A)| \gtrsim |A|^{\frac{3}{2} + \frac{1}{34}}$ ,  $|A(A + A)| \gtrsim |A|^{\frac{3}{2} + \frac{5}{242}}$ ,  $|{(a_1 + a_2 + a_3 + a_4)^2 + \log a_5 : a_i \in A}| \gg \frac{|A|^2}{\log |A|}$ .

**Key words.** sum-product estimates, expanders, additive combinatorics, discrete geometry

**AMS subject classifications.** 52C10, 11B30, 11B75

**DOI.** 10.1137/17M112316X

**1. Introduction.** Throughout this paper, the standard notation  $\ll, \gg$  is applied to positive quantities in the usual way. Saying  $X \gg Y$  means that  $X \geq cY$  for some absolute constant  $c > 0$ . The notation  $X \approx Y$  denotes that  $X \gg Y$  and  $X \ll Y$  occur simultaneously. All logarithms in the paper are base 2. We use the symbols  $\lesssim, \gtrsim$  to suppress both constant and logarithmic factors. To be precise, we write  $X \gtrsim Y$  if there is some absolute constant  $c > 0$  such that  $X \gg Y/(\log X)^c$ .

This paper considers several variations on the sum-product problem, all of which follow a common theme. The story of the sum-product problem begins with the Erdős–Szemerédi conjecture, which states that for any finite  $A \subset \mathbb{Z}$  and for all  $\epsilon > 0$ ,

$$\max\{|A + A|, |AA|\} \geq c_\epsilon |A|^{2-\epsilon},$$

where

$$A + A := \{a + b : a, b \in A\}$$

is the *sum set* and

$$AA := \{ab : a, b \in A\}$$

is the *product set*. Although the conjecture was originally stated for subsets of the integers, it is also widely believed to be true over the reals.

\*Received by the editors March 28, 2017; accepted for publication (in revised form) June 23, 2017; published electronically August 30, 2017.

<http://www.siam.org/journals/sidma/31-3/M112316.html>

**Funding:** The second author is grateful for the hospitality of the Rényi Institute, where part of this work was conducted with support from grant ERC-AdG. 321104, for the support of the Austrian Science Fund FWF Project F5511-N26, which is part of the Special Research Program “Quasi-Monte Carlo Methods: Theory and Applications,” and also for the support of FWF Project P30405-N32.

<sup>†</sup>School of Mathematics, University of Bristol, Bristol, BS8 1TW, UK, and Heilbronn Institute of Mathematical Research, Bristol, UK (brendan.murphy@bristol.ac.uk).

<sup>‡</sup>Johannes Kepler Universität, Linz, Austria (o.rochenewton@gmail.com).

<sup>§</sup>Steklov Mathematical Institute, ul. Gubkina, 8, Moscow, 119991, Russia, IITP RAS, Bolshoy Karetny per. 19, Moscow, 127994, Russia, and MIPT, Institutskii per. 9, Dolgoprudnii, 141701, Russia (ilya.shkredov@gmail.com).

Modern literature on the sum-product problem often focuses on the problem of proving that certain sets defined by a combination of different arithmetic operations on elements of a set  $A$  are always significantly larger than  $|A|$ . Growth results of this type are often referred to as *expanders*.

In [7], the authors considered several expander problems. The aim of this sequel is to improve the main results from [7].

One result that was established in [7] is that, for any  $A \subset \mathbb{R}$ ,

$$(1.1) \quad |A(A + A)| \gtrsim |A|^{\frac{3}{2} + \frac{1}{178}}.$$

This result gave a small quantitative improvement on the inequality  $|A(A + A)| \gg |A|^{3/2}$ , which follows from a simple application of the Szemerédi–Trotter theorem (see [10, Lemma 3.2] for a formal proof).

The exponent  $3/2$  arises often as a threshold for what is achievable in sum-product type problems by using the Szemerédi–Trotter theorem in the simplest way that we know of, essentially copying the arguments of Elekes [1]. To improve on these fundamental results, refinements of these techniques have been required; see, for example, [6] and [11]. Most of these basic sum-product results with exponent  $3/2$  have by now been improved in the Euclidean setting, but some remain out of reach. For example, a simple application of the Szemerédi–Trotter theorem (see, for example, Exercise 8.3.3 in [16]) yields the bound  $|AA + A| \gg |A|^{3/2}$ , and no improvement is known.

The main new theorem that we prove in this paper is another example of a result that breaks the  $3/2$  threshold.

**THEOREM 1.1.** *Let  $A \subset \mathbb{R}$  be finite. Then, there exists  $a \in A$  such that*

$$|A(A + a)| \gtrsim |A|^{\frac{3}{2} + \frac{1}{186}}.$$

This improves on [7, Theorem 2.9], in which it was established that there is some  $a \in A$  such that  $|A(A + a)| \gg |A|^{3/2}$ .

**1.1. Outline of the proof of Theorem 1.1.** There are two main new lemmas which go into the proof of Theorem 1.1. The first of these is a lemma which states that there exists  $a \in A$  such that  $|A(A + a)| \geq |A|^{\frac{3}{2} + c}$ , where  $c > 0$  is an absolute constant, unless the multiplicative energy<sup>1</sup> of  $A$  is essentially as large as possible. This lemma is proved using the Szemerédi–Trotter theorem. In proving such a result, we improve qualitatively and quantitatively on some of the main lemmas from [7]. See the forthcoming Lemma 3.2.

The second new lemma (see the forthcoming Lemma 4.1 for a more precise statement) proves that if the product set of  $A$  is very small, then the bound  $|A(A + \alpha)| \gg |A|^{\frac{3}{2} + c}$  holds for any nonzero  $\alpha \in \mathbb{R}$ . The proof of this is a little more involved, using techniques from additive combinatorics, and is closely related to the work of the third author in [14]. A nontrivial result bounding the additive energy for sets with small product set, also due to the third author in [13], is used as a black box in the proof of this lemma. See the forthcoming Theorem 4.3 for the statement.

<sup>1</sup>The multiplicative energy of  $A$  is the number of solutions to the equation

$$a_1 a_2 = a_3 a_4, \quad (a_1, a_2, a_3, a_4) \in A.$$

See section 2 for more on the multiplicative energy and other types of energy.

The Balog–Szemerédi–Gowers theorem tells us that if a set  $A$  has large multiplicative energy, then there is a large subset  $A' \subset A$  such that  $|A'A'|$  is small. Therefore, one can then use the Balog–Szemerédi–Gowers theorem and combine these two lemmas together to conclude the proof. However, we instead use a technical argument, building on the work of Konyagin and Shkredov [3], which allows us to avoid an application of the Balog–Szemerédi–Gowers theorem and thus to improve the aforementioned constant  $c$ .

**1.2. Further new results.** The proof of inequality (1.1) in [7] followed a structure similar to the above sketch, and the Balog–Szemerédi–Gowers theorem was used to conclude the argument. Once again, we are able to make this argument more efficient by using tools from [3] to avoid using the Balog–Szemerédi–Gowers theorem, resulting in the following two results.

**THEOREM 1.2.** *Let  $A \subset \mathbb{R}$  be finite. Then,*

$$|A(A + A)| \gtrsim |A|^{\frac{3}{2} + \frac{5}{242}}.$$

**THEOREM 1.3.** *Let  $A \subset \mathbb{R}$  be finite. Then,*

$$|A(A - A)| \gtrsim |A|^{\frac{3}{2} + \frac{1}{34}}.$$

Theorem 1.2 gives an improvement on (1.1), while Theorem 1.3 gives an improvement on equation (49) in [7].

Another of the main results in [7] was the bound

$$(1.2) \quad |A(A + A + A + A)| \gg \frac{|A|^2}{\log |A|}.$$

Note that (1.2) is optimal, up to finding the correct power of the logarithmic factor, as can be seen by taking  $A = \{1, 2, \dots, N\}$ . In the last of our new theorems, we follow a similar argument to prove the following, admittedly curious, expander bound.

**THEOREM 1.4.** *Let  $A \subset \mathbb{R}^+$  be finite. Then,*

$$|\{(a_1 + a_2 + a_3 + a_4)^2 + \log a_5 : a_i \in A\}| \gg \frac{|A|^2}{\log |A|}.$$

Note that the simple example whereby  $A = \{1, 2, \dots, N\}$  illustrates that Theorem 1.4 is also optimal up to the logarithmic factor.

**1.3. The structure of the rest of this paper.** The rest of the paper will be structured as follows. Section 2 will be used to introduce some notation and preliminary results that will be used throughout the paper. As mentioned above, there are two main new lemmas in this paper. Section 3 is devoted to proving the first of these, and section 4 proves the second. Section 5 is used to conclude the proof of Theorem 1.1. In section 6 the proofs of Theorem 1.2 and 1.3 are concluded. Section 7 is devoted to the proof of Theorem 1.4.

**1.4. A note on an earlier preprint [9].** This paper supersedes the preprint [9] by the second author.

**2. Notation and preliminary results.** Given finite sets  $A, B \subset \mathbb{R}$ , the additive energy of  $A$  and  $B$  is the number of solutions to the equation

$$a_1 - b_1 = a_2 - b_2.$$

The additive energy is denoted  $E^+(A, B)$ . Let

$$r_{A-B}(x) := |\{(a, b) \in A \times B : a - b = x\}|.$$

Note that  $r_{A-B}(x) = |A \cap (B + x)|$ . The notation of the representation function  $r$  will be used with flexibility throughout this paper, with the information about the kind of representations it counts being contained in a subscript. For example,

$$r_{(A-A)^2+(A-A)^2}(x) = |\{(a_1, a_2, a_3, a_4) \in A^4 : (a_1 - a_2)^2 + (a_3 - a_4)^2 = x\}|.$$

Note that

$$E^+(A, B) = \sum_{x \in A-B} r_{A-B}^2(x).$$

The shorthand  $E^+(A) = E^+(A, A)$  is used.

The notion of energy can be extended to an arbitrary power  $k$ . We define  $E_k^+(A)$  by the formula

$$E_k^+(A) = \sum_{x \in A-A} r_{A-A}^k(x).$$

Similarly, the *multiplicative energy* of  $A$  and  $B$ , denoted  $E^\times(A, B)$ , is the number of solutions to the equation

$$\frac{a_1}{b_1} = \frac{a_2}{b_2}$$

such that  $a_1, a_2 \in A$  and  $b_1, b_2 \in B$ . For  $x \neq 0$ , let  $A_x$  denote the set  $A_x = A \cap x^{-1}A$  and note that  $r_{A/A}(x) = |A_x|$ .

A simple but important feature of energies is that the Cauchy–Schwarz inequality can be used to convert an upper bound for energy into a lower bound for the cardinality of a set. In particular, it follows from the Cauchy–Schwarz inequality that

$$(2.1) \quad E^\times(A, B) \geq \frac{|A|^2|B|^2}{|AB|}.$$

The Cauchy–Schwarz inequality can also be used to prove the bounds

$$(2.2) \quad E^\times(A, B) \leq (E^\times(A))^{1/2}(E^\times(B))^{1/2} \quad \text{and} \quad E^+(A, B) \leq (E^+(A))^{1/2}(E^+(B))^{1/2}.$$

The notions of additive and multiplicative energy have been central in the literature on sum-product estimates. For example, the key ingredient in the beautiful work of Solymosi [15], which until recently held the record for the best known sum-product estimate, is the following bound.

**THEOREM 2.1.** *For any finite  $A \subset \mathbb{R}$ ,*

$$E^\times(A) \ll |A + A|^2 \log |A|.$$

A major tool that is used in this paper several times, both explicitly and implicitly, is the Szemerédi–Trotter theorem. In particular, we will need the following result, which follows from a simple application of the Szemerédi–Trotter theorem. See, for example, Corollary 8.8 in [16].

LEMMA 2.2. *Let  $P \subset \mathbb{R}$  be a finite point set such that no line contains more than  $|P|^{1/2}$  points from  $P$ . Then there are  $O(|P|^2 \log |P|)$  collinear triples.*

*In particular, let  $A \subset \mathbb{R}$  be a finite set, and set  $P = A \times A$ . Then it follows that no line contains more than  $|A| = |P|^{1/2}$  points from  $P$ , and hence there are  $O(|A|^4 \log |A|)$  collinear triples in  $A \times A$ .*

In a recent paper of Konyagin and Shkredov [3], a new characteristic for a finite set of real numbers  $A$  was considered. Define  $d_*(A)$  by the formula

$$d_*(A) = \min_{t>0} \min_{\emptyset \neq Q, R \subset \mathbb{R} \setminus \{0\}} \frac{|Q|^2 |R|^2}{|A| t^3},$$

where the second minimum is taken over all  $Q$  and  $R$  such that  $\max\{|Q|, |R|\} \geq |A|$  and such that for every  $a \in A$ , the bound  $|Q \cap aR^{-1}| \geq t$  holds. Konyagin and Shkredov proved the following lemma.

LEMMA 2.3 (Lemma 13, [3]). *For any  $A, B \subset \mathbb{R}$  and any  $\tau \geq 1$*

$$(2.3) \quad |\{x : r_{A-B}(x) \geq \tau\}| \ll \frac{|A||B|^2}{\tau^3} d_*(A).$$

Using the language introduced in [12] and also used in [3],  $A$  is said to be a Szemerédi–Trotter set with parameter  $M$  if the bound  $|\{x : r_{A-B}(x) \geq \tau\}| \leq \frac{|A||B|^2}{\tau^3} M$  holds for any  $B \subset \mathbb{R}$ . Using this terminology, Lemma 2.3 states that every set  $A$  is a Szemerédi–Trotter set with parameter  $O(d_*(A))$ . The main theoretical tool in the proof of Lemma 2.3 is the Szemerédi–Trotter theorem. Lemma 2.3 generalizes an earlier result in which the bound

$$(2.4) \quad |\{x : r_{A-B}(x) \geq \tau\}| \ll \frac{|A||B|^2}{\tau^3} d(A)$$

was established, where  $d(A) = \min_{C \neq \emptyset} \frac{|AC|^2}{|A||C|}$ . See [8, Lemma 7] for a proof. As pointed out in [3],  $d_*(A) \leq d(A)$ , since for any nonempty  $C$  we can take  $t = |C|$ ,  $Q = AC$ , and  $R = C^{-1}$  in the definition of  $d_*(A)$ .

We conclude this section by attempting to give some heuristic behind the meaning of these technical parameters  $d(A)$  and  $d_*(A)$ . Roughly speaking, these parameters give some measure of how multiplicatively structured  $A$  is; if these parameters are small, then the set  $A$  has some multiplicative structure. One classical measure of this vague property in additive combinatorics is the *doubling constant*  $K = K(A)$ , where  $K = \frac{|AA|}{|A|}$ . Note that

$$(2.5) \quad d_*(A) \leq d(A) \leq K^2.$$

The theory of sum-product estimates tells us that if  $K$  is small, then  $A + A$  is large. Estimates (2.3) and (2.4) give us similar information if  $d_*(A)$  and  $d(A)$ , respectively, are small. The advantage of working with the parameters  $d(A)$  and  $d_*(A)$  instead of the doubling constant is that, as (2.5) indicates, the property of these parameters being small is more frequently attained.

**3. A bound on sums of multiplicative energies with shifts.** The aim of this section is to prove the first of the two main new lemmas of this paper. This is the

following lemma, which gives an improvement of Lemma 2.4 in [7], in the case when the sets involved are approximately the same size, unless the multiplicative energy is essentially as large as possible.

LEMMA 3.1. *Let  $A, B, C \subset \mathbb{R}$  be a finite sets such that  $|A| \approx |C|$ . Then*

$$\sum_{a \in A} E^\times(B, C - a) \ll E^\times(B)^{1/2} |A|^2 \log^{1/2} |A| + |A|^3 + |A||B|^2.$$

*Proof.* We have

$$\begin{aligned} \sum_{a \in A} E^\times(B, C - a) &= |\{(a, b, b', c, c') \in A \times B \times B \times C \times C : b(c - a) = b'(c' - a)\}| \\ &\leq |\{(a, b, b', c, c') \in A \times B \times B \times C \times C : b(c - a) = b'(c' - a) \neq 0\}| \\ &\quad + |A|^3 + |A|^2|B| + |A||B|^2 \\ &\leq \left| \left\{ (a, b, b', c, c') \in A \times B \times B \times C \times C, : \frac{b}{b'} = \frac{c' - a}{c - a} \neq 0 \right\} \right| \\ &\quad + |A|^3 + E^\times(B)^{1/2} |A|^2 + |A||B|^2. \end{aligned}$$

The remaining task is to bound the main term. Applying the Cauchy–Schwarz inequality yields

$$\begin{aligned} &\left| \left\{ (a, b, b', c, c') \in A \times B \times B \times C \times C : \frac{b}{b'} = \frac{c' - a}{c - a} \neq 0 \right\} \right| \\ &= \sum_{x \neq 0} r_{B/B}(x) n(x) \\ &\leq \left( \sum_x r_{B/B}^2(x) \right)^{1/2} \left( \sum_{x \neq 0} n^2(x) \right)^{1/2} \\ (3.1) \quad &= E^\times(B)^{1/2} \left( \sum_{x \neq 0} n^2(x) \right)^{1/2}, \end{aligned}$$

where

$$n(x) = \left| \left\{ (a, c, c') \in A \times C \times C : x = \frac{c' - a}{c - a} \right\} \right|.$$

Note that

$$\begin{aligned} \sum_x n^2(x) &\leq \left| \left\{ (a_1, a_2, c_1, c_2, c'_1, c'_2) : \frac{c'_1 - a_1}{c_1 - a_1} = \frac{c'_2 - a_2}{c_2 - a_2} \right\} \right| \\ &= \left| \left\{ (a_1, a_2, c_1, c_2, c'_1, c'_2) : \frac{c_2 - a_2}{c_1 - a_1} = \frac{c'_2 - a_2}{c'_1 - a_1} \right\} \right|. \quad \square \end{aligned}$$

The identity  $\frac{c_2 - a_2}{c_1 - a_1} = \frac{c'_2 - a_2}{c'_1 - a_1}$  occurs only if the three points  $(a_1, a_2), (c_1, c_2), (c'_1, c'_2) \in (A \cup C) \times (A \cup C)$  are collinear. By Lemma 2.2, there are  $O(|A \cup C|^4 \log |A|)$  such collinear triples, and so

$$\sum_x n^2(x) \ll |A|^4 \log |A|.$$

Combining this with (3.1), we have

$$\sum_{a \in A} E^\times(B, C - a) \ll E^\times(B)^{1/2} |A|^2 \log^{1/2} |A| + |A|^3 + |A||B|^2.$$

COROLLARY 3.2. *Let  $A \subset \mathbb{R}$  be finite. Then there exists  $a \in A$  such that*

$$(3.2) \quad E^\times(A) |A(A + a)|^2 \gg \frac{|A|^6}{\log |A|}.$$

Similarly, there exists  $b \in A$  such that

$$(3.3) \quad E^\times(A) |A(A - b)|^2 \gg \frac{|A|^6}{\log |A|}.$$

*Proof.* By Lemma 3.1, we have

$$\sum_{a \in A} E^\times(A, A + a) \ll E^\times(A)^{1/2} |A|^2 \log^{1/2} |A| + |A|^3 \ll E^\times(A)^{1/2} |A|^2 \log^{1/2} |A|.$$

Therefore, by the pigeonhole principle, there exists  $a \in A$  such that

$$E^\times(A, A + a) \ll E^\times(A)^{1/2} |A| \log^{1/2} |A|.$$

By the Cauchy–Schwarz inequality

$$\frac{|A|^4}{|A(A + a)|} \leq E^\times(A, A + a) \ll E^\times(A)^{1/2} |A| \log^{1/2} |A|.$$

A rearrangement of this inequality completes the proof. The proof of (3.3) is essentially identical.  $\square$

Corollary 3.2 provides a pinned version of the following result, which was the main lemma from [7].

LEMMA 3.3. *For any finite sets  $A, B, C \in \mathbb{R}$ ,*

$$E^\times(A) |A(B + C)|^2 \gg \frac{|A|^4 |B| |C|}{\log |A|}.$$

**4. A conditional lower bound on  $|A(A + \alpha)|$ .** The second main new lemma of this paper is the following.

LEMMA 4.1. *Let  $A \subset \mathbb{R}$  and  $\alpha \in \mathbb{R} \setminus \{0\}$ . Then*

$$|A(A + \alpha)| \gtrsim \frac{E^\times(A)^2}{|A|^{\frac{58}{13}} d_*^{\frac{7}{13}}(A)}.$$

In particular, this result implies the following statement:<sup>2</sup>

$$|AA| \leq M|A| \Rightarrow |A(A + \alpha)| \gg_M |A|^{\frac{3}{2} + c}.$$

<sup>2</sup>The notation  $X \gg_M Y$  means that  $X \geq cY$ , where the value  $c > 0$  depends on a parameter  $M$ .



Indeed, since by the Cauchy–Schwarz inequality  $E^\times(A) \geq |A|^3/M$  and since  $d_*(A) \leq d(A) \leq M^2$ , we have

$$|A(A + \alpha)| \gtrsim \frac{|A|^{\frac{20}{13}}}{M^{\frac{40}{13}}}.$$

In the proof of Lemma 4.1 we will need the following simple lemma. See [14, Lemma 4].

LEMMA 4.2. *Let  $G$  be an abelian group and  $A \subset G$  be a finite set. Then there is  $z$  such that*

$$\sum_{x \in zA} |(zA) \cap x(zA)| \gg \frac{E^\times(A)}{|A| \log |A|}.$$

We will also need the following result of the third author [13], which tells us that a set  $A$  such that the parameter  $d_*(A)$  is small has small additive energy. One can obtain similar but weaker results using the Szemerédi–Trotter theorem in a more elementary way (see, for example, Corollary 8 in [8]), but the important thing for our application of this result is that the exponent  $32/13$  is smaller than  $5/2$ .

THEOREM 4.3 ([12], Theorem 5.4). *Let  $A \subset \mathbb{R}$  be a finite set. Then*

$$E^+(A) \ll d_*(A)^{7/13} |A|^{32/13} \log^{71/65} |A|.$$

The result in [13] did not use the quantity  $d_*(A)$  because it was introduced later in [4], but one can check that the arguments of [13] work for any Szemerédi–Trotter set. The only fact we need is an upper bound for the size of a set  $\{x : r_{A-B}(x) \geq \tau\}$ , so Lemma 2.3 is enough for us.

*Proof of Lemma 4.1.* Without loss of generality, we may assume that  $0 \notin A$ . Applying Lemma 4.2 and writing  $B = zA$ , we have

$$(4.1) \quad \sum_{\lambda \in B} |B \cap \lambda^{-1}B| \gg \frac{E^\times(A)}{|A| \log |A|}.$$

Next we double-count the number of solutions to the equation

$$(4.2) \quad b_1(b'_1 + \alpha z) = b_2(b'_2 + \alpha z)$$

such that  $b_1, b_2 \in B$ ,  $b'_1 \in B_{b_1}$ , and  $b'_2 \in B_{b_2}$ . We recall here that  $B_x := B \cap x^{-1}B$ .

Let  $S$  denote the number of solutions to (4.2). Suppose that we have such a solution. Then  $b'_1 = b''_1/b_1$  and  $b'_2 = b''_2/b_2$  for some  $b''_1, b''_2 \in B$ . Therefore,

$$\alpha z b_1 + b''_1 = \alpha z b_2 + b''_2,$$

and it follows that  $S \leq E^+(B, \alpha z B)$ . An application of (2.2) then gives  $E^+(B, \alpha z B) \leq E^+(B)^{1/2} E^+(\alpha z B)^{1/2} = E^+(B) = E^+(A)$ . So, by Theorem 4.3

$$(4.3) \quad S \leq E^+(A) \lesssim d_*(A)^{7/13} |A|^{32/13}.$$

On the other hand, denote

$$m(t) = |\{(b, b') \in B \times B_b : b(b' + \alpha z) = t\}|$$

and note that  $S = \sum_t m^2(t)$ . Also,  $m(t) > 0$  only if  $t \in B(B + \alpha z)$ . Then, by (4.1), the Cauchy–Schwarz inequality, and (4.3)

$$(4.4) \quad \frac{E^\times(A)^2}{|A|^2 \log^2 |A|} \ll \left( \sum_{\lambda \in B} |B \cap \lambda^{-1} B| \right)^2$$

$$(4.5) \quad = \left( \sum_t m(t) \right)^2$$

$$(4.6) \quad \leq |B(B + \alpha z)| \sum_t m^2(t)$$

$$(4.7) \quad \lesssim |B(B + \alpha z)| d_*(A)^{7/13} |A|^{32/13}.$$

Finally, note that  $|B(B + \alpha z)| = |A(A + \alpha)|$ . We conclude that

$$|A(A + \alpha)| \gtrsim \frac{E^\times(A)^2}{|A|^{58/13} d_*(A)^{7/13}}. \quad \square$$

Actually, one can see that we have proved the inequality  $|A|^2 |A(A + \alpha)| E^+(A) \gtrsim (E^\times(A))^2$  for any finite subset of reals.

**5. Proof of Theorem 1.1.** Before proving Theorem 1.1, we need one more lemma.

LEMMA 5.1. *Let  $A \subset \mathbb{R}$  and suppose that  $E^\times(A) \geq \frac{|A|^3}{K}$ . Then for any  $\alpha \in \mathbb{R} \setminus \{0\}$*

$$|A(A + \alpha)| \gtrsim \frac{|A|^{20/13}}{K^{40/13}}.$$

*Proof.* We claim that for any  $A \subset \mathbb{R}$  such that  $E^\times(A) \geq \frac{|A|^3}{K}$  there is a subset  $A' \subseteq A$  such that

$$(5.1) \quad E^\times(A') \gtrsim E^\times(A) \geq \frac{|A|^3}{K},$$

$$(5.2) \quad d_*(A') \lesssim \frac{K^2 |A|}{|A'|}.$$

Given such a subset  $A'$ , we may apply Lemma 4.1 to find that

$$\begin{aligned} |A(A + \alpha)| \geq |A'(A' + \alpha)| &\gtrsim \frac{E^\times(A')^2}{|A'|^{58/13} d_*^{7/13}(A')} \\ &\gtrsim \frac{|A|^6}{K^2 |A'|^{58/13}} \cdot \frac{|A'|^{7/13}}{|A|^{7/13} K^{14/13}} \\ &= \frac{|A|^{71/13}}{K^{40/13} |A'|^{51/13}} \geq \frac{|A|^{20/13}}{K^{40/13}}. \end{aligned}$$

It remains to prove (5.1) and (5.2). By the popularity principle and dyadic pigeonholing there is a subset  $P \subseteq A/A$  and a number  $\Delta \geq |A|/2K$  such that for all  $x$  in  $P$

$$(5.3) \quad \Delta \leq |A \cap xA| < 2\Delta,$$

and

$$(5.4) \quad \sum_{x \in P} |A \cap xA|^2 \gtrsim E^\times(A).$$

Now we perform an additional refinement step. Let  $A' \subseteq A$  denote the set of  $x$  such that

$$|P \cap xA^{-1}| \geq \frac{\Delta|P|}{4|A|}.$$

Since

$$(5.5) \quad \sum_{x \in A} |P \cap xA^{-1}| = \sum_{x \in P} |A \cap xA| \geq \Delta|P|,$$

by the popularity principle we have

$$\sum_{x \in A'} |P \cap xA^{-1}| \geq \frac{3\Delta|P|}{4}.$$

If  $x \notin A'$ , then

$$|P \cap x(A')^{-1}| \leq |P \cap xA^{-1}| < \frac{\Delta|P|}{4|A|}.$$

Thus

$$\frac{3\Delta|P|}{4} \leq \sum_{x \in A'} |P \cap xA^{-1}| = \sum_{x \in A} |P \cap x(A')^{-1}| \leq \frac{\Delta|P|}{4} + \sum_{x \in A'} |P \cap x(A')^{-1}|,$$

which yields

$$\frac{\Delta|P|}{2} \leq \sum_{x \in A'} |P \cap x(A')^{-1}| = \sum_{x \in P} |A' \cap xA'|.$$

By Cauchy–Schwarz, we have

$$E^\times(A') \gg \Delta^2|P| \gtrsim E^\times(A).$$

Setting  $Q = P$ ,  $R = A$ , and  $t = (\Delta|P|)/(4|A|)$  in the definition of the quantity  $d_*(A')$ , we obtain

$$d_*(A') \ll \frac{|P|^2|A|^2}{\left(\frac{\Delta|P|}{4|A|}\right)^3|A'|} \ll \frac{|A|^5}{|A'| |P| \Delta^3} \lesssim \frac{|A|^5}{|A'| E^\times(A) \Delta} \ll \frac{K^2|A|}{|A'|},$$

where the last inequality follows from the lower bounds for  $E^\times(A)$  and  $\Delta$ . □

*Proof of Theorem 1.1.* Write  $E^\times(A) = \frac{|A|^3}{K}$ . By Corollary 3.2 there is some  $a \in A$  such that

$$(5.6) \quad |A(A+a)| \gg K^{1/2}|A|^{3/2} \log^{-1/2}|A| \gtrsim K^{1/2}|A|^{3/2}.$$

On the other hand, for any  $a \in A \setminus \{0\}$ , Lemma 5.1 implies that

$$(5.7) \quad |A(A+a)| \gtrsim \frac{|A|^{20/13}}{K^{40/13}}.$$

Optimizing over (5.6) and (5.7), we obtain

$$|A(A+a)| \gtrsim |A|^{3/2+1/186}$$

as required. □

In fact, by taking more care with the pigeonholing argument in the proof of Corollary 3.2 it follows that the bound  $|A(A+a)| \gtrsim |A|^{3/2+1/186}$  holds for at least half of the elements  $a \in A$ .

**6. Three variable expanders.** The proofs of Theorems 1.2 and 1.3 follow an argument similar to that of the proof of Theorem 1.1. We can use Corollary 3.2 to get an exponent better than  $3/2$  in the case when  $E^\times(A)$  is not too large. However, in the case when  $E^\times(A)$  is large, we need analogues of Lemma 5.1 that are quantitatively better for the purposes of these problems. These bounds are given by the following lemma.

LEMMA 6.1. *Let  $A \subset \mathbb{R}$  and suppose that  $E^\times(A) \geq \frac{|A|^3}{K}$ . Then*

$$(6.1) \quad |A - A| \gtrsim \frac{|A|^{8/5}}{K^{6/5}}$$

and

$$(6.2) \quad |A + A| \gtrsim \frac{|A|^{58/37}}{K^{42/37}}.$$

In order to prove (6.1), we will need the following lemma.

LEMMA 6.2. *For any finite set  $A \subset \mathbb{R}$ ,*

$$|A - A| \gg \frac{|A|^{8/5}}{d_*^{3/5}(A) \log^{2/5} |A|}.$$

Although this result has not appeared explicitly in the literature, it can be proved by essentially copying the arguments from [8] and predecessors with the stronger Lemma 2.3 in place of the bound (2.4). The proof is included in the appendix for completeness.

The following similar result for sum sets follows from a combination of the work in [12] and [3].

LEMMA 6.3. *For any finite set  $A \subset \mathbb{R}$ ,*

$$|A + A| \gtrsim \frac{|A|^{58/37}}{d_*^{21/37}(A)}.$$

To be more precise, it was proven in [12] that if  $A$  is a Szemerédi–Trotter set with parameter  $D(A)$ , then  $|A + A| \gtrsim \frac{|A|^{58/37}}{D^{21/37}(A)}$ , and it was subsequently established in [3] that any set  $A$  is a Szemerédi–Trotter set with  $O(d_*(A))$ .

In addition, we need the following lemma, which uses the hypothesis that the energy is large in order to find a large subset  $A' \subset A$  such that  $d_*(A)$  is small.

LEMMA 6.4 (double pigeonholing argument). *Let  $A \subset \mathbb{R}$  and suppose that  $E^\times(A) \geq \frac{|A|^3}{K}$ . Then there is a subset  $A' \subseteq A$  and a number  $\Delta \gg |A|/K$  such that*

$$(6.3) \quad |A'| \gtrsim \frac{|A|^2}{K\Delta},$$

$$(6.4) \quad d_*(A') \lesssim \frac{K|A'|^2}{|A|\Delta}.$$

The proof of Lemma 6.4 is similar to the refinement step in the proof of Lemma 5.1.

*Proof.* As in (5.3) and (5.4), by the popularity principle and dyadic pigeonholing there is a subset  $P \subseteq A/A$  and a number  $\Delta \geq |A|/2K$  such that for all  $x$  in  $P$

$$\Delta \leq |A \cap xA| < 2\Delta$$

and

$$\sum_{x \in P} |A \cap xA|^2 \gtrsim E^\times(A).$$

Now we perform a second dyadic pigeonholing argument. As in (5.5) we have

$$\sum_{a \in A} |A \cap aP| = \sum_{x \in P} |A \cap xA| \geq |P|\Delta.$$

Thus there exist a subset  $A' \subseteq A$  and a number  $0 < t \leq |A|$  such that for all  $a$  in  $A'$

$$t \leq |A \cap aP| < 2t$$

and

$$\sum_{a \in A'} |A \cap aP| \gtrsim |P|\Delta;$$

hence

$$|A'|t \gtrsim |P|\Delta.$$

Since  $t \leq |A|$  and  $|P|\Delta^2 \gtrsim |A|^3/K$  we have

$$|A'| \gtrsim \frac{|P|\Delta}{|A|} \gtrsim \frac{|A|^2}{K\Delta},$$

which proves (6.3).

For every  $a \in A'$  we have  $|A \cap aP| \geq t$ . Therefore, we can take

$$t = t, \quad Q = A, \quad \text{and} \quad R = P^{-1}$$

in the definition of  $d_*(A')$ . We then have

$$\begin{aligned} d_*(A') &\leq \frac{|A|^2|P|^2}{|A'|t^3} \\ &\lesssim \frac{|A|^2|P|^2}{|A'|t^3} \cdot \left(\frac{|A'|t}{|P|\Delta}\right)^3 \\ &= \frac{|A|^2|A'|^2}{|P|\Delta^3} = \frac{|A|^2|A'|^2}{(|P|\Delta^2)\Delta} \lesssim \frac{K|A'|^2}{|A|\Delta}, \end{aligned}$$

which proves (6.4). □

*Proof of Lemma 6.1.* Similar to the proof of Lemma 5.1, the idea here is to use the double pigeonholing argument (Lemma 6.4) to find a large subset  $A' \subset A$  such that  $d_*(A)$  is small, and to then apply Lemmas 6.2 and 6.3 to complete the proof.

Since  $E^\times(A) \geq |A|^3/K$ , by Lemma 6.4 there is a subset  $A' \subseteq A$  and a number  $\Delta \gg |A|/K$  such that

$$\begin{aligned} |A'| &\gtrsim \frac{|A|^2}{K\Delta}, \\ d_*(A') &\lesssim \frac{K|A'|^2}{|A|\Delta}. \end{aligned}$$

Applying Lemma 6.2 yields

$$\begin{aligned} |A - A| &\geq |A' - A'| \gtrsim \frac{|A'|^{8/5}}{d_*(A')^{3/5}} \\ &\gtrsim |A'|^{8/5} \cdot \frac{|A|^{3/5} \Delta^{3/5}}{|A'|^{6/5} K^{3/5}} = \frac{|A'|^{2/5} |A|^{3/5} \Delta^{3/5}}{K^{3/5}} \\ &\gtrsim \frac{|A|^{4/5}}{K^{2/5} \Delta^{2/5}} \cdot \frac{|A|^{3/5} \Delta^{3/5}}{K^{3/5}} = \frac{|A|^{7/5}}{K} \Delta^{1/5} \\ &\gg \frac{|A|^{8/5}}{K^{6/5}}. \end{aligned}$$

Applying Lemma 6.3 yields

$$\begin{aligned} |A + A| &\geq |A' + A'| \gtrsim \frac{|A'|^{58/37}}{d_*^{21/37}(A')} \\ &\gtrsim |A'|^{58/37} \cdot \frac{|A|^{21/37} \Delta^{21/37}}{|A'|^{42/37} K^{21/37}} = \frac{|A'|^{16/37} |A|^{21/37} \Delta^{21/37}}{K^{21/37}} \\ &\gtrsim \frac{|A|^{32/37}}{K^{16/37} \Delta^{16/37}} \cdot \frac{|A|^{21/37} \Delta^{21/37}}{K^{21/37}} = \frac{|A|^{53/37}}{K} \Delta^{5/37} \\ &\gg \frac{|A|^{58/37}}{K^{42/37}}. \quad \square \end{aligned}$$

We are now ready to prove the new lower bounds for  $A(A - A)$  and  $A(A + A)$ .

*Proof of Theorem 1.2.* Write  $E^\times(A) = \frac{|A|^3}{K}$ . By Corollary 3.2

$$(6.5) \quad |A(A + A)| \gg K^{1/2} |A|^{3/2} \log^{-1/2} |A| \gtrsim K^{1/2} |A|^{3/2}.$$

On the other hand, Lemma 6.1 implies that

$$(6.6) \quad |A(A + A)| \geq |A + A| \gtrsim \frac{|A|^{58/37}}{K^{42/37}}.$$

Optimizing over (6.5) and (6.6), we obtain

$$|A(A + A)| \gtrsim |A|^{3/2+5/242}$$

as required.  $\square$

*Proof of Theorem 1.3.* Write  $E^\times(A) = \frac{|A|^3}{K}$ . By Corollary 3.2

$$(6.7) \quad |A(A - A)| \gg K^{1/2} |A|^{3/2} \log^{-1/2} |A| \gtrsim K^{1/2} |A|^{3/2}.$$

On the other hand, Lemma 6.1 implies that

$$(6.8) \quad |A(A - A)| \geq |A - A| \gtrsim \frac{|A|^{8/5}}{K^{6/5}}.$$

Optimizing over (6.7) and (6.8), we obtain

$$|A(A - A)| \gtrsim |A|^{3/2+1/34}$$

as required. □

**7. Five variable expander.** In this section we will prove Theorem 1.4 based on the proof in [7] of the inequality

$$(7.1) \quad |A(A + A + A + A)| \gg \frac{|A|^2}{\log |A|}.$$

The proof of (7.1) in [7] follows from comparing the upper bound on the multiplicative energy in Theorem 2.1 with the lower bound in Lemma 3.3. Here, we need a suitable analogue of Lemma 3.3, the proof of which relies on the following celebrated result of Guth and Katz [2].

**THEOREM 7.1.** *For any finite set  $A \subset \mathbb{R}$ , the number of solutions to the equation*

$$(a_1 - a_2)^2 + (a_3 - a_4)^4 = (a_5 - a_6)^2 + (a_7 - a_8)^2$$

*such that  $a_1, \dots, a_8 \in A$  is at most  $O(|A|^6 \log |A|)$ .*

Theorem 7.1 is a special case of a more general geometric result which immediately implies a resolution of the Erdős distinct distances problem up to logarithmic factors, but here it is stated only in the form in which it will be used in this paper.

Theorem 7.1 can be used to prove the following variation of Lemma 3.3.

**LEMMA 7.2.** *For any finite sets  $A, B \subset \mathbb{R}$ ,*

$$E^+(A) |\{a + (b_1 + b_2)^2 : a \in A, b_1, b_2 \in B\}|^2 \gg \frac{|A|^4 |B|^2}{\log |B|}.$$

*Proof.* The proof proceeds by the familiar method of double-counting the number of solutions to the equation

$$(7.2) \quad a_1 + (b_1 + b_2)^2 = a_2 + (b_3 + b_4)^2$$

such that  $a_i \in A$  and  $b_i \in B$ . Let  $S$  denote the number of solutions to (7.2) and write

$$A + (B + B)^2 := \{a + (b_1 + b_2)^2 : a \in A, b_1, b_2 \in B\}.$$

By the Cauchy–Schwarz inequality

$$S \geq \frac{|A|^2 |B|^4}{|A + (B + B)^2|}.$$

On the other hand, also by the Cauchy–Schwarz inequality,

$$\begin{aligned} S^2 &= \left( \sum_x r_{A-A}(x) r_{(B+B)^2-(B+B)^2}(x) \right)^2 \\ &\leq \left( \sum_x r_{A-A}^2(x) \right) \left( \sum_x r_{(B+B)^2-(B+B)^2}^2(x) \right) \\ &= E^+(A) \left( \sum_x r_{(B+B)^2-(B+B)^2}^2(x) \right). \end{aligned}$$

Theorem 7.1 tells us that  $(\sum_x r_{(B+B)^2-(B+B)^2}^2(x)) = O(|B|^6 \log |B|)$ . Therefore,

$$\begin{aligned} |A|^4 |B|^8 &\leq |A + (B + B)^2|^2 S^2 \\ &\ll |A + (B + B)^2|^2 E^+(A) |B|^6 \log |B|. \end{aligned}$$

After rearranging this inequality, we obtain the desired result.  $\square$

Unfortunately, we are not aware of a proof of Lemma 7.2 which does not use the deep results from [2].

We are now ready to prove Theorem 1.4.

*Proof of Theorem 1.4.* Apply Lemma 7.2 with  $A = \log A$  and  $B = A + A$ . We have

$$E^+(\log A) |\{(a_1 + a_2 + a_3 + a_4)^2 + \log a_5 : a_i \in A\}|^2 \gg \frac{|A|^4 |A + A|^2}{\log |A|}.$$

Note that  $\log a_1 + \log a_2 = \log a_3 + \log a_4$  if and only if  $a_1 a_2 = a_3 a_4$ , and so  $E^+(\log A) = E^\times(A)$ . We can apply Theorem 2.1 to deduce that

$$E^+(\log A) \ll |A + A|^2 \log |A|.$$

It then follows that

$$|\{(a_1 + a_2 + a_3 + a_4)^2 + \log a_5 : a_i \in A\}|^2 \gg \frac{|A|^4}{\log^2 |A|},$$

which completes the proof.  $\square$

**Appendix.** The purpose of this appendix is to present a formal proof of Lemma 6.2. We will call upon the following result on the relationship between different types of energy.

LEMMA 7.3 ([5], Lemma 2.4 and 2.5). *For any finite sets  $A, B \subset \mathbb{R}$*

$$|A|^2 (E_{1.5}^+(A))^2 \leq (E_3^+(A))^{2/3} (E_3^+(B))^{1/3} E(A, A - B).$$

In fact, Lemma 7.3 holds for any abelian group.

*Proof of Lemma 6.2.* Recall that Lemma 6.2 states that for any finite set  $A \subset \mathbb{R}$ ,

$$|A - A| \gg \frac{|A|^{8/5}}{d_*^{3/5}(A) \log^{2/5} |A|}.$$



In order to prove this, we will first prove two energy bounds. Note that, by Lemma 2.3,

$$\begin{aligned}
 E_3^+(A) &= \sum_x r_{A-A}^3(x) \\
 (7.3) \quad &= \sum_{j \geq 1} \sum_{x: 2^{j-1} \leq r_{A-A}(x) < 2^j} r_{A-A}^3(x) \\
 &\ll |A|^3 d_*(A) \log |A|.
 \end{aligned}$$

Similarly, for any  $F \subset \mathbb{R}$  and a parameter  $\Delta > 0$

$$\begin{aligned}
 E^+(A, F) &= \sum_x r_{A-F}^2(x) \\
 (7.4) \quad &= \sum_{x: r_{A-F}(x) \leq \Delta} r_{A-F}^2(x) + \sum_{j \geq 1} \sum_{x: 2^{j-1} \Delta \leq r_{A-A}(x) < 2^j \Delta} r_{A-F}^2(x) \\
 &\ll \Delta |A| |F| + \frac{|A| |F|^2 d_*(A)}{\Delta}.
 \end{aligned}$$

We choose  $\Delta = (|F| d_*(A))^{1/2}$  and thus conclude that

$$(7.5) \quad E(A, F) \ll |A| |F|^{3/2} d_*(A)^{1/2}.$$

Now, by Hölder’s inequality

$$\begin{aligned}
 |A|^6 &= \left( \sum_{x \in A-A} r_{A-A}(x) \right)^3 \\
 (7.6) \quad &\leq |A - A| \left( \sum_x r_{A-A}^{3/2}(x) \right)^2 \\
 &= |A - A| (E_{1.5}^+(A))^2.
 \end{aligned}$$

Applying (7.6) with Lemma 7.3, as well as inequalities (7.3) and (7.5), we have

$$\begin{aligned}
 |A|^8 &\leq |A - A| (E_{1.5}^+(A))^2 |A|^2 \\
 &\leq |A - A| E_3^+(A) E(A, A - A) \\
 &= |A - A|^{5/2} |A|^4 d_*(A)^{3/2} \log |A|.
 \end{aligned}$$

Rearranging this inequality completes the proof. □

Finally, we note that a similar method can be used to prove a quantitatively weaker version of Lemma 6.3, in the form of the following result:

$$(7.7) \quad |A + A| \gg \frac{|A|^{14/9}}{d_*^{5/9}(A) \log^{2/9} |A|}.$$

To see how this works, one can repeat the arguments from the proof of Theorem 1.2 in [6], but using Lemma 2.3 in place of Lemma 3.2 from [6]. This is worth noting, since the proofs of the main results in [3] and [4], that is, the bound

$$\max\{|A + A|, |AA|\} \gg |A|^{4/3+c}$$

for some  $c > 0$ , both include applications of Lemma 6.3. One can also obtain this sum-product estimate, albeit with a smaller positive value  $c$ , by using the bound (7.7) instead of Lemma 6.3.

**Acknowledgments.** We thank Antal Balog, Orit Raz, and Endre Szemerédi for helpful discussions.

## REFERENCES

- [1] G. ELEKES, *On the number of sums and products*, Acta Arith., 81 (1997), pp. 365–367.
- [2] L. GUTH AND N. H. KATZ, *On the Erdős distinct distance problem in the plane*, Ann. of Math. (2), 181 (2015), pp. 155–190.
- [3] S. V. KONYAGIN AND I. D. SHKREDOV, *New results on sums and products in  $\mathbb{R}$* , Proc. Steklov Inst. Math., 294 (2016), pp. 78–88.
- [4] S. V. KONYAGIN AND I. D. SHKREDOV, *On sum sets of sets, having small product set*, Proc. Steklov Inst. Math., 290 (2015), pp. 288–299.
- [5] L. LI, *On a Theorem of Schoen and Shkredov on Sumsets of Convex Sets*, preprint, <https://arxiv.org/abs/1108.4382>, 2011.
- [6] L. LI AND O. ROCHE-NEWTON, *Convexity and a sum-product type estimate*, Acta Arith., 156 (2012), pp. 247–255.
- [7] B. MURPHY, O. ROCHE-NEWTON, AND I. D. SHKREDOV, *Variations on the sum-product problem*, SIAM J. Discrete Math., 29 (2015), pp. 514–540, <https://doi.org/10.1137/140952004>.
- [8] O. RAZ, O. ROCHE-NEWTON, AND M. SHARIR, *Sets with few distinct distances do not have heavy lines*, Discrete Math., 338 (2015), pp. 1484–1492.
- [9] O. ROCHE-NEWTON, *A New Expander and Improved Bounds for  $A(A + A)$* , preprint, <https://arxiv.org/abs/1603.06827>, 2016.
- [10] O. ROCHE-NEWTON AND D. ZHELEZOV, *A bound on the multiplicative energy of a sum set and extremal sum-product problems*, Mosc. J. Comb. Number Theory, 5 (2015), pp. 52–69.
- [11] T. SCHOEN AND I. D. SHKREDOV, *On sumsets of convex sets*, Combin. Probab. Comput., 20 (2011), pp. 793–798.
- [12] I. D. SHKREDOV, *On sums of Szemerédi-Trotter sets*, Trans. Steklov Math. Inst., 289 (2015), pp. 300–309.
- [13] I. D. SHKREDOV, *Some new results on higher energies*, Trans. Moscow Math. Soc., 2013 (2013), pp. 31–63.
- [14] I. D. SHKREDOV, *Some remarks on sets with small quotient set*, Mat. Sbornik, accepted; preprint, <https://arxiv.org/abs/1603.04948>, 2016.
- [15] J. SOLYMOSI, *Bounding multiplicative energy by the sumset*, Adv. Math., 222 (2009), pp. 402–408.
- [16] T. TAO AND V. VU, *Additive Combinatorics*, Cambridge University Press, Cambridge, UK, 2006.