



University of Dundee

## Antitrypanosomatid Pharmacomodulation at Position 3 of the 8-Nitroquinolin-2(1H)-one Scaffold Using Palladium-Catalysed Cross-Coupling Reactions

Pedron, Julien; Boudot, Clotilde; Bourgeade-Delmas, Sandra; Sournia-Saquet, Alix; Paloque, Lucie; Rastegari, Maryam

*Published in:*  
ChemMedChem

*DOI:*  
[10.1002/cmdc.201800456](https://doi.org/10.1002/cmdc.201800456)

*Publication date:*  
2018

*Document Version*  
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

*Citation for published version (APA):*

Pedron, J., Boudot, C., Bourgeade-Delmas, S., Sournia-Saquet, A., Paloque, L., Rastegari, M., ... Valentin, A. (2018). Antitrypanosomatid Pharmacomodulation at Position 3 of the 8-Nitroquinolin-2(1H)-one Scaffold Using Palladium-Catalysed Cross-Coupling Reactions. *ChemMedChem*, 13(20), 2217-2228. <https://doi.org/10.1002/cmdc.201800456>

### General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Antitrypanosomatid pharmacomodulation at position 3 of the 8-nitroquinolin-2(1H)-one scaffold using pallado-catalyzed cross coupling reactions.

Julien Pedron<sup>[a]#</sup>, Clotilde Boudot<sup>[b]#</sup>, Sandra Bourgeade-Delmas<sup>[c]</sup>, Alix Sournia-Saquet<sup>[a]</sup>, Lucie Paloque<sup>[a]</sup>, Maryam Rastegari<sup>[a]</sup>, Mansour Abdoulaye<sup>[a]</sup>, Hussein El-Kashef<sup>[d]</sup>, Colin Bonduelle<sup>[a]</sup>, Geneviève Prativiel<sup>[a]</sup>, Susan Wyllie<sup>[e]</sup>, Alan Fairlamb<sup>[e]</sup>, Bertrand Courtioux<sup>[b]</sup>, Pierre Verhaeghe<sup>[a]\*</sup> and Alexis Valentin<sup>[c]</sup>.

[a] J. Pedron, Dr A. Saquet, Dr L. Paloque (<https://orcid.org/0000-0003-0359-3959>), Dr C. Bonduelle (<https://orcid.org/0000-0002-7213-7861>), Dr G. Prativiel (<https://orcid.org/0000-0003-0688-2107>) and Pr P. Verhaeghe\* (<https://orcid.org/0000-0003-0238-2447>). LCC-CNRS, Université de Toulouse, CNRS, UPS, 205 route de Narbonne, 31077 Toulouse, France. E-mail: pierre.verhaeghe@lcc-toulouse.fr

[b] C. Boudot and Dr B. Courtioux.

Université de Limoges, UMR INSERM 1094, Neuroépidémiologie Tropicale, Faculté de Pharmacie, 2 rue du Dr Marcland, 87025 Limoges, France.

[c] Dr S. Bourgeade-Delmas and Pr A. Valentin (<https://orcid.org/0000-0002-2133-112X>).

UMR 152 PharmaDev, Université de Toulouse, IRD, UPS, 35 Chemin des Maraîchers, 31400 Toulouse, France.

[d] Pr H. El-Kashef.

Assiut University, Faculty of Science, Department of Chemistry, 71516 Assiut, Egypt.

[e] Dr S. Wyllie (<https://orcid.org/0000-0001-8810-5605>) and Pr A. Fairlamb (<https://orcid.org/0000-0001-5134-0329>).

University of Dundee, School of Life Sciences, Division of Biological Chemistry and Drug Discovery, Dow Street, Dundee DD1 5EH, Scotland, United Kingdom

# Co-first-authors

Supporting information for this article is given via a link at the end of the document.

**Abstract:** An antikinoplastid pharmacomodulation study at position 3 of the recently described 3-bromo-8-nitroquinolin-2(1H)-one hit molecule was conducted. 24 derivatives were synthesized using the Suzuki-Miyaura cross-coupling reaction and evaluated *in vitro* toward both *L. infantum* axenic amastigotes and *T. brucei brucei* trypomastigotes. The results show that the introduction of a *para*-carboxyphenyl group at position 3 of the scaffold leads to a selective antitrypanosomal hit molecule (**21**) with a lower reduction potential (-0.56 V) than the initial hit (-0.45 V). Compound **21** displays a micromolar antitrypanosomal activity (IC<sub>50</sub> = 1.5 μM) and a low cytotoxicity on the human HepG2 cell line (CC<sub>50</sub> = 120 μM), reaching higher selectivity index (IS = 80) than the reference drug eflornithine. Contrary to the results previously obtained in this series, hit compound **21** is not active toward *L. infantum* and is not efficiently bioactivated by *T. brucei brucei* type I nitroreductase, which suggests the existence of an additional mechanism of action.

## Introduction


Kinetoplastids are flagellated protozoan parasites responsible for lethal neglected tropical diseases such as human African trypanosomiasis (HAT) and visceral leishmaniasis (VL). They are characterised by the presence of a circular DNA called kinetoplast, adjacent to the flagellar basal body. *Trypanosoma* parasites are transmitted by the bite of an infected "tse tse fly" and are the causative agents of HAT, also known as "sleeping sickness". There are many species of *Trypanosoma*, but only *T. brucei gambiense* and *T. brucei rhodesiense* are pathogenic for humans.[1] During the infection, metacyclic trypomastigotes enter in the blood circulation via the bite of the "tse tse fly" and disseminate in the whole organism where they differentiate into bloodstream trypomastigotes: this is the

hemolymphatic stage. After several weeks or months including symptoms such as headaches, anaemia and hepatosplenomegaly, trypomastigotes cross the blood-brain barrier (BBB) and cause damages to the central nervous system, leading to sleeping disorders, behavioural disorders, seizure, coma and finally death: this is the meningoencephalic stage.[2] *Leishmania* is another protozoan parasite, at the origin of leishmaniasis and transmitted by the bite of an infected sandfly. *L. donovani* and *L. infantum* are the two major species causing the most severe form of the disease: visceral leishmaniasis (VL).[3] Briefly, metacyclic promastigotes penetrate into the skin during the blood meal of an infected sandfly. They are internalized by mononuclear phagocytic cells such as macrophages where they differentiate into amastigotes. Parasites multiply in these cells until their destruction and disseminate in many organs such as liver and spleen, leading to death.[4]

HAT and VL represent more than 1 billion people at risk and are responsible for more than 55.000 new cases and 25.000 deaths each year.[5,6] These numbers are surely underestimated because of the difficulty to access to some rural areas and the unspecific symptoms in the early stages of the diseases. Currently, there are very few efficient and safe drugs available on the market against these neglected tropical diseases. Pentamidine and suramin are used for the treatment of the first stage of HAT but these molecules are highly toxic and need hospitalisation for the IV administration.[7] Melarsoprol, an arsenic containing-drug and a combination of eflornithine and nifurtimox are suitable for the second stage of HAT (Fig. 1).[8]. The same observation can be made with VL for which only Amphotericin B, miltefosine, antimonial derivatives, pentamidine and paromomycine are available. Among these drugs, miltefosine is the only orally available drug. These drugs are either expensive (liposomal amphotericin B), present severe side effects (nephrotoxicity of amphotericin B, teratogenicity of miltefosine...) or


This is the peer reviewed version of the following article: Pedron, J., et al. (2019) "Antitrypanosomatid Pharmacomodulation at Position 3 of the 8-Nitroquinolin-2(1H)-one Scaffold Using Palladium-Catalysed Cross-Coupling Reactions", *ChemMedChem* 13:20, pp. 2217-2228, which has been published in final form at <https://doi.org/10.1002/cmdc.201800456>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

show an increasing lack of efficacy due to the emergence of resistant parasites (antimony derivatives and miltefosine).[9] This global context calls for the discovery of new antikinoplastid molecules.



**Figure 1.** Structure, indication and route of administration of the antitrypanosomal drugs on the market.

Unfortunately, today, there are only two new chemical entities in clinical trials against HAT and none against VL (Fig 2.).[10] Acoziborole is an orally-active benzoxaborole in phase IIb/III of clinical trials, this molecule is active against both stages of HAT.[11] Fexinidazole, a 5-nitroimidazole, was recently in phase II of clinical trials against VL but showed a lack of efficacy whereas it succeeded to a phase IIIb study against HAT.[12,13] This molecule is rapidly metabolised *in vivo* into two metabolites (a sulfoxide and a sulfone derivative) which are still active against the *trypanosoma* parasites. Fexinidazole is selectively bioactivated by type I parasitic nitroreductases (NTR) leading to cytotoxic electrophilic metabolites, through a successive 2 electron reduction, such as nitroso and hydroxylamine derivatives.[14] There are two NTR identified in *Leishmania* (NTR1 and NTR2)[15,16] and only one in *Trypanosoma*. [17] These nitroreductases are absent from mammalian cells. Consequently, substrates of these enzymes can afford selective antikinoplastid candidates. Unfortunately, no X-ray structure of these parasitic NTRs is available, which prevents the use of the classical rational medicinal chemistry approaches, such as docking, to design new substrates of these enzymes.




**Figure 2.** Molecular structure of drug candidates acoziborole and fexinidazole (with its active metabolites).

Our research team has been working on the synthesis of new antikinoplastid molecules for several years. Starting from a

chemical study about 2-substituted nitroquinoline derivatives with antiparasitic potential,[18] we identified a new antileishmanial hit: the 8-nitroquinolin-2(1*H*)-one.[19] Pharmacomodulation studies at position 4 of this scaffold were then realized.[20,21] Recently, we described a comprehensive study of this pharmacophore *via* an electrochemistry-guided work and the development of a computational model, able to predict the redox potentials of each molecule in the series.[22] Thus, a new antikinoplastid hit molecule was identified (Fig. 3). This molecule was not genotoxic in the comet assay and was selectively bioactivated by type 1 NTRs of *L. donovani* and *T. brucei brucei*. [22]

Here, we present a pharmacomodulation study at position 3 of the scaffold, using the Suzuki-Miyaura or Sonogashira cross-coupling reaction. To explore SARs, twenty six molecules were synthesized and evaluated on both *L. infantum axenic* amastigotes and *T. brucei brucei* trypomastigotes. All molecules were also assessed for their cytotoxicity on the HepG2 human cell line.




**Figure 3.** Structure and biological profile of the previously identified antikinoplastid hit 3-bromo-8-nitroquinolin-2(1*H*)-one.

## Results and Discussion


### Chemistry

In a first time, 3-bromo-8-nitroquinolin-2(1*H*)-one was prepared in 3 steps, as presented in Scheme 1. [22] The nitration of the 2-chloroquinoline mainly led to the 2-chloro-8-nitroquinoline intermediate which was transformed in a second step into the corresponding lactam, according to a previously reported procedure.[23] The 8-nitroquinolin-2(1*H*)-one was finally selectively halogenated at position 3 by refluxing in HBr (48%) in the presence of sodium bromate, as reported by O'Brien and co-workers.[24]



**Scheme 1.** Synthesis of the initial antikinoplastid hit. a) H<sub>2</sub>SO<sub>4</sub>, HNO<sub>3</sub>, rt, 2 h; b) CH<sub>3</sub>CN, HClO<sub>4</sub>, 100 °C, 72 h; c) NaBrO<sub>3</sub>, HBr, 100 °C, 5 h.

Then, the microwave-assisted Suzuki-Miyaura cross coupling reaction between 3-bromo-8-nitroquinolin-2(1*H*)-one and *para*-methoxyphenylboronic acid was studied, looking for optimal conditions. (Table 1). Several parameters were then investigated such as the nature of the solvent, base, and Pd-catalyst or the reaction temperature.

**Table 1.** Optimization of the Suzuki-Miyaura cross-coupling reaction between 3-bromo-8-nitroquinolin-2(1H)-one and *para*-methoxyphenylboronic acid.


Entry	Solvent	Base, 3 equiv.	Catalyst	Temperature (°C)	Time (h)	Yield (%) <sup>[a]</sup>
1	DMF	Na <sub>2</sub> CO <sub>3</sub>	Pd(OAc) <sub>2</sub> , 0.1 equiv.	150	1	43
2	DMSO	Na <sub>2</sub> CO <sub>3</sub>	Pd(OAc) <sub>2</sub> , 0.1 equiv.	150	1	- <sup>[b]</sup>
3	Toluene	Na <sub>2</sub> CO <sub>3</sub>	Pd(OAc) <sub>2</sub> , 0.1 equiv.	110	8	- <sup>[c]</sup>
4	Dioxane	Na <sub>2</sub> CO <sub>3</sub>	Pd(OAc) <sub>2</sub> , 0.1 equiv.	100	8	- <sup>[c]</sup>
5	THF	Na <sub>2</sub> CO <sub>3</sub>	Pd(OAc) <sub>2</sub> , 0.1 equiv.	66	8	- <sup>[c]</sup>
6	DME	Na <sub>2</sub> CO <sub>3</sub>	Pd(OAc) <sub>2</sub> , 0.1 equiv.	85	8	- <sup>[c]</sup>
7	DMF/H <sub>2</sub> O (8/2)	Na <sub>2</sub> CO <sub>3</sub>	Pd(OAc) <sub>2</sub> , 0.1 equiv.	100	1	46
8	DMF/H <sub>2</sub> O (8/2)	K <sub>2</sub> CO <sub>3</sub>	Pd(OAc) <sub>2</sub> , 0.1 equiv.	100	1	- <sup>[b]</sup>
9	DMF/H <sub>2</sub> O (8/2)	Cs <sub>2</sub> CO <sub>3</sub>	Pd(OAc) <sub>2</sub> , 0.1 equiv.	100	1	- <sup>[b]</sup>
10	DMF	Na <sub>2</sub> CO <sub>3</sub>	Pd(PPh <sub>3</sub> ) <sub>4</sub> , 0.1 equiv.	150	1	26
11	DME	Na <sub>2</sub> CO <sub>3</sub>	Pd(PPh <sub>3</sub> ) <sub>4</sub> , 0.1 equiv.	85	3	52
12	DME	K <sub>2</sub> CO <sub>3</sub>	Pd(PPh <sub>3</sub> ) <sub>4</sub> , 0.1 equiv.	85	4	84
13	DME	K <sub>2</sub> CO <sub>3</sub>	Pd(PPh <sub>2</sub> )Cl <sub>2</sub> , 0.1 equiv.	85	8	- <sup>[c]</sup>
14	DME	K <sub>2</sub> CO <sub>3</sub>	Pd(dppf)Cl <sub>2</sub> , 0.1 equiv.	85	8	- <sup>[c]</sup>
15	DME	K <sub>2</sub> CO <sub>3</sub>	Pd(OAc) <sub>2</sub> , 0.1 equiv.	85	8	- <sup>[c]</sup>
16	DME	Cs <sub>2</sub> CO <sub>3</sub>	Pd(PPh <sub>3</sub> ) <sub>4</sub> , 0.1 equiv.	85	2	88
17	DME	CsF	Pd(PPh <sub>3</sub> ) <sub>4</sub> , 0.1 equiv.	85	2	88
18	DME	Cs <sub>2</sub> CO <sub>3</sub>	Pd(PPh <sub>3</sub> ) <sub>4</sub> , 0.05 equiv.	85	4	56

[a] The yield was calculated after purification by chromatography on silica gel with adapted eluent; [b] Degradation of the substrate was observed on TLC [c] Only partial conversion of the substrate was observed on TLC.

The first Suzuki-Miyaura reaction conditions (entry 1) were inspired from a previously described protocol to introduce aryl moiety at position 3 of the quinolinone ring.<sup>[25]</sup> This reaction was realized in DMF under microwave (MW) heating, using 3 equiv. of Na<sub>2</sub>CO<sub>3</sub> as a base, 0.1 equiv. of Pd(OAc)<sub>2</sub> as a catalyst and 1.2 equiv of *p*-methoxyphenylboronic acid in a sealed tube, to afford the desired compound in 43 % yield. The first parameter studied was the nature of the solvent (entries 1-7). By using toluene, dioxane, THF, DME or DMSO, partial conversion or degradation of the substrate was observed whereas the use of a DMF/H<sub>2</sub>O mixture led to a 46% yield. Then, two others bases were studied (entries 8-9) but led to the degradation of the substrate within 1 h. In DMF, the replacement of Pd(OAc)<sub>2</sub> by Pd(PPh<sub>3</sub>)<sub>4</sub> decreased the yield of the reaction from 43% to 26% (entry 10). Then, adapting another previously described protocol using DME and Pd(PPh<sub>3</sub>)<sub>4</sub> (entry 11), a slightly better yield was obtained.<sup>[26]</sup> This result was then improved by replacing Na<sub>2</sub>CO<sub>3</sub> by K<sub>2</sub>CO<sub>3</sub> which afford a yield of 84%. The nature of the


catalyst was then studied (entries 13-15) but none of these assays led to an improvement in the reaction yield and only a few conversion was observed. Finally, the best results were obtained by using Cs<sub>2</sub>CO<sub>3</sub> or CsF as a base (entries 16-17), which afford an efficient procedure of a Suzuki-Miyaura cross coupling reaction on this substrate. In the entry 18, the decrease of the amount of catalyst to 0.05 equiv. led to a lower reaction yield (56%) in comparison to entry 16, which was chosen as the best reaction conditions.

Then, the procedure was extended to 19 other arylboronic acids, to afford new derivatives bearing a phenyl, thiophene, furane or pyridine moiety at position 3 of the scaffold (Scheme 2). The reactions yields were generally superior to 65% (for 14 derivatives) but were lower for 4-hydroxyphenylboronic acid and 4-aminophenylboronic acid with respectively 37% and 41% yields.



**Scheme 2.** General procedure for the Suzuki-Miyaura cross coupling reaction between 3-bromo-8-nitroquinolin-2(1*H*)-one and various arylboronic acids.


Three additional molecules (**21-23**) were synthesized by the saponification of compounds **18-20** into the carboxylic acid derivatives, using an excess of sodium hydroxide in an ethanol/water mixture (Scheme 3). Finally, compound **24** was obtained in two steps: *O*-methylation of **18**, using methyl iodide in DMF under inert atmosphere, followed by a saponification with the same procedure as described for compounds **21-23** (Scheme 3).



**Scheme 3.** Preparation of compounds **21-24**.

In parallel, in a view to enlarge the chemical diversity at position 3 of the scaffold and introduce alkynyl moieties, a Sonogashira cross-coupling reaction was set up. The initial conditions were adapted from a previously described protocol. [27] As presented in Table 2, after a brief optimization of the reaction between 3-bromo-8-nitroquinolinone and phenylacetylene, DME was chosen as solvent, and the reaction was carried out at 15°C (entry 6). Interestingly, it was noted that temperature had a strong influence on the reaction. At 0 °C in DME, the reaction did not take place whereas it worked at 15 and 25 °C, affording awaited compound **25**. Surprisingly, it led to the unexpected 8-nitro-2-phenylfuro[2,3-*b*]quinoline **26** when heating at 40 °C in DME or DMF. Compound **27** was also obtained when trying to recrystallize **26** in acetonitrile, underlining the instability of this product. Such a consecutive Sonogashira coupling and cyclization reaction, leading to furo[2,3-*b*]pyridine derivatives, was already reported in the literature [28,29] The optimization of both Suzuki-Miyaura and Sonogashira cross-coupling reactions led to the synthesis of 26 new molecules with an aryl or an alkynyl group at position 3 of the scaffold. Apart molecule **25**, which was considered too unstable, all these molecules were then evaluated *in vitro* to determine their antikinoplastid potential. Compound **26** could not be evaluated because of a very poor aqueous solubility. Thus, only the Suzuki-Miyaura coupling products **1-23** could be assessed.

**Table 2.** Optimization of the Sonogashira cross-coupling reaction on 3-bromo-8-nitroquinolin-2(1*H*)-one substrate.




Entry	Solvent	Temperature (°C)	Yield <b>25</b> (%) <sup>[a]</sup>	Yield <b>26</b> (%) <sup>[a]</sup>	Time (h)
1	DMF	25°C	..[b]	..[b]	48
2	THF	25°C	..[b]	..[b]	48
3	Et <sub>3</sub> N	25°C	..[b]	..[b]	48
4	DME	25°C	<b>56</b>	0	0.25
5	DME	0	..[b]	..[b]	48
6	DME	15	<b>72</b>	0	1.5
7	DME	40	0	<b>51</b>	0.5
8	DMF	40	0	<b>56</b>	36

[a] The yield was calculated after purification by chromatography on silica gel with adapted eluent; [b] Only partial conversion of the substrate was observed on TLC.

## Compound evaluation

In a first time, the cytotoxicity of these molecules was assessed *in vitro* on the HepG2 human cell line and the corresponding cytotoxic concentration 50% (CC<sub>50</sub>) were compared to the one of the reference drug doxorubicin (Table 3). The biological results showed that compounds **1-11** presented a lack of solubility in aqueous medium and could not be tested. The water-solubility was improved with compounds **10, 12-17** and **21-23**, bearing either a pyridine-3-yl or a phenyl moiety at position 3 of the scaffold, this latter being substituted by a hydrophilic group such as hydroxymethyl and aldehyde or an ionized carboxylic group. These molecules displayed a low cytotoxicity on HepG2 human cell line with CC<sub>50</sub> > 25 μM, the compound bearing an aldehyde group in *para* position of the phenyl ring (**15**) was the most cytotoxic of this series with a CC<sub>50</sub> = 30 μM.

Then, all synthesized compounds were tested *in vitro* against *Leishmania infantum* axenic amastigotes. Their inhibitory concentration 50% (IC<sub>50</sub>) were determined and compared to the ones of two antileishmanial reference drugs (amphotericin B and miltefosine) and to the drug candidate fexinidazole.


**Table 3.** *In vitro* antileishmanial, antitrypanosomal and cytotoxic activities of the synthesized compounds **1-25** and reference standards.

Compound	R (Yield %)	IC <sub>50</sub> <i>L. infantum</i> axenic amastigotes (μM)	IC <sub>50</sub> <i>T. brucei brucei</i> trypomastigotes (μM)	CC <sub>50</sub> HepG2 (μM)	Antitrypanosomal selectivity Index <sup>[g]</sup>
1	Phenyl (90)	>12 <sup>[a]</sup>	4.7 +/- 2.7	>12 <sup>[a]</sup>	>2
2	4-OCH <sub>3</sub> -phenyl (88)	>6 <sup>[a]</sup>	-	>6 <sup>[a]</sup>	-
3	4-OH-phenyl (37)	>3 <sup>[a]</sup>	-	>3 <sup>[a]</sup>	-
4	4-NH <sub>2</sub> -phenyl (41)	>3 <sup>[a]</sup>	-	>3 <sup>[a]</sup>	-
5	4-Cl-phenyl (73)	>3 <sup>[a]</sup>	-	>3 <sup>[a]</sup>	-
6	4-F-phenyl (70)	>6 <sup>[a]</sup>	-	>6 <sup>[a]</sup>	-
7	4-CF <sub>3</sub> -phenyl (60)	>6 <sup>[a]</sup>	-	>6 <sup>[a]</sup>	-
8	Thiophen-3-yl (92)	>6 <sup>[a]</sup>	-	>6 <sup>[a]</sup>	-
9	Furan-2-yl (65)	>3 <sup>[a]</sup>	-	>3 <sup>[a]</sup>	-
10	Pyridin-3-yl (62)	>50 <sup>[a]</sup>	2.8 +/- 0.8	>25 <sup>[a]</sup>	>9
11	Pyridin-4-yl (71)	NS <sup>[b]</sup>	-	NS <sup>[b]</sup>	-
12	4-CH <sub>2</sub> OH-phenyl (77)	>25 <sup>[a]</sup>	1.9 +/- 0.3	>25 <sup>[a]</sup>	>13
13	3-CH <sub>2</sub> OH-phenyl (87)	29.3 +/- 4.2	1.5 +/- 0.3	>25 <sup>[a]</sup>	>17
14	2-CH <sub>2</sub> OH-phenyl (72)	22 +/- 2.0	7.2 +/- 0.6	>100 <sup>[c]</sup>	>14
15	4-CHO-phenyl (51)	35 +/- 1.7	0.5 +/- 0.1	30 +/- 3.7	60
16	3-CHO-phenyl (55)	10.2 +/- 0.6	5.6 +/- 0.4	>100 <sup>[c]</sup>	>18
17	2-CHO-phenyl (65)	9.8 +/- 1.2	7.5 +/- 0.4	>50 <sup>[a]</sup>	>7
18	4-COOCH <sub>3</sub> -phenyl (72)	>12.5 <sup>[a]</sup>	-	>12.5 <sup>[a]</sup>	-
19	3-COOCH <sub>3</sub> -phenyl (71)	NS <sup>[b]</sup>	-	NS <sup>[b]</sup>	-
20	2-COOCH <sub>3</sub> -phenyl (71)	NS <sup>[b]</sup>	-	NS <sup>[b]</sup>	-
<b>21</b>	<b>4-COOH-phenyl (65)</b>	<b>&gt;100<sup>[c]</sup></b>	<b>1.5 +/- 0.2</b>	<b>120 +/- 7</b>	<b>80</b>
22	3-COOH-phenyl (60)	>100 <sup>[c]</sup>	7.5 +/- 1.0	>100 <sup>[c]</sup>	>13
23	2-COOH-phenyl (66)	>100 <sup>[c]</sup>	>50 <sup>[a]</sup>	>100 <sup>[c]</sup>	-
Initial Hit <sup>[22]</sup>	Br	7.1 +/- 1.5	1.9 +/- 0.44	92 +/- 13.0	48
8-nitroquinolinone <sup>[22]</sup>	H	15.5 +/- 0.5	23.4 +/- 5.7	164 +/- 28	7
Doxorubicin <sup>[d]</sup>		-	-	0.2 +/- 0.02	-
Amphotericin B <sup>[e]</sup>		0.06 +/- 0.001	-	7.0 +/- 0.25	-
Miltefosine <sup>[e]</sup>		0.8 +/- 0.2	-	84.5 +/- 8.8	-
Fexinidazole <sup>[e, f]</sup>		3.3 +/- 0.7	0.4 +/- 0.18	>100 <sup>[c]</sup>	>250
Suramin <sup>[f]</sup>		-	0.03 +/- 0.009	>100 <sup>[c]</sup>	>3333
Eflornithine <sup>[f]</sup>		-	15.8 +/- 2.1	>100 <sup>[c]</sup>	>6

[a] The product could not be tested at higher concentrations due to a poor solubility in aqueous medium; [b] The product was not soluble at any tested concentration; [c] The IC<sub>50</sub> or CC<sub>50</sub> value was not reached at the highest tested concentration; [d] Doxorubicin was used as a cytotoxic reference drug; [e] Amphotericin B, Miltefosine and Fexinidazole were used as antileishmanial reference drugs or drug candidate; [f] Fexinidazole, Suramin and Eflornithine were used as anti-trypanosomal reference drugs or drug candidate; [g] Antitrypanosomal selectivity index was calculated according to the following formula : CC<sub>50</sub> HepG2 / IC<sub>50</sub> *T. brucei brucei*

Regarding antileishmanial activity, apart for aldehyde-containing compounds **16** and **17**, the tested series appeared either poorly active ( $IC_{50} = 22 - 35 \mu M$ ) or even totally inactive ( $IC_{50} > 100 \mu M$  for carboxylic group containing derivatives) toward *L. infantum*, in comparison with the reference drugs. Thus, introducing an aryl moiety at position 3 of the pharmacophore did not seem to favour antileishmanial activity.

In a second time, only the compounds with appropriate aqueous soluble solubility were tested *in vitro* against *T. brucei brucei* trypomastigotes and compared to reference antitrypanosomal drugs (suramin and eflornithine) and the drug candidate fexinidazole. All tested molecules displayed good antitrypanosomal activity ( $0.5 \mu M \leq IC_{50} \leq 7.5 \mu M$ ), better than the one of eflornithine ( $IC_{50} = 15.8 \mu M$ ) and approaching the one of fexinidazole ( $IC_{50} = 0.4 \mu M$ ). Interestingly, compounds substituted in *ortho* position of the phenyl ring (**14**, **17**, **23**) displayed higher  $IC_{50}$  values than their analogues substituted in *meta* and *para* positions. Compounds with a *para*-substituted phenyl ring generally showed lower  $IC_{50}$  than their *meta*-substituted analogues, **15** being 11 times more potent than **16**, and **21** 5 times more potent than **22**. Compounds **15** and **21** appeared as the most promising antitrypanosomal molecules in this series with respective  $IC_{50}$  values of 0.5 and 1.5  $\mu M$ . Compound **21** appears as a new antitrypanosomal hit, with an activity against *T. brucei brucei* close to the one of the parent compound ( $IC_{50} = 1.9 \mu M$ ) but with a better cytotoxicity profile, leading to a better selectivity index (80 versus 48 for the initial hit). By comparison with reference drugs, compound **21** appears less active than suramin but more active than eflornithine. Regarding fexinidazole, another nitroheterocycle, compound **21** presents the same cytotoxic profile with  $CC_{50} = 120 \mu M$  and is three times less active than this drug candidate. Indeed, unlike fexinidazole and 3-bromo-8-nitroquinolin-2(1*H*)-one, compound **21** displayed a selective antitrypanosomal activity, being inactive ( $IC_{50} > 100 \mu M$ ) against *L. infantum*. This is a first element indicating that molecule **21** shows a specific antiparasitic profile in the studied series.



**Figure 4.** *In vitro* antiparasitic activities and cytotoxicity of compound **24**.

Finally, the *O*-methylated analogue **24** of hit compound **21** was synthesized and tested *in vitro* against both *L. infantum* and *T. brucei brucei* (Fig. 4). It was as active against *T. brucei brucei* as **21** ( $IC_{50} = 2.2 \mu M$ ). Molecule **24** was also active against *L. infantum* ( $IC_{50} = 12.8 \mu M$ ) whereas **21** was not ( $IC_{50} > 100 \mu M$ ). These results are surprising, considering that the hydrogen bond between the lactam function and the nitro group appeared mandatory for providing antileishmanial activity in 8-nitroquinolin-2(1*H*)-one series by increasing the reduction potential value [22]. This is a second element indicating that the introduction of an aryl group at position 3 of the 8-nitroquinolin-2(1*H*)-one scaffold could lead to a new antikineto-plasid mechanism of action, in comparison with 3-bromo-8-nitroquinolin-2(1*H*)-one.

To assess if the antitrypanosomal nitroheterocycle **21** was bioactivated by the NTR of *T. brucei brucei*, its  $IC_{50}$  value was measured on both a wild type *T. brucei brucei* trypomastigotes strain

and a NTR-overexpressing one (Table 4). The results were compared with the ones obtained for the initial hit.[22] This latter is clearly bioactivated by the trypanosomal NTR, being 4.5 more active against the strain overexpressing the NTR than on the wild type one, whereas **21** is only 1.3 time more active on the strain overexpressing the NTR. This result is a third element that suggests that **21**, which is less intensively bioactivated by the trypanosomal NTR than the initial hit, could present another parasitic target. This assay also showed that compound **21** presents the same level of activity toward *T. brucei brucei* than the drug nifurtimox, used as a bioactivation control.








**Table 4.** Study of the bioactivation of **21** by trypanosomal nitroreductase NTR.

Compound	$IC_{50}$ <i>T. brucei brucei</i> trypomastigotes wild type strain ( $\mu M$ )	$IC_{50}$ <i>T. brucei brucei</i> trypomastigotes NTR over-expressing strain ( $\mu M$ ).	Fold change
<b>21</b>	5.4 +/- 0.12	4.2 +/- 0.2	1.3
<b>Initial Hit</b>	17.7 +/- 1.0	3.9 +/- 0.1	4.5
<b>Nifurtimox</b>	1.9 +/- 0.05	0.6 +/- 0.05	3.1

In parallel, an electrochemistry study was carried out by measuring in DMSO the reduction potentials of five 8-nitroquinolin-2(1*H*)-one derivatives bearing an aryl group at position 3, using cyclic voltammetry (Table 5). For all compounds, a reversible single-electron reduction was observed (formation of an anion radical). The redox potentials of the new compounds bearing an aryl group at position 3 ranged between -0.53 V and -0.59 V, higher than for the initial hit (-0.45 V) but quite close to the one of 8-nitroquinolin-2(1*H*)-one (-0.54 V). As previously noted in the studied series [22], the *O*-methylation of compound **21** leading to compound **24**, is responsible for an important decrease in the redox potential value from -0.56 V to -0.93 V. This shift is mainly due to the removal of the intramolecular hydrogen bond between the lactam function and the nitro group. It can be concluded that the introduction of a phenyl ring at the position 3 of the scaffold has no significant impact on the redox potential of the studied series but that it allows to reach novel antitrypanosomal molecules that display lower reduction potential than the initial hit with the same level of efficacy.

Thus, to understand the selective antitrypanosomal activity of **21**, other parasitic targets should be explored. Among the parasitic targets that were recently reported in the literature concerning the antitrypanosomal activity of new diverse nitroheterocycles, the *S*-adenosylmethionine decarboxylase (AdoMetDC) was highlighted by a high-throughput mass spectrometry-based assay conducted on 400.000 molecules [30] and is to consider.

**Table 5.** Effects of the substitution of the phenyl ring on reduction potentials in Suzuki-miyaura series.

Compound	Structure	E° (V) <sup>[a]</sup>
<b>1</b>		-0.59
<b>7</b>		-0.53
<b>13</b>		-0.53
<b>21</b>		-0.56
<b>24</b>		-0.93
<b>Initial Hit</b>		-0.45
<b>8-nitroquinolin-2(1H)-one</b>		-0.54

[a] Cyclic voltammetry conditions: DMSO/TBAPF<sub>6</sub>, SCE/GC, 1 electron reversible reduction, values are given in V versus NHE

## Conclusions

An optimized Suzuki-Miyaura reaction at position 3 of 3-bromo-8-nitroquinolin-2(1H)-one led to the synthesis of 24 new derivatives. These molecules were screened *in vitro* against *L. infantum* and *T. brucei brucei* to evaluate their antikinetoplastid potential. Among these molecules, a new selective antitrypanosomal hit **21**, bearing a *para*-carboxyphenyl group, was identified. Compound **21** was not cytotoxic on the HepG2 human cell line (CC<sub>50</sub> = 120 μM), displayed a good antitrypanosomal activity (IC<sub>50</sub> = 1.5 μM, SI = 80), better than the one of the drug eflornithine and approaching the one of fexinidazole, a 5-nitroimidazole in phase III of clinical trials against HAT. Interestingly, **21** was inactive against *L. infantum* showing a parasitic selectivity among kinetoplastids. Unlike fexinidazole and the previously identified hit molecule in the series, this molecule was not efficiently metabolized by the type I trypanosomal NTR, suggesting a probable additional mechanism of action in this series.

## Experimental Section

### Chemistry

All reagents and solvents were obtained from commercial sources (Fluorochem®, Sigma-Aldrich® or Alfa Aesar®) and used as received. The progress of the reactions was monitored by pre-coated thin layer chromatography (TLC) sheets ALUGRAM® SIL G/UV<sub>254</sub> from Macherey-Nagel and were visualized by ultraviolet light at 254 nm. The <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on a Bruker UltraShield 300 MHz, a Bruker IconNMR 400 MHz, or a Bruker Avance NEO 600 MHz instrument, at the Laboratoire de Chimie de Coordination and data are presented as follows: chemical shifts in parts per million (δ) using tetramethylsilane (TMS) as reference, coupling constant in Hertz (Hz), multiplicity by abbreviations: (s) singlet, (d) doublet, (t) triplet, (q) quartet, (dd) doublets of doublets, (m) multiplet and (br s) broad singlet. Melting points are uncorrected and were measured on a Stuart Melting Point SMP3 instrument. High-resolution mass measurements were done on a GCT Premier Spectrometer (DCI, CH<sub>4</sub>, HRMS) or Xevo G2 QTOF (Waters, ESI+, HRMS) instrument at the Université Paul Sabatier, Toulouse. Microwave reactions were realized in a CEM Discover® microwave reactor.

3-bromo-8-nitroquinolin-2(1H)-one was prepared according to a previously reported procedure.[22]

General procedure for the preparation of compounds **1-20** and **25**

One equiv. of 3-bromo-8-nitroquinolin-2(1H)-one (1.1 mmol, 300 mg), 3 equiv. of cesium carbonate (3.3 mmol, 1.1 g), 0.1 equiv. of Pd(PPh<sub>3</sub>)<sub>4</sub> (0.12 mmol, 127 mg) and 1.2 equiv. of the appropriate phenylboronic acid were added in a sealed flask of 25 mL. Under Ar atmosphere, 10 mL of dry dimethoxyethane were then added. The reaction mixture was heated at 85°C in a microwave reactor during 2 h. The reaction mixture was poured into water and extracted 3 times with dichloromethane (3x100 mL). The organic layer was washed with water, dried over anhydrous MgSO<sub>4</sub> and evaporated *in vacuo*. The crude residues were purified by chromatography on silica gel using adapted eluent and recrystallized if necessary to give compounds **1-20** and **25**.

**8-nitro-3-phenylquinolin-2(1H)-one 1** (C<sub>15</sub>H<sub>10</sub>N<sub>2</sub>O<sub>3</sub>) was purified by chromatography on silica gel using dichloromethane/ethyl acetate (97/3) as eluent, isolated and recrystallized in acetonitrile to yield a yellow solid (264 mg, 0.99 mmol, 90%); Tdec: 177 °C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ:



7.31-7.35 (m, 1H, H6), 7.41-7.50 (m, 3H, H3', H4' and H5'), 7.75-7.77 (m, 2H, H2' and H6'), 7.91 (s, 1H, H4), 7.94 (dd, 1H, J=7.6 and 1.4 Hz, H5), 8.49 (dd, 1H, J=8.4 and 1.4 Hz, H7), 11.40 (br s, 1H, NH). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ: 121.4 (CH), 122.7 (C), 127.3 (CH), 128.5 (2xCH), 128.7 (2xCH), 129.1 (CH), 132.7 (C), 133.1 (C), 134.5 (C), 134.6 (C), 135.6 (CH), 136.7 (CH), 161.0 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>15</sub>H<sub>11</sub>N<sub>2</sub>O<sub>3</sub> [M+H]<sup>+</sup> = 267.0770, found 267.0762.

**3-(4-methoxyphenyl)-8-nitroquinolin-2(1H)-one 2** (C<sub>16</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>) was purified by chromatography on silica gel using dichloromethane/ethyl acetate (95/5) as eluent, isolated and recrystallized in acetonitrile to yield an orange solid (287 mg, 0.97 mmol, 88%); m.p. 229 °C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ: 3.87 (s, 3H, CH<sub>3</sub>), 6.99-7.02 (m, 2H, H3' and H5'), 7.29-7.33 (m, 1H, H6), 7.73-7.77 (m, 2H, H2' and H6'), 7.87 (s, 1H, H4), 7.91 (dd, J=7.6 and 1.4 Hz, 1H, H5), 8.47 (dd, J=8.3 and 1.4 Hz, 1H, H7), 11.38 (br s, 1H, NH). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz) δ: 55.4 (CH<sub>3</sub>), 114.0 (2xCH), 121.3 (CH), 122.9 (C), 125.6 (C), 126.9 (CH), 130.0 (2xCH), 132.7 (C), 132.9 (C), 134.0 (C), 135.3 (CH), 135.4 (CH), 160.3 (C), 161.2 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>16</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub> [M+H]<sup>+</sup> = 297.0875, found 297.0864.

**3-(4-hydroxyphenyl)-8-nitroquinolin-2(1H)-one 3** (C<sub>15</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>) was purified by chromatography on silica gel using dichloromethane/ethyl acetate (75/25) as eluent and isolated to yield an orange solid (115 mg, 0.41 mmol, 37%); m.p. 266 °C, <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz) δ: 6.84-6.88 (m, 2H, H3' and H5'), 7.40-7.44 (m, 1H, H6), 7.67-7.71 (m, 2H, H2' and H6'), 8.20 (dd, J= 7.7 and 1.4 Hz, 1H, H5), 8.26 (s, 1H, H4), 8.40 (dd, J= 8.3 and 1.4 Hz, 1H, H7), 9.74 (s, 1H, OH), 11.12 (br s, 1H, NH). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>, 75 MHz) δ: 115.4 (2xCH), 122.0 (CH), 122.8 (C), 125.8 (C), 127.1 (CH), 130.4 (2xCH), 132.3 (C), 132.5 (C), 133.5 (C), 136.0 (CH), 136.1 (CH), 158.5 (C), 160.8 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>15</sub>H<sub>11</sub>N<sub>2</sub>O<sub>4</sub> [M+H]<sup>+</sup> = 283.0719, found 283.0706.

**3-(4-aminophenyl)-8-nitroquinolin-2(1H)-one 4** (C<sub>15</sub>H<sub>11</sub>N<sub>3</sub>O<sub>3</sub>) was purified by chromatography on silica gel using dichloromethane/ethyl acetate (95/5) as eluent and isolated to yield an orange solid (128 mg, 0.45 mmol, 41%); m.p. 280 °C, <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz) δ: 5.44 (s, 2H, NH<sub>2</sub>), 6.61-6.65 (m, 2H, H3' and H5'), 7.37-7.41 (m, 1H, H6), 7.58-7.62 (m, 2H, H2' and H6'), 8.15-8.18 (m, 2H, H4 and H5), 8.36 (dd, 1H, J= 8.3 and 1.4 Hz, H7), 11.07 (br s, 1H, NH). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>, 100 MHz) δ: 113.6 (2xCH), 121.9 (CH), 122.1 (C), 123.1 (C), 126.5 (C), 129.9 (2xCH), 131.9 (C), 132.9 (C), 133.4 (C), 134.1 (CH), 135.8 (CH), 150.0

(CH), 161.0 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>15</sub>H<sub>12</sub>N<sub>3</sub>O<sub>3</sub> [M+H]<sup>+</sup> = 282.0879, found 282.0874.

**3-(4-chlorophenyl)-8-nitroquinolin-2(1H)-one 5** (C<sub>15</sub>H<sub>9</sub>N<sub>2</sub>O<sub>3</sub>Cl) was purified by chromatography on silica gel using dichloromethane/ethyl acetate (97/3) as eluent and isolated to yield a yellow solid (241 mg, 0.80 mmol, 73%); m.p. 222 °C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ: 7.33-7.37 (m, 1H, H6), 7.40-7.42 (m, 2H, H3' and H5'), 7.66-7.75 (m, 2H, H2' and H6'), 7.92 (s, 1H, H4), 7.95 (dd, 1H, J=7.6 and 1.5 Hz, H5), 8.51 (dd, 1H, J=8.4 and 1.5 Hz, H7), 11.42 (br s, 1H, NH). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100 MHz) δ: 121.6 (CH), 122.4 (C), 126.9 (CH), 127.7 (CH), 128.7 (CH), 129.1 (CH), 129.7 (CH), 132.8 (C), 133.1 (C), 133.2 (C), 134.4 (C), 135.7 (CH), 136.2 (C), 137.3 (CH), 160.6 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>15</sub>H<sub>10</sub>ClN<sub>2</sub>O<sub>3</sub> [M+H]<sup>+</sup> = 301.0380, found 301.0373.

**3-(4-fluorophenyl)-8-nitroquinolin-2(1H)-one 6** (C<sub>15</sub>H<sub>9</sub>N<sub>2</sub>O<sub>3</sub>F) was purified by chromatography on silica gel using dichloromethane/ethyl acetate (98/2) as eluent and isolated to yield a yellow solid (219 mg, 0.77 mmol, 70%); m.p. 217 °C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ: 7.12-7.20 (m, 2H, H2' and H6'), 7.31-7.36 (m, 1H, H6), 7.73-7.80 (m, 2H, H3' and H5'), 7.89 (s, 1H, H4), 7.94 (dd, 1H, J= 7.6 et 1.5 Hz, H5), 8.50 (dd, 1H, J= 8.4 and 1.5 Hz, H7), 11.41 (br s, 1H, NH). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100 MHz) δ: 115.5 (d, J= 21.6 Hz, 2xCH), 121.5 (CH), 122.6 (C), 124.4 (C), 127.4 (CH), 130.6 (d, J= 8.3 Hz, 2xCH), 132.7 (C), 133.1 (C), 133.5 (C), 135.5 (CH), 136.5 (CH), 160.9 (C), 163.2 (d, J= 249.2 Hz, C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>15</sub>H<sub>10</sub>FN<sub>2</sub>O<sub>3</sub> [M+H]<sup>+</sup> = 285.0675, found 285.0676.

**8-nitro-3-(4-trifluoromethylphenyl)quinolin-2(1H)-one 7** (C<sub>16</sub>H<sub>9</sub>N<sub>2</sub>O<sub>3</sub>F<sub>3</sub>) was purified by chromatography on silica gel using dichloromethane/ethyl acetate (98/2) as eluent and isolated to yield a yellow solid (220 mg, 0.66 mmol, 60%); m.p. 168 °C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ: 7.34-7.38 (m, 1H, H6), 7.72-7.75 (m, 2H, H2' and H6'), 7.88 (m, 2H, H3' and H5'), 7.95-7.98 (m, 2H, H4 and H5), 8.53 (dd, 1H, J= 8.4 and 1.4 Hz, H7), 11.45 (br s, 1H, NH). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 150 MHz) δ: 121.7 (CH), 122.4 (C), 124.0 (q, J= 272.3 Hz, C), 125.4 (q, J= 3.7 Hz, 2xCH), 127.9 (CH), 129.1 (2xCH), 130.9 (q, J= 32.5 Hz, C), 132.8 (C), 133.2 (C), 133.4 (C), 135.8 (CH), 137.6 (CH), 138.0 (C), 160.5 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>16</sub>H<sub>10</sub>N<sub>2</sub>O<sub>3</sub>F<sub>3</sub> [M+H]<sup>+</sup> = 335.0644, found 335.0629.

**8-nitro-3-(thiophen-3-yl)quinolin-2(1H)-one 8** (C<sub>13</sub>H<sub>8</sub>N<sub>2</sub>SO<sub>3</sub>) was purified by chromatography on silica gel using dichloromethane/ethyl acetate (98/2) as eluent and isolated to yield a yellow solid (275 mg, 1.01 mmol, 92%); m.p. 199 °C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ: 7.31-7.35 (m, 1H,

H6), 7.41-7.43 (m, 1H, H4'), 7.59 (dd, 1H, J= 5.1 Hz and 1.3 Hz, H5'), 7.94 (dd, 1H, J= 7.7 Hz and 1.4 Hz, H5), 8.05 (s, 1H, H4), 8.39 (dd, 1H, J=3.0 and 1.3 Hz, H2'), 8.48 (dd, 1H, J=8.4 and 1.4 Hz, H7), 11.42 (br s, 1H, NH). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz) δ: 121.4 (CH), 122.6 (C), 123.6 (C), 125.6 (CH), 126.2 (CH), 127.0 (CH), 127.1 (CH), 128.6 (C), 132.4 (C), 134.1 (CH), 134.3 (C), 135.4 (CH), 160.6 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>13</sub>H<sub>9</sub>N<sub>2</sub>SO<sub>3</sub> [M+H]<sup>+</sup> = 273.0334, found 273.0330.

**3-(2-furanyl)-8-nitroquinolin-2(1H)-one 9** (C<sub>13</sub>H<sub>8</sub>N<sub>2</sub>O<sub>4</sub>) was purified by chromatography on silica gel using dichloromethane as eluent and isolated to yield a dark red solid (183 mg, 0.71 mmol, 65%); m.p. 257 °C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ: 6.58 (dd, 1H, J= 3.3 Hz and 1.9 Hz, H4'), 7.30-7.35 (m, 1H, H6), 7.55-7.56 (m, 2H, H3' and H5'), 7.96 (dd, 1H, J= 7.6 and 1.4 Hz, H5), 8.23 (s, 1H, H4), 8.46 (dd, 1H, J= 8.3 and 1.4 Hz, H7), 11.42 (br s, 1H, NH). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100 MHz) δ: 112.6 (CH), 114.2 (CH), 121.6 (CH), 122.5 (C), 123.6 (C), 126.9 (CH), 130.7 (CH), 132.0 (C), 132.8 (C), 135.5 (CH), 143.3 (CH), 147.7 (C), 158.7 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>13</sub>H<sub>8</sub>N<sub>2</sub>O<sub>4</sub> [M+H]<sup>+</sup> = 257.0562, found 257.0558.

**8-nitro-3-(pyridine-3-yl)quinolin-2(1H)-one 10** (C<sub>14</sub>H<sub>9</sub>N<sub>3</sub>O<sub>3</sub>) was purified by chromatography on silica gel using dichloromethane/ethyl acetate as eluent and isolated to yield a yellow solid (182 mg, 0.68 mmol, 62%); m.p. 237 °C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ: 7.35-7.39 (m, 1H, H6), 7.40-7.43 (m, 1H, H5'), 7.97-7.99 (m, 2H, H4 and H5), 8.20-8.23 (m, 1H, H6'), 8.53 (dd, 1H, J=8.4 and 1.4 Hz, H7), 8.66 (dd, 1H, J=4.9 and 1.7 Hz, H4'), 8.90 (dd, 1H, J=2.3 and 0.9 Hz, H2'), 11.45 (br s, 1H, NH). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100 MHz) δ: 121.7 (CH), 122.3 (C), 123.1 (CH), 127.9 (CH), 130.5 (C), 131.4 (C), 132.8 (C), 133.3 (C), 135.8 (CH), 136.4 (CH), 137.2 (CH), 149.0 (CH), 150.0 (CH), 160.6 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>14</sub>H<sub>9</sub>N<sub>3</sub>O<sub>3</sub> [M+H]<sup>+</sup> = 268.0722, found 268.0712.

**8-nitro-3-(pyridine-4-yl)quinolin-2(1H)-one 11** (C<sub>14</sub>H<sub>9</sub>N<sub>3</sub>O<sub>3</sub>) was purified by chromatography on silica gel using dichloromethane/acetone (80/20) as eluent and isolated to yield a yellow solid (209 mg, 0.78 mmol, 71%); m.p. 297 °C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ: 7.36-7.40 (m, 1H, H6), 7.70-7.72 (m, 2H, H2' and H6'), 7.98 (dd, 1H, J=7.9 and 1.4 Hz, H5), 8.03 (s, 1H, H4), 8.55 (dd, 1H, J= 8.4 and 1.4 Hz, H7), 8.73-8.74 (m, 2H, H3' and H5'), 11.46 (br s, 1H, NH). <sup>13</sup>C NMR (TFA-d, 100 MHz) δ: 121.6 (C), 123.7 (CH), 126.1 (C), 126.7 (2xCH), 130.7 (CH), 132.4 (C), 133.4 (C), 137.5 (CH), 140.6 (2xCH), 144.1 (CH), 153.5 (C), 161.1 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>14</sub>H<sub>9</sub>N<sub>3</sub>O<sub>3</sub> [M+H]<sup>+</sup> = 268.0722, found 268.0713.

**3-(4-hydroxymethylphenyl)-8-nitroquinolin-2(1H)-one 12**

(C<sub>16</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>) was purified by chromatography on silica gel using dichloromethane/acetone (75/25) as eluent and isolated to yield a yellow solid (251 mg, 0.85 mmol, 77%); m.p. 219 °C, <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz) δ: 4.56 (d, 2H, J=5.6, CH<sub>2</sub>), 5.26 (t, 1H, J= 5.6, OH), 7.41-7.46 (m, 3H, H6, H3' and H5'), 7.76-7.79 (m, 2H, H2' and H6'), 8.23 (dd, 1H, J=7.6 and 1.4 Hz, H5), 8.36 (s, 1H, H4), 8.43 (dd, 1H, J= 8.3 and 1.4 Hz, H7), 11.17 (br s, 1H, NH). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>, 100 MHz) δ: 63.1 (CH<sub>2</sub>), 122.1 (CH), 122.6 (C), 126.6 (2xCH), 127.6 (CH), 128.8 (2xCH), 132.6 (C), 132.7 (C), 133.6 (C), 134.0 (C), 136.4 (CH), 137.6 (CH), 143.6 (C), 160.8 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>16</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub> [M+H]<sup>+</sup> = 297.0875, found 297.0878.

**3-(3-hydroxymethylphenyl)-8-nitroquinolin-2(1H)-one 13**

(C<sub>16</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>) was purified by chromatography on silica gel using dichloromethane/acetone (75/25) as eluent and isolated to yield a yellow solid (267 mg, 0.90 mmol, 82%); m.p. 155 °C, <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz) δ: 4.58 (d, 2H, J=5.6 Hz, CH<sub>2</sub>), 5.27 (t, 1H, J= 5.6 Hz, OH), 7.37-7.47 (m, 3H, H6, H4' and H5'), 7.66-7.68 (m, 1H, H6'), 7.73-7.75 (m, 1H, H2'), 8.25 (dd, 1H, J=7.8 and 1.4 Hz, H5), 8.35 (s, 1H, H4), 8.44 (dd, 1H, J= 8.3 and 1.4 Hz, H7), 11.17 (br s, 1H, NH). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>, 75 MHz) δ: 63.3 (CH<sub>2</sub>), 122.1 (CH), 122.6 (C), 127.1 (CH), 127.2 (CH), 127.5 (CH), 127.7 (CH), 128.4 (CH), 132.7 (C), 133.0 (C), 133.7 (C), 135.1 (C), 136.5 (CH), 138.0 (CH), 142.9 (C), 160.6 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>16</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub> [M+H]<sup>+</sup> = 297.0875, found 297.0861.

**3-(2-hydroxymethylphenyl)-8-nitroquinolin-2(1H)-one 14**

(C<sub>16</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>) was purified by chromatography on silica gel using dichloromethane/ethyl acetate (50/50) as eluent and isolated to yield a yellow solid (235 mg, 0.79 mmol, 72%); m.p. 208 °C, <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz) δ: 4.44 (d, J=5.5 Hz, 2H, CH<sub>2</sub>), 5.09 (sl, 1H, OH), 7.28 (dd, 1H, J= 7.5 and 1.4 Hz, H3'), 7.33-7.37 (m, 1H, H6), 7.42-7.46 (m, 2H, H4' and H5'), 7.57 (dd, 1H, J=7.5 and 1.4 Hz, H6'), 8.12 (s, 1H, H4), 8.19 (dd, 1H, J= 7.6 Hz and 1.4 Hz, H5), 8.45 (dd, 1H, J= 8.3 and 1.4 Hz, H7), 11.19 (br s, 1H, NH). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100 MHz) δ: 63.7 (CH<sub>2</sub>), 121.9 (CH), 122.5 (C), 127.8 (CH), 128.4 (CH), 129.8 (CH), 130.1 (CH), 130.6 (CH), 132.7 (C), 133.1 (C), 133.9 (C), 135.7 (CH), 135.8 (C), 139.2 (CH), 139.9 (C), 162.2 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>16</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub> [M+H]<sup>+</sup> = 297.0875, found 297.0869.

**3-(4-formylphenyl)-8-nitroquinolin-2(1H)-one 15** (C<sub>16</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>) was purified by chromatography on silica gel using dichloromethane/cyclohexane (80/20) as eluent, isolated and

recrystallized in acetonitrile to yield a yellow solid (165 mg, 0.56 mmol, 51%); m.p. 243 °C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ: 7.34-7.39 (m, 1H, H<sub>6</sub>), 7.93-8.00 (m, 6H, H<sub>4</sub>, H<sub>5</sub>, H<sub>2</sub>', H<sub>3</sub>', H<sub>5</sub>' and H<sub>6</sub>'), 8.53 (dd, 1H, J= 8.5 Hz and 1.4 Hz, H<sub>7</sub>), 10.08 (s, 1H, CHO), 11.45 (br s, 1H, NH). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz) δ: 121.7 (CH), 122.3 (C), 128.0 (CH), 129.4 (2xCH), 129.8 (2xCH), 132.8 (C), 133.2 (C), 133.4 (C), 135.9 (CH), 136.4 (C), 137.9 (CH), 140.5 (C), 160.5 (C), 191.7 (CH). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>16</sub>H<sub>11</sub>N<sub>2</sub>O<sub>4</sub> [M+H]<sup>+</sup> = 295.0719, found 295.0710.

**3-(3-formylphenyl)-8-nitroquinolin-2(1H)-one 16** (C<sub>16</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>) was purified by chromatography on silica gel using dichloromethane/acetone (75/25) as eluent and isolated to yield a yellow solid (104 mg, 0.35 mmol, 32%); m.p. 209 °C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ: 7.34-7.39 (m, 1H, H<sub>6</sub>), 7.63-7.69 (m, 1H, H<sub>5</sub>'), 7.94-7.99 (m, 2H, H<sub>5</sub> and H<sub>6</sub>'), 8.01 (s, 1H, H<sub>4</sub>), 8.10-8.13 (m, 1H, H<sub>4</sub>'), 8.25-8.26 (m, 1H, H<sub>2</sub>'), 8.53 (dd, 1H, J= 8.4 and 1.4 Hz, H<sub>7</sub>), 10.10 (s, 1H, CHO), 11.5 (br s, 1H, NH). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100 MHz) δ: 121.7 (CH), 122.4 (C), 127.8 (CH), 129.2 (CH), 129.7 (CH), 130.2 (CH), 132.8 (C), 133.1 (C), 133.3 (C), 134.7 (CH), 135.6 (C), 135.8 (CH), 136.6 (C), 137.5 (CH), 160.7 (C), 192.0 (CH). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>16</sub>H<sub>11</sub>N<sub>2</sub>O<sub>4</sub> [M+H]<sup>+</sup> = 295.0719, found 295.0713.

**3-(2-formylphenyl)-8-nitroquinolin-2(1H)-one 17** (C<sub>16</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>) was purified by chromatography on silica gel using cyclohexane/acetone (60/40) as eluent and isolated to yield a yellow solid (210 mg, 0.71 mmol, 65%); m.p. 230-231 °C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ: 7.33-7.37 (m, 1H, H<sub>6</sub>), 7.44 (dd, 1H, J= 7.5 Hz and 1.3 Hz, H<sub>6</sub>'), 7.61-7.65 (m, 1H, H<sub>4</sub>'), 7.68-7.72 (m, 1H, H<sub>5</sub>'), 7.82 (s, 1H, H<sub>4</sub>), 7.93 (dd, 1H, J= 7.6 Hz and 1.4 Hz, H<sub>3</sub>'), 7.99 (dd, 1H, J= 7.8 Hz and 1.4 Hz, H<sub>5</sub>), 8.53 (dd, 1H, J= 8.4 Hz and 1.4 Hz, H<sub>7</sub>), 10.03 (s, 1H, CHO), 11.42 (br s, 1H, NH). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100 MHz) δ: 121.6 (CH), 122.3 (C), 127.8 (CH), 129.5 (CH), 130.8 (CH), 131.0 (CH), 133.0 (C), 133.7 (C), 133.9 (CH), 134.5 (C), 135.0 (C), 135.7 (CH), 136.1 (C), 138.0 (CH), 160.8 (C), 191.2 (CH). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>16</sub>H<sub>11</sub>N<sub>2</sub>O<sub>4</sub> [M+H]<sup>+</sup> = 295.0719, found 295.0718.

**3-(4-methoxycarbonylphenyl)-8-nitroquinolin-2(1H)-one 18** (C<sub>17</sub>H<sub>12</sub>N<sub>2</sub>O<sub>5</sub>) was purified by chromatography on silica gel using dichloromethane as eluent, isolated and recrystallized to yield a yellow solid (257 mg, 0.79 mmol, 72%); m.p. 245 °C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ: 3.95 (s, 3H, CH<sub>3</sub>), 7.33-7.38 (m, 1H, H<sub>6</sub>), 7.84-7.87 (m, 2H, H<sub>2</sub>' and H<sub>6</sub>'), 7.95-7.97 (m, 2H, H<sub>4</sub> and H<sub>5</sub>), 8.12-8.15 (m, 2H, H<sub>3</sub>' and H<sub>5</sub>'), 8.52 (dd, 1H, J= 8.3 and 1.4 Hz, H<sub>7</sub>), 11.43 (br s, 1H, NH). <sup>13</sup>C NMR (CDCl<sub>3</sub>,

100 MHz) δ: 52.3 (CH<sub>3</sub>), 121.6 (CH), 122.4 (C), 127.8 (CH), 128.7 (2xCH), 129.7 (2xCH), 130.4 (C), 132.8 (C), 133.3 (C), 133.5 (C), 135.8 (CH), 137.6 (CH), 139.0 (C), 160.6 (C), 166.7 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>17</sub>H<sub>13</sub>N<sub>2</sub>O<sub>5</sub> [M+H]<sup>+</sup> = 325.0824, found 325.0818.

**3-(3-methoxycarbonylphenyl)-8-nitroquinolin-2(1H)-one 19**

(C<sub>17</sub>H<sub>12</sub>N<sub>2</sub>O<sub>5</sub>) was purified by chromatography on silica gel using dichloromethane/cyclohexane (90/10) as eluent and isolated to yield a yellow solid (253 mg, 0.78 mmol, 71%); m.p. 212 °C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ: 3.95 (s, 3H, CH<sub>3</sub>); 7.33-7.37 (m, 1H, H<sub>6</sub>), 7.53-7.58 (m, 1H, H<sub>5</sub>'), 7.96 (dd, 1H, J= 7.7 Hz and 1.4 Hz, H<sub>5</sub>), 7.98 (s, 1H, H<sub>4</sub>), 8.05-8.11 (m, 2H, H<sub>4</sub>' and H<sub>6</sub>'), 8.36-8.37 (m, 1H, H<sub>2</sub>'), 8.51 (dd, 1H, J= 8.4 Hz and 1.4 Hz, H<sub>7</sub>), 11.42 (br s, 1H, NH). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100 MHz) δ: 52.3 (CH<sub>3</sub>), 121.6 (CH), 122.5 (C), 127.7 (CH), 128.6 (CH), 129.6 (CH), 130.0 (CH), 130.5 (C), 132.8 (C), 133.2 (C), 133.4 (CH), 133.5 (C), 134.8 (C), 135.7 (CH), 137.3 (CH), 160.7 (C), 166.7 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>17</sub>H<sub>13</sub>N<sub>2</sub>O<sub>5</sub> [M+H]<sup>+</sup> = 325.0824, found 325.0809.

**3-(2-methoxycarbonylphenyl)-8-nitroquinolin-2(1H)-one 20**

(C<sub>17</sub>H<sub>12</sub>N<sub>2</sub>O<sub>5</sub>) was purified by chromatography on silica gel using cyclohexane/acetone (70/30) as eluent and isolated to yield a yellow solid (257 mg, 0.79 mmol, 72%); m.p. 212 °C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ: 3.82 (s, 3H, CH<sub>3</sub>), 7.29-7.33 (m, 1H, H<sub>6</sub>), 7.37-7.39 (m, 1H, H<sub>6</sub>'), 7.49-7.53 (m, 1H, H<sub>4</sub>'), 7.60-7.64 (m, 1H, H<sub>5</sub>'), 7.77 (s, 1H, H<sub>4</sub>), 7.91 (d, 1H, J=7.65 Hz, H<sub>3</sub>'), 8.02 (d, 1H, J= 7.8 Hz, H<sub>5</sub>), 8.48 (d, 1H, J= 8.4 Hz, H<sub>7</sub>), 11.34 (br s, 1H, NH). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100 MHz) δ: 52.3 (CH<sub>3</sub>), 121.4 (CH), 122.7 (C), 127.2 (CH), 129.0 (CH), 130.1 (CH), 130.6 (CH), 131.0 (C), 132.4 (CH), 132.9 (C), 133.4 (C), 135.5 (CH), 135.6 (CH), 135.8 (C), 137.0 (C), 160.9 (C), 167.5 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>17</sub>H<sub>13</sub>N<sub>2</sub>O<sub>5</sub> [M+H]<sup>+</sup> = 325.0824, found 325.0819.

**General procedure for the preparation of compounds 21-23**

A mixture of 40 mL of H<sub>2</sub>O/Ethanol (2/8) was added onto 1 equiv. (200 mg) of the corresponding 3-(methoxycarbonylphenyl)-8-nitroquinolin-2(1H)-one derivative (**18-20**). Then, 5 equiv. of NaOH were added and the reaction mixture was stirred at 80°C for 3 h. The reaction mixture was then poured into water, acidified to pH=1 with HCl 37%, extracted twice with dichloromethane (2x50 mL) and twice with ethyl acetate (2x50 mL). The combined organic layers were washed with water, dried over anhydrous MgSO<sub>4</sub> and evaporated *in vacuo*. The crude residues were purified by chromatography on silica gel using adapted eluent and recrystallized if necessary to give compounds **21-23**.

**3-(4-carboxyphenyl)-8-nitroquinolin-2(1H)-one 21** (C<sub>16</sub>H<sub>10</sub>N<sub>2</sub>O<sub>5</sub>) was washed with acid water and recrystallized in acetonitrile to yield a yellow solid (124 mg, 0.40 mmol, 65%); m.p. > 310 °C, <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz) δ: 7.45-7.49 (m, 1H, H<sub>6</sub>), 7.93-7.96 (m, 2H, H<sub>2</sub>' and H<sub>6</sub>'), 8.03-8.06 (m, 2H, H<sub>3</sub>' and H<sub>5</sub>'), 8.26 (dd, 1H, J=7.9 and 1.4 Hz, H<sub>5</sub>), 8.46 (dd, 1H, J=8.3 and 1.4 Hz, H<sub>7</sub>), 8.49 (s, 1H, H<sub>4</sub>), 11.23 (br s, 1H, NH), 13.07 (br s, 1H, COOH). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>, 100 MHz) δ: 122.3 (CH), 122.35 (C), 128.1 (CH), 129.2 (2xCH), 129.6 (2xCH), 131.0 (C), 131.7 (C), 132.8 (C), 133.8 (C), 136.7 (CH), 139.1 (CH), 139.5 (C), 160.4 (C), 167.5 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>16</sub>H<sub>11</sub>N<sub>2</sub>O<sub>5</sub> [M+H]<sup>+</sup> 311.0668, found 311.0669.

**3-(3-carboxyphenyl)-8-nitroquinolin-2(1H)-one 22** (C<sub>16</sub>H<sub>10</sub>N<sub>2</sub>O<sub>5</sub>) was washed with acid water and recrystallized in acetonitrile to yield a brown solid (88 mg, 0.28 mmol, 42%); Tdec. 305-306 °C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ: 7.44-7.48 (m, 1H, H<sub>6</sub>), 7.60-7.64 (m, 1H, H<sub>5</sub>'), 8.00-8.02 (m, 1H, H<sub>6</sub>'), 8.03-8.06 (m, 1H, H<sub>4</sub>'), 8.27 (dd, 1H, J=7.8 and 1.4 Hz, H<sub>5</sub>), 8.39-8.40 (m, 1H, H<sub>2</sub>'), 8.46 (dd, 1H, J=8.3 and 1.4 Hz, H<sub>7</sub>), 8.47 (s, 1H, H<sub>4</sub>), 11.22 (br s, 1H, NH), 13.05 (br s, 1H, COOH). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>, 100 MHz) δ: 122.2 (CH), 122.5 (C), 127.9 (CH), 129.0 (CH), 129.8 (CH), 129.9 (CH), 131.3 (C), 131.8 (C), 132.8 (C), 133.4 (CH), 133.7 (C), 135.6 (C), 136.7 (CH), 138.6 (CH), 160.5 (C), 167.6 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>16</sub>H<sub>11</sub>N<sub>2</sub>O<sub>4</sub> [M+H]<sup>+</sup> = 311.0668, found 311.0672.

**3-(2-carboxyphenyl)-8-nitroquinolin-2(1H)-one 23** (C<sub>16</sub>H<sub>10</sub>N<sub>2</sub>O<sub>5</sub>) was purified by chromatography on silica gel using dichloromethane/ ethyl acetate (50/50) as eluent and isolated to yield a yellow solid (109 mg, 0.35 mmol, 57%); T dec. 282-285 °C, <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz) δ: 7.43-7.48 (m, 2H, H<sub>6</sub> et H<sub>6</sub>'), 7.54-7.58 (m, 1H, H<sub>4</sub>'), 7.66-7.70 (m, 1H, H<sub>5</sub>'), 7.91 (dd, 1H, J=7.7 and 1.4 Hz, H<sub>5</sub>), 8.13 (s, 1H, H<sub>4</sub>), 8.20 (dd, 1H, J= 7.8 and 1.4 Hz, H<sub>3</sub>'), 8.42 (dd, 1H, J=8.3 and 1.4 Hz, H<sub>7</sub>) ; 11.08 (br s, 1H, NH), 12.72 (br s, 1H, COOH). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>, 100 MHz) δ: 122.1 (CH), 122.6 (C), 127.4 (CH), 129.1 (CH), 129.8 (CH), 131.3 (CH), 132.4 (CH), 132.5 (C), 132.7 (C), 133.9 (C), 136.1 (CH), 136.2 (C), 136.3 (CH), 136.4 (C), 160.7 (C), 168.4 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>16</sub>H<sub>11</sub>N<sub>2</sub>O<sub>5</sub> [M+H]<sup>+</sup> 311.0668, found 311.0673.

#### Preparation of 3-(4-carboxyphenyl)-2-methoxy-8-nitroquinoline 24

Under Ar atmosphere, 260 mg of 3-(4-methoxycarbonylphenyl)-8-nitroquinolin-2(1H)-one (0.80 mmol, 1 equiv.) were solubilized in 5 mL of dry DMF. This solution was then added onto 64 mg of 60% sodium hydride (1.6 mmol, 2 equiv.), solubilized in DMF (5 mL). After 10 min of stirring at rt, 100 μL of methyl iodide (1.6 mmol, 2 equiv.) were added

dropwise. The reaction mixture was stirred at rt overnight, before being poured into ice. A precipitate was formed, filtered off and washed with water. The precipitate was solubilized into dichloromethane. The solution was dried over anhydrous MgSO<sub>4</sub> and evaporated *in vacuo*. The crude residue was purified by chromatography on silica gel using dichloromethane as eluent. Compound **24** was isolated to yield a pale white solid (76%, 0.61 mmol, 206 mg). Then, 2-methoxy-3-(4-methoxycarbonylphenyl)-8-nitroquinoline (0.44 mmol, 150 mg) was solubilized into 40 mL of a mixture of H<sub>2</sub>O/Ethanol (2/8). 88 mg of NaOH (2.2 mmol, 5 equiv.) were then added and the reaction mixture was stirred at 80°C for 2 h. The reaction mixture was then poured into water, acidified to pH=1 with HCl 37%, extracted twice with dichloromethane (2x50 mL) and twice with ethyl acetate (2x50 mL). The combined organic layers were washed with water, dried over anhydrous MgSO<sub>4</sub> and evaporated *in vacuo*. Compound **24** was washed with basic water to yield a white solid (87 mg, 0.27 mmol, 61%)

Compound **24** (C<sub>17</sub>H<sub>12</sub>N<sub>2</sub>O<sub>5</sub>); Tdec. 270-274 °C, <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz) δ: 4.00 (s, 3H, CH<sub>3</sub>), 7.61-7.65 (m, 1H, H<sub>6</sub>), 7.80-7.82 (m, 2H, H<sub>2</sub>' and H<sub>6</sub>'), 8.05-8.07 (m, 2H, H<sub>3</sub>' and H<sub>5</sub>'), 8.23 (dd, 1H, J=7.8 and 1.4 Hz, H<sub>5</sub>), 8.27 (dd, 1H, J=8.1 and 1.4 Hz, H<sub>7</sub>), 8.56 (s, 1H, H<sub>4</sub>), 13.07 (br s, 1H, NH). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>, 100 MHz) δ: 54.5 (CH<sub>3</sub>), 124.1 (CH), 124.7 (C), 126.6 (CH), 127.0 (C), 129.7 (2xCH), 130.0 (2xCH), 130.9 (C), 132.7 (CH), 136.7 (C), 139.3 (CH), 140.0 (C), 146.2 (C), 160.8 (C), 167.5 (C). HRMS (DCI-CH<sub>4</sub>) calcd for C<sub>17</sub>H<sub>13</sub>N<sub>2</sub>O<sub>5</sub> [M+H]<sup>+</sup> = 325.0824, found 325.0822.

#### Preparation of 8-nitro-3-phenylethynylquinolin-2(1H)-one 25

100 mg of 3-bromo-8-nitroquinolin-2(1H)-one previously synthesized in our team (0.37 mmol, 1 equiv.), 7 mg of CuI (0.037 mmol, 0.1 equiv.) and 43 mg of Pd(PPh<sub>3</sub>)<sub>4</sub> (0.037 mmol, 0.1 equiv.) were added in a sealed flask of 10 mL. Under Argon atmosphere, 5 mL of dry dimethoxyethane 155 μL of Et<sub>3</sub>N (1.11 mmol, 3 equiv.), 61 μL of phenylacetylene (0.56 mmol, 1.5 equiv.) were successively added. The reaction mixture was cooled at 15°C during 1.5 hours. The reaction mixture was poured into water and extracted three times with dichloromethane. The organic layer was washed with water, dried over anhydrous MgSO<sub>4</sub> and evaporated *in vacuo*. The crude residue was purified by chromatography on silica gel using dichloromethane as an eluent and isolated to yield a yellow solid (78 mg, 0.27 mmol, 72%).

1 Compound **25** (C<sub>17</sub>H<sub>10</sub>N<sub>2</sub>O<sub>3</sub>); Pf: 186-187 °C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz)  
2 δ: 7.31-7.35 (m, 1H, H6), 7.37-7.38 (m, 2H, H3', H4' and H5'), 7.59-7.63  
3 (m, 2H, H2' and H6'), 7.87 (dd, 1H, J= 7.6 and 1.5 Hz, H5), 8.06 (s, 1H,  
4 H4), 8.49 (dd, 1H, J= 8.4 and 1.4 Hz, H7), 11.40 (br s, 1H, NH). <sup>13</sup>C NMR  
5 (CDCl<sub>3</sub>-d<sub>6</sub>, 100 MHz) δ: 83.4 (C), 97.3 (C), 119.6 (C), 121.8 (CH), 122.0  
6 (C), 122.1 (C), 128.0 (CH), 128.4 (2xCH), 129.2 (CH), 132.1 (2xCH),  
7 133.0 (C), 133.0 (C), 135.3 (CH), 141.6 (CH), 159.9 (C). HRMS (DCI-  
8 CH<sub>4</sub>) calcd for C<sub>17</sub>H<sub>13</sub>N<sub>2</sub>O<sub>5</sub> [M+H]<sup>+</sup> 291.0770, found 291.0776.

#### 14 Preparation of 8-nitro-2-phenyl-furo[2,3-*b*]quinoline **26**

15 100 mg of 3-bromo-8-nitroquinolin-2(1*H*)-one previously synthesized in  
16 our team (0.37 mmol, 1 equiv.), 7 mg of CuI (0.037 mmol, 0.1 equiv.) and  
17 43 mg of Pd(PPh<sub>3</sub>)<sub>4</sub> (0.037 mmol, 0.1 equiv.) were added in a sealed  
18 flask of 10 mL. Under Argon atmosphere, 5 mL of dry dimethoxyethane  
19 155 μL of Et<sub>3</sub>N (1.11 mmol, 3 equiv.), 61 μL of phenylacetylene (0.56  
20 mmol, 1.5 equiv.) were successively added. The reaction mixture was  
21 heated at 40°C during 30 min. The reaction mixture was poured into  
22 water and extracted three times with dichloromethane. The organic layer  
23 was washed with water, dried over anhydrous MgSO<sub>4</sub> and evaporated *in*  
24 *vacuo*. The crude residue was purified by chromatography on silica gel  
25 using dichloromethane as an eluent and isolated to yield a grey solid (55  
26 mg, 0.19 mmol, 51%).

27 Compound **26** (C<sub>17</sub>H<sub>10</sub>N<sub>2</sub>O<sub>3</sub>); Pf: 226 °C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ:  
28 7.16 (s, 1H, H3), 7.45-7.55 (m, 3H, H3', H4' and H5'), 7.56-7.59 (m, 1H,  
29 H6), 7.99-8.02 (m, 2H, H2' and H6'), 8.08 (dd, 1H, J= 7.5 and 1.4 Hz, H5),  
30 8.14 (dd, 1H, J= 8.3 and 1.4 Hz, H7), 8.41 (s, 1H, H4). <sup>13</sup>C NMR (CDCl<sub>3</sub>,  
31 100 MHz) δ: 99.2 (CH), 123.3 (CH), 123.4 (CH), 124.0 (C), 125.9 (2xCH),  
32 127.7 (C), 128.2 (CH), 128.6 (C), 129.1 (2xCH), 130.5 (CH), 132.0 (CH),  
33 136.0 (C), 147.7 (C), 159.6 (C), 162.4 (C). HRMS (DCI-CH<sub>4</sub>) calcd for  
34 C<sub>17</sub>H<sub>11</sub>N<sub>2</sub>O<sub>3</sub> [M+H]<sup>+</sup> 291.0770, found 291.0768.

#### 34 Electrochemistry

35 Voltammetric measurements were carried out with a potentiostat Autolab  
36 PGSTAT100 (ECO Chemie, The Netherlands) controlled by GPES 4.09  
37 software. Experiments were performed at room temperature in a  
38 homemade airtight three-electrode cell connected to a vacuum/argon  
39 line. The reference electrode consisted of a saturated calomel electrode  
40 (SCE) separated from the solution by a bridge compartment. The counter  
41 electrode was a platinum wire of approximately 1 cm<sup>2</sup> apparent surface.  
42 The working electrode was GC microdisk (1.0 mm of diameter – Bio-logic

SAS). The supporting electrolyte (nBu<sub>4</sub>N)[PF<sub>6</sub>] (Fluka, 99% puriss electrochemical grade) and the solvent DMSO (Sigma-Aldrich puriss p.a. dried < 0.02% water) were used as received and simply degassed under argon. The solutions used during the electrochemical studies were typically 10<sup>-3</sup> M in compound and 0.1 M in supporting electrolyte. Before each measurement, the solutions were degassed by bubbling Ar and the working electrode was polished with a polishing machine (Presi P230). Under these experimental conditions employed in this work, the half-wave potential (E<sub>1/2</sub>) of the ferrocene Fc<sup>+/0</sup> couple in DMSO was E<sub>1/2</sub> = 0.45 V vs SCE. Experimental peak potentials have been measured versus SCE and converted to NHE by adding 0.241 V.

#### 34 Biology

##### Antileishmanial activity on *L. infantum axenic amastigotes*. [31]

*L. infantum* promastigotes (MHOM/MA/67/ITMAP-263, CNR Leishmania, Montpellier, France, expressing luciferase activity) in logarithmic phase cultivated in RPMI 1640 medium supplemented with 5% foetal calf serum (FCS), 2 mM L-glutamine and antibiotics (100 U/mL penicillin and 100 μg/mL streptomycin), were centrifuged at 900 g for 10 min. The supernatant was removed carefully and was replaced by the same volume of RPMI 1640 complete medium at pH 5.4 and incubated for 24 h at 24 °C. The acidified promastigotes were incubated for 24 h at 37 °C in a ventilated flask. Promastigotes were then transformed into axenic amastigotes. The effects of the tested compounds on the growth of *L. infantum* axenic amastigotes were assessed as follows. *L. infantum* amastigotes were incubated at a density of 2.10<sup>6</sup> parasites/mL in sterile 96-well plates with various concentrations of compounds dissolved in DMSO (final concentration less than 0.5% v/v), in duplicate. Appropriate controls DMSO, amphotericin B, miltefosine and fexinidazole (reference drugs purchased from Sigma Aldrich) were added to each set of experiments. After a 48 h incubation period at 37 °C, each plate-well was then microscope-examined for detecting any precipitate formation. To estimate the luciferase activity of axenic amastigotes, 80 μl of each well are transferred to white 96-well plates, Steady Glow® reagent (Promega) was added according to manufacturer's instructions, and plates were incubated for 2 min. The luminescence was measured in Microbeta Luminescence Counter (PerkinElmer). Inhibitory concentration 50% (IC<sub>50</sub>) was defined as the concentration of drug required to inhibit by 50% the metabolic activity of *L. infantum* amastigotes compared to control. IC<sub>50</sub> values were calculated by non-linear regression analysis processed

1 on dose response curves, using TableCurve 2D V5 software. IC<sub>50</sub> values  
2 represent the mean of three independent experiments.

#### 3 *Antitrypanosomal activity on T. brucei brucei trypomastigotes.*

4  
5 Assays were performed on *Trypanosoma brucei brucei* AnTat 1.9 strain  
6 (IMTA, Antwerpen, Belgium). It was cultured in MEM with Earle's salts,  
7 supplemented according to the protocol of Baltz et al.[32] with the  
8 following modifications, 0.5 mM mercaptoethanol (Sigma Aldrich®,  
9 France), 1.5 mM L-cysteine (Sigma Aldrich®), 0.05 mM bathocuproïne  
10 sulfate (Sigma Aldrich®) and 20% heat-inactivated horse serum (Gibco®,  
11 France), at 37°C in an atmosphere of 5% CO<sub>2</sub>. The parasites were  
12 incubated at an average density of 2000 parasites/well in sterile 96-wells  
13 plates (Mc2®, France) with various concentrations of compounds  
14 dissolved in DMSO (Sigma Aldrich®), in duplicate. Reference drugs  
15 suramin, eflornithine, and fexinidazole (purchased from Sigma Aldrich,  
16 France and Fluorochem, UK) suspended in NaCl 0.9% or DMSO, were  
17 added to each set of experiments. The effects of the tested compounds  
18 were assessed by the viability marker Alamar Blue® (Fisher, France)  
19 assay described by Ráz et al.[33] After a 69 h incubation period at 37 °C,  
20 10 µL of Alamar Blue® was then added to each well, and the plates were  
21 incubated for 5h.[34] The plates were read in a PerkinElmer ENSPIRE  
22 (Germany) microplate reader using an excitation wavelength of 530 nm  
23 and an emission wavelength of 590 nm. IC<sub>50</sub> were calculated by  
24 nonlinear regression analysis processed on dose-response curves, using  
25 GraphPad Prism software (USA). IC<sub>50</sub> was defined as the concentration  
26 of drug necessary to inhibit by 50% the viability of *T. brucei brucei*  
27 compared to the control. IC<sub>50</sub> values were calculated from three  
28 independent experiments in duplicate.

#### 29 *Antitrypanosomal activity on T. brucei trypomastigotes overexpressing* 30 *the nitroreductase (NTR1).*

31 *Trypanosoma brucei* bloodstream-form 'single marker' S427 (T7RPOL  
32 TETR NEO) and drug-resistant cell lines were cultured at 37°C in HMI9-T  
33 medium [35] supplemented with 2.5 µg ml<sup>-1</sup> G418 to maintain expression  
34 of T7 RNA polymerase and the tetracycline repressor protein.  
35 Bloodstream trypanosomes overexpressing the *T. brucei* nitroreductase  
36 (NTR1).[36] were grown in medium supplemented with 2.5 µg ml<sup>-1</sup>  
37 phleomycin and expression of NTR was induced by the addition of 1 µg  
38 ml<sup>-1</sup> tetracycline. Cultures were initiated with 1 × 10<sup>5</sup> cells ml<sup>-1</sup> and sub-  
39 cultured when cell densities approached 1–2 (× 10<sup>6</sup>) ml<sup>-1</sup>. In order to

40 examine the effects of inhibitors on the growth of these parasites,  
41 triplicate cultures containing the inhibitor were seeded at 1 × 10<sup>5</sup>  
42 trypanosomes ml<sup>-1</sup>. Cells overexpressing NTR were induced with  
43 tetracycline 48 h prior to EC<sub>50</sub> analysis. Cell densities were determined  
44 after culture for 72 h, as previously described [37]. EC<sub>50</sub> values were  
45 determined using the following two-parameter equation by non-linear

$$y = \frac{100}{1 + \left( \frac{[I]}{EC_{50}} \right)^m}$$

46 regression using GraFit: where the  
47 experimental data were corrected for background cell density and  
48 expressed as a percentage of the uninhibited control cell density. In this  
49 equation [I] represents inhibitor concentration and m is the slope factor.

#### 50 *Cytotoxic evaluation on HepG2 cell line.*

51 The evaluation of the tested molecules cytotoxicity on the HepG2  
52 (hepatocarcinoma cell line from ECACC purchased from Sigma-Aldrich,  
53 ref 85011430-1VL certificated without mycoplasma) was done according  
54 to the method of Mosman with slight modifications.[38] Briefly, cells (1 ×  
55 10<sup>5</sup> cells/mL) in 100 µL of complete medium, [Alpha MEM Eagle from  
56 PAN BIOTECH supplemented with 10% foetal bovine serum, 2 mM L-  
57 glutamine and antibiotics (100 U/mL penicillin and 100 µg/mL  
58 streptomycin)] were seeded into each well of 96-well plates and  
59 incubated at 37 °C and 5% CO<sub>2</sub>. After a 24 h incubation, 100 µL of  
60 medium with various product concentrations and appropriate controls  
61 were added and the plates were incubated for 72 h at 37 °C and 5% CO<sub>2</sub>.  
62 Each plate-well was then microscope-examined for detecting possible  
63 precipitate formation before the medium was aspirated from the wells.  
64 100 µL of MTT solution (0.5 mg/mL in Alpha MEM Eagle) were then  
65 added to each well. Cells were incubated for 2 h at 37 °C and 5% CO<sub>2</sub>.  
66 After this time, the MTT solution was removed and DMSO (100 µL) was  
67 added to dissolve the resulting formazan crystals. Plates were shaken  
68 vigorously (300 rpm) for 5 min. The absorbance was measured at 570  
69 nm with a microplate spectrophotometer (Eon BioTek). DMSO was used  
70 as blank and doxorubicin (purchased from Sigma Aldrich) as positive  
71 control. CC<sub>50</sub> were calculated by non-linear regression analysis  
72 processed on dose response curves, using TableCurve 2D V5 software.  
73 CC<sub>50</sub> values represent the mean value calculated from three independent  
74 experiments.

## Acknowledgements


The authors thank the Université Paul Sabatier de Toulouse and the Région Occitanie / Pyrénées-Méditerranée for funding this work. A. Fairlamb and S. Wyllie are supported by funding from Wellcome Trust (WT105021). C. Bijani from the NMR facility Laboratoire de Chimie de Coordination de Toulouse, C. Claparols, N. Martins-Froment, V. Bourdon and E. Leroy from the mass spectrometry service of Institut de Chimie de Toulouse are also acknowledged for their support, respectively for NMR and HRMS experiments.

**Keywords:** Trypanosomatids • 8-nitroquinolin-2(1*H*)-one series • Antikinetoplastid pharmacomodulation • Parasitic nitroreductases • Pallado-catalyzed cross-coupling reactions.

## References:


- [1] P. Capewell, N. J. Veitch, C. M. Turner, C. M. R. Turner, J. Raper, M. Berriman, S. L. Hadjuk and A. MacLeod, *PLoS Negl. Trop. Dis.* **2011**, *5*, e1287.
- [2] P. Büscher, G. Cecchi, V. Jamonneau, and G. Priotto *Lancet*, **2017**, *390*, 2397-2405.
- [3] D. Pace, *J. Infect.* **2014**, *69*, S10-S18.
- [4] M. M. Ashkan and K. M. Rahim, *Trop. Doct.* **2008**, *38*, 186-188.
- [4] [http://www.who.int/neglected\\_diseases/diseases/en/](http://www.who.int/neglected_diseases/diseases/en/)
- [5] World Health Organization: <http://www.who.int/news-room/fact-sheets/detail/leishmaniasis> updated 03/2018
- [6] World Health Organization: [http://www.who.int/news-room/fact-sheets/detail/trypanosomiasis-human-african-\(sleeping-sickness\)](http://www.who.int/news-room/fact-sheets/detail/trypanosomiasis-human-african-(sleeping-sickness)), updated 02/2018
- [7] M. C. Field, D. Horn, A. H. Fairlamb, M. A. J. Ferguson, D. W. Gray, K. D. Read, M. De Rycker, L. S. Torrie, P. G. Wyatt, S. Wyllie, and I. H. Gilbert *Nat. Rev. Microbiol.* **2017**, *15*, 217-231.
- [8] G. Eperon, M. Balasegaram, J. Potet, C. Mowbray, O. Valverde and F. Chappuis, *Expert Rev. Anti Infect., Ther.* **2014**, *12*, 1407-1417.
- [9] B. Zulfiqar, T. B. Shelper and V. M. Avery, *Drug Discov. Today*, **2017**, *22*, 1516-1531.
- [10] S. Patterson and A. H. Fairlamb, *Curr. Med. Chem.* **2018**, *in press*, DOI: 10.2174/0929867325666180426164352.
- [11] R. T. Jacobs, B. Nare, S. A. Wring, M. D. Orr, D. Chen, J. M. Sligar, M. X. Jenks, R. A. Noe, T. S. Bowling, L. T. Mercer, C. Rewerts, E. Gaukel, J. Owens, R. Parham, R. Randolph, B. Beaudet, C. J. Bacchi, N. Yarett, J. J. Plattner, Y. Freund, C. Ding, T. Akama, Y. K. Zhang, R. Brun, M. Kaiser, I. Scandale, and R. Don, *PLoS Negl. Trop. Dis.* **2011**, *5*, e1151.
- [12] S. Wyllie, S. Patterson, L. Stojanovski, F. R. C. Simeons, S. Norval, R. Kime, K. D. Read and A. H. Fairlamb, *Sci. Transl. Med.* **2012**, *4*, e119.
- [13] V. K. B. K. Mesu, W. M. Kalonji, C. Bardonneau, O. V. Mordt, S. Blesson, F. Simon, S. Delhomme, S. Bernhard, W. Kuziena, J. F. Lubaki, S. L. Vuvu, P. N. Ngima, H. M. Mbembo, M. Ilunga, A. K. Bonama, J. A. Heradi, J. L. L. Salomo, G. Mandula, L. K. Badibabi, F. R. Dama, P. K. Lukula, D. N. Tete, C. Lumbala, B. Scherrer, N. Strub-Wourgaft and A. Tarral, *Lancet* **2018**, 144-154.
- [14] S. Patterson, S. Wyllie *Trends Parasitol.* **2014**, *30*, 289-298. Nitro drugs
- [15] S. Wyllie, S. Patterson and A. H. Fairlamb, *Antimicrob. Agents Chemother.* **2013**, *57*, 901-906.
- [16] S. Wyllie, A. J. Roberts, S. Norval, S. Patterson, B. Foth, M. Berriman, K. Read and A. H. Fairlamb, *PLoS Pathog.* **2016**, *12*, e1005971/22.
- [17] B. S. Hall, X. Wu, L. Hu and S. R. Wilkinson, *Antimicrob. Agents Chemother.* **2010**, *54*, 1193-1199.
- [18] P. Verhaeghe, P. Rathelot, S. Rault and P. Vanelle, *Lett. Org. Chem.* **2006**, *3*, 891-897.
- [19] L. Paloque, P. Verhaeghe, M. Casanova, C. Castera-Ducros, A. Dumètre, L. Mbatchi, S. Hutter, M. Kraiem-M'Rabet, M. Laget, V. Remusat, P. Rathelot, N. Azas and P. Vanelle, *Eur. J. Med. Chem.* **2012**, *54*, 75-86.
- [20] C. Kieffer, A. Cohen, P. Verhaeghe, S. Hutter, C. Castera-Ducros, M. Laget, V. Remusat, M. Kraiem-M'Rabet, S. Rault, P. Rathelot, N. Azas and P. Vanelle, *Eur. J. Med. Chem.* **2015**, *92*, 282-294.
- [21] C. Kieffer, A. Cohen, P. Verhaeghe, L. Paloque, S. Hutter, C. Castera-Ducros, M. Laget, S. Rault, A. Valentin, P. Rathelot, N. Azas and P. Vanelle, *Bioorg. Med. Chem.* **2015**, *23*, 2377-2386.
- [22] J. Pedron, C. Boudot, S. Hutter, S. Bourgeade-Delmas, J.-L. Stigliani, A. Sournia-Saquet, A. Moreau, E. Boutet-Robinet, L. Paloque, E. Mothes, M. Laget, L. Vendier, G. Pratiel, S. Wyllie, A. H. Fairlamb, N. Azas, B. Courtioux, A. Valentin and P. Verhaeghe, *Eur. J. Med. Chem.* **2018**, *155*, 135-152.
- [23] V. P. Andreev, Y. P. Nizhnik, *Russ. J. Org. Chem.* **2002**, *38*, 137-138.
- [24] N. J. O'Brien, M. Brzozowski, D. J. D. Wilson, L. W. Deady and B. M. Abbott, *Bioorg. Med. Chem.* **2014**, *22*, 3781-3790.
- [25] M. S. Tremblay, M. Halim and D. Sames, *JACS*, **2007**, *129*, 7570-7577.
- [26] E. Chong and L. L. Schafer, *Org. Lett.* **2013**, *15*, 6002-6005.
- [27] S. D. Kuduk, J. W. Skudlarek, C. N. Di Marco, J. G. Bruno, M. A. Pausch, J. A. O'Brien, T. D. Cabalu, J. Stevens, J. Brunner, P. L. Tannenbaum, A. L. Gotter, C. J. Winrow, J. J. Renger and P. J. Coleman, *Bioorg. Med. Chem. Lett.* **2014**, *24*, 1784-1789.
- [28] R. Zhou, W. Wang, Z. Jiang, K. Wang, X. Zheng, H. Fu, H. Chen and R. Li, *Chem. Commun.* **2014**, *50*, 6023-6026.
- [29] H. Van de Poël, G. Guillaumet, M.C. Viaux-Massuard, *Heterocycles* **2002**, *57*, 55-71.
- [30] O. A. Volkov, C. C. Cosner, A. J. Brockway, M. Kramer, M. Booker, S. Zhong, A. Ketcherside, S. Wei, J. Longgood, M. K. McCoy, T. E. Richardson, S. A. Wring, M. Peel, J. D. Klinger, B. A. Posner, J. K. De Brabander and M. A. Phillips, *ACS Infect. Dis.*, **2017**, *3*, 512-526.
- [31] C. Zhang, S. Bourgeade-Delmas, A. F. Alvarez, A. Valentin, C. Hemmert and H. Gornitzka, *Eur. J. Med. Chem.* **2018**, *143*, 1635-1643.
- [32] T. Baltz, D. Baltz C. Giroud and J. Crockett, *EMBO J.* **1985**, *4*, 1273-1277.
- [33] B. Rätz, M. Iten, Y. Grether-Bühler and R. Kaminsky, *Acta. Trop.* **1997**, *68*, 139-147.
- [34] J. Guillon, A. Cohen, R. Nath Das, C. Boudot, N. Gueddouda, S. Moreau, L. Ronga, S. Savrimoutou, S. Rubio, S. Amaziane, A. Dassonville-Klimpt, N. Azas, B. Courtioux, J. L. Mergny, C. Mullié and P. Sonnet, *Chem. Biol. Drug. Des.* **2018**, *68*, 1-22.
- [35] N. Greig, S. Wyllie, S. Patterson and A. H. Fairlamb, *FEBS J.* **2009**, *276*, 376-386.
- [36] S. Wyllie, B. J. Foth, A. Kelner, A. Y. Sokolova, M. Berriman, A. H. Fairlamb, *Antimicrob. Agents Chemother.* **2016**, *71*, 625-634.
- [37] D. C. Jones, I. Hallyburton, L. Stojanovski, K. D. Read, J. A. Frearson and A. H. Fairlamb, *Biochem. Pharmacol.*, **2010**, *80*, 1478-1486.
- [38] T. J. Mossman, *J. Immunol. Methods* **1983**, *65*, 55-63.

## Entry for the Table of Contents



An antikinetoplastid pharmacomodulation at position 3 of the scaffold was conducted using an optimized Suzuki-Miyaura cross-coupling reaction. Among the 24 molecules synthesized, compound **21** was identified as a new selective antitrypanosomal hit with a lower reduction potential. Unlike the initial hit, compound **21** is not efficiently bioactivated by the type I trypanosomal nitroreductase which suggests an additional mechanism of action.





[Click here to access/download](#)

**Supporting Information**

Supporting info ChemMedChem Pedron July 2018.docx

