



University of Dundee

Real-world effectiveness and safety of Vedolizumab for the treatment of Inflammatory Bowel Disease

Plevris, N.; Chuah, C. S.; Allen, R. M.; Arnott, I. D.; Brennan, P. N. ; Chaudhary, S.

Published in:

Journal of Crohn's and Colitis

DOI:

[10.1093/ecco-jcc/jjz042](https://doi.org/10.1093/ecco-jcc/jjz042)

Publication date:

2019

Document Version

Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Plevris, N., Chuah, C. S., Allen, R. M., Arnott, I. D., Brennan, P. N., Chaudhary, S., Churchhouse, A. M. D., Din, S., Donoghue, E., Gaya, D. R., Groome, M., Jafferbhoy, H. M., Jenkinson, P. W., Lam, W. L., Lyons, M., Macdonald, J. C., MacMaster, M., Mowat, C., Naismith, G. D., ... Lees, C. W. (2019). Real-world effectiveness and safety of Vedolizumab for the treatment of Inflammatory Bowel Disease: The Scottish Vedolizumab Cohort. *Journal of Crohn's and Colitis*, 13(9), 1111-1120. <https://doi.org/10.1093/ecco-jcc/jjz042>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Title: Real-world effectiveness and safety of Vedolizumab for the treatment of Inflammatory Bowel Disease: The Scottish Vedolizumab Cohort

Short title: Real-world effectiveness and safety of vedolizumab for Inflammatory Bowel Disease

N Plevris¹, C S Chuah¹, R M Allen², I D Arnott¹, P N Brennan³, S Chaudhary⁴, A MD Churchhouse⁵, S Din¹, E Donoghue⁶, D R Gaya², M Groome³, H M Jafferbhoy⁵, P W Jenkinson^{1,7}, W L Lam², M Lyons¹, J C Macdonald⁸, M MacMaster², C Mowat³, G D Naismith⁹, L F Potts⁷, E Saffouri², J P Seenan⁸, A Sengupta⁵, P Shasi³, D I Sutherland⁴, J A Todd³, J Veryan², A JM Watson⁷, D A Watts⁶, G R Jones¹, C W Lees¹

Author Affiliations:

1. The Edinburgh IBD Unit, Western General Hospital, Edinburgh, EH4 2XU, UK
2. Department of Gastroenterology, Glasgow Royal Infirmary, Glasgow, G4 0SF
3. Department of Gastroenterology, Ninewells Hospital, Dundee, DD2 9SY,
4. Department of Gastroenterology, University Hospital Hairmyres, East Kilbride, G75 8RG
5. Department of Gastroenterology, Victoria Hospital, Kirkcaldy, KY2 5AH
6. Department of Gastroenterology, Forth Valley Royal Hospital, Larbert, FK5 4WR
7. Department of Colorectal Surgery, Raigmore Hospital, Inverness, IV2 3UJ
8. Department of Gastroenterology, Queen Elizabeth University Hospital, Glasgow, G51 4TF
9. Department of Gastroenterology, Royal Alexandra Hospital, Paisley, PA2 9PN
10. Department of Gastroenterology, Raigmore Hospital, Inverness, IV2 3UJ

1
2
3
4 **Corresponding Author:** Dr N Plevris, Clinical Research Fellow, the Edinburgh IBD Unit,
5
6 Western General Hospital, Edinburgh, EH4 2XU, UK. Tel: +44 07702230863, Email:
7
8 nikolas.plevris@nhslothian.scot.nhs.uk
9

10 11 12 13 **Disclosures**

14
15 *Declaration of funding interests:* None
16

17
18 *Declaration of personal interests:* CSC, GRJ, RMA, PB, PNB, SC, AMDC, ED, MG, HMJ,
19
20 PWJ, WLL, ML, JCM, MM, CM, JDN, LFP, ES, JPS, AS, PS, DIS, JAT, JV, AJMW, DAW
21
22 have no personal interests to declare in relation to this work. NP has received consultancy
23
24 fees from Takeda, speaker fees and/or travel support from Abbvie, Takeda, Norgine. IDA
25
26 has received consultancy fees from Vifor and travel support from Shire. SD has received
27
28 travel support from Dr Falk and Abbvie. DRG has received speaker fees and/or travel
29
30 support from Ferring, Takeda, Pfizer, Abbvie, Vifor. CWL has received research support
31
32 from Abbvie and Shire, consultancy fees from Abbvie, Pfizer, Dr Falk, Hospira, MSD,
33
34 Pharmacosmos, Takeda and Vifor, speaker fees and/or travel support from Abbvie, Pfizer,
35
36 Dr Falk, Ferring, Hospira, MSD, Shire, Takeda and Warner-Chilcott.
37
38
39
40
41
42

43 **Authorship contributions:**

44
45 NP study design, data collection, analysis and writing of manuscript. CSC data collection
46
47 and writing of manuscript. GRJ, CWL study design, analysis and critical revision of
48
49 manuscript. RMA, IDA, PB, PNB, SC, AMDC, SD, ED, DRG, MG, HMJ, PWJ, WLL, ML,
50
51 JCM, MM, CM, GDN, LFP, ES, JPS, AS, PS, DIS, JAT, JV, AJMW, DAW data collection
52
53 and critical revision of manuscript. All authors approved the final version of the manuscript.
54
55
56
57
58
59
60

ABSTRACT

Introduction

Vedolizumab is an anti-a4b7 monoclonal antibody that is licensed for the treatment of moderate to severe Crohn's disease and ulcerative colitis.

Aims

To establish the real-world efficacy and safety of vedolizumab in the treatment of inflammatory bowel disease.

Methods

Retrospective study involving six NHS health boards in Scotland between June 2015 and November 2017. Inclusion criteria included a diagnosis ulcerative colitis or Crohn's disease with objective evidence of active inflammation at baseline (HBI \geq 5 / Partial Mayo \geq 2 plus CRP >5 g/L or faecal calprotectin \geq 250 μ g/g or inflammation on endoscopy / MRI) and at least one clinical follow up by 12 months. Kaplan-Meier survival analysis was used to establish 12-month cumulative rates of clinical remission (HBI <5 or Partial Mayo <2), mucosal healing (absence of ulceration/erosions on colonoscopy/MRI or Mayo score 0) and deep remission (clinical remission plus mucosal healing). Rates of serious adverse events were described quantitatively.

Results

Our cohort consisted of 180 patients with ulcerative colitis and 260 with Crohn's disease. Combined median follow up was 52 weeks (IQR 26-52 weeks). In ulcerative colitis, 12-month cumulative rates of clinical remission, mucosal healing and deep remission were 57.4%, 47.3% and 38.5% respectively. In Crohn's disease, 12-month cumulative rates of

1
2 clinical remission, mucosal healing and deep remission were 58.4%, 38.9% and 28.3%
3
4 respectively. The serious adverse event rate was 15.6 per 100 patient years of follow up.
5
6
7

8 **Conclusion**

9
10 Vedolizumab is a safe and effective treatment for achieving both clinical remission and
11
12 mucosal healing in ulcerative colitis and Crohn's disease.
13
14
15
16
17

18 **Keywords:** Vedolizumab, real-world, Crohn's disease, Ulcerative colitis, mucosal healing.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

1. INTRODUCTION

The anti-TNF monoclonal antibodies have revolutionized the treatment of patients with inflammatory bowel disease (IBD). Clinical trials have clearly demonstrated their ability to induce and maintain clinical remission but also achieve mucosal healing in patients with IBD.^{1,2,3} However, up to 30% and 45% of patients will experience a primary loss or secondary loss of response respectively.⁴ Furthermore, the systemic immunosuppressive effect of anti-TNF agents is associated with significant morbidity.⁵ Therefore, alternative treatments with different mechanisms of actions are required.

Vedolizumab is a human monoclonal antibody that targets $\alpha 4\beta 7$ integrin and blocks lymphocyte trafficking to the gut via the MADCAM pathway.⁶ Following the phase 3 GEMINI trials, that demonstrated efficacy in induction and maintenance of clinical remission in both ulcerative colitis (UC) and Crohn's disease (CD), vedolizumab was approved as the first gut selective biologic for the treatment of moderate to severe inflammatory bowel disease (IBD).^{7,8,9} In GEMINI 1 and 2, 41.8% and 39.0% of UC and CD responders were in clinical remission at week 52 respectively.^{7,8} However, because of the strict inclusion and exclusion criteria adopted in the seminal trials (e.g. age restrictions, exclusion of patients with stomas, stricturing disease or multiple small bowel resections) the populations treated are not always representative of those encountered in routine clinical practice.¹⁰ Therefore, establishing the effectiveness and safety of vedolizumab in real-world settings is essential, especially when considering how it should be positioned in treatment algorithms.

To this date a number of real world studies have published data on the effectiveness and safety of vedolizumab.¹¹ However, limitations of several studies include their small sample size, small anti-TNF naive numbers, lack of long-term follow up and confinement to specialist tertiary centres.¹¹ Furthermore, with the paradigm shift of treating beyond clinical symptoms

1 to normalisation of faecal calprotectin (FC) and mucosal healing, increasing data is required
2
3
4 on the ability of vedolizumab to achieve these objective end-points in the real world.¹²
5
6 Therefore, in this study we aimed to assess the real-world effectiveness and safety of
7
8 vedolizumab utilising a large multicentre Scottish cohort of patients with IBD, primarily
9
10 looking to assess cumulative rates of clinical remission and mucosal healing by 12-months.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

2. METHODS

2.1 Study Design

This was a multi-centre retrospective cohort analysis involving thirteen hospitals within six NHS health-boards across Scotland (NHS Lothian [co-ordinating site], NHS Greater Glasgow and Clyde, NHS Lanarkshire, NHS Fife, NHS Forth Valley, NHS Highlands). Data were retrospectively collected at each hospital by review of electronic medical records.

2.2 Data Collection

A standardised electronic data collection pro-forma was used at all sites. Baseline parameters for patient phenotyping were obtained. Follow up data on clinical disease activity (Harvey-Bradshaw Index [HBI] / Partial Mayo score/ Physician's global assessment [PGA]), faecal calprotectin (FC), endoscopy, radiology, surgery, steroid use and adverse events were collected where possible via review of electronic medical records. All data had been prospectively collected as part of routine clinical care.

2.3 Participants

Patients receiving vedolizumab for IBD between June 2015 and November 2017 were identified following review of prescribing, infusion suite and electronic medical records at each site. Inclusion criteria for our analysis were: (a) confirmed diagnosis of CD or UC (based on standard clinical, radiological, endoscopic and histological criteria); (b) active disease prior to starting vedolizumab defined by either a HBI ≥ 5 or Partial Mayo score ≥ 2 plus CRP ≥ 5 mg/L and/or endoscopic / radiographic assessment and/or FC ≥ 250 $\mu\text{g/g}$ (c) completion of standard induction dosing (0, 2, 6 weeks); (d) at least one clinical follow up (assessment of disease activity score) within 12-months of initiation of vedolizumab. Patients with inflammatory bowel disease unknown (IBDU) were included in the UC analysis unless pathology favoured CD. Patients who did not complete induction were excluded.

1
2 Patients on vedolizumab but no objective evidence of active disease at baseline or receiving
3
4 treatment for the indication of pouchitis and microscopic colitis were also excluded from the
5
6 analysis.
7
8
9

10 11 **2.4 Primary and Secondary Outcomes**

12
13 Our primary outcomes were the proportion of patients achieving clinical remission and
14
15 mucosal healing. Secondary outcomes included: proportion of patients achieving deep
16
17 remission (clinical remission plus mucosal healing); change in FC during follow up; rates of
18
19 resectional surgery; **baseline predictors of outcome;** and serious adverse events.
20
21 Achievement of mucosal healing was determined from endoscopy or radiology reports.
22
23 Serious adverse events were defined as any event leading to disruption or discontinuation
24
25 of therapy, hospitalisation or death.
26
27
28
29
30
31

32 33 **2.5 Ulcerative Colitis Definitions**

34
35 Clinical remission was defined as a Partial Mayo score <2 and complete tapering of steroids.
36
37 Mucosal healing was assessed either by flexible sigmoidoscopy or ileo-colonoscopy and
38
39 defined as an Endoscopic Mayo score of 0 with complete tapering of steroids. Where a score
40
41 was not documented it was determined by review of endoscopy reports / photos. At least
42
43 one assessment of mucosal healing by flexible sigmoidoscopy or ileo-colonoscopy was
44
45 available in 101/180 UC patients (56.1%).
46
47
48
49

50 51 **2.6 Crohn's Disease Definitions**

52
53 Clinical remission was defined as a HBI <5 off steroids (PGA used in patients with stoma).
54
55 Mucosal healing was defined as the absence of mucosal ulceration / erosions on ileo-
56
57 colonoscopy and off steroids. In patients where ileo-colonoscopy was not possible mucosal
58
59 healing was assessed by MRI (defined according to local site radiologist) or capsule
60

1
2 endoscopy (defined according to local site physician) where available. We were unable to
3
4 apply any validated MRI scoring indices, therefore a sensitivity analysis was performed
5
6 excluding individuals who had a mucosal assessment based on radiology alone. At least
7
8 one assessment of mucosal healing was available in 140/260 CD patients (53.8%),
9
10 performed via; ileo-colonoscopy in 114 (81.4%); MRI in 25 (17.9%) because of stricturing
11
12 disease (n=21) or proximal small bowel disease (n=4); and capsule endoscopy in 1 (0.7%)
13
14 because of proximal small bowel disease.
15
16
17
18
19

20 **2.7 Statistical Analysis**

21
22 SPSS Version 24 (IBM Inc., Chicago, IL, USA) and Prism Version 7.0 (Graphpad Software,
23
24 San Diego, CA, USA) were used for statistical analyses and generation of graphs.
25
26 Descriptive statistics are presented as medians with interquartile range (IQR) for continuous
27
28 variables and frequencies with percentages for categorical variables. For comparison of
29
30 non-parametric continuous variables, the Mann-Whitney U or Kruskal-Wallis test was used
31
32 where appropriate. For comparison of categorical variables, the Chi-squared test was used.
33
34 Primary and secondary outcomes were assessed using Kaplan-Meier survival analysis and
35
36 comparisons made using the log-rank test. Patients were censored at failure, last follow up
37
38 (if less than 365 days) or at 365 days follow up. To try and account for some of the possible
39
40 bias introduced from retrospective observational data with varying follow up, we performed
41
42 a non-responder imputation (NRI) analysis to obtain a conservative estimate of outcomes at
43
44 12 months. Cox proportional hazard regression analyses were carried out to identify
45
46 possible baseline predictors of clinical remission, mucosal healing and deep remission.
47
48 Variables for analysis were chosen a-priori and are listed in Supplementary table 1.
49
50 Variables from the univariable analysis with a p-value <0.20 were fitted and a stepwise
51
52 backward selection approach was adopted to identify significant predictors. A p-value <0.05
53
54 was considered significant for all statistical tests.
55
56
57
58
59
60

2.8 Ethics

This study was considered a retrospective audit by the local ethics committee as all data was collected as part of routine clinical care; hence no formal ethical approval was necessary. Caldicott guardian approval (NHS Lothian) was granted for data collection, storage and submission for publication (Application ID: 1845).

For Review Only

3. RESULTS

3.1 Patient Cohort

A total of 481 patients receiving vedolizumab were identified. Of these, 180 UC and 260 CD patients fulfilled our inclusion criteria for analysis (Figure 1, Table 1). The median follow up (time from initiation to last infusion or discontinuation) was 51 weeks (IQR 26-52 weeks) for CD and 52 weeks (IQR 26-52 weeks) for UC. Patient demographics and disease characteristics are shown in Table 1. Amongst the CD patients, 230 (88.5%) had been previously exposed to anti-TNF with 62.7% receiving vedolizumab as either 3rd or 4th line biologic. Amongst the UC patients, 111 (61.7%) had been previously exposed to anti-TNF.

3.2 Ulcerative colitis outcomes

The cumulative rates of clinical remission (Partial Mayo score <2) were 20.0%, 43.3% and 57.4% at 3, 6 and 12 months respectively. The median time to first endoscopic assessment was 13 weeks (IQR 13-51 weeks). Cumulative rates of mucosal healing (Endoscopic Mayo score of 0) were 17.9% and 47.3% after 6 and 12 months of treatment respectively. **In those achieving outcome, median time to clinical remission and mucosal healing was 14 weeks (IQR 13-26 weeks) and 26 weeks (IQR 22-51 weeks) respectively.** The cumulative rates of deep remission (mucosal healing plus clinical remission) at 6 and 12 months were 14.3% and 38.5% respectively. Utilising a NRI analysis at 12 months; clinical remission, mucosal healing and deep remission was achieved in 19.4% (n=35/180), 15.8% (n=16/101) and 10.9% (n=11/101) of patients respectively. Cumulative clinical remission rates were significantly higher in patients without endoscopic follow up compared to those with (74.6% vs 43.6% at 12 months, log-rank p<0.01) (Supplementary Figure 1). A significant drop in faecal calprotectin levels was observed after 3, 6 and 12 months of treatment compared to baseline (Figure 2).

Cumulative rates of colectomy were 6.0% and 12.4% after 6 and 12 months of vedolizumab treatment. Amongst these patients, n=7 had received IV steroids for severe disease (n=2/7 also started on concomitant cyclosporine) and underwent colectomy within 4 months of starting vedolizumab due to non-response.

3.3 Predictors of treatment outcomes in ulcerative colitis

Multivariable Cox regression analysis identified receiving vedolizumab as first line treatment as the only independent predictor of clinical remission (HR 1.9, 95% CI 1.24-2.91, $p<0.01$), mucosal healing (HR 2.32, 95% CI 1.11-4.89, $p=0.03$) and deep remission (HR 3.78, 95% CI 1.55-9.22, $p<0.01$) (Table 2). Stratifying patients by previous TNF exposure revealed significantly higher cumulative rates of clinical remission (71.4% vs 48.0% at 12 months, log-rank $p<0.01$), mucosal healing (61.6% vs 34.3% at 12 months, log-rank $p=0.03$) and deep remission (58.4% vs 20.0% at 12 months, log-rank $p<0.01$) in patients who were anti-TNF naïve (Table 2, Figure 3). No significant difference was observed in cumulative rates of colectomy between anti-TNF naïve and exposed patients (7.2% vs 15.5%, $p=0.15$) (Figure 2). Comparison of baseline disease activity measures showed no difference in median partial Mayo scores (6 vs 6, $p=0.33$) (*partial Mayo scores available in n=120*) or median CRP levels (4.0 vs 5.0 mg/L, $p=0.11$) (*CRP available in n=164*) between anti-TNF naïve and exposed patients. However, anti-TNF patients had significantly lower median FC levels at baseline (745 vs 1050 $\mu\text{g/g}$) (*FC available in n=116*) but higher rates of concomitant steroid use (62.3% vs 43.2%, $p=0.01$) compared to those who were anti-TNF exposed.

3.3 Crohn's disease outcomes

Cumulative rates of clinical remission were 21.9%, 39.9% and 58.4% after 3, 6 and 12 months of vedolizumab treatment respectively. The median time to first mucosal assessment was 28 weeks (IQR 14-48 weeks). Cumulative rates of mucosal healing were

1
2 10.1% and 38.9% after 6 and 12 months of treatment respectively. In those achieving
3
4 remission, median time to clinical remission and mucosal healing was 14 weeks (IQR 12-26
5
6 weeks) and 39 weeks (IQR 21-50 weeks), respectively.
7
8
9

10
11 Analysis of mucosal healing, including only patients with endoscopic follow up (n=114),
12
13 revealed cumulative rates of 12.2% and 42.6% at 6 and 12 months respectively.
14
15 Furthermore, cumulative rates of deep remission at 6 and 12 months were 7.4% and 28.3%
16
17 respectively. Utilising a NRI analysis at 12 months; clinical remission, mucosal healing and
18
19 deep remission was achieved in 16.2% (n=42/260), 9.3% (n=13/140) and 7.1% (n=10/140)
20
21 of patients respectively. Cumulative clinical remission rates were similar in patients without
22
23 follow up for mucosal healing compared to those with (62.5% vs 55.3% at 12 months, log-
24
25 rank p=0.25) (Supplementary Figure 1). We observed a significant drop in faecal calprotectin
26
27 levels after 3, 6 and 12 months of treatment compared to baseline (Figure 2).
28
29
30
31
32

33
34 Cumulative rates of resectional surgery were 5.4% and 13.1% after 6 and 12 months of
35
36 vedolizumab respectively (pan-proctocolectomy with ileostomy n=10; subtotal colectomy
37
38 n=1; ileocaecal resection n=8; ileocaecal resection for stricturing disease n=2; ileocaecal
39
40 resection for penetrating disease n=3; ileocaecal resection for perforation n=2). Surgical
41
42 procedures for perianal disease were performed in 11 patients (fistulotomy n=1; fistulectomy
43
44 n=2; examination under anaesthesia with drainage of abscess and seton insertion n=8).
45
46
47
48
49

50 **3.4 Predictors of treatment outcomes in Crohn's disease**

51
52 Multivariable Cox regression analysis revealed individuals with an IBD related
53
54 hospitalisation in the last 12 months were less likely to achieve clinical remission (HR 0.59,
55
56 0.40-0.86, p<0.01) (Table 3). However, no predictive factors were identified for achieving
57
58 mucosal healing or deep remission (Table 3). Stratifying patients by prior IBD admissions
59
60

1
2 revealed significantly higher cumulative rates of clinical remission in those that had not been
3
4 hospitalised due to their IBD in the 12 months prior to vedolizumab (65.3% vs 49.1% at 12
5
6 months, log-rank $p < 0.01$). However, no difference was seen in cumulative mucosal healing
7
8 (61.6% vs 34.3% at 12 months, log-rank $p = 0.03$), deep remission (58.4% vs 20.0% at 12
9
10 months, log-rank $p = 0.003$) or resectional surgery. Comparison of baseline disease activity
11
12 measures showed no difference in median HBI (7 vs 6, $p = 0.07$) (HBI available in $n = 130$)
13
14 and FC levels (540 vs 780 $\mu\text{g/g}$, $p = 0.30$) (FC available in $n = 130$) but did show higher CRP
15
16 levels (7.5 vs 7.0 mg/L , $p = 0.02$) (CRP available in $n = 256$) between patients with previous
17
18 IBD related hospitalisations versus those without.
19
20
21
22
23
24

25 Stratifying patients by previous TNF exposure revealed no significant difference for
26
27 cumulative clinical remission (61.3% vs 57.8% at 12 months, log-rank $p = 0.27$), mucosal
28
29 healing (37.0% vs 39.4% at 12 months, log-rank $p = 0.92$), deep remission (27.3% vs 28.5%
30
31 at 12 months, log-rank $p = 0.91$) and resectional surgery (3.0% vs 14.6% at 12 months, log-
32
33 rank $p = 0.13$) between patients who were anti-TNF naïve and exposed (Supplementary
34
35 Figure 3).
36
37
38
39
40

41 3.4 Safety

42
43 The combined CD and UC cohort had over 333 patient years of follow-up (PYF). Dose
44
45 escalation to Q4 vedolizumab was carried out in 9 UC patients (after 3 months due to sub-
46
47 optimal response) and 6 CD patients (after 6 months due to loss of response [$n = 3$] and sub-
48
49 optimal response [$n = 3$]). The rate of serious adverse events was 15.6 per 100 PYF (Table
50
51 2). The rate of serious infections was 6.3 per 100 PYF, however the majority continued
52
53 vedolizumab following treatment of their infection. There was a total of 4 (0.9%) infusion
54
55 reactions and 2 (0.5%) delayed hypersensitivity reactions, all requiring discontinuation of
56
57 vedolizumab. Arthralgia was the most common non-infective severe adverse event (3.3 per
58
59
60

1
2 100 PYF). No deaths occurred during follow-up. Three malignancies were reported during
3
4 treatment. One patient was diagnosed with a cholangiocarcinoma following 4 doses of
5
6 vedolizumab. However, he was thought to also have concurrent PSC. Another patient was
7
8 diagnosed with metastatic ovarian cancer after 3 doses of vedolizumab. The final patient
9
10 underwent a colectomy due to non-response to vedolizumab and upon review of his
11
12 pathology was found to also have a colonic neuro-endocrine tumour. In patients undergoing
13
14 surgery following vedolizumab, post-operative complications (30-day) were observed in
15
16 22.2% (Clavien Dindo Grade 1: n=2; Grade 2: n=2) and 19.2% (Clavien Dindo Grade 1: n=1;
17
18 Grade 2: n=2; Grade 3: n=2) of the UC and CD patients respectively.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

4. DISCUSSION

Here we present results from a large, multi-centre, real-world cohort investigating the effectiveness and safety of vedolizumab in the treatment of patients with IBD. To our knowledge, this is the largest UK cohort published to date, with data not only on clinical outcomes but also assessment of mucosal healing. In our cohort, we observed 12-month cumulative clinical remission, mucosal healing and deep remission rates of 57.4%, 47.3%, 38.5% in UC respectively; and 58.4%, 38.9%, 28.3% in CD respectively. We were also able to show a significant drop in FC for both CD and UC after 3-months of treatment. Furthermore, with over 333 patient years of follow-up no new safety signals were observed. These results are in keeping with other previously reported experiences and add to the growing body of literature on the effectiveness and safety of vedolizumab in the real world.

Our results support the use and effectiveness of vedolizumab for the treatment of both UC and CD. Despite both cohorts being mature with high anti-TNF exposure (61.7% in UC; 88.5% in CD) by 12-months over 50% of patients were able to achieve clinical remission. Adopting a NRI analysis, clinical remission rates of 19.4% and 16.2%; and mucosal healing rates of 15.8% and 9.3% were observed in UC and CD at 12 months respectively. However, these estimates are conservative and several patients who had missing data at 12 months or had less than 12 months follow up, had already achieved the end point of clinical remission and/or mucosal healing. Furthermore, we did observe in both cohorts that FC dropped significantly by 3-months (Figure 2). When assessing cumulative mucosal healing, superior rates were observed in UC, possibly due to the greater number of anti-TNF naive patients (Table 1). For both cohorts, the greatest benefit was observed after >6 months of treatment. This observation has been reported by others and suggests that in many patients

1
2 vedolizumab may have a slower onset of action and persistence to >6 months of therapy is
3
4 required to achieve optimum outcomes.¹²⁻¹⁵
5
6
7
8
9

10
11 In UC, our results are in keeping with other real-world cohorts from the US (12-month
12 cumulative clinical remission, 51.0%; 12-month cumulative endoscopic remission, 41.0%;
13 NRI 12-month clinical remission, 20.0%; NRI 12-month mucosal healing, 17.0%), France
14 (12-month clinical remission, 42.1%), Sweden (12-month clinical remission, 64.0%), Canada
15 (12-month: clinical remission 62.0%; endoscopic remission 47.8%) and Germany (NRI 12-
16 month clinical remission, 25.0%).¹³⁻¹⁸ Again in CD our results are similar to those reported
17 by others, although there appears to be more variation in published outcomes.^{11,13-18} For
18 example, in the US VICTORY consortium (n=221) which was analysed in a similar way to
19 our study, cumulative 12-month clinical remission was 35.0% and mucosal healing much
20 higher at 63.0%.¹⁴ Their higher mucosal healing rates compared to clinical remission may
21 suggest that a number of the patients had symptoms unrelated to active IBD, especially as
22 patients could be included based on symptoms alone.¹⁴ Individuals may not always have
23 active IBD driving their symptoms (e.g. bile salt malabsorption in patient with small bowel
24 resection, irritable bowel syndrome) or have disproportionate symptoms for the degree of
25 mucosal inflammation. As such, treatments aimed at treating inflammation may show poor
26 efficacy for achieving clinical remission but superior mucosal healing as baseline
27 inflammation is not as severe as expected. In contrast to this our inclusion criteria
28 necessitated objective evidence of inflammation (HBI \geq 5 or Partial Mayo score \geq 2 plus CRP
29 \geq 5 mg/L and/or endoscopic / radiographic assessment and/or FC \geq 250 μ g/g) which might
30 explain some of the differences observed. Other reasons for differences in reported
31 outcomes between our study and the US VICTORY consortium include; differences in the
32 patient populations examined; the use of the PGA to assess efficacy rather than validated
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2 disease activity scores; and the definition of mucosal healing necessitating being off
3
4 steroids.¹⁴
5
6
7
8
9

10 Optimum positioning of vedolizumab for the treatment of IBD is still unknown and highly
11 debated. Currently in the UK, NICE guidelines allow the use of vedolizumab first line in
12 patients with moderate to severe UC but in CD stipulate its use only after anti-TNF failure or
13 if there are contraindications to anti-TNF use.^{19,20} Results from the GEMINI trials indicated
14 higher efficacy in bio-naïve patients for both UC and CD, something that has been echoed
15 in other real-world cohorts.^{7-9,11} In the VICTORY consortium study they showed that previous
16 anti-TNF exposure was associated with a reduced probability of achieving clinical remission
17 (HR 0.53 [95% CI 0.38-0.75] for UC; HR 0.40 [95% CI 0.20-0.81] for CD) and mucosal
18 healing (HR 0.51 [95% CI 0.29-0.88] for UC; HR 0.29 [95% CI 0.12-0.73] for CD).^{13,14} We
19 observed similar results in patients with UC, where Cox regression analysis identified first
20 line use of vedolizumab as the only independent predictor of clinical remission (HR 1.9, 95%
21 CI 1.24-2.91, p<0.01), mucosal healing (HR 2.32, 95% CI 1.11-4.89, p=0.03) and deep
22 remission (HR 3.78, 95% CI 1.55-9.22, p<0.01) (Table 2). Although, anti-TNF exposed
23 patients had a higher baseline FC; partial Mayo scores and CRP were similar between the
24 two groups. Furthermore, the proportion of patients on concomitant steroids was higher in
25 the anti-TNF naïve group suggesting significant disease. Interestingly, in a recent network
26 meta-analysis of randomised controlled trials published by Singh et al., vedolizumab and
27 infliximab were ranked highest as first line agents for induction of remission and mucosal
28 healing in patients with moderate to severe UC.²¹ No difference was observed in our CD
29 cohort but our analysis was likely underpowered due to the small number of bio-naïve
30 patients (Supplementary Figure 2). Despite data showing superior efficacy with first line use
31 for both CD and UC, in the absence of head to head trials no definite conclusions can be
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2 made as to its positioning compared to anti-TNF therapy. However, factors such as safety
3
4 and cost, especially in the advent of biosimilar anti-TNF agents, may also influence health
5
6 trusts and providers.
7
8
9

10
11
12 In our CD cohort, we were unable to identify any independent predictors of mucosal healing
13
14 or deep remission (Table 3). However, we did find that in patients who had an IBD related
15
16 hospitalisation in the prior 12 months to vedolizumab, there was a significantly lower
17
18 probability of achieving clinical remission (HR 0.59, 0.40-0.86, $p < 0.01$) (Table 3). This
19
20 observation may be due to more severe disease in those with prior hospitalisation. Other
21
22 studies have also identified severe disease, active perianal disease and smoking history as
23
24 independent predictors of clinical remission in CD.¹⁴
25
26
27
28
29
30
31
32

33 The mechanism of action of vedolizumab theoretically results in reduced systemic
34
35 immunosuppression and thus superior safety profile when compared to other agents like
36
37 anti-TNF or azathioprine.⁶ Recently, Colombel et al. published safety data from six
38
39 vedolizumab trials including 2830 patients with over 4800 patient years of follow up.²²
40
41 Authors found no increased risk of any infection or serious infection with vedolizumab
42
43 exposure compared to placebo.²² Furthermore, infusion reactions were described in $\leq 5\%$ of
44
45 patients and reported malignancy rates of $< 1\%$, in keeping with background risk.²² In our
46
47 cohort, rates of SAEs were low (15.6 per 100 PYF) and in keeping with the published
48
49 literature. Infusion reactions were also uncommon and no deaths occurred during follow up.
50
51 Three malignancies were reported but these all occurred soon after therapy was initiated
52
53 and thought to be unrelated to vedolizumab.
54
55
56
57
58
59
60

1
2 There are several strengths to our study. Firstly, we present results from the largest CD and
3
4 second largest UC cohort published to date with data gathered from multiple centres within
5
6 different NHS health boards in Scotland. This allows representation of a diverse patient
7
8 population, capturing practices from both tertiary referral centres and district general
9
10 hospitals treating patients with IBD. In contrast to others, we have also used standardised
11
12 and validated disease activity scores to assess clinical remission. Furthermore, in the
13
14 paradigm shift of treating to normalisation of FC and mucosal healing, we present data on
15
16 the ability of vedolizumab to achieve these important objective end points. However, we also
17
18 acknowledge certain limitations within our study. The retrospective nature of patient
19
20 identification and data collection may have influenced our results. We have tried to
21
22 overcome this by creating a standardised data collection proforma and using objective
23
24 outcome measures where possible. Furthermore, variability in follow-up intervals and both
25
26 assessment and reporting of mucosal healing may have introduced bias. To account for
27
28 differences in follow-up, Kaplan-Meier analysis was used although our results may have
29
30 been influenced by heavy censoring and / or lack of independence between censoring and
31
32 events. Furthermore, we performed a NRI analysis to provide conservative outcomes
33
34 following the use of vedolizumab. In addition, when we compared cumulative clinical
35
36 remission rates depending on whether patients had a follow up for mucosal healing we found
37
38 that in UC cumulative clinical remission rates were significantly higher in patients who did
39
40 not have endoscopic follow up (Supplementary Figure 1). This suggests a possible selection
41
42 bias for performing endoscopy in more unwell patients, therefore outcomes may be
43
44 underestimated. Finally, as our study was multi-centre, encompassing different types of
45
46 hospitals, we would expect some variations in patient populations and practice patterns.
47
48 Although, as previously stated, our aims were to assess real-world outcomes that are
49
50 generalisable to all practice.
51
52
53
54
55
56
57
58
59
60

1
2 In conclusion, we have shown in a large, real-world, multi-centre cohort that vedolizumab is
3
4 effective at achieving not only clinical remission but mucosal healing at 12-months for both
5
6 UC and CD. In addition, we add to the body of evidence that in UC, first line use results in
7
8 superior outcomes. Finally, our safety data further supports the favourable safety profile of
9
10 vedolizumab in the treatment of IBD. Further real-world prospective as well as head to head
11
12 trials are now required.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

REFERENCES

1. Colombel JF, Sandborn WJ, Reinisch W, et al. Infliximab, azathioprine, or combination therapy for Crohn's disease. *N Engl J Med.* 2010;362(15):1383-1395.
Doi:10.1056/NEJMoa0904492
2. Rutgeerts P, Sandborn WJ, Feagan BG, et al. Infliximab for Induction and Maintenance Therapy for Ulcerative Colitis. *N Engl J Med.* 2005;353(23):2462-2476.
Doi:10.1056/NEJMoa050516
3. Papi C, Fasci-Spurio F, Rogai F, et al. Mucosal healing in inflammatory bowel disease: Treatment efficacy and predictive factors. *Dig Liver Dis.* 2013;45(12):978-985.
Doi:10.1016/j.dld.2013.07.006
4. Roda G, Jharap B, Neeraj N, et al. Loss of Response to Anti-TNFs: Definition, Epidemiology, and Management. *Clin Transl Gastroenterol.* 2016;7.
Doi:10.1038/ctg.2015.63
5. Hoentjen F, van Bodegraven AA. Safety of anti-tumor necrosis factor therapy in inflammatory bowel disease. *World J Gastroenterol.* 2009;15(17):2067-2073.
Doi:10.3748/wjg.15.2067
6. Wyant T, Fedyk E, Abhyankar B. An overview of the mechanism of action of the monoclonal antibody vedolizumab. *J Crohn's Colitis.* 2016;10(12):1437-1444.
Doi:10.1093/ecco-jcc/jjw092

- 1
2 7. Sandborn WJ, Feagan BG, Rutgeerts P, et al. Vedolizumab as induction and maintenance
3
4 therapy for Crohn's disease. *N Engl J Med.* 2013;369(8):711-721.
5
6 Doi:10.1056/NEJMoa1215739
7
8
9
10
11 8. Feagan BG, Rutgeerts P, Sands BE, et al. Vedolizumab as Induction and Maintenance
12
13 Therapy for Ulcerative Colitis. *N Engl J Med.* 2013;369(8):699-710.
14
15 Doi:10.1056/NEJMoa1215734
16
17
18
19
20 9. Sands BE, Feagan BG, Rutgeerts P, et al. Effects of Vedolizumab Induction Therapy for
21
22 Patients With Crohn's Disease in Whom Tumor Necrosis Factor Antagonist Treatment
23
24 Failed. *Gastroenterology.* 2014;147(3):618-627.e3. doi:10.1053/j.gastro.2014.05.008
25
26
27
28
29 10. Ha C, Ullman TA, Siegel CA, et al. Patients enrolled in randomized controlled trials do
30
31 not represent the inflammatory bowel disease patient population. *Clin Gastroenterol*
32
33 *Hepatol.* 2012;10(9):1002-1e78. Doi:10.1016/j.cgh.2012.02.004
34
35
36
37
38 11. Schreiber S, Dignass A, Peyrin-Biroulet L, et al. Systematic review with meta-analysis:
39
40 real-world effectiveness and safety of vedolizumab in patients with inflammatory bowel
41
42 disease. *Journal of Gastroenterology.* 2018:1-17.
43
44
45
46
47 12. Peyrin-Biroulet L, Sandborn W, Sands BE, et al. Selecting Therapeutic Targets in
48
49 Inflammatory Bowel Disease (STRIDE): Determining Therapeutic Goals for Treat-to-Target.
50
51 *Am J Gastroenterol.* 2015;110(9):1324-1338. Doi:10.1038/ajg.2015.233
52
53
54
55
56
57
58
59
60

- 1
2 13. Narula N, Peerani F, Meserve J, et al. Vedolizumab for Ulcerative Colitis: Treatment
3
4 Outcomes from the VICTORY Consortium. *American Journal of Gastroenterology*. 2018:1-
5
6 10.
7
8
9
10
11 14. Dulai PS, Singh S, Jiang X, et al. The real-world effectiveness and safety of vedolizumab
12
13 for moderate-severe Crohn's disease: Results from the US VICTORY consortium. *Am J*
14
15 *Gastroenterol*. 2016;111(8):1147-1155. Doi:10.1038/ajg.2016.236
16
17
18
19
20 15. Amiot A, Serrero M, Peyrin-Biroulet L, et al. One-year effectiveness and safety of
21
22 vedolizumab therapy for inflammatory bowel disease: a prospective multicentre cohort
23
24 study. *Aliment Pharmacol Ther*. 2017;46(3):310-321. Doi:10.1111/apt.14167
25
26
27
28
29 16. Eriksson C, Marsal J, Bergemalm D, et al. Long-term effectiveness of vedolizumab in
30
31 inflammatory bowel disease: a national study based on the Swedish National Quality
32
33 Registry for Inflammatory Bowel Disease (SWIBREG). *Scand J Gastroenterol*. 2017;52(6-
34
35 7):722-729. Doi:10.1080/00365521.2017.1304987
36
37
38
39
40 17. Kotze PG, Ma C, Almutairdi A, et al. Real-world clinical, endoscopic and radiographic
41
42 efficacy of vedolizumab for the treatment of inflammatory bowel disease. *Aliment Pharmacol*
43
44 *Ther*. 2018. Doi:10.1111/apt.14919
45
46
47
48
49 18. Stallmach A, Langbein C, Atreya R, et al. Vedolizumab provides clinical benefit over
50
51 1 year in patients with active inflammatory bowel disease - a prospective multicenter
52
53 observational study. *Aliment Pharmacol Ther*. 2016;44:1199–212.
54
55
56
57
58
59
60

- 1
2 19. National Institute for Health and Care Excellence (2015) Vedolizumab for treating
3
4 moderately to severely active Crohn's disease after prior therapy. Technology appraisal
5
6 guidance (TA352)
7
8
9
10
11
12
13 20. National Institute for Health and Care Excellence (2015) Vedolizumab for treating
14
15 moderately to severely active ulcerative colitis. Technology appraisal guidance (TA342)
16
17
18
19
20 21. Singh S, Fumery M, Sandborn WJ, Murad MH. Systematic review with network meta-
21
22 analysis: first- and second-line pharmacotherapy for moderate-severe ulcerative colitis.
23
24 *Aliment Pharmacol Ther.* 2018;47(2):162-175. Doi:10.1111/apt.14422
25
26
27
28
29 22. Colombel J-F, Sands BE, Rutgeerts P, et al. The safety of vedolizumab for ulcerative
30
31 colitis and Crohn's disease. *Gut.* 2017;66(5):839-851. Doi:10.1136/gutjnl-2015-311079
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Patient demographics and disease characteristics of cohort. CD, Crohn's disease; UC, ulcerative colitis; IFX, infliximab; ADA, adalimumab; UST, ustekinumab.

| | CD (n=260) | | UC (n=180) | |
|--|-----------------------|------------|-----------------------|-----------|
| Median age, years (IQR) | 39 (29-54) | | 41 (30-56) | |
| Median disease duration, years (IQR) | 9 (5-14) | | 6 (3-11) | |
| Female gender, n (%) | 160 (61.5) | | 77 (42.8) | |
| <i>Smoking, n (%)</i> | | | | |
| Current | 23 (8.8) | | 7 (3.9) | |
| Ex | 33 (12.7) | | 30 (16.7) | |
| <i>Disease location / extent, n (%)</i> | L1 | 50 (19.2) | E1 | 15 (8.3) |
| | L2 | 83 (31.9) | E2 | 68 (37.8) |
| | L3 | 127 (48.8) | E3 | 97 (53.9) |
| <i>Disease behaviour n (%)</i> | B1 | 150 (57.7) | - | |
| | B2 | 57 (21.9) | - | |
| | B3 | 53 (20.4) | - | |
| Perianal disease, n (%) | 52 (20.0) | | - | |
| Extra-intestinal manifestations, n (%) | 83 (31.9) | | 44 (24.4) | |
| IBD hospitalisations in last year, n (%) | 106 (40.8) | | 61 (33.9) | |
| Previous Resectional Surgery, n (%) | 108 (41.5) | | - | |
| Previous anti-TNF exposure, n (%) | 226 (86.9) | | 111 (61.7) | |
| <i>Position of vedolizumab, n (%)</i> | | | | |
| 1 st Line | 34 (13.1) | | 69 (38.3) | |
| 2 nd Line | 86 (33.1) | | 86 (47.8) | |
| 3 rd Line | 132 (50.8) | | 24 (13.3) | |
| 4 th Line | 8 (3.1) | | 1 (0.6) | |
| <i>Reason for vedolizumab, n (%)</i> | | | | |
| -Failure of IFX and/or ADA | 199 (76.5) | | 100 (55.6) | |
| -Failure of IFX and/or ADA plus UST | 8 (3.1) | | - | |
| -Contraindication to anti-TNF | 32 (12.3) | | 30 (16.7) | |

| | | |
|---|-------------------|------------------|
| -Intolerance of anti-TNF | 24 (9.2) | 6 (3.3) |
| -Patient preference | 3 (1.2) | 15 (8.3) |
| -Clinician preference | - | 29 (16.1) |
| Vedolizumab dosing, n (%) | | |
| Week 10 dose | | |
| 8-weekly | 53 (20.4) | 4 (2.2) |
| 4-weekly | 253 (97.3) | 180 (100) |
| | 7 (2.7) | 0 |
| Concomitant immunosuppressant, n (%) | 77 (13.0) | 58 (32.2) |
| Concomitant steroids, n (%) | 100 (38.5) | 91 (50.6) |
| Median Harvey-Bradshaw Index, (IQR)* | 6 (5-10) | - |
| Median Partial Mayo Score, (IQR)** | - | 6 (3-8) |
| Median CRP, mg/L (IQR) | 7 (4-15) | 5 (2-10)** |
| Median Faecal Calprotectin*, µg/g (IQR) | 690 (250-1065)* | 840 (500-1500)** |

*Harvey Bradshaw Index, CRP and faecal calprotectin available in n= 130, 256 and 130 at baseline respectively

**Partial Mayo Score, CRP and faecal calprotectin available in n= 120, 164 and 116 at baseline respectively

Table 2. Univariable and multivariable predictors of outcomes in ulcerative colitis. HR, hazard ratio; CI, confidence interval

| | Univariable analyses | | | Multivariable analyses | | |
|--|----------------------|-----------|---------|------------------------|-----------|---------|
| | HR | 95% CI | p-value | HR | 95% CI | p-value |
| Clinical Remission | | | | | | |
| First line treatment | 1.99 | 1.30-3.04 | <0.01 | 1.90 | 1.24-2.91 | <0.01 |
| Age >60 years | 0.51 | 0.27-1.00 | 0.04 | | | |
| Mucosal Healing | | | | | | |
| First line treatment | 2.19 | 1.05-4.59 | 0.04 | 2.32 | 1.11-4.89 | 0.03 |
| IBD related hospitalisation in the prior 12 months | 0.51 | 0.21-1.27 | 0.15 | | | |
| Deep Remission | | | | | | |
| First line treatment | 3.52 | 1.45-8.58 | <0.01 | 3.78 | 1.55-9.22 | <0.01 |
| Female | 0.46 | 0.17-1.24 | 0.12 | | | |
| Age >60 years | 1.96 | 0.77-4.98 | 0.16 | | | |
| IBD related hospitalisation in the prior 12 months | 0.42 | 0.14-1.23 | 0.11 | | | |
| Left sided disease (E2) | 0.41 | 0.11-1.54 | 0.18 | | | |

Table 3. Univariable and multivariable predictors of outcomes in Crohn's disease. HR, hazard ratio; CI, confidence interval

| | Univariable analyses | | | Multivariable analyses | | |
|--|----------------------|-----------|---------|------------------------|-----------|---------|
| | HR | 95% CI | p-value | HR | 95% CI | p-value |
| Clinical Remission | | | | | | |
| Female | 0.73 | 0.51-1.05 | 0.09 | | | |
| Age >60 years | 1.39 | 0.90-2.14 | 0.14 | | | |
| IBD related hospitalisation in the prior 12 months | 0.60 | 0.41-0.88 | <0.01 | 0.59 | 0.40-0.86 | <0.01 |
| Disease duration >5 years | 0.74 | 0.49-1.12 | 0.15 | | | |
| Smoking | 0.57 | 0.27-1.23 | 0.15 | | | |
| Ileo-colonic disease (L3) | 1.76 | 1.03-2.99 | 0.04 | | | |
| Mucosal Healing | | | | | | |
| Age >60 years | 2.01 | 0.95-4.26 | 0.07 | | | |
| Disease duration >5 years | 2.54 | 0.77-8.36 | 0.12 | | | |
| Penetrating disease (B3) | 0.44 | 0.16-1.26 | 0.13 | | | |
| Deep Remission | | | | | | |
| Age >60 years | 1.83 | 0.80-4.21 | 0.16 | | | |
| Concomitant steroids | 1.96 | 0.90-4.27 | 0.09 | | | |
| Penetrating disease (B3) | 0.31 | 0.07-1.30 | 0.11 | | | |

Table 4. Cumulative rates of clinical remission, mucosal healing and deep remission stratified by **(A)** previous TNF exposure in ulcerative colitis **(B)** IBD related hospitalisations in the 12 months prior to vedolizumab in Crohn's disease.

A.

| | Overall | | | anti-TNF naive | | | anti-TNF exposed | | |
|---------------------------|---------|-------|-------|----------------|-------|-------|------------------|-------|-------|
| | 3M | 6M | 12M | 3M | 6M | 12M | 3M | 6M | 12M |
| Clinical Remission | 20.0% | 43.3% | 57.4% | 23.8% | 53.1% | 71.4% | 17.6% | 29.7% | 48.0% |
| Mucosal Healing | 4.2% | 17.9% | 47.3% | 8.6% | 20.9% | 61.6% | 1.6% | 11.8% | 34.3% |
| Deep Remission | 4.2% | 14.3% | 38.5% | 8.6% | 20.9% | 58.4% | 1.6% | 8.0% | 20.0% |

B.

| | Overall | | | Hospitalised | | | Not Hospitalised | | |
|---------------------------|---------|-------|-------|--------------|-------|-------|------------------|-------|-------|
| | 3M | 6M | 12M | 3M | 6M | 12M | 3M | 6M | 12M |
| Clinical Remission | 21.9% | 39.9% | 58.4% | 13.7% | 28.0% | 49.1% | 27.4% | 40.3% | 65.3% |
| Mucosal Healing | 2.3% | 10.1% | 38.9% | 0% | 4.4% | 31.1% | 4.0% | 12.6% | 43.0% |
| Deep Remission | 1.5% | 7.4% | 28.3% | 0% | 4.2% | 27.6% | 2.6% | 9.7% | 28.6% |

Table 5. Serious adverse events. PYF, patient years of follow-up.

| Serious Adverse Events | Incidence, n (%) | Event rate per 100 PYF |
|-----------------------------|------------------|------------------------|
| <i>Serious Infection</i> | 21 (4.7) | 6.3 per 100 PYF |
| -Intra-abdominal collection | 1 (0.2) | |
| -Tonsillitis | 1 (0.2) | |
| -Nasopharyngitis | 2 (0.5) | |
| -Cellulitis | 2 (0.5) | |
| -Shingles | 1 (0.2) | |
| -Pneumonia | 5 (1.1) | |
| -Cholecystitis | 2 (0.5) | |
| -Urogenital Infection | 4 (1.0) | |
| -Septic Arthritis | 1 (0.2) | |
| -Dental Abscess | 1 (0.2) | |
| -Clostridium Difficile | 1 (0.2) | |
| Arthralgia | 11 (2.5) | 3.3 per 100 PYF |
| Infusion reaction | 4 (1.0) | 1.2 per 100 PYF |
| Deranged LFTs | 4 (1.0) | 1.2 per 100 PYF |
| Delayed Hypersensitivity | 2 (0.4) | 0.6 per 100 PYF |
| Malignancy | 3 (0.7) | 0.9 per 100 PYF |
| Bowel Perforation | 2 (0.4) | 0.6 per 100 PYF |
| Headache | 3 (0.7) | 0.9 per 100 PYF |
| Depression | 1 (0.2) | 0.3 per 100 PYF |
| Neutropenia | 1 (0.2) | 0.3 per 100 PYF |

Figure 1. Study flowchart of patients included in analysis. IBD, inflammatory bowel disease; UC, ulcerative colitis; CD, Crohn's disease.




Figure 2. Effect of vedolizumab on faecal calprotectin after 3, 6 and 12 months of therapy in (A) ulcerative colitis and (B) Crohn's disease (faecal calprotectin value used if within +/- 1 month of specified time interval). Dotted line indicates faecal calprotectin cut-off of 250 µg/g.




Figure 3. Kaplan-Meier curves for ulcerative colitis treatment outcomes stratified by previous anti-TNF exposure. **A**, cumulative rates of clinical remission; **B**, cumulative rates of mucosal healing; **C**, cumulative rates of deep remission; **D**, cumulative rates of colectomy.





Figure 4. Kaplan-Meier curves for Crohn's disease treatment outcomes stratified by IBD related hospitalisations in the 12 months prior to vedolizumab. **A**, cumulative rates of clinical remission; **B**, cumulative rates of mucosal healing; **C**, cumulative rates of deep remission; **D**, cumulative rates of surgery.



Supplementary Table 1. (A) Covariates selected for Cox regression model in UC. (B)

Covariates selected for Cox regression model in CD.

A.



| | |
|----------|--|
| 1 | First line treatment |
| 2 | Female |
| 3 | Age >60 years |
| 4 | Disease duration >5 years |
| 5 | Disease extent (E1, E2, E3; reference E1) |
| 6 | Concomitant steroids |
| 7 | Concomitant immunosuppressant |
| 8 | IBD related hospitalisation in the prior 12 months |

Smoking was not included as a covariate as there were only n=7 documented active smokers in the UC cohort.

B.

| | |
|-----------|--|
| 1 | First line treatment |
| 2 | Female |
| 3 | Age >60 years |
| 4 | Disease duration >5 years |
| 5 | Disease distribution (L1, L2, L3; reference L1) |
| 6 | Concomitant steroids |
| 7 | Concomitant immunomodulator |
| 8 | IBD related hospitalisation in the prior 12 months |
| 9 | Peri-anal disease |
| 10 | Week 10 dose |
| 11 | Previous surgery |
| 12 | Smoking |
| 13 | Penetrating disease (B3) |

1
2 **Supplementary Figure 1.** Cumulative clinical remission rates in UC (A) and CD (B).
3
4 depending on whether endoscopic / MRI follow up was available. MA, mucosal
5
6 assessment.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Supplementary Figure 2. Kaplan-Meier curves for Crohn's disease treatment outcomes stratified by previous anti-TNF exposure. **A**, cumulative rates of clinical remission; **B**, cumulative rates of mucosal healing; **C**, cumulative rates of deep remission; **D**, cumulative rates of surgery.

