

Boumpouli, M. and Danton, M. and Gourlay, T. and Kazakidi, A. (2018) Blood flow simulations in models of the pulmonary bifurcation to facilitate treatment of adults with congenital heart disease. In: 31st Scottish Fluid Mechanics Meeting 2018, 2018-05-29 - 2018-05-29, University of Aberdeen.

This version is available at https://strathprints.strath.ac.uk/65500/

Strathprints is designed to allow users to access the research output of the University of Strathclyde. Unless otherwise explicitly stated on the manuscript, Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Please check the manuscript for details of any other licences that may have been applied. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<u>https://strathprints.strath.ac.uk/</u>) and the content of this paper for research or private study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to the Strathprints administrator: strathprints@strath.ac.uk

The Strathprints institutional repository (https://strathprints.strath.ac.uk) is a digital archive of University of Strathclyde research outputs. It has been developed to disseminate open access research outputs, expose data about those outputs, and enable the management and persistent access to Strathclyde's intellectual output.

Blood flow simulations in models of the pulmonary bifurcation to facilitate treatment of adults with congenital heart disease

M.Boumpouli1, M. Danton 1,2, T. Gourlay 1, A. Kazakidi 1 1 Department of Biomedical Engineering, University of Strathclyde, G4 ONW, Glasgow. 2 Scottish Adult Congenital Cardiac Service, Golden Jubilee National Hospital, Clydebank, G81 4DY, Glasgow.

Tetralogy of Fallot (ToF) is the most common cyanotic congenital heart disease, for which patients require surgical intervention at a very young age. Although these patients have long survival rates, they are at risk of chronic complications and frequently require re-operations with the most common being pulmonary valve replacement (PVR). However, the decision for surgical intervention is currently based on clinical indications and the right timing for PVR remains ambiguous [1,2]. The overall objective of this work is to identify a computational metric that will help assess the right timing for surgical intervention in adults with repaired tetralogy of Fallot. This current study concerns a preliminary computational analysis of blood flow in simplified geometries of the pulmonary bifurcation. The focus lies on the effect of geometric and haemodynamic parameters on the wall shear stress patterns around the pulmonary bifurcation.

Different models of the pulmonary bifurcation were created with varied angles and origins of the branches, representing normal and diseased cases. Blood flow simulations were performed based on steady and incompressible flow, governed by the Newtonian Navier-Stokes equations. Physiological vessel dimensions and boundary conditions were applied in all models and different Reynolds numbers were tested.

Blood flow in the pulmonary bifurcation is strongly dependent on geometric and haemodynamic conditions. An increase in the angle of the pulmonary branches cannot be directly correlated with increased regions of recirculation zones downstream of the junction, but more extended regions were observed for higher Reynolds numbers. The origin of the branches also appeared to influence flow separation and recirculation, for the same branching angle. Changes in velocities and shear stresses developed on the vessels walls were observed for each model. Evaluation of these parameters can provide an insight into the underlying flow mechanisms of more complex 3D geometries. Future work will involve the reconstruction of 3D patient-specific geometries from medical images of adult patients with repaired tetralogy of Fallot, pre- and post- operatively. Time-dependent effects will also be tested. These results contribute towards the identification of new reliable approaches for the assessment of PVR and may help towards better medical care.

Acknowledgments

This work is supported in part from the University of Strathclyde Research Studentship Scheme (SRSS) Research Excellence Awards (REA), Project No 1208 and the European Union's Horizon 2020 research and innovation programme, under the Marie Skłodowska-Curie grant agreement No 749185.

References:

- 1. Buechel, E.R.V., et al. European Heart Journal. 2005; 26: 2721-2727.
- 2. Geva T. Circulation. 2013; 128: 1855-1857.