



Heath, Michael and Benkort, Deborah and Brierley, Andrew and Daewel, Ute and Hofmeister, Richard and Proud, Roland and Schrum, Corinna and Speirs, Douglas (2018) Modelling the effects of changes in sea-ice extent on Arctic marine food webs. In: MASTS Annual Science Conference, 2018-10-31 - 2018-11-01, TIC conferenec centre. ,

This version is available at <https://strathprints.strath.ac.uk/67247/>

Strathprints is designed to allow users to access the research output of the University of Strathclyde. Unless otherwise explicitly stated on the manuscript, Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Please check the manuscript for details of any other licences that may have been applied. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<https://strathprints.strath.ac.uk/>) and the content of this paper for research or private study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to the Strathprints administrator: strathprints@strath.ac.uk

Modelling the effects of changes in sea-ice extent on Arctic marine food webs

Michael Heath¹, Deborah Benkort², Andrew Brierley³, Ute Daewel², Richard Hofmeister², Roland Proud³, Corinna Schrum² and Douglas Speirs¹

¹Department of Mathematics and Statistics, University of Strathclyde, UK

²Institute of Coastal Research, Helmholtz-Zentrum Geesthacht, Germany

³School of Biology, University of St Andrews, UK




Retreating Arctic sea-ice cover and increasing primary production

1979





2016



Barents Sea

April
(winter maximum ice extent)




September
(summer minimum ice extent)

Total extent (million km²)

	1979	2016	decrease
winter	15.4	13.7	11%
summer	7.1	4.5	37%

Climate projections indicate continuing loss of ice cover in the future


Increasing Arctic primary production linked to loss of ice cover



Arrigo and van Dijken (2011) JGR Oceans

Images: National Snow and Ice Data Centre, Univ. Colorado

Atlantification of the Barents Sea



Winter ice concentration 1979-2010 (colours), and ice extent (15% concⁿ)


From: Arthun et al. (2012). *Journal of Climate* 25, 4736-4743
See also: Lind et al. (2018). *Nature Climate Change* 8, 634-639

Ice-edge, Svalbard, June 2008



Borealization of Barents Sea fish communities

Shifting distributions of fish communities




Redrawn from Fossheim et al. 2015 *Nature Climate Change* DOI:10.1038/NCLIMATE2647



Retreating ice and the prospect of fisheries in polar waters

On 30 November 2017, the EU plus 9 major fishing nations agreed not to develop fisheries in the Central Arctic Ocean for at least the next 16 years, to give time for development of scientific understanding.

Science, 1 Dec 2017. doi:10.1126/science.aar6437



How will increased primary production propagate up the food web in the absence of sea-ice?



Depending on details of the processes, we might see:

Enhanced pelagic system




Enhanced benthic system

StrathE2E food web model - predicting the ecological effects of multiple types of perturbations



Dynamic, mass conserving network model, driven by input data on hydrodynamics, nutrient fluxes, and fishery harvesting rates

“Middle-out” cascade

Harvesting of top predator species



- Harvesting of mid-trophic level species,
- Benthos mortality



- Nutrient inputs
- Light intensity
- Sediment disturbance
- Mixing and transport


Heath (2012) Prog. Oceanogr. 102, 42-66

Heath et al. (2014) Ecology Letters 17, 101-114

Heath et al. (2014).Nature Communications 5:3893

Four interconnected compartments: Pelagic, Benthic, Scavenging, and Sediment

Birds & mammals (B/M)
 Demersal fish (DF)
 Pelagic fish (PF)
 Carnivorous zooplankton (CZ)
 Carnivorous/scavenging benthos (CSB)
 Omnivorous zooplankton (OZ)
 Susp/deposit feeding benthos (SDB)
 Phytoplankton (P)
 Nitrate (Nit)
 Ammonia (Am)
 Detritus (Det)
 Corpses (Corp)
 Discards (Disc)



Network of coupled ordinary differential equations



- Models flow rates of nitrogen between the living, dead and inorganic components of the food web
- Output at daily intervals
- Computational fitting to observed data

Barents Sea model – annual cycles of physical driving data

Compare two StrathE2E models for an Arctic shelf sea (Barents Sea)



- (1) Baseline : **December-June sea-ice cover**; (2) Scenario : **year-round ice-free**
- Both models – initially, no fishing

Temperature
(0.5°C warmer in the ice-free model)



Vertical mixing
(ice shields the water column from wind mixing)





Light intensity
(ice attenuates incoming light)



Freshwater input
(earlier river runoff in warmer conditions)

Ice-cover baseline model driving data based on 1980's Barents Sea

Comparison of stationary annual cycle in the ice-cover baseline model with 1980's observations from Svalbard




Solid lines – Model simulation with ice-cover December - June

Symbols – Monthly averaged 1980's data from Hornsund, Svalbard

(Węslawski et al. (1988). *Polar Research* 6, 185-189)

Differences between annual average masses of ecosystem components in the ice-free model and the baseline



The increase in phytoplankton biomass is amplified as it cascades up the food web.

BUT – sensitive to:

- *Temperature*
- *Attenuation of light and vertical mixing by ice-cover*
- *Preference parameters defining the accessibility of prey to predators vertically and horizontally*



Annual primary production:

(gC.m⁻².y⁻¹)

Ice-cover model: **50.9**

Ice-free model: **55.4**

How will increased primary production propagate up the food web in the absence of sea-ice?



Ice-free scenario:



Increased draw-down of nitrate and more primary production, as anticipated, but this leads to:

Enhanced benthic system

Increased mixing carries spring primary production deeper into the water column where it becomes accessible to filter feeding benthos



Sensitivity of steady state fisheries yield to ice cover



Modelled fish catches in relation to annual harvest ratio (\equiv fishing mortality rate)

- **Demersal fish yields are greater in the ice-free model than in the baseline – due to enhanced benthos production**
- **Pelagic fish yields are smaller in the ice-free model than in the baseline – due to enhanced predation from demersal fish and mammals**

Take-home messages

- Loss of winter sea-ice leads to increased primary production
- In this model, increased primary production leads to enhancement of the benthos/demersal fish, rather than the pelagic system.
- Sensitivity analysis and more investigation needed, to better represent the physical and biological effects of ice cover
- Potential impact - strategic guidance related to general policy on future exploitation of Arctic living resources
- Scope for public engagement – digestible results to illustrate the threats and opportunities posed by climate change in the Arctic





**Changing
Arctic
Ocean** NERC



**Helmholtz-Zentrum
Geesthacht**

Centre for Materials and Coastal Research



University of
St Andrews



MiMeMo Microbes to Megafauna Modelling of Arctic Seas

www.changing-arctic-ocean.ac.uk/project/mimemo/

Fishing village with cod-drying racks in the Norwegian Arctic