

Pain as a threat to the social self: a motivational account

Kai Karos¹, Amanda C. de C. Williams², Ann Meulders^{1,3,4}, & Johan W.S. Vlaeyen^{1,3,4}

¹ Research Group of Health Psychology, KU Leuven, Belgium

² Research Department of Clinical, Health and Educational Psychology, University College London, London, United Kingdom

³ Center for Excellence on Generalization Research in Health and Psychopathology, KU Leuven, Belgium

⁴ Department of Clinical Psychological Sciences, Maastricht University, The Netherlands

Number of pages: 23

Number of figures: 1

Number of tables: 0

Correspondence concerning this article should be addressed to Kai Karos, Department of Psychology, University of Leuven, Tiensestraat 102, box 3726, 3000 Leuven, Belgium. E-mail: Kai.Karos@kuleuven.be, T: +32 (0)16 32 57 82, F: +32 (0)16 32 61 44.

Keywords: interpersonal; social; evolutionary psychology; needs; motivations; social threat; pain communication; pain behavior; goals

1. Introduction

It has been proposed that the definition of pain ideally recognizes not only sensory, cognitive, and emotional dimensions, but also a *social* dimension [110]. Although it is widely acknowledged that interpersonal context modulates pain experience and communication [39,56], we still fail to understand why and how this modulation occurs. Drawing from evolutionary, social, and health psychology, we argue that pain is a fundamentally social and threatening human experience because it challenges several basic needs (see Fig. 1): (1) the need for autonomy, (2) the need to belong, and (3) the need for justice/fairness. Examining how pain interferes with these basic human needs can help us better understand the dynamic interplay between social context and pain. Here, we (1) define these fundamental human needs, (2) outline how pain threatens these needs, and (3) describe the consequences of such threats, especially for pain itself.

2. The need for autonomy

Even though humans are an inherently social species, they possess a fundamental need for autonomy and a *sense of agency*, a subjective feeling of control over their own actions and their outcomes [13]. To predict and control the environment is at the center of survival fitness, especially in the case of aversive experiences like pain [17,18,100]. We argue that in the context of pain (and illness), the need for autonomy has a fundamentally interpersonal connotation because agency shifts from the person with pain to others.

From an evolutionary perspective, pain can be conceptualized as a homeostatic emotion [16] or an “awareness of a need state” that is communicated to conspecifics [109]. In contrast to more solitary animal species, humans evolved as a reciprocally altruistic species, prioritizing the ability to communicate need states to elicit help from or warn others. This may help to allow survival in otherwise fatal situations [31]. Illness symptoms in general can have

a signaling function [94] and the same is true for facial pain expression [108]. Therefore, a state of illness in general and pain in particular can place humans into a state of dependency on others.

Feelings of helplessness and uncontrollability are common when suffering from pain, especially when it is chronic [83]. Moreover, Western society emphasizes individual function and autonomy, which is severely hampered by chronic illness. Consequently, many people with chronic pain experience shame, embarrassment, and humiliation [88]. These emotions are often fueled by interpersonal worries such as being a burden or whether pain is taken seriously [97]. Last, pain can be the result of victimization at the hand of others, such as in the cases of torture, bullying, or physical assault [3,101]. In the latter, the difference in power and control is paramount, as it places the victim at the mercy of a hostile assailant.

Feelings of uncontrollability and helplessness have adverse effects on physical and psychological health and pain. The debilitating consequences of uncontrollable events have been shown across different species and led to the “learned helplessness hypothesis” of depression [1]. Regarding pain, experimental research has demonstrated that predictability and controllability of pain are major determinants of cognitive pain processing [12,68,70,83,105]. Uncontrollable pain stimuli are perceived as more intense, harmful, and unpleasant than controllable ones [12,61], and subjective feelings of helplessness have been associated with increased pain intensity [69]. Losing control over pain is even worse than never having had control over pain in the first place [17]. Also, feelings of helplessness [83], and shame, guilt, and fear of negative evaluation predict pain intensity levels [97]. Although this research does not emphasize uncontrollability and helplessness in a social domain, pain places the individual in a state where s/he relies on others for support, transferring control to others such as family, friends, and most importantly, healthcare providers. Additionally, doctor-patient relationships are often characterized by authoritarian and paternalistic

physicians and problematic treatment compliance, and many people with chronic pain are dissatisfied with their care [47].

In the case of victimization, consequences are even direr. Victims of bullying are at increased risk of developing chronic pain complaints [30,99,101] and the same is true for victims of torture [3,11], and interpersonal abuse [6]. Pain intentionally inflicted by others is perceived as more intense than incidental pain [37] and is associated with reduced pain communication [51,74,108].

3. The need to belong

Humans as social animals have a pervasive drive to form and maintain at least some lasting, positive, and significant interpersonal relationships [7]. This motivation has been demonstrated across cultures, affects a broad variety of behaviors and leads to serious adverse effects when thwarted [41,111].

Pain threatens the need to belong in numerous ways. First, pain interferes with social activities such as work and hobbies [35]. People with chronic pain experience stigmatization and invalidation (i.e., ignoring, rejecting, or negatively evaluating one's thoughts or feelings) [14,44,78,103,104,107], especially when medical pathology is not confirmed [22,79,80,82]. People in pain are judged as less warm and less competent by pain-free individuals: effectively, a derogated out-group [5]. More broadly, chronic pain conflicts with societal norms that emphasize health, autonomy and functionality until old age and the dominant norm that pain should be short-lived, diagnostically meaningful, and denote a fixable problem [23,73,110].

Individuals with chronic pain are more socially isolated and excluded than healthy controls across the lifespan [33,71,75], especially when pain has no medical explanation [77]. While it has been argued that one of the 'key objects of fear' in chronic pain is threat to self-

identity [65], this threat also directly affects the ‘social self’ [88]. Individuals with chronic pain internalize stigma regarding alienation and discrimination [104]. The link between illness and social isolation is supported by evolutionary theories. Ostracism of ill members has been observed in most social species and across time and culture, increasing group cohesiveness for those not ostracized and enhancing fitness for survival [38,112], for instance by limiting the spread of infectious diseases [87].

Thwarting the need to belong has a host of detrimental consequences for physical and psychological health and for pain. Chronic deprivation of the sense of belonging has been associated with increased stress levels, reduced immune functioning, and increased mortality. Similarly, loneliness is strongly associated with increased risk for morbidity and mortality [2,7,41]. In regard to pain, social exclusion is associated with increased pain reports and vice versa [8,28,112]. Social stress and pain also share overlapping neural pathways, often interpreted as evidence that the human pain detection system has been coopted to detect and react to social threat [26,27,29,53,72]. Social exclusion has been associated with increased [8] but also decreased [20] pain reports. Pain tolerance is positively related to social network size [45], and perceived social isolation in individuals with low-back pain predicts future disability [46,71]. In rheumatoid arthritis pain, perceived stigmatization is related to poorer physical and psychological well-being. Importantly, perceptions of social isolation and stigmatization seem to be more important than actual size of the social network [55,78].

4. The need for justice / fairness

Humans have a universal investment in justice and fairness [10]. Justice and fairness have been of interest for hundreds of years in philosophy, economy, law, and increasingly, psychology [63]. The literature commonly distinguishes between *outcome fairness/justice*, which refers to equality/fairness in outcome and *procedural fairness/justice*, which refers to equality/fairness in the procedure leading to a certain outcome. A preference for fairness and

justice has been demonstrated in humans but also in other species including apes, monkeys, dogs, and birds, providing evidence for an evolutionary basis, a mechanism for payoff evaluation in species that cooperate with others and rely on reciprocal altruism (e.g., tit-for-tat) [9,10,93]. In pain, perceived injustice has been conceptualized as a set of cognitions comprising attributions of blame, magnitude of loss, and irreparability of loss [89].

Perceptions of injustice are both common and detrimental in pain sufferers [89,91], especially when injury or pain can be blamed on someone else's actions or negligence (e.g., a car accident) and when pain is perceived as unnecessary. In such cases, responsibility is attributed by the person with pain to someone else: attribution of blame and responsibility are key processes in making sense of chronic pain [24]. In the case of stigmatization and invalidation of individuals seeking care for persistent pain, it is often assumed that the person with pain has a vested interest in exaggerating or even fabricating pain and other symptoms [80,82,107]. Perceptions of unfair behavior by the person in pain can directly lead to pain underestimation in observers [48,49], and this is particularly marked where medical causes are absent, leading to suspicions of malingering [49]. Overall, perceptions of injustice often arise from the behavior of close others such as colleagues, family, and health-care providers, particularly where the person with pain believes that those others contributed to the injury, inadequately assessed or treated pain, or responded punitively to expression of pain [85]. Some research has also highlighted the importance of "just world" beliefs in perceptions of injustice: a belief that the "environment is a just and orderly place where people usually get what they deserve" (p. 1030) [58]. Pain, especially when chronic, fundamentally challenges this belief, and blame functions to restore just world beliefs. Furthermore, injustice is not merely "perceived" by individuals suffering from pain, but there are a range of systemic and societal injustices in the assessment and treatment of pain based on gender [84], race [21,25,43], ethnicity [66], age [42], and other characteristics of the person in pain. Perceptions

of injustice also play a crucial role in the violations of the needs discussed previously. Experiences of exclusion are often associated with perceptions of injustice [112] and so are experiences of helplessness and uncontrollability, especially when others take advantage of such a situation [36,37,74].

Violations of justice and fairness have a range of detrimental consequences, often categorized as *reactions of withdrawal* and *reactions of attack* [63]. Reactions of withdrawal involve exclusion and discontinuation of prior cooperation; reactions of attack include feelings of anger and retributive behavior that is intended to punish the offender [9]. This set of responses is observed in human and non-human animals and is evolutionarily adaptive, as it promotes future cooperation and deters cheaters and freeloaders. In animals, violations of justice might be met with temporary social exclusion and discontinuation of cooperative relationships as well as displays of anger [10]. In humans, discontinued cooperation can take the form of not complying with authority that is perceived as unjust. Moreover, the link between injustice and anger is well established [62], and perceptions of injustice might be the single greatest cause for feelings of anger, retribution and aggression [4,52,63]. According to the social interaction theory of aggression, the primary function of aggression is to produce some change in others' behavior (e.g., to exact retributive justice for perceived wrongs). Similarly, social exclusion frequently leads to aggression and punishment when it is perceived as unjustified [98,102] and punishment for injustice often occurs even at the cost to the punisher [9].

In pain, perceptions of injustice have been associated with increased pain intensity and unpleasantness [60,95]. In individuals with chronic pain, perceptions of injustice have been associated with a range of problematic outcomes [91], including prolonged work disability and impaired physical functioning [89], increased pain catastrophizing, persistence of post-traumatic stress symptoms [92], heightened protective pain behavior [90], and anger [59].

Anger itself has also been associated with increased pain reports and decreased pain tolerance [96], disability and treatment non-adherence [86,91] and interpersonal consequences, such as social exclusion and reduced social support [32]. Moreover, if the person in pain is perceived as someone who is deceptive and unfairly tries to gain an advantage (e.g., care or financial compensation), this may result in invalidation, stigmatization, pain underestimation, and negative evaluation of the person with pain [48,49,80,81].

5. Future directions

Pain is an inherently social experience and threatens our social needs in three fundamental ways: (1) it shifts control away from the person with pain to others, (2) it excludes, and (3) it is often associated with (perceived) injustice. This framework has direct clinical implications, since social needs are often considered secondary to physical health-related needs. While specific recommendations are premature, the recognition that pain directly affects the social self may help clinicians to better customize their interventions to the needs of those seeking help. This awareness also highlights the responsibility of the healthcare professional not only to meet an individual's pain-related needs, but also to address interpersonal needs, most effectively achieved with a multidisciplinary effort [107].

Specifically, clinical practice should involve (1) standard assessment of interpersonal needs and motivations, (2) recognition of threats to these needs, and (3) attempt to resolve thwarted needs as far as possible, and facilitates active involvement of the social network such as family members. Crucially, considering that threats to these needs also frequently originate on a systematic and societal level, interventions should target these levels as well [78,107].

Critically, the relationship between social needs and pain is *bidirectional*: Pain does thwart social needs but thwarted social needs also have detrimental consequences for health *and* pain (e.g., by affecting the perceived harmfulness of pain [56]). Psychological experiences,

including pain, are powerfully shaped by social context [34,40,67]. Thus, managing social context *is* managing pain.

This framework also generates several paths for further empirical investigation. *First*, there is a need to accurately assess interpersonal needs and motivations within pain research. Assessment methods for these could be borrowed from other fields where available (e.g., [7,112]), or developed from first principles. We have a wide variety of instruments available to assess pain-management motivations, but other motivations (especially interpersonal motivations) are rarely assessed [50]. *Second*, and in line with recent motivational accounts of pain [18,19,106], we need to better understand how pain-related goals and motivations compete or interact with concurrent goals, social or otherwise, and how they affect (pain) behavior [15,39]. It is relevant to know when and how an individual prioritizes certain pain-related goals over others. *Third*, the expansion and integration of different disciplines, levels of analyses, and populations could substantially advance understanding. Specifically, the evolutionary framework underlying this work emphasizes the importance of translating and integrating research findings from different species and disciplines, and to identify universal mechanisms by which social needs affect behavior, and variants [57,64,109]. Similarly, while most empirical research focuses on the individual level (up to small groups), the phenomena highlighted in the present account originate from, or are influenced by, the macro-level as well (e.g., the health-care system, societal and cultural norms) [107]. Several existing lines of research such as gender and sex differences in pain [54], perceptions of injustice [91], or stigma and invalidation [78] would profit from a macro-approach. *Fourth*, there are several other ways by which pain can threaten interpersonal human needs (e.g., challenging the self-concept [76]) which are beyond the scope of this review but deserve further scrutiny.

In sum, dissemination of the biopsychosocial model of pain to health-care professionals, patients, scientists and the general public is crucial to understand that meeting the interpersonal needs of an individual with pain is fundamentally associated with increased physical and psychological well-being of the individual in general, and their experience of pain in particular.

Acknowledgements

The authors report no conflict of interest. Kai Karos is a doctoral researcher supported by the Research Foundation, Flanders, Belgium (grant 1111015N). Ann Meulders is a postdoctoral researcher of the Research Foundation Flanders (FWO-Vlaanderen), Belgium, (grant ID 12E3717N) and is supported by a Vidi grant from the Netherlands Organization for Scientific Research (NWO), The Netherlands (grant ID 452-17-002).

6. References

- [1] Abramson LY, Seligman ME, Teasdale JD. Learned helplessness in humans: Critique and reformulation. *J. Abnorm. Psychol.* 1978;87:49–74. doi:10.1037/0021-843X.87.1.49.
- [2] Allen NB, Badcock PBT. The Social Risk Hypothesis of Depressed Mood: Evolutionary, Psychosocial, and Neurobiological Perspectives. *Psychol. Bull.* 2003;129:887–913. doi:10.1037/0033-2909.129.6.887.
- [3] Amris K, Williams AC de C. Managing chronic pain in survivors of torture. *Pain Manag.* 2015;5:5–12. doi:10.2217/pmt.14.50.
- [4] Anderson C a, Bushman BJ. Human Aggression. *Annu. Rev. Psychol.* 2002;53:27–51. doi:10.1146/annurev.psych.53.100901.135231.

- [5] Ashton-james CE, Richardson DC, Williams ACDC, Bianchi-berthouze N, Dekker PH. Impact of pain behaviors on evaluations of warmth and competence. *Pain* 2014;155:2656–2661.
- [6] Balousek S, Plane MB, Fleming M. Prevalence of Interpersonal Abuse in Primary Care Patients Prescribed Opioids for Chronic Pain. *J. Gen. Intern. Med.* 2007;22:1268–1273. doi:10.1007/s11606-007-0257-6.
- [7] Baumeister RF, Leary MR. The need to belong: desire for interpersonal attachments as a fundamental human motivation. *Psychol. Bull.* 1995;117:497–529.
- [8] Bernstein MJ, Claypool HM. Social exclusion and pain sensitivity: why exclusion sometimes hurts and sometimes numbs. *Pers. Soc. Psychol. Bull.* 2012;38:185–96. doi:10.1177/0146167211422449.
- [9] Brosnan SF, Waal FBM De. Evolution of responses to (un)fairness. *Science* (80-.). 2014;346.
- [10] Brosnan SF, Waal FBM De. Fairness in Animals : Where to from Here ? 2012:336–351.
- [11] Carinci AJ, Mehta P, Christo PJ. Chronic pain in torture victims. *Curr. Pain Headache Rep.* 2010;14:73–79.
- [12] Carlsson K, Andersson J, Petrovic P, Petersson KM, ?hman A, Ingvar M. Predictability modulates the affective and sensory-discriminative neural processing of pain. *Neuroimage* 2006;32:1804–1814. doi:10.1016/j.neuroimage.2006.05.027.
- [13] Caspar EA, Desantis A, Dienes Z, Cleeremans A, Haggard P. The sense of agency as tracking control. *PLoS One* 2016;11:1–16.

- [14] Cohen M, Quintner J, Buchanan D, Nielsen M, Guy L. Stigmatization of Patients with Chronic Pain: The Extinction of Empathy. *Pain Med.* 2011;12:1637–1643. doi:10.1111/j.1526-4637.2011.01264.x.
- [15] Craig KD. Social communication model of pain. *Pain* 2015;156:1198–1199. doi:10.1097/j.pain.0000000000000185.
- [16] Craig KD. The social communication model of pain. *Can. Psychol. Can.* 2009;50:22–32. doi:10.1037/a0014772.
- [17] Crombez G, Eccleston C, De Vlieger P, Van Damme S, De Clercq A. Is it better to have controlled and lost than never to have controlled at all? An experimental investigation of control over pain. *Pain* 2008;137:631–9. doi:10.1016/j.pain.2007.10.028.
- [18] Van Damme S, Crombez G, Eccleston C. Coping with pain: a motivational perspective. *Pain* 2008;139:1–4. doi:10.1016/j.pain.2008.07.022.
- [19] Van Damme S, Legrain V, Vogt J, Crombez G. Keeping pain in mind: a motivational account of attention to pain. *Neurosci. Biobehav. Rev.* 2010;34:204–13. doi:10.1016/j.neubiorev.2009.01.005.
- [20] DeWall CN, Baumeister RF. Alone but Feeling No Pain: Effects of Social Exclusion on Physical Pain Tolerance and Pain Threshold, Affective Forecasting, and Interpersonal Empathy. *J. Pers. Soc. Psychol.* 2006;91:1–15. doi:10.1037/0022-3514.91.1.1.
- [21] Drwecki BB, Moore CF, Ward SE, Prkachin KM. Reducing racial disparities in pain treatment : The role of empathy and perspective-taking. *Pain* 2011;152:1001–1006.

- [22] Dumit J. Illnesses you have to fight to get: Facts as forces in uncertain, emergent illnesses. *Soc. Sci. Med.* 2006;62:577–590.
- [23] Eccleston C. *Embodied: the psychology of physical sensation*. 1st ed. Oxford: Oxford University Press, 2015 p.
- [24] Eccleston C, Williams ACDC, Stainton Rogers W. Patients' and professionals' understandings of the causes of chronic pain: Blame, responsibility and identity protection. *Soc. Sci. Med.* 1997;45:699–709.
- [25] Edwards RR, Moric M, Husfeldt B, Buvanendran A, Ivankovich O. Ethnic Similarities and Differences in the Chronic Pain Experience : A Comparison of African American , Hispanic , and White Patients. 2005;6.
- [26] Eisenberger NI. Social Pain and the Brain: Controversies, Questions, and Where to Go from Here. *Annu. Rev. Psychol.* 2015;66:601–629. doi:10.1146/annurev-psych-010213-115146.
- [27] Eisenberger NI. The pain of social disconnection: examining the shared neural underpinnings of physical and social pain. *Nat. Rev. Neurosci.* 2012;13:421–34. doi:10.1038/nrn3231.
- [28] Eisenberger NI, Jarcho JM, Lieberman MD, Naliboff BD. An experimental study of shared sensitivity to physical pain and social rejection. *Pain* 2006;126:132–8. doi:10.1016/j.pain.2006.06.024.
- [29] Eisenberger NI, Lieberman MD, Williams KD. Does rejection hurt? An fMRI study of social exclusion. *Science* 2003;302:290–2. doi:10.1126/science.1089134.
- [30] Fekkes M, Pijpers FIM, Fredriks AM, Vogels T, Verloove-Vanhorick SP. Do bullied children get ill, or do ill children get bullied? A prospective cohort study on the

- relationship between bullying and health-related symptoms. *Pediatrics* 2006;117:1568–74. doi:10.1542/peds.2005-0187.
- [31] Finlay BL, Syal S. The pain of altruism. *Trends Cogn. Sci.* 2014;18:615–617. doi:10.1016/j.tics.2014.08.002.
- [32] Fitzgerald ST, Haythornthwaite JA, Suchday S, Ewart CK. Anger in Young Black and White Workers: Effects of Job Control, Dissatisfaction, and Support. *J. Behav. Med.* 2003;26:283–296. doi:10.1023/A:1024228026022.
- [33] Forgeron PA, King S, Stinson JN, McGrath PJ, MacDonald AJ, Chambers CT. Social functioning and peer relationships in children and adolescents with chronic pain: A systematic review. *Pain Res. Manag.* 2010;15:27–41. doi:10.1155/2010/820407.
- [34] Friston K. The free-energy principle: A unified brain theory? *Nat. Rev. Neurosci.* 2010;11:127–138. doi:10.1038/nrn2787.
- [35] Gatchel RJ, Peng YB, Peters ML, Fuchs PN, Turk DC. The biopsychosocial approach to chronic pain: scientific advances and future directions. *Psychol. Bull.* 2007;133:581–624. doi:10.1037/0033-2909.133.4.581.
- [36] Gray K. The Power of Good Intentions: Perceived Benevolence Soothes Pain, Increases Pleasure, and Improves Taste. *Soc. Psychol. Personal. Sci.* 2012;3:639–645. doi:10.1177/1948550611433470.
- [37] Gray K, Wegner DM. The sting of intentional pain. *Psychol. Sci.* 2008;19:1260–2. doi:10.1111/j.1467-9280.2008.02208.x.
- [38] Gruter M, Masters RD. Ostracism as a social and biological phenomenon: An introduction. *Ethol. Sociobiol.* 1986;7:149–158. doi:10.1016/0162-3095(86)90043-9.

- [39] Hadjistavropoulos T, Craig KD, Duck S, Cano A, Goubert L, Jackson PL, Mogil JS, Rainville P, Sullivan MJL, de C Williams AC, Vervoort T, Fitzgerald TD. A biopsychosocial formulation of pain communication. *Psychol. Bull.* 2011;137:910–39. doi:10.1037/a0023876.
- [40] Hari R, Henriksson L, Malinen S, Parkkonen L. Centrality of Social Interaction in Human Brain Function. *Neuron* 2015;88:181–193. doi:10.1016/j.neuron.2015.09.022.
- [41] Hawkley LC, Cacioppo JT. Loneliness matters: A theoretical and empirical review of consequences and mechanisms. *Ann. Behav. Med.* 2010;40:218–227.
- [42] Hirsh AT, Alqudah AF, Stutts LA, Robinson ME. Virtual human technology : Capturing sex , race , and age influences in individual pain decision policies. *Pain* 2009;140:231–238.
- [43] Hoffman KM, Trawalter S, Axt JR, Oliver MN. Racial bias in pain assessment and treatment recommendations , and false beliefs about biological differences between blacks and whites. 2016.
- [44] Iglor EC, Defenderfer EK, Lang AC, Bauer K, Uihlein J, Davies WH. Gender differences in the experience of pain dismissal in adolescence. *J. Child Heal. Care* 2017;21:381–391. doi:10.1177/1367493517727132.
- [45] Johnson KV-A, Dunbar RIM. Pain tolerance predicts human social network size. *Sci. Rep.* 2016;6:25267. doi:10.1038/srep25267.
- [46] de Jong Gierveld J, van Tilburg T. A shortened scale for overall, emotional and social loneliness. *Tijdschr. Gerontol. Geriatr.* 2008;39:4–15.
- [47] Kaba R, Sooriakumaran P. The evolution of the doctor- patient relationship. 2016;5:57–65.

- [48] Kappesser J, de C Williams AC. Pain judgements of patients' relatives: examining the use of social contract theory as theoretical framework. *J. Behav. Med.* 2008;31:309–317.
- [49] Kappesser J, Williams ACDC, Prkachin KM. Testing two accounts of pain underestimation. *Pain* 2006;124:109–116.
- [50] Karoly P. Motivation, broadly construed, matters. *Pain* 2015;156:1375–1376.
- [51] Karos K, Meulders A, Goubert L, Vlaeyen JWS. The Influence of Social Threat on Pain, Aggression, and Empathy in Women. *J. Pain* 2018;19:291–300. doi:10.1016/j.jpain.2017.11.003.
- [52] Karos K, Meulders A, Goubert L, Vlaeyen JWS. The Influence of Social Threat on Pain, Aggression, and Empathy in Women. *J. Pain* 2017.
- [53] Karos K, Meulders A, Vlaeyen JWS. Threatening Social Context Facilitates Pain-Related Fear Learning. *J. Pain* 2015;16:214–225. doi:10.1016/j.jpain.2014.11.014.
- [54] Keogh E. Gender differences in the nonverbal communication of pain: A new direction for sex, gender, and pain research? *Pain* 2014;155:1927–1931. doi:10.1016/j.pain.2014.06.024.
- [55] Kool MB, van Middendorp H, Lumley MA, Bijlsma JW, Geenen R. Social support and invalidation by others contribute uniquely to the understanding of physical and mental health of patients with rheumatic diseases. *J. Health Psychol.* 2013;18:86–95. doi:10.1177/1359105312436438.
- [56] Krahé C, Springer A, Weinman J a, Fotopoulou A. The social modulation of pain: others as predictive signals of salience - a systematic review. *Front. Hum. Neurosci.* 2013;7:386. doi:10.3389/fnhum.2013.00386.

- [57] Langford DJ. Social Modulation of Pain as Evidence for Empathy in Mice. *Science* (80-.). 2006;312:1967–1970. doi:10.1126/science.1128322.
- [58] Lerner MJ, Miller DT. Just World Research and the Attribution Process : Looking Back and Ahead. 1978;85:1030–1051.
- [59] McCracken LM. Anger, injustice, and the continuing search for psychological mechanisms of pain, suffering, and disability. *Pain* 2013;154:1495–6. doi:10.1016/j.pain.2013.05.042.
- [60] McParland JL, Knussen C, Murray J. The effects of a recalled injustice on the experience of experimentally induced pain and anxiety in relation to just-world beliefs. *Eur. J. Pain* 2016:n/a-n/a. doi:10.1002/ejp.862.
- [61] Meulders A, Vervliet B, Fonteyne R, Baeyens F, Hermans D, Vansteenwegen D. Preexposure to (un)predictable shock modulates discriminative fear learning between cue and context: an investigation of the interaction between fear and anxiety. *Int. J. Psychophysiol.* 2012;84:180–7. doi:10.1016/j.ijpsycho.2012.02.004.
- [62] Miller DT. Disrespect and the experience of injustice. *Annu. Rev. Psychol.* 2001;52:527–553.
- [63] Miller DT. Disrespect and the Experience of Injustice. *Annu. Rev. Psychol.* 2001;52:527–553. doi:10.1146/annurev.psych.52.1.527.
- [64] Mogil JS. Social modulation of and by pain in humans and rodents. *Pain* 2015;156:S33–S41.
- [65] Morley S, Eccleston C. “The object of fear in pain.” In: Asmundsen J, Vlaeyen J, Crombez G, editors. *Understanding and treating fear of pain*. NY: Oxford University Press, 2004. pp. 163–188.

- [66] Mossey JM. Defining Racial and Ethnic Disparities in Pain Management. 2011;1859–1870.
- [67] Moutoussis M, Fearon P, El-Dereedy W, Dolan RJ, Friston KJ. Bayesian inferences about the self (and others): A review. *Conscious. Cogn.* 2014;25:67–76. doi:10.1016/j.concog.2014.01.009.
- [68] Müller MJ. Will it hurt less if I believe I can control it? Influence of actual and perceived control on perceived pain intensity in healthy male individuals: A randomized controlled study. *J. Behav. Med.* 2012;35:529–537.
- [69] Müller MJ, Netter P. Relationship of subjective helplessness and pain perception after electric skin stimuli. *Stress Med.* 2000;16:109–115.
- [70] Oka S, Chapman CR, Kim B, Shimizu O, Noma N, Takeichi O, Imamura Y, Oi Y. Predictability of Painful Stimulation Modulates Subjective and Physiological Responses. *J. Pain* 2010;11:239–246. doi:10.1016/j.jpain.2009.07.009.
- [71] Oliveira VC, Ferreira ML, Morso L, Albert HB, Refshauge KM, Ferreira PH. Patients' perceived level of social isolation affects the prognosis of low back pain. *Eur. J. Pain* 2014;1–8. doi:10.1002/ejp.578.
- [72] Panksepp J. *Affective Neuroscience: The Foundations of Human and Animal Emotions*. Oxford University Press, 1998 p. Available: <https://books.google.nl/books?id=qqcRGagyEuAC>.
- [73] Patsavas A. Recovering a Cripistemology of Pain. *J. Lit. Cult. Disabil. Stud.* 2014;8:203–218. doi:10.3828/jlcds.2014.16.
- [74] Peeters P a M, Vlaeyen JWS. Feeling more pain, yet showing less: the influence of social threat on pain. *J. Pain* 2011;12:1255–61. doi:10.1016/j.jpain.2011.07.007.

- [75] Piotrowski C. Chronic pain patients and loneliness: A systematic review of the literature. *Loneliness in life: Education, business, society*. NY: McGraw Hill, 2014.
- [76] Purdie F, Morley S. Self-compassion, pain, and breaking a social contract. *Pain* 2015;156:2354–2363. doi:10.1097/j.pain.0000000000000287.
- [77] De Ruddere L, Bosmans M, Crombez G, Goubert L. Patients are socially excluded when their pain has no medical explanation. *J. Pain* 2016;17:1028–1035. doi:10.1016/j.jpain.2016.06.005.
- [78] De Ruddere L, Craig KD. Understanding stigma and chronic pain. *Pain* 2016;157:1. doi:10.1097/j.pain.0000000000000512.
- [79] De Ruddere L, Goubert L, Stevens MAL, Deveugele M, Craig KD, Crombez G. Health Care Professionals' Reactions to Patient Pain: Impact of Knowledge About Medical Evidence and Psychosocial Influences. *J. Pain* 2014;15:262–270. doi:10.1016/j.jpain.2013.11.002.
- [80] Ruddere L De, Goubert L, Stevens M, Williams AC, Crombez G. Discounting pain in the absence of medical evidence is explained by negative evaluation of the patient. *Pain* 2013;154:669–676. doi:10.1016/j.pain.2012.12.018.
- [81] De Ruddere L, Goubert L, Vervoort T, Kappesser J, Crombez G. Impact of being primed with social deception upon observer responses to others' pain. *Pain* 2013;154:221–6. doi:10.1016/j.pain.2012.10.002.
- [82] De Ruddere L, Goubert L, Vervoort T, Prkachin KM, Crombez G. We discount the pain of others when pain has no medical explanation. *J. Pain* 2012;13:1198–1205. doi:10.1016/j.jpain.2012.09.002.

- [83] Samwel HJA, Evers AWM, Crul BJP, Kraaimaat FW. The Role of Helplessness, Fear of Pain, and Passive Pain-Coping in Chronic Pain Patients. *Clin. J. Pain* 2006;22:245–251. doi:10.1097/01.ajp.0000173019.72365.f5.
- [84] Scd LL. Defining Gender Disparities in Pain Management. 2011:1871–1877.
- [85] Scott W, Mcevoy A, Garland R, Bernier E, Milioto M, Trost Z, Sullivan M, Sullivan M. Sources of injustice among individuals with persistent pain following musculoskeletal injury. 2016:6–15.
- [86] Scott W, Trost Z, Bernier E, Sullivan MJL. Anger differentially mediates the relationship between perceived injustice and chronic pain outcomes. *Pain* 2013;154:1691–1698. doi:10.1016/j.pain.2013.05.015.
- [87] Shakhar K, Shakhar G. Why Do We Feel Sick When Infected??? Can Altruism Play a Role? *PLoS Biol.* 2015;13:1–15.
- [88] Smith JA, Osborn M. Pain as an assault on the self: An interpretative phenomenological analysis of the psychological impact of chronic benign low back pain. *Psychol. Health* 2007;22:517–534. doi:10.1080/14768320600941756.
- [89] Sullivan MJL, Adams H, Horan S, Maher D, Boland D, Gross R. The role of perceived injustice in the experience of chronic pain and disability: scale development and validation. *J. Occup. Rehabil.* 2008;18:249–61. doi:10.1007/s10926-008-9140-5.
- [90] Sullivan MJL, Davidson N, Scott W. Perceived Injustice is Associated with Heightened Pain Behavior and Disability in Individuals with Whiplash Injuries. 2009:238–247.
- [91] Sullivan MJL, Scott W, Trost Z. Perceived injustice: a risk factor for problematic pain outcomes. *Clin. J. Pain* 2012;28:484–8. doi:10.1097/AJP.0b013e3182527d13.

- [92] Sullivan MJL, Thibault P, Simmonds MJ, Milioto M, Cantin A-P, Velly AM. Pain, perceived injustice and the persistence of post-traumatic stress symptoms during the course of rehabilitation for whiplash injuries. *Pain* 2009;145:325–31. doi:10.1016/j.pain.2009.06.031.
- [93] Taylor AJW. Justice as a basic human need. *New Ideas Psychol.* 2003;21:209–219. doi:10.1016/j.newideapsych.2003.09.004.
- [94] Tiokhin L. Do Symptoms of Illness Serve Signaling. *Q. Rev. Biol.* 2016;91:1–19.
- [95] Trost Z, Scott W, Lange JM, Manganelli L, Bernier E, Sullivan MJ. An experimental investigation of the effect of a justice violation on pain experience and expression among individuals with high and low just world beliefs. *Eur. J. Pain (United Kingdom)* 2014;18:415–423.
- [96] Trost Z, Vangronsveld K, Linton SJ, Quartana PJ, Sullivan MJL. Cognitive dimensions of anger in chronic pain. *Pain* 2012;153:515–7. doi:10.1016/j.pain.2011.10.023.
- [97] Turner-Cobb JM, Michalaki M, Osborn M. Self-conscious emotions in patients suffering from chronic musculoskeletal pain: A brief report. *Psychol. Health* 2015;30:495–501. doi:10.1080/08870446.2014.991735.
- [98] Twenge JM, Baumeister RF, Tice DM, Stucke TS. If you can't join them, beat them: effects of social exclusion on aggressive behavior. *J. Pers. Soc. Psychol.* 2001;81:1058–1069.
- [99] Vanderbilt D, Augustyn M. The effects of bullying. *Paediatr. Child Health (Oxford)*. 2010;20:315–320. doi:10.1016/j.paed.2010.03.008.
- [100] Vlaeyen JWS. Learning to predict and control harmful events: chronic pain and conditioning. *Pain* 2015;156 Suppl:S86-93. doi:10.1097/j.pain.000000000000107.

- [101] Voerman JS, Vogel I, De Waart F, Westendorp T, Timman R, Busschbach JJ V, Van De Looij-Jansen P, De Klerk C. Bullying, abuse and family conflict as risk factors for chronic pain among Dutch adolescents. *Eur. J. Pain (United Kingdom)* 2015;19:1544–1551.
- [102] Warburton W a., Williams KD, Cairns DR. When ostracism leads to aggression: The moderating effects of control deprivation. *J. Exp. Soc. Psychol.* 2006;42:213–220.
- [103] Waugh OC, Byrne DG, Nicholas MK. Internalised Stigma in People Living with Chronic Pain. *J. Pain* 2014. doi:10.1016/j.jpain.2014.02.001.
- [104] Waugh OC, Byrne DG, Nicholas MK. Internalized stigma in people living with chronic pain. *J. Pain* 2014;15:550.e1-550.e10. doi:10.1016/j.jpain.2014.02.001.
- [105] Wiech K, Kalisch R, Weiskopf N, Pleger B, Stephan KE, Dolan RJ. Anterolateral Prefrontal Cortex Mediates the Analgesic Effect of Expected and Perceived Control over Pain. *J. Neurosci.* 2006;26:11501–11509. doi:10.1523/JNEUROSCI.2568-06.2006.
- [106] Wiech K, Tracey I. Pain, decisions, and actions: a motivational perspective. *Front. Neurosci.* 2013;7:1–12. doi:10.3389/fnins.2013.00046.
- [107] Williams AC de C. Defeating the stigma of chronic pain. *Pain* 2016;157:1. doi:10.1097/j.pain.0000000000000530.
- [108] Williams ACDC. Facial expression of pain: an evolutionary account. *Behav. Brain Sci.* 2002;25:439-55-88. Available: <http://www.ncbi.nlm.nih.gov/pubmed/12879700>.
- [109] Williams ACDC. What can evolutionary theory tell us about chronic pain? *Pain* 2016;157:788–790. doi:10.1097/j.pain.0000000000000464.

[110] Williams ACDC, Craig KD. Updating the definition of pain. *Pain* 2016;157:2420–2423. doi:10.1097/j.pain.0000000000000613.

[111] Williams KD. *Ostracism: The power of silence*. New York: Guilford Press, 2002 p.

[112] Williams KD. Ostracism. *Annu. Rev. Psychol.* 2007;58:425–52.
doi:10.1146/annurev.psych.58.110405.085641.

Figure legend

Figure 1: A schematic representation of the three fundamental interpersonal needs at the nexus of an individuals' pain experience. Pain threatens these needs in a myriad of ways but is in turn affected by thwarted needs as well. Note that pain in this context refers to both pain appraisals (e.g., pain intensity and unpleasantness), but also pain behavior (e.g., pain expression, disability, treatment adherence). Different needs may overlap and influence each other (e.g., an experience of ostracism may be excluding but at the same time lead to perceptions of injustice and helplessness). While these needs are private to the individual (level 1), they are inherently influenced by the social context, be it the interpersonal context (level 2, i.e., partner, family, health-care professionals) or the socio-cultural context (level 3, i.e., groups, culture, social norms). Conversely, thwarted interpersonal needs can affect the relationship between the individual and the social context as well (e.g., withdrawal from social circles or attack in the form of antisocial behavior). Lastly, individual factors (e.g., personality traits) might function as a buffer or facilitator for the flow between these different levels.

