

# Low Cerebrospinal Fluid A $\beta$ <sub>42</sub> and A $\beta$ <sub>40</sub> are Related to White Matter Lesions in Cognitively Normal Elderly

Ingmar Skoog<sup>a,1,\*</sup>, Silke Kern<sup>a,b,1</sup>, Henrik Zetterberg<sup>b,c</sup>, Svante Östling<sup>a</sup>, Anne Börjesson-Hanson<sup>a</sup>, Xinxin Guo<sup>a</sup> and Kaj Blennow<sup>b</sup>

<sup>a</sup>Department of Psychiatry and Neurochemistry, Neuropsychiatric Epidemiology Unit, Institute of Neuroscience

View metadata, citation and similar papers at [core.ac.uk](http://core.ac.uk)

brought to you by  CORE  
provided by UCL Discovery

and Physiology, Sahlgrenska Academy at the University of Gothenburg, Gothenburg, Sweden

<sup>c</sup>UCL Institute of Neurology, Queen Square, London, UK

Accepted 15 January 2018

## Abstract.

**Background:** Low cerebrospinal fluid (CSF) levels of A $\beta$ <sub>42</sub> may be the earliest manifestation of Alzheimer's disease (AD). Knowledge on how CSF A $\beta$  interacts with different brain pathologies early in the disease process is limited. We examined how CSF A $\beta$  markers relate to brain atrophy and white matter lesions (WMLs) in octogenarians with and without dementia to explore the earliest pathogenetic pathways of AD in the oldest old.

**Objective:** To study CSF amyloid biomarkers in relation to brain atrophy and WMLs in 85-year-olds with and without dementia.

**Methods:** 53 octogenarians took part in neuropsychiatric examinations and underwent both a lumbar puncture and a brain CT scan. CSF levels of A $\beta$ <sub>42</sub> and A $\beta$ <sub>40</sub> were examined in relation to cerebral atrophy and WMLs. Dementia was diagnosed.

**Results:** In 85-year-olds without dementia, lower levels of both CSF A $\beta$ <sub>42</sub> and CSF A $\beta$ <sub>40</sub> were associated with WMLs. CSF A $\beta$ <sub>42</sub> also correlated with measures of central atrophy, but not with cortical atrophy. In participants with dementia, lower CSF levels of A $\beta$ <sub>42</sub> were related to frontal, temporal, and parietal cortical atrophy but not to WMLs.

**Conclusions:** Our findings may suggest that there is an interrelationship between A $\beta$  and subcortical WMLs in older persons without dementia. After onset of dementia, low CSF A $\beta$ <sub>42</sub>, probably representing amyloid deposition in plaques, is associated with cortical atrophy. WMLs may be an earlier manifestation of A $\beta$  deposition than cortical degeneration.

**Keywords:** Alzheimer's disease, amyloid- $\beta$ , biomarkers, cerebrospinal fluid, computerized tomography, dementia, epidemiological methods, population-based, vascular dementia, white matter lesions

## INTRODUCTION

Alzheimer's disease (AD), alone or in combination with other brain disorders, is the most common cause of dementia in old age. AD is characterized by

synaptic and neuronal degeneration and the presence of extensive amounts of plaques and tangles in cortical brain regions [1]. A central finding in AD is the deposition of amyloid- $\beta$  (A $\beta$ ) in the brain, especially in the core of the plaques and in the microvessels [2]. According to the "amyloid cascade hypothesis", A $\beta$  oligomerization with subsequent deposition of the peptide in plaques start a cascade of events that results in neuronal and synaptic degeneration and dementia [3–6].

<sup>1</sup>These authors contributed equally to the work.

\*Correspondence to: Ingmar Skoog, Neuropsychiatric Epidemiology Unit, Sahlgrenska University Hospital/Mölndal, Wallingsgatan 6, SE 43141 Mölndal, Sweden. Tel.: +46 31 342 2164; Fax: +46 31 828163; E-mail: [ingmar.skoog@neuro.gu.se](mailto:ingmar.skoog@neuro.gu.se).

Biomarkers are valuable tools to study the temporal evolution of the central pathogenetic processes in AD [5, 7]. Low levels of cerebrospinal fluid (CSF) A $\beta$ <sub>42</sub> reflect cortical A $\beta$  deposition [8, 9], which is supported by the finding that it is related to high cortical retention of the positron-emission tomography (PET) tracer Pittsburgh Compound B (PIB) that binds fibrillar A $\beta$  [10–12]. Low CSF A $\beta$ <sub>42</sub> levels are also strongly associated with AD with dementia and incipient AD in MCI [7], as well as with future cognitive impairment in asymptomatic individuals [13–17].

Brain atrophy on magnetic resonance imaging (MRI) or computerized tomography (CT) of the brain reflects neurodegenerative processes [5, 18, 19]. The earliest changes in the evolution of A $\beta$  deposits in AD are thought to occur in the neocortex and from there spread to allocortical regions, brainstem, and finally to the cerebellum [20].

The temporal sequence in the pathophysiological processes of AD has received increased interest recently. As suggested by Jack and co-workers [4, 5], it is supposed that the initiating event is abnormal processing of A $\beta$ , reflected by changes in CSF A $\beta$  [4, 21], which occurs many years before symptoms of cognitive disturbances [22]. The second event is related to tau-mediated injury and dysfunction, and the third to neurodegeneration of brain structures, i.e., cerebral atrophy ([4, 5, 23]). Cognitive symptoms are supposed to relate closely to the evolution of neuronal degeneration [5]. The hypothetical model of the pathophysiological process was intended to model pure AD [5]. However, in late-onset AD, especially in cases with very late onset, there is a high prevalence of co-existing pathological processes [4]. One important type of concomitant pathology is ischemic white matter lesions (WMLs), which are frequent both in clinical and autopsied AD cases, as well as in normal elderly [24–26]. WMLs have been shown to aggravate clinical symptoms of AD [27]. These changes are associated with lipohyalinosis causing narrowing of the lumen of the small perforating arteries and arterioles that nourish the deep white matter. The underlying cause of WMLs is not established, but the main hypothesis is that they are due to long-standing hypertension that cause lipohyalinosis and thickening of the vessel walls [28], which leads to ischemia in vulnerable areas. Animal experiments have shown that brain ischemia also may cause an acute increase in A $\beta$  secretion through induced BACE1 protein expression, with accumulation of oligomeric A $\beta$  in the hippocampus [29–31]. Furthermore,

neuropathological studies in patients who died after acute stroke or after resuscitation from cardiac arrest have shown a marked increase in A $\beta$  expression in hippocampus together with a marked increase in non-fibrillar A $\beta$  plaques in cortex [32, 33]. Thus, ischemia may trigger A $\beta$  deposition and plaque formation. In addition, a study on normal elderly individuals without dementia but with extensive AD neuropathology suggested that WMLs precede the cortical atrophy in the evolution of AD pathology [25]. WMLs have also been related to deposition of A $\beta$  in the vessels [34], further suggesting that interactions between A $\beta$  and WMLs may occur early in the disease process. We aimed to examine the relationship between amyloid metabolism (measured by CSF A $\beta$ ), stage of neuronal degeneration (measured as atrophy on CT), and small vessel disease (measured as WMLs on CT) in very old persons from a population-based sample of 85-year-olds with and without dementia, to provide further knowledge about the relationship between these pathogenetic markers in older persons.

## STUDY POPULATION AND METHODS

### *Study population*

A representative sample of 85-year-olds and registered for census purposes in Gothenburg, was invited to take part in a health survey. Both people living in the community and those in institutions were included. A neuropsychiatric examination was performed on a systematic sub-sample ( $n=494$ , response rate 63%). The sample has been described in detail previously [35].

The first 165 individuals were invited to undergo a lumbar puncture. Sixty-nine (31 with dementia and 38 without dementia) accepted. Of these, five individuals were excluded due to technical reasons and another two due to hemorrhagic spinal taps, leaving 62 individuals (27 with dementia and 35 without, 40 women and 22 men) for the present study. AD was diagnosed in 12 individuals (four men and eight women), vascular dementia (VaD) in 13 (two men and 11 women) and other types of dementia in two individuals (one man and one woman). All had onset of dementia after age 65 years. The cohort that underwent lumbar puncture has been described in detail previously [13]. All 85-year-olds with a lumbar puncture were invited to undergo a CT scan of the head. Fifty-three accepted both examinations: 23 with

Table 1  
Demographics and CSF biomarker levels in a population sample of 85-year-olds with and without dementia

Characteristic	Without Dementia (n = 30)	With Dementia (n = 23)	p-value <sup>1</sup>	Alzheimer's dementia (n = 9)	Vascular dementia (n = 12)	p-value <sup>2</sup>
No. (%)						
Women	16 (53.3)	17 (73.9)	p = 0.16	6 (66.7)	10 (83.3)	p = 0.61
Current/Ex-smokers	18 (60.0)	6 (33.4)	p = 0.14	2 (28.6)	2 (22.2)	p = 1.0
Living in care	0 (0)	16 (69.6)	p < 0.00	6 (66.6)	9 (75.0)	p = 1.0
Stroke	4 (13.3)	9 (39.1)	p = 0.05	0	9 (75)	p = 0.001
Mean (SD)						
CSF A $\beta$ <sub>42</sub>	632 (280)	420 (200)	p < 0.01	381 (161)	431 (198)	p = 0.55
CSF A $\beta$ <sub>40</sub>	1975 (684)	1393 (438)	p < 0.001	1225 (408)	1530 (414)	p = 0.11
Mean age at interview	85.4 (0.14)	85.5 (0.18)	p < 0.05	85.1 (0.86)	85.1 (27.5)	p = 0.35
Systolic blood pressure	163.5 (30.1)	146.1 (24.9)	p = 0.05	145.6 (21.7)	148.2 (28.5)	p = 0.82
MMSE	28.3 (1.5)	12.6 (8.9)	p < 0.0001	11.7 (9.5)	13.5 (8.2)	p = 0.66
No. (%)						
Frontal atrophy severe	12 (40.0)	15 (65.2)	p = 0.15	6 (66.7)	7 (58.3)	p = 1.0
Temporal atrophy	14 (60.9)	6 (20.0)	p < 0.01	7 (77.8)	6 (50.0)	p = 0.37
Parietal atrophy	11 (36.7)	14 (60.9)	p = 0.08	4 (44.4)	9 (75.0)	p = 0.20
Occipital atrophy	4 (17.4)	6 (20.0)	p = 0.92	2 (22.2)	2 (16.7)	p = 1.0
White matter lesions severe	3 (10)	7 (30.4)	p < 0.01	7 (77.78)	8 (66.7)	p = 0.66

Values are given as number (percent) or mean (standard deviation). CSF biomarker levels are given as pg/mL. In the statistical analyses, Fisher's exact test was used for discrete variables and Student's *t*-test for continuous variables. <sup>1</sup>Test between participants with dementia and participants without. <sup>2</sup>Test between participants with Vascular dementia and participants with Alzheimer's dementia. CSF, cerebrospinal fluid; MMSE, Mini-Mental State Examination.

dementia [AD was diagnosed in 9 individuals (3 men and 6 women), 12 with VaD (2 men and 10 women), and 2 with other types of dementia (one man and one woman)] and 30 without dementia. The demographic of this sample is given in Table 1. Mean age of onset of dementia in the 23 participants with dementia was 79.8 (SD 4.12) years (13 were diagnosed with dementia between 80 and 85 years, 8 between 75 and 80 years and 2 persons before age 75).

Follow-up data on this sample has been published previously [13]. During the following 3 years (age 85-88), 7 of the 35 85-year-olds without dementia developed dementia, while 28 remained dementia-free.

The study was approved by the Ethics Committee at the University of Gothenburg. All subjects (or their closest relatives) gave their informed consent to participate in the study.

#### General examinations

The examinations included medical history and physical examinations performed by a geriatrician, neuropsychiatric examinations and a telephone interview with a close informant performed by a psychiatrist, an electrocardiogram, a chest X-ray, a battery of blood tests, CT of the brain, and a lumbar puncture [35]. All clinical examinations were made without knowledge of the brain imaging and biochemical results and vice versa.

#### CSF sampling and analysis

Lumbar punctures were performed in the morning, under standardized conditions, through the L3/L4 or L4/L5 interspace. The first 12 ml of CSF was collected in polypropylene tubes and gently mixed to avoid possible gradient effects. CSF samples with more than 500 erythrocytes per  $\mu$ L were excluded.

CSF A $\beta$ <sub>42</sub> was determined using a sandwich ELISA which is based on the 3D6 and 21F12 monoclonal antibodies, making it specific for A $\beta$  1-42, as previously described [36]. CSF A $\beta$ <sub>40</sub> was determined using an in-house research sandwich ELISA in which an A $\beta$ <sub>40</sub>-specific polyclonal antibody (Quality Control Biochemicals, Hopkinton, MA, USA) was used as capturing antibody, and an N-terminal specific monoclonal antibody (3D6) was used as detecting antibody. High performance liquid chromatography purified A $\beta$ <sub>40</sub> (Bachem, Bubendorf, Switzerland), was used as standard. CSF samples were analyzed in duplicates. Details of the ELISA characteristics, including intra- and inter-day variability have been presented in detail [37].

#### Computerized tomography evaluations

All CT scans were performed without contrast enhancement and the CT scans were examined by two experienced radiologists, who were blinded to the clinical and CSF results [26].

The occipital, parietal, frontal, and temporal lobes were categorized using a four-point scale (normal, mild, moderate and severe) according to the estimated extent of brain atrophy [18]. The rating procedure was carried out separately for the cortical and ventricular studies.

The following linear distances were determined using a transparent metric ruler as described by de Leon and co-workers [18]: (a) the bifrontal span of the lateral ventricle; (b) the width of the lateral ventricles at the head of the caudate nucleus; (c) the sum of the separate widths of the left and right Sylvian fissures; (d) the minimal width of the bodies of the lateral ventricles at the waist, and (e) the width of the third ventricle. Ratios were determined by dividing the obtained values by the width of the brain at the level of the measurement, giving the following ratios: (a) bifrontal ratio, (b) bicaudate ratio, (c) Sylvian fissure ratio, (d) cella media ratio.

WMLs were defined as low-density areas in periventricular or subcortical white matter. The changes were always diffusely distributed within the white matter. Decreased density was subjectively rated as no, mild, moderate, or severe in relation to the attenuation of normal white matter [26, 38]. In cases of disagreement a consensus conference was performed.

#### *Diagnostic methods*

Dementia and its severity were diagnosed according to the DSM-III-R criteria, using information from the psychiatric examination and the close informant interview, as described previously [35]. Subjects with dementia were classified into diagnostic subgroups: AD according to the NINCDS-ADRDA criteria [39] and VaD according to the possible NINDS-AIREN criteria [40]. In the diagnostic classification, information from the psychiatric examinations, close informant interview, laboratory tests, ECG, and case records were used [35].

#### *Statistical analyses*

Fisher's exact test and independent *t*-test were used to test the hypothesis of no differences between the groups. Pitman's permutation test was used to test correlations ( $r$ ) between CSF  $A\beta$  and CT variables (atrophy, WMLs), adjusted for the presence of dementia. Before stratification by dementia status, we tested if there was an interaction with dementia status, i.e., if the strength of correlations between CSF  $A\beta$

and CT measures (atrophy, WMLs) differed between 85-year-olds with and without dementia. Interactions were calculated with a permutation test as both groups were relatively small, and as the variables were not normally distributed. In cases of interactions  $p < 0.20$ , we stratified analyses by dementia status. Results were retested with robust regression by using STATA procedure "regress" with the parameter vce (robust).  $p$ -values  $< 0.05$  (two-sided) were regarded as statistically significant.

## RESULTS

*Description of the sample is presented in Table 1*

Table 2 gives the correlations between CSF  $A\beta_{42}$  and CT measures of brain atrophy and WMLs in the total group. Lower CSF  $A\beta_{42}$  levels correlated with higher degree of temporal atrophy, bifrontal ratio, bicaudate ratio, and third ventricular width. There were interaction effects with dementia status for the associations between CSF  $A\beta_{42}$  and frontal, temporal, and parietal atrophy, and WMLs. Analyses of these CT measures were therefore stratified by dementia status in further analyses. In the stratified analyses (Tables 3 and 4), lower CSF  $A\beta_{42}$  was related to severity of WMLs in the group without dementia, but not in the group with dementia. In contrast, lower CSF  $A\beta_{42}$  levels were related to more severe frontal and temporal lobe atrophy in the group with dementia, while no such associations were found in the group without dementia.

Table 2 also gives the correlations between CSF  $A\beta_{40}$  and CT measures of brain atrophy and WMLs in the total group. Lower CSF  $A\beta_{40}$  was related to higher bifrontal ratio, sella media ratio, and severity of WMLs. There were interactions with dementia status for the associations between CSF  $A\beta_{40}$  and frontal atrophy and WMLs. Analyses of these CT measures were therefore stratified by dementia status in further analyses. In these analyses, lower CSF  $A\beta_{40}$  was related to severity of WMLs in the group without dementia, but not in those with dementia (Table 3). The main results were unchanged using a robust regression analysis.

## DISCUSSION

We examined biomarkers for  $A\beta$  metabolism, neuronal degeneration, and WMLs in 85-year-olds

Table 2

Cerebrospinal fluid Aβ<sub>40</sub> and Aβ<sub>42</sub> in relation to measures of cerebral atrophy and white matter lesions in a population sample of 85-year-olds (n = 53). Pooled correlation coefficient adjusted for dementia

	Aβ <sub>42</sub>		Aβ <sub>40</sub>		Aβ <sub>42/40</sub>	
	Correlation r <sup>a</sup>	Interaction with dementia p value	Correlation r <sup>a</sup>	Interaction with dementia p value	Correlation r <sup>a</sup>	Interaction with dementia p value
Cortical atrophy						
Frontal	-0.10	0.02	-0.05	0.11	-0.21	0.51
Temporal	-0.31 <sup>b</sup>	0.03	-0.26	0.26	-0.26	0.27
Parietal	-0.27	0.11	-0.11	0.75	-0.37 <sup>c</sup>	0.43
Occipital	-0.12	0.89	-0.09	0.68	-0.01	0.53
Central atrophy						
Bifrontal ratio	-0.41 <sup>c</sup>	0.47	-0.41 <sup>c</sup>	0.41	-0.19	0.57
Bicaudate ratio	-0.28 <sup>b</sup>	0.97	-0.23	0.80	-0.12	0.17
Third ventricle width	-0.30 <sup>b</sup>	0.73	-0.25	0.66	-0.18	0.40
Sylvian fissure ratio	-0.03	0.99	-0.04	0.15	-0.10	0.52
Sella media ratio	-0.25	0.44	-0.30 <sup>b</sup>	0.49	-0.08	0.35
White Matter Lesions	-0.22	0.10	-0.31 <sup>b</sup>	0.17	0.05	0.63

<sup>a</sup>Pitman's permutation test was used to test correlations (r) adjusted for dementia status, 30 participants without dementia diagnosis and 23 with a dementia diagnosis. <sup>b</sup>p < 0.05. <sup>c</sup>p < 0.01.

Table 3

Stratified analyses in those correlations between brain atrophy/white matter lesions and CSF Aβ where interaction effect by dementia status had a p-value of < 0.20 and the correlation was significantly not equal in any subgroup

	CSF Aβ <sub>42</sub>			CSF Aβ <sub>40</sub>			CSF Aβ <sub>42/40</sub>		
	Without Dementia (n = 29)	With Dementia (n = 22)	Test of Interaction*	Without Dementia (n = 29)	With Dementia (n = 22)	Test of Interaction*	Without Dementia (n = 29)	With Dementia (n = 22)	Test of Interaction*
Cortical atrophy									
Frontal	0.12	-0.52 <sup>b</sup>	p = 0.02	0.11	-0.37	p = 0.11			
Temporal	-0.15	-0.70 <sup>d</sup>	p = 0.03			p = 0.26			
Parietal	-0.13	-0.59 <sup>c</sup>	p = 0.11			p = 0.75			
Central atrophy									
Bicaudate ratio							0.10	-0.25	p = 0.234
White Matter Lesions	-0.39 <sup>b</sup>	0.03	p = 0.10	-0.46 <sup>b</sup>	-0.08	p = 0.17			

\*Significance of interaction evaluated by a permutation test. <sup>b</sup>p < 0.05. <sup>c</sup>p < 0.01. <sup>d</sup>p < 0.001.

Table 4

Stratified analyses between brain atrophy/white matter lesions and CSF Aβ according to Alzheimer's dementia and vascular dementia status

	CSF Aβ <sub>42</sub>			CSF Aβ <sub>40</sub>			CSF Aβ <sub>42/40</sub>		
	Alzheimer's Dementia (n = 9)	Vascular Dementia (n = 12)	Test of Interaction*	Alzheimer's Dementia (n = 9)	Vascular Dementia (n = 12)	Test of Interaction*	Alzheimer's Dementia (n = 9)	Vascular Dementia (n = 12)	Test of Interaction*
Cortical atrophy									
Frontal	-0.61	-0.67 <sup>b</sup>	p = 0.84	-0.74 <sup>b</sup>	-0.15	p = 0.14	-0.11	-0.47	p = 0.44
Temporal	-0.61	-0.67 <sup>b</sup>	p = 0.83	-0.54	-0.23	p = 0.49	-0.35	-0.51	p = 0.71
Parietal	-0.48	-0.70 <sup>b</sup>	p = 0.49	-0.35	-0.13	p = 0.65	-0.29	-0.64 <sup>b</sup>	p = 0.37
White Matter Lesions	-0.016	0.13	p = 0.82	0.23	-0.18	p = 0.43	-0.22	0.27	p = 0.39

\*Significance of interaction evaluated by a permutation test. <sup>b</sup>p < 0.05. <sup>c</sup>p < 0.01. <sup>d</sup>p < 0.001.

with and without dementia to learn more about the early pathogenetic processes in very late-life sporadic dementia. Lower CSF levels of Aβ<sub>42</sub> are supposed to be the earliest event in the evolution of AD, occurring approximately 15–20 years before brain atrophy and clinical symptoms [5]. We found that lower CSF levels of Aβ<sub>42</sub> and Aβ<sub>40</sub> were related to WMLs in

85-year-olds without dementia, and Aβ<sub>42</sub> to degree of frontal, parietal, and temporal cortical brain atrophy in those with dementia. This may indicate that disturbed metabolism of Aβ is related to cerebrovascular disease, such as WMLs, in the very early phases of the disease, and to the stage of neuronal degeneration, as measured by brain atrophy on CT, only in

the later disease stages with manifest dementia. The latter is in line with the hypothesis of Jack et al. [5], that symptoms of dementia do not occur until the start of neurodegeneration.

Lower CSF levels of both A $\beta_{42}$  and A $\beta_{40}$  correlated with severity of WMLs in individuals without dementia, but not in those with dementia. Amyloid PET studies show a correlation between cortical amyloid load and CSF A $\beta_{42}$ , but no relation with CSF A $\beta_{40}$  [14, 41, 42]. Thus, the correlation between low CSF levels of both A $\beta_{40}$  and A $\beta_{42}$  and WMLs suggest that our finding might reflect a disturbance in A $\beta$  metabolism or clearance rather than amyloid deposition in plaques. Indeed, neuropathological studies have reported that the periventricular white-matter in individuals without dementia and with WMLs contain deposits of A $\beta$ , while the white matter in those without WMLs does not [43]. Other studies also report increased quantities of parenchymal A $\beta$  in AD white matter [44, 45], strengthening the hypothesis that WMLs may be connected to the amyloidogenic process early in AD. There are also data supporting a relation between A $\beta$  and WMLs in preclinical AD. One neuropathology study on individuals without dementia but with extensive neurofibrillary tangles and plaques in the cerebral cortex suggested that WMLs precede the cortical atrophy in AD [25]. Further support for our findings comes from a recent clinical study reporting that white matter hyperintensities on MRI were related to lower CSF levels of A $\beta_{42}$  in controls and in individuals with VAD, but not in individuals with manifest AD dementia [46]. Also other studies suggest that WMLs may cause or precede brain atrophy [47]. WMLs have previously been related to AD in both clinical and autopsy studies [24–26]. These data, together with our findings that CSF measures of A $\beta$  neuropathology are associated with WMLs in 85-year-olds without dementia and with cortical atrophy among individuals with dementia, suggest that there is an interrelationship between A $\beta$  and WMLs in those without dementia. The timing of the event is uncertain, but WMLs may be an even earlier manifestation than cortical neurodegeneration. However, this finding needs to be further elucidated using the more sensitive MRI technique in a larger sample. It must be emphasized that in octogenarians it may be hard to disentangle age-related pathology, preclinical dementia pathology and co-morbidities.

In line with our study, van Westen et al. [48] reported that lower A $\beta_{40}$  was related to WMLs in non-demented persons. However, in contrast to our

findings, they found that WMLs were only associated with lower CSF A $\beta_{42}$  in AD dementia, and not in the cognitively healthy. One reason for the latter may be that study participants in that study were younger.

The association between CSF A $\beta_{42}$  and A $\beta_{40}$ , and WMLs in individuals without dementia should also be seen in light of the hypothesis that vascular factors may be involved in the pathogenesis of AD [49]. The metabolism of A $\beta$ PP and generation of A $\beta$  may be affected by intracerebral small-vessel arteriolosclerosis [50], which may cause hypoperfusion and ischemia [51] that may further exacerbate A $\beta$  oligomerization and deposition [52]. This hypothesis is supported by both experimental animal studies and human post-mortem studies suggesting that ischemia may cause an upregulation of A $\beta$  production with subsequent accumulation and oligomerization of A $\beta$  and formation of non-fibrillar plaques [29, 30, 33]. The vessel changes related to WMLs may also lead to disturbance in clearance of A $\beta$  from the brain, leading to lower levels of CSF A $\beta$ . High blood pressure has been reported to precede AD onset by 10–15 years [53]. Hypertension is also a risk factor for WMLs [54]. Furthermore, middle-aged hypertensive individuals without dementia have an increased amount of senile plaques in their brains [55], which should translate into lower CSF A $\beta_{42}$  values [8, 9]. Recently, a PET study in individuals free from dementia and aged 83 years and above reported that arterial stiffness was associated with progressive deposition of A $\beta$  in the brain [56]. The direction of the association is not clear. These findings, as well as those in the present study, may either reflect that similar mechanisms may be involved in the pathogenesis of both disorders or that vascular diseases may exacerbate the AD process, by for instance inducing blood-brain barrier dysfunction, oxidative stress and ischemia [51, 52, 57]. Finally, recent studies suggest that WMLs may influence the expression and/or processing of A $\beta$ PP, which may cause a downstream reduction in the levels of A $\beta_{42}$  and A $\beta_{40}$ , not related to deposition of the peptides in the brain parenchyma [58, 59]. Whether this change contributes to neurodegeneration is unclear.

The cortical atrophy in AD is due to neuronal and synaptic degeneration and loss. In the present study, the strongest correlation between atrophy and lower CSF A $\beta_{42}$  was found for the temporal lobe, which is in line with studies reporting a relation between CSF A $\beta_{42}$  and temporal lobe atrophy visualized on MRI [60–62]. We could not find any correlation between CSF A $\beta_{42}$  and cortical atrophy in

85-year-olds without dementia, despite that CSF A $\beta$ <sub>42</sub> levels were lower in cognitively healthy individuals who later developed dementia in this sample [13, 15] and in women who developed cognitive impairment in another of our population samples [15]. These findings support the hypothesis that a disturbance in brain A $\beta$  metabolism occurs before cortical neuronal degeneration in AD [4, 6]. Furthermore, a PET study found that A $\beta$  burden was not associated with brain atrophy measured over 10 years in elderly individuals without dementia, indicating that A $\beta$  load has not yet affected brain volume in individuals without dementia [63, 64], and that clinical symptoms of dementia start to occur when A $\beta$  gives rise to brain atrophy and neurodegeneration. In contrast, another PET study reported that lower levels of CSF A $\beta$ <sub>42</sub> at baseline were related to brain volume reductions during 1 year of follow-up in cognitively healthy elderly controls [65].

Some limitations of the study have to be considered. First, the number of individuals in some of the groups was relatively small. However, the findings were in the expected direction, even when the sample was divided into several subgroups, and may be considered biologically plausible. Second, although the sample is drawn from the general population, only 30% consented to a lumbar puncture. However, it seems unlikely that this should have affected the associations between CSF levels of A $\beta$  and brain atrophy. Third, linear measurements and visual ratings of cortical atrophy and WMLs on CT is a rather crude method. However, if anything, this should decrease the possibility of finding differences between the groups. In addition, we have previously shown a high interrater reliability in the CT measures [66]. Furthermore, CT is less sensitive than MRI in detecting WMLs and is more severely affected by bone hardening artifacts, which is particularly severe in the region of the medial temporal lobe [67]. However, CT is better in delineating clinically relevant WMLs [68] and comparable to MRI in detecting brain atrophy [67]. On the other hand, if very mild changes are not detected, it may have influenced the results. Also, CT may be more suitable than MRI for the elderly, as it is less sensitive to motion artefacts. CT is also the most used brain imaging tool world-wide. Fourth, all scans were rated by a single person which may increase the risk for systematic error. Fifth, we cannot exclude the possibility that some cases of AD in this very old population might have mixed dementia. Finally, the cross-sectional design limits conclusions regarding cause-effect relations.

In summary, we found that low levels of CSF A $\beta$ <sub>42</sub> and A $\beta$ <sub>40</sub> correlated with WMLs in individuals without dementia, suggesting that this might be an early event in the dementia process, maybe related to deficient clearance of amyloid or ischemia. In contrast, CSF A $\beta$ <sub>42</sub> correlated with cerebral cortical atrophy only in individuals with dementia. These findings suggest that A $\beta$  deposition manifests itself earlier in subcortical white matter compared with cortical brain regions and that subcortical arteriolosclerosis may play a role in the pathogenetic process of AD.

## ACKNOWLEDGMENTS

This study was supported by grants from The Swedish Research Council (11267, 2005-8460, 825-2007-7462, 825-2012-5041, 2013-8717, 2013-61X-14002, 2015-02830), the Torsten Söderbergs Stiftelse at the Royal Swedish Academy of Sciences, Swedish Research Council for Health, Working Life and Welfare (no 2001-2646, 2001-2835, 2003-0234, 2004-0150, 2006-0020, 2008-1229, 2004-0145, 2006-0596, 2008-1111, 2010-0870, AGECAP 2013-2300, 2013-2496, Epilife 2006-1506), Swedish Alzheimer foundation, Hjärnfonden, Swedish Brain Power, The Alzheimer's Association Zenith Award (ZEN-01-3151), The Alzheimer's Association Stephanie B. Overstreet Scholars (IIRG-00-2159), The Knut and Alice Wallenberg Foundation, Sahlgrenska University Hospital (ALF), The Emil and Maria Palm Foundation, The Bank of Sweden Tercentenary Foundation, EU FP7 project LipiDiDiet, Grant Agreement N° 211696, Konung Gustaf V:s och Drottning Victorias Frimurarestiftelse, Eivind och Elsa K:son Sylvans stiftelse, Stiftelsen Söderström-Königska Sjukhemmet, Stiftelsen för Gamla Tjänarinnor, Handlanden Hjalmar Svenssons Forskningsfond, Stiftelsen Längmanska Kulturfonden, Epilife Small Grant, Stiftelsen Demensfonden, Stiftelsen Wilhelm and Martina Lundgrens Vetenskapsfond. None of the funders were involved in the design or interpretation of the study. The funding sources had no involvement in study design; the collection, analysis, and interpretation of data; in the writing of the paper; and in the decision to submit the paper for publication.

Authors' disclosures available online (<https://www.j-alz.com/manuscript-disclosures/17-0950r1>).

## REFERENCES

- [1] Blennow K, de Leon MJ, Zetterberg H (2006) Alzheimer's disease. *Lancet* **368**, 387-403.

- [2] Querfurth HW, LaFerla FM (2010) Alzheimer's disease. *N Engl J Med* **362**, 329-344.
- [3] Hardy J, Selkoe DJ (2002) The amyloid hypothesis of Alzheimer's disease: Progress and problems on the road to therapeutics. *Science* **297**, 353-356.
- [4] Jack Jr CR, Knopman DS, Jagust WJ, Petersen RC, Weiner MW, Aisen PS, Shaw LM, Vemuri P, Wiste HJ, Weigand SD, Lesnick TG, Pankratz VS, Donohue MC, Trojanowski JQ (2013) Tracking pathophysiological processes in Alzheimer's disease: An updated hypothetical model of dynamic biomarkers. *Lancet Neurol* **12**, 207-216.
- [5] Jack Jr CR, Knopman DS, Jagust WJ, Shaw LM, Aisen PS, Weiner MW, Petersen RC, Trojanowski JQ (2010) Hypothetical model of dynamic biomarkers of the Alzheimer's pathological cascade. *Lancet Neurol* **9**, 119-128.
- [6] Weiner MW, Veitch DP, Aisen PS, Beckett LA, Cairns NJ, Green RC, Harvey D, Jack CR, Jagust W, Liu E, Morris JC, Petersen RC, Saykin AJ, Schmidt ME, Shaw L, Siu-ciak JA, Soares H, Toga AW, Trojanowski JQ (2012) The Alzheimer's Disease Neuroimaging Initiative: A review of papers published since its inception. *Alzheimers Dement* **8**, S1-68.
- [7] Blennow K, Hampel H, Weiner M, Zetterberg H (2010) Cerebrospinal fluid and plasma biomarkers in Alzheimer disease. *Nat Rev Neurol* **6**, 131-144.
- [8] Strozzyk D, Blennow K, White LR, Launer LJ (2003) CSF Abeta 42 levels correlate with amyloid-neuropathology in a population-based autopsy study. *Neurology* **60**, 652-656.
- [9] Tapiola T, Alafuzoff I, Herukka SK, Parkkinen L, Hartikainen P, Soinen H, Pirttila T (2009) Cerebrospinal fluid {beta}-amyloid 42 and tau proteins as biomarkers of Alzheimer-type pathologic changes in the brain. *Arch Neurol* **66**, 382-389.
- [10] Fagan AM, Mintun MA, Mach RH, Lee SY, Dence CS, Shah AR, LaRossa GN, Spinner ML, Klunk WE, Mathis CA, DeKosky ST, Morris JC, Holtzman DM (2006) Inverse relation between *in vivo* amyloid imaging load and cerebrospinal fluid Abeta42 in humans. *Ann Neurol* **59**, 512-519.
- [11] Knopman DS, Jack CR, Wiste HJ, Weigand SD, Vemuri P, Lowe VJ, Kantarci K, Gunter JL, Senjem ML, Mielke MM, Roberts RO, Boeve BF, Petersen RC (2013) Selective worsening of brain injury biomarker abnormalities in cognitively normal elderly persons with beta-amyloidosis. *JAMA Neurol* **70**, 1030-1038.
- [12] Palmqvist S, Zetterberg H, Blennow K, Vestberg S, Andreasson U, Brooks DJ, Owenius R, Hagerstrom D, Wollmer P, Minthon L, Hansson O (2014) Accuracy of brain amyloid detection in clinical practice using cerebrospinal fluid beta-amyloid 42: A cross-validation study against amyloid positron emission tomography. *JAMA Neurol* **71**, 1282-1289.
- [13] Skoog I, Davidsson P, Aevarsson O, Vanderstichele H, Vanmechelen E, Blennow K (2003) Cerebrospinal fluid beta-amyloid 42 is reduced before the onset of sporadic dementia: A population-based study in 85-year-olds. *Dement Geriatr Cogn Disord* **15**, 169-176.
- [14] Fagan AM, Roe CM, Xiong C, Mintun MA, Morris JC, Holtzman DM (2007) Cerebrospinal fluid tau/beta-amyloid(42) ratio as a prediction of cognitive decline in nondemented older adults. *Arch Neurol* **64**, 343-349.
- [15] Gustafson DR, Skoog I, Rosengren L, Zetterberg H, Blennow K (2007) Cerebrospinal fluid beta-amyloid 1-42 concentration may predict cognitive decline in older women. *J Neurol Neurosurg Psychiatry* **78**, 461-464.
- [16] Stomrud E, Hansson O, Blennow K, Minthon L, Londo E (2007) Cerebrospinal fluid biomarkers predict decline in subjective cognitive function over 3 years in healthy elderly. *Dement Geriatr Cogn Disord* **24**, 118-124.
- [17] Handoko M, Grant M, Kuskowski M, Zahs KR, Wallin A, Blennow K, Ashe KH (2013) Correlation of specific amyloid-beta oligomers with tau in cerebrospinal fluid from cognitively normal older adults. *JAMA Neurol* **70**, 594-599.
- [18] De Leon MJ, Ferris SH, George AE, Reisberg B, Kricheff II, Gershon S (1980) Computed tomography evaluations of brain-behavior relationships in senile dementia of the Alzheimer's type. *Neurobiol Aging* **1**, 69-79.
- [19] Zhang Y, Londo E, Minthon L, Wattmo C, Liu H, Aspelin P, Wahlund LO (2008) Usefulness of computed tomography linear measurements in diagnosing Alzheimer's disease. *Acta Radiol* **49**, 91-97.
- [20] Thal DR, Rub U, Orantes M, Braak H (2002) Phases of A beta-deposition in the human brain and its relevance for the development of AD. *Neurology* **58**, 1791-1800.
- [21] Buchhave P, Minthon L, Zetterberg H, Wallin AK, Blennow K, Hansson O (2012) Cerebrospinal fluid levels of beta-amyloid 1-42, but not of tau, are fully changed already 5 to 10 years before the onset of Alzheimer dementia. *Arch Gen Psychiatry* **69**, 98-106.
- [22] Selkoe DJ, Hardy J (2016) The amyloid hypothesis of Alzheimer's disease at 25 years. *EMBO Mol Med* **8**, 595-608.
- [23] Jack Jr CR, Lowe VJ, Weigand SD, Wiste HJ, Senjem ML, Knopman DS, Shiung MM, Gunter JL, Boeve BF, Kemp BJ, Weiner M, Petersen RC, Alzheimer's Disease Neuroimaging Initiative (2009) Serial PIB and MRI in normal, mild cognitive impairment and Alzheimer's disease: Implications for sequence of pathological events in Alzheimer's disease. *Brain* **132**, 1355-1365.
- [24] Brun A, Englund E (1986) A white matter disorder in dementia of the Alzheimer type: A pathoanatomical study. *Ann Neurol* **19**, 253-262.
- [25] de la Monte SM (1989) Quantitation of cerebral atrophy in preclinical and end-stage Alzheimer's disease. *Ann Neurol* **25**, 450-459.
- [26] Skoog I, Palmertz B, Andreasson LA (1994) The prevalence of white-matter lesions on computed tomography of the brain in demented and nondemented 85-year-olds. *J Geriatr Psychiatry Neurol* **7**, 169-175.
- [27] Jellinger KA (2002) Alzheimer disease and cerebrovascular pathology: An update. *J Neural Transm* **109**, 813-836.
- [28] Skoog I (1998) A review on blood pressure and ischaemic white matter lesions. *Dement Geriatr Cogn Disord* **9**(Suppl 1), 13-19.
- [29] Guglielmotto M, Aragno M, Autelli R, Giliberto L, Novo E, Colombatto S, Danni O, Parola M, Smith M, Perry G, Tamagno E, Tabaton M (2009) The up-regulation of BACE1 mediated by hypoxia and ischemic injury: Role of oxidative stress and HIF1alpha. *J Neurochem* **108**, 1045-1056.
- [30] Koike MA, Green KN, Blurton-Jones M, Laferla FM (2010) Oligemic hypoperfusion differentially affects tau and amyloid-{beta}. *Am J Pathol* **177**, 300-310.
- [31] Wang X, Xing A, Xu C, Cai Q, Liu H, Li L (2010) Cerebrovascular hypoperfusion induces spatial memory impairment, synaptic changes, and amyloid-beta oligomerization in rats. *J Alzheimers Dis* **21**, 813-822.
- [32] Qi J, Wu H, Yang Y, Wang D, Chen Y, Gu Y, Liu T (2007) Cerebral ischemia and Alzheimer's disease: The expression of amyloid-beta and apolipoprotein E in human hippocampus. *J Alzheimers Dis* **12**, 335-341.



- [33] Wiśniewski HM, Maślińska D (1996) Beta-protein immunoreactivity in the human brain after cardiac arrest. *Folia Neuropathol* **34**, 65-71.
- [34] Grinberg LT, Thal DR (2010) Vascular pathology in the aged human brain. *Acta Neuropathol* **119**, 277-290.
- [35] Skoog I, Nilsson L, Palmertz B, Andreasson LA, Svanborg A (1993) A population-based study of dementia in 85-year-olds. *N Engl J Med* **328**, 153-158.
- [36] Vanderstichele H, Van Kerschaver E, Hesse C, Davidsson P, Buyse MA, Andreasen N, Minthon L, Wallin A, Blennow K, Vanmechelen E (2000) Standardization of measurement of beta-amyloid(1-42) in cerebrospinal fluid and plasma. *Amyloid* **7**, 245-258.
- [37] Andreasen N, Hesse C, Davidsson P, Minthon L, Wallin A, Winblad B, Vanderstichele H, Vanmechelen E, Blennow K (1999) Cerebrospinal fluid beta-amyloid(1-42) in Alzheimer disease: Differences between early- and late-onset Alzheimer disease and stability during the course of disease. *Arch Neurol* **56**, 673-680.
- [38] Simoni M, Pantoni L, Pracucci G, Palmertz B, Guo X, Gustafson D, Skoog I (2008) Prevalence of CT-detected cerebral abnormalities in an elderly Swedish population sample. *Acta Neurol Scand* **118**, 260-267.
- [39] McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM (1984) Clinical diagnosis of Alzheimer's disease: Report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology* **34**, 939-944.
- [40] Roman GC, Tatemichi TK, Erkinjuntti T, Cummings JL, Masdeu JC, Garcia JH, Amaducci L, Orgogozo JM, Brun A, Hofman A, et al. (1993) Vascular dementia: Diagnostic criteria for research studies. Report of the NINDS-AIREN International Workshop. *Neurology* **43**, 250-260.
- [41] Kadir A, Andreasen N, Almkvist O, Wall A, Forsberg A, Engler H, Hagman G, Larksater M, Winblad B, Zetterberg H, Blennow K, Langstrom B, Nordberg A (2008) Effect of phenserine treatment on brain functional activity and amyloid in Alzheimer's disease. *Ann Neurol* **63**, 621-631.
- [42] Cairns NJ, Ikonomic MD, Benzinger T, Storandt M, Fagan AM, Shah AR, Reinwald LT, Carter D, Felton A, Holtzman DM, Mintun MA, Klunk WE, Morris JC (2009) Absence of Pittsburgh compound B detection of cerebral amyloid beta in a patient with clinical, cognitive, and cerebrospinal fluid markers of Alzheimer disease. *Arch Neurol* **66**, 1557-1562.
- [43] Tabaton M, Caponnetto C, Mancardi G, Loeb C (1991) Amyloid beta protein deposition in brains from elderly subjects with leukoaraiosis. *J Neurol Sci* **106**, 123-127.
- [44] Roher AE, Weiss N, Kokjohn TA, Kuo YM, Kalback W, Anthony J, Watson D, Luehrs DC, Sue L, Walker D, Emmerling M, Goux W, Beach T (2002) Increased A beta peptides and reduced cholesterol and myelin proteins characterize white matter degeneration in Alzheimer's disease. *Biochemistry* **41**, 11080-11090.
- [45] Chalmers K, Wilcock G, Love S (2005) Contributors to white matter damage in the frontal lobe in Alzheimer's disease. *Neuropathol Appl Neurobiol* **31**, 623-631.
- [46] Kester MI, Goos JD, Teunissen CE, Benedictus MR, Bouwman FH, Wattjes MP, Barkhof F, Scheltens P, van der Flier WM (2014) Associations between cerebral small-vessel disease and Alzheimer disease pathology as measured by cerebrospinal fluid biomarkers. *JAMA Neurol* **71**, 855-862.
- [47] Appelman AP, Exalto LG, van der Graaf Y, Biessels GJ, Mali WP, Geerlings MI (2009) White matter lesions and brain atrophy: More than shared risk factors? A systematic review. *Cerebrovasc Dis* **28**, 227-242.
- [48] van Westen D, Lindqvist D, Blennow K, Minthon L, Nagga K, Stomrud E, Zetterberg H, Hansson O (2016) Cerebral white matter lesions - associations with Abeta isoforms and amyloid PET. *Sci Rep* **6**, 20709.
- [49] Knopman DS, Roberts R (2010) Vascular risk factors: Imaging and neuropathologic correlates. *J Alzheimers Dis* **20**, 699-709.
- [50] Stenset V, Hofoss D, Johnsen L, Berstad AE, Negaard A, Skinningsrud A, Gjerstad L, Fladby T (2011) White matter lesion load increases the risk of low CSF Abeta42 in apolipoprotein E-varepsilon4 carriers attending a memory clinic. *J Neuroimaging* **21**, e78-82.
- [51] Skoog I (1997) The relationship between blood pressure and dementia: A review. *Biomed Pharmacother* **51**, 367-375.
- [52] FINDER VH, GLOCKSHUBER R (2007) Amyloid-beta aggregation. *Neurodegener Dis* **4**, 13-27.
- [53] Skoog I, Lernfelt B, Landahl S, Palmertz B, Andreasson LA, Nilsson L, Persson G, Oden A, Svanborg A (1996) 15-year longitudinal study of blood pressure and dementia. *Lancet* **347**, 1141-1145.
- [54] Guo X, Pantoni L, Simoni M, Bengtsson C, Bjorkelund C, Lissner L, Gustafson D, Skoog I (2009) Blood pressure components and changes in relation to white matter lesions: A 32-year prospective population study. *Hypertension* **54**, 57-62.
- [55] Sparks DL, Scheff SW, Liu H, Landers TM, Coyne CM, Hunsaker 3rd JC (1995) Increased incidence of neurofibrillary tangles (NFT) in non-demented individuals with hypertension. *J Neurol Sci* **131**, 162-169.
- [56] Hughes TM, Kuller LH, Barinas-Mitchell EJ, McDade EM, Klunk WE, Cohen AD, Mathis CA, Dekosky ST, Price JC, Lopez OL (2014) Arterial stiffness and beta-amyloid progression in nondemented elderly adults. *JAMA Neurol* **71**, 562-568.
- [57] Zhang X, Zhou K, Wang R, Cui J, Lipton SA, Liao FF, Xu H, Zhang YW (2007) Hypoxia-inducible factor 1alpha (HIF-1alpha)-mediated hypoxia increases BACE1 expression and beta-amyloid generation. *J Biol Chem* **282**, 10873-10880.
- [58] Jonsson M, Zetterberg H, van Straaten E, Lind K, Syversen S, Edman A, Blennow K, Rosengren L, Pantoni L, Inzitari D, Wallin A (2010) Cerebrospinal fluid biomarkers of white matter lesions - cross-sectional results from the LADIS study. *Eur J Neurol* **17**, 377-382.
- [59] Selnes P, Blennow K, Zetterberg H, Grambaite R, Rosengren L, Johnsen L, Stenset V, Fladby T (2010) Effects of cerebrovascular disease on amyloid precursor protein metabolites in cerebrospinal fluid. *Cerebrospinal Fluid Res* **7**, 10.
- [60] Schroder J, Pantel J, Ida N, Essig M, Hartmann T, Knopp MV, Schad LR, Sandbrink R, Sauer H, Masters CL, Beyreuther K (1997) Cerebral changes and cerebrospinal fluid beta-amyloid in Alzheimer's disease: A study with quantitative magnetic resonance imaging. *Mol Psychiatry* **2**, 505-507.
- [61] Bouwman FH, Schoonenboom SN, van der Flier WM, van Elk EJ, Kok A, Barkhof F, Blankenstein MA, Scheltens P (2007) CSF biomarkers and medial temporal lobe atrophy predict dementia in mild cognitive impairment. *Neurobiol Aging* **28**, 1070-1074.
- [62] Schuff N, Woerner N, Boreta L, Kornfield T, Shaw LM, Trojanowski JQ, Thompson PM, Jack Jr CR, Weiner MW (2009) MRI of hippocampal volume loss in early

- Alzheimer's disease in relation to ApoE genotype and biomarkers. *Brain* **132**, 1067-1077.
- [63] Driscoll I, Zhou Y, An Y, Sojkova J, Davatzikos C, Kraut MA, Ye W, Ferrucci L, Mathis CA, Klunk WE, Wong DF, Resnick SM (2011) Lack of association between 11C-PiB and longitudinal brain atrophy in non-demented older individuals. *Neurobiol Aging* **32**, 2123-2130.
- [64] Ewers M, Insel P, Jagust WJ, Shaw L, Trojanowski JQ, Aisen P, Petersen RC, Schuff N, Weiner MW, Alzheimer's Disease Neuroimaging Initiative (2012) CSF biomarker and PIB-PET-derived beta-amyloid signature predicts metabolic, gray matter, and cognitive changes in nondemented subjects. *Cereb Cortex* **22**, 1993-2004.
- [65] Fjell AM, Walhovd KB, Fennema-Notestine C, McEvoy LK, Hagler DJ, Holland D, Blennow K, Brewer JB, Dale AM (2010) Brain atrophy in healthy aging is related to CSF levels of Abeta1-42. *Cereb Cortex* **20**, 2069-2079.
- [66] Gustafson D, Lissner L, Bengtsson C, Bjorkelund C, Skoog I (2004) A 24-year follow-up of body mass index and cerebral atrophy. *Neurology* **63**, 1876-1881.
- [67] Frisoni GB (2001) Structural imaging in the clinical diagnosis of Alzheimer's disease: Problems and tools. *J Neurol Neurosurg Psychiatry* **70**, 711-718.
- [68] Lopez OL, Becker JT, Jungreis CA, Rezek D, Estol C, Boller F, DeKosky ST (1995) Computed tomography—but not magnetic resonance imaging—identified periventricular white-matter lesions predict symptomatic cerebrovascular disease in probable Alzheimer's disease. *Arch Neurol* **52**, 659-664.