

CSF neurofilament light concentration is increased in presymptomatic *CHMP2B* mutation carriers

Nina Rostgaard, MSc, Peter Roos, MD, Erik Portelius, PhD, Kaj Blennow, Henrik Zetterberg, Anja H. Simonsen, PhD, and Jørgen E. Nielsen, MD

Neurology® 2018;90:e157-163. doi:10.1212/WNL.0000000000004799

Correspondence

Dr. Simonsen
anja.hviid.simonsen@regionh.dk

Abstract

Objective

A rare cause of familial frontotemporal dementia (FTD) is a mutation in the *CHMP2B* gene on chromosome 3 (FTD-3), described in a Danish family. Here we examine whether CSF biomarkers change in the preclinical phase of the disease.

Methods

In this cross-sectional explorative study, we analyzed CSF samples from 16 mutation carriers and 14 noncarriers from the Danish FTD-3 family. CSF biomarkers included total tau (t-tau) and neurofilament light chain (NfL) as a marker for neurodegeneration, phosphorylated tau (p-tau) as a marker for tau pathology, β -amyloid ($A\beta$) 38, 40, and 42 ($A\beta_{38}$, $A\beta_{40}$, and $A\beta_{42}$) to monitor $A\beta$ metabolism, and YKL-40 as a marker of neuroinflammation. $A\beta$ isoform concentrations were measured using a multiplexed immunoassay; t-tau, p-tau, NfL, and YKL-40 concentrations were measured using sandwich ELISAs.

Results

CSF NfL concentration was significantly increased in mutation carriers vs noncarriers. Further, CSF NfL concentration was significantly higher in symptomatic mutation carriers compared to presymptomatic carriers, and also significantly higher in presymptomatic carriers compared to noncarriers. No differences in t-tau and p-tau and YKL-40 concentrations between controls and mutation carriers were observed. CSF concentrations of the $A\beta$ peptides $A\beta_{38}$ and $A\beta_{40}$ but not $A\beta_{42}$ were significantly lower in mutation carriers compared to noncarriers.

Conclusions

Increased NfL levels in presymptomatic individuals and in symptomatic patients with FTD-3 indicate a continuous process of neurodegeneration from the presymptomatic to symptomatic state. Although not specific for FTD-3 pathology, our data suggest that CSF NfL could serve as a valuable biomarker to detect onset of neurodegeneration in FTD-3 mutation carriers.

From the Danish Dementia Research Centre (N.R., P.R., A.H.S., J.E.N.), Department of Neurology, Rigshospitalet, University of Copenhagen, Denmark; Clinical Neurochemistry Laboratory (E.P., K.B., H.Z.), Sahlgrenska University Hospital; Institute of Neuroscience and Physiology (E.P., K.B., H.Z.), Department of Psychiatry and Neurochemistry, the Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden; and Department of Molecular Neuroscience and UK Dementia Research Institute (H.Z.), UCL Institute of Neurology, Queen Square, London, UK.

Go to Neurology.org/N for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

The Article Processing Charge was funded by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Glossary

AAO = age at onset; **A β ₄₂** = β -amyloid 42; **AD** = Alzheimer disease; **FTD** = frontotemporal dementia; **FTD-3** = chromosome 3–linked frontotemporal dementia; **GLM** = general linear modeling; **NfL** = neurofilament light; **p-tau** = phosphorylated tau; **t-tau** = total tau; **TMB** = tetramethyl benzidine.

A truncating mutation in the *CHMP2B* gene (c.532-1G>C) on chromosome 3 results in early-onset frontotemporal dementia (FTD) (chromosome 3–linked FTD [FTD-3] or *CHMP2B*-FTD).¹ This rare cause of FTD was first described in a large Danish family² and later a different *CHMP2B* mutation was identified in an unrelated Belgian patient with familial FTD.³ The disease is mapped through 6 generations from the first known case in 1876 and the family now counts more than 500 individuals. FTD-3 is characterized by progressive cognitive deficits with behavioral changes. FTD-3 brains present primarily with frontal degeneration as well as temporal and dominant parietal lobe dysfunction.²

The histopathologic hallmarks of *CHMP2B*-FTD include ubiquitin-positive inclusions, autofluorescent aggregates, and p62-positive inclusions, while there is no tau, TAR DNA-binding protein 43, or fused in sarcoma pathology.⁴

The core CSF biomarkers used for dementia diagnosis are β -amyloid 42 (A β ₄₂), total tau (t-tau), and phosphorylated tau (p-tau).⁵ A large subset of FTD cases have pathologic tau changes with robust increases in CSF t-tau and p-tau levels but normal levels of A β ₄₂.⁶ CSF biomarkers in familial FTD have been investigated in individuals with mutations in *MAPT*, *GRN*, or the *C9ORF79* repeat expansion⁷ but not in *CHMP2B* mutation carriers.

In this cross-sectional explorative study, we analyzed the CSF biomarkers A β ₄₂, t-tau, and p-tau181 in 30 individuals from the Danish FTD-3 family. We further included neurofilament light (NfL) as a marker of neuronal injury and YKL-40 as a marker of neuroinflammation in our analysis.

Methods

Study population

The FTD-3 family has been subject to extensive studies over more than 20 years within the Frontotemporal Dementia Research in Jutland Association (FReJA) collaboration, and biological material has been collected during the years for linkage analyses, gene identification, and functional studies.² The disease has been tracked through 6 generations, and clinical characteristics have been recorded in 45 cases of disease, providing information about natural history, clinical characteristics, and age at onset (AAO).

Samples from a total of 30 individuals were included: 10 affected *CHMP2B* mutation carriers, 6 presymptomatic *CHMP2B* mutation carriers, and 14 noncarrier family

members (table). In the affected FTD-3 patients, the mean AAO was 60.5 years (SD 4.8 years).

Standard protocol approvals, registrations, and patient consents

The study was approved by the Ethics Committee of the Capital Region of Denmark (H-1-2012-041), and written informed consent was obtained from each participant before enrollment. Subject age was recorded as the age of the individual on the day of lumbar puncture.

Collection of CSF

In brief, CSF samples were obtained by lumbar puncture and collected in polypropylene tubes, centrifuged at 2,000 g for 10 minutes, aliquoted into cryo tubes, and stored at -80°C until further use.⁸

Immunoassays

A β triplex

CSF concentrations of A β ₃₈, A β ₄₀, and A β ₄₂ were measured using V-plex Peptide Panel 1 Kits A β ₃₈, A β ₄₀, and A β ₄₂ (Meso Scale Discovery system, Rockville, MD) according to the manufacturer's protocol: 60 μL of CSF was diluted 2-fold in diluent provided with the kit. Calibrator samples and controls were prepared according to manufacturer's protocol. On 96-well MSD plates precoated with capture antibodies, 25 μL of sample, calibrator, or controls were added to each well followed by addition of 25 μL of detection antibody and incubation at room temperature for 2 hours. Finally, the plate was washed in kit washing buffer and read in a Meso Scale Discovery imager at appropriate wavelength.

Total tau

CSF t-tau concentration was measured by the hTAU Ag ELISA assay (INNOTEST, Fujirebio, Japan) according to the manufacturer's protocol: on a 96-well microtiter plate precoated with anti-human tau antibody, 25- μL samples, controls, and standards ranging from 50 to 2,500 pg/mL were added followed by incubation with a biotinylated detection antibody to tau and addition of peroxidase-conjugated streptavidin. The reaction was developed with tetramethyl benzidine (TMB) chromogen solution and subsequently stopped with 0.9 M sulfuric acid and quantified at 450 nm in a microplate reader.

Phosphorylated tau

The CSF concentration of tau phosphorylated at amino acid 181 (p-tau) was measured by the p-tau (181p) ELISA assay (INNOTEST, Fujirebio, Japan) according to the manufacturer's protocol: on a 96-well microtiter plate precoated

Table Levels of CSF biomarkers

	Mutation carriers		Noncarriers		p Value
	Mean	95% CL, mean	Mean	95% CL, mean	
Age, y, mean (min-max)	57.8 (32.7– 73.3)	—	59.7 (38.1– 71.1)	—	0.92
Sex, F/M	7/9	—	7/7	—	0.59
A β ₃₈ , pg/mL	1,868.5	1,528.0–2,208.9	2,405.6	2,103.3–2,707.8	0.019
A β ₄₀ , pg/mL	4,874.1	4,109.1–5,639.2	5,849.5	5,319.6–6,379.4	0.038
A β ₄₂ , pg/mL	422.8	332.3–513.2	504.6	414.7–594.6	0.18
YKL-40, pg/mL	132,618.4	108,464–156,772	119,549.9	100,558–138,542	0.38
t-tau, pg/mL	258.7	216.6–301.2	273.4	209.6–337.3	0.68
p-tau, pg/mL	31.3	26.6–36.1	40.1	31.9–48.2	0.049
NfL, pg/mL	2,473.0	1,933.9–3,012.0	686.5	456.6–916.3	<0.0001
A β ₄₂ /A β ₃₈	0.23	0.20–0.26	0.21	0.18–0.25	0.53
A β ₄₂ /A β ₄₀	0.09	0.08–0.10	0.09	0.07–0.10	0.87
t-tau/A β ₄₂	0.69	0.51–0.87	0.63	0.36–0.90	0.68
p-tau/A β ₄₂	0.09	0.06–0.11	0.09	0.06–0.13	0.77

Abbreviations: A β = β -amyloid; CL = confidence level; NfL = neurofilament light; p-tau = phosphorylated tau; t-tau = total tau. CSF concentrations of A β ₃₈, A β ₄₀, A β ₄₂, t-tau, p-tau, YKL-40, and NfL were measured with immunoassays in 30 individuals from the Danish chromosome 3-linked frontotemporal dementia family. Results are given as mean and 95% CL.

with anti-human p-tau antibody, 75- μ L samples, controls, and standards ranging from 15.6 to 1,000 pg/mL were added followed by incubation with a biotinylated detection antibody to p-tau and addition of peroxidase-conjugated streptavidin. The reaction was developed with TMB chromogen solution and subsequently stopped with 0.9 M sulfuric acid and quantified at 450 nm in a microplate reader.

Neurofilament light chain

CSF NfL concentration was measured by sandwich ELISA with the antibody NfL21 used for coating the plates (monoclonal mouse antibody against NfL [in-house]). The coating concentration was 0.5 μ g/mL in carbonate buffer (pH 9.6).

Briefly, 50 μ L CSF (1:2 dilution), 100 μ L calibrator, and 50 μ L PBST (0.05% tween) were added to the coated plates and incubated 1 hour while shaking at room temperature followed by overnight incubation at +4°C. For detection, a biotinylated NfL23 mouse monoclonal antibody against NfL (in-house) was used together with streptavidin–horseradish peroxidase diluted 1:20,000 (100 μ L/well).

Following wash, 100 μ L TMB substrate (TMB One ready-to-use substrate, Kem En Tec Diagnostic, Taastrup, Denmark) was added to each well and plates were incubated for 20 minutes in the dark at room temperature. The reaction was stopped with 100 μ L/well of 0.2 M H₂SO₄ and absorbance was quantified at 450/650 nm in a microplate reader.

YKL-40

CSF YKL-40 concentration was measured using a solid-phase sandwich ELISA (Human Chitinase 3-like 1 Quantikine ELISA kit, R&D Systems, Oxon, UK) according to the manufacturer's protocol: 50 μ L of CSF was diluted 2-fold in diluent provided with the kit. Calibrators and controls were prepared according to protocol.

On a capture-antibody precoated 96-well microtiter plate, 50 μ L of CSF sample, calibrator, or controls were added followed by addition of 200 μ L of conjugate and incubation at room temperature for 2 hours. Finally, each well was washed in the provided washing buffer, followed by addition of 200 μ L substrate solution, and subsequently stopped with 50 μ L of the provided stop solution. Quantification was carried out at 540/570 nm in a microplate reader.

All biochemical measurements were performed by board-certified laboratory technicians who were blinded to clinical information. The measurements were performed in one round of experiments using one batch of reagents. Intra-assay coefficients of variation were below 10%.

Statistical methods

Data were analyzed using SAS software (Enterprise Guide 7.1, 2014, SAS Institute Inc, Cary, NC). Concentrations of each biomarker were compared between controls and *CHMP2B* mutation carriers, and between affected individuals and pre-symptomatic mutation carriers. As some biomarkers were

found to be age-dependent, participant age was included in a subsequent analysis using general linear modeling (GLM). Where it was necessary, logarithmic transformation was performed to obtain normal distribution.

Results

CSF data for all measured markers are shown in the table.

Participant age and sex distribution did not differ significantly between mutation carriers and noncarriers ($t[29] = -0.63, p = 0.53$ and $\chi^2[1] = 0.28, p = 0.59$, respectively).

In our dataset, t-tau, p-tau, and YKL-40 levels were found to be age-dependent (GLM, $F_{1,28} = 6.68, p = 0.0153$; $F_{1,28} = 5.14, p = 0.0314$; $F_{1,28} = 6.37, p = 0.0176$, respectively) while $A\beta_{38}$, $A\beta_{40}$, $A\beta_{42}$, and NfL were not (GLM, $F_{1,28} = 2.45, p = 0.1288$; $F_{1,28} = 1.80, p = 0.1906$; $F_{1,28} = 0.22, p = 0.6450$; $F_{1,27} = 0.38, p = 0.4935$, respectively).

When adjusted for age-dependency, statistical analysis showed no difference in t-tau and p-tau levels between controls and mutation carriers (GLM, $F_{3,26} = 3.29, p = 0.5173$, and $F_{3,26} = 3.86, p = 0.69$, respectively). Levels of $A\beta_{38}$ and $A\beta_{40}$ but not $A\beta_{42}$ were significantly decreased in mutation carriers compared to noncarriers (GLM, $F_{1,28} = 6.23, p = 0.02$; $F_{1,28} = 4.75, p = 0.04$; and $F_{1,28} = 1.87, p = 0.18$, respectively) (figure 1). $A\beta_{42}/A\beta_{38}$, $A\beta_{42}/A\beta_{40}$, t-tau/ $A\beta_{42}$, and p-tau/ $A\beta_{42}$ ratios were not found to be significantly different between the 2 groups (GLM, $F_{1,28} = 0.23, p = 0.63$; $F_{1,28} = 0.05, p = 0.83$; $F_{3,25} = 2.37, p = 0.68$; and $F_{3,25} = 1.96, p = 0.73$, respectively).

YKL-40 CSF levels did not differ significantly between mutation carriers and noncarriers after adjustment for age (GLM, $F_{3,26} = 2.64, p = 0.57$) (figure 1).

CSF levels of NfL were significantly higher in mutation carriers compared to noncarriers (GLM, $F_{1,28} = 53.92, p \leq 0.0001$) (figure 2). When adjusting for participant age, this difference persisted (GLM, $F_{3,26} = 29.44, p = 0.021$). Symptomatic mutation carriers had significantly higher NfL levels than presymptomatic mutation carriers (GLM, $F_{1,14} = 8.61, p = 0.0109$), while presymptomatic carriers had significantly increased NfL levels compared to noncarriers (GLM, $F_{1,17} = 10.97, p = 0.004$).

In the age-corrected model, levels of NfL were found to be significantly increased in symptomatic mutation carriers when compared to presymptomatic carriers (GLM, $F_{3,12} = 7.02, p = 0.0064$), while the significant difference between presymptomatic mutation carriers and noncarriers disappeared (GLM, $F_{3,15} = 9.94, p = 26.22$).

As illustrated (figure 2), NfL levels were high in mutation carriers years prior to the expected clinical onset of 58 years.⁹

Even when adjusting for participant age, this elevation was significant (GLM, $F_{3,26} = 29.44, p = 0.021$).

Discussion


In this explorative study, we analyzed CSF samples from 10 affected *CHMP2B* mutation carriers, 6 presymptomatic *CHMP2B* mutation carriers, and 14 noncarriers from the Danish FTD-3 family. Further investigations on the diagnostic performance of the markers are needed to clarify the influence of age on the levels of the investigated biomarkers.

In our FTD-3 cohort, mutation carriers had significantly increased levels of CSF NfL compared to controls. Interestingly, this increase could be detected in presymptomatic individuals several years prior to expected AAO. Even the youngest presymptomatic mutation carrier (participant age 32, which is 26 years prior to expected clinical onset) had higher CSF NfL levels than some of the oldest noncarriers (figure 2). This suggests that the neurodegenerative process is ongoing in mutation carriers several decades before onset of symptoms, as was also reported in individuals with autosomal inherited Alzheimer disease (AD) in the Dominantly Inherited Alzheimer Network (DIAN) cohort.¹⁰

In contrast to our findings, a recent publication from the GENFI consortium found that CSF NfL levels were significantly increased in symptomatic carriers of mutations in other FTD-causing risk genes, *GRN*, *MAPT*, or *C9ORF27*, when compared to presymptomatic carriers and that presymptomatic carriers and control participants had similar NfL levels.¹¹ In that study, the symptomatic mutation carriers' NfL levels correlated with disease severity, brain atrophy, and survival.¹¹ In recent years, neurofilaments have been investigated in several neurodegenerative diseases¹² and a meta-analysis has concluded that both NfL and neurofilament heavy are increased overall in patients with sporadic AD, FTD, and vascular dementia.¹³ Other studies have shown that CSF NfL levels were increased in sporadic FTD when compared to other dementia groups or healthy controls.^{14–19} Our results and the results from the studies of sporadic FTD cases suggest that NfL may be applied as biomarker of neurodegenerative progression and disease severity.

We detected a significant decrease in CSF $A\beta_{38}$ and $A\beta_{40}$ levels, but not in $A\beta_{42}$ levels, in mutation carriers compared to noncarriers. The $A\beta_{42}/A\beta_{38}$ and $A\beta_{42}/A\beta_{40}$ ratios, however, did not differ between mutation carriers and controls. These results concur with the hypothesis that FTD-3 is not an amyloid disease, as no amyloid pathology has been identified.^{2,20} Substantiating this further, amyloid ligand PET scans in 4 affected patients with FTD-3 have shown no accumulation of amyloid (unpublished material). Lower $A\beta_{38}$ and $A\beta_{40}$ concentrations in CSF have been noted in *PSEN1* mutation carriers and may indicate impaired γ -secretase processing of APP.²¹ Lower $A\beta_{38}$ and $A\beta_{40}$ have also been seen in

Figure 1 CSF biomarkers in *CHMP2B* mutation carriers compared to noncarriers



Box plots show CSF concentrations of β -amyloid 42 ($A\beta_{38}$, $A\beta_{40}$, $A\beta_{42}$), total tau (t-tau), phosphorylated tau (p-tau), and YKL-40 analyzed by immunoassays. Levels of the $A\beta$ peptides $A\beta_{38}$ (A) and $A\beta_{40}$ (C) were significantly decreased in mutation carriers compared to noncarriers ($p = 0.02$ and $p = 0.04$, respectively). Levels of $A\beta_{42}$, t-tau, p-tau, and YKL-40 (B, D–F) were not found to differ significantly between the 2 groups.

neuroinflammatory conditions.²² Finally, there are some indications that reduced neuronal/synaptic activity may lead to less $A\beta$ production in general.^{23,24}


In *MAPT*, *GRN*, and *C9ORF72* mutation carriers, levels of $A\beta_{42}$ are generally found to be within normal range while levels of t-tau and p-tau are not well-characterized due to lack of consistent CSF profile for these patients.⁷

We did not observe a difference in CSF p-tau levels in mutation carriers compared to noncarriers in the current study. This finding is not surprising as patients with FTD-3 do not display the hyperphosphorylated tau pathology that is otherwise observed in other FTDs.^{2,20} Also, we found no

significant increase in CSF t-tau levels. This was surprising as t-tau is known to be a general marker of neurodegeneration and it is well-described that patients with FTD-3 have brain atrophy even early in disease.^{20,25,26}

The CSF level of the inflammation marker YKL-40 was found to increase with age in both mutation carriers and noncarriers in our FTD-3 cohort. This finding concurs with another study showing that CSF YKL-40 levels did not differ between patients with FTD and controls.²⁷ YKL-40 is a secreted 40-kDa glycoprotein belonging to the member of the human chitinase activity family.²⁸ YKL-40 is expressed in several tissues including astrocytes and microglia in the brain. It has been shown that YKL-40 plays an important role in astrocyte

Figure 2 CSF concentrations of neurofilament light (NfL) in symptomatic and presymptomatic *CHMP2B* mutation carriers (MC) and noncarriers



Box plots (A, B) and scatterplot (C) show CSF concentrations of NfL. (A) CSF levels of NfL were significantly higher in MCs compared to noncarriers ($p \leq 0.0001$), also when adjusted for age correlation ($p = 0.021$). (B) Symptomatic MCs had significantly higher NfL levels than presymptomatic MCs ($p = 0.0109$), while presymptomatic carriers had significantly increased NfL levels compared to noncarriers ($p = 0.0002$). (C) Even the youngest presymptomatic MCs had higher NfL levels than aged noncarriers.

and microglial response to neuroinflammation.^{29,30} Our finding suggests that the observed increase is a consequence of age rather than mutation status and therefore YKL-40 may not be a valuable marker of disease in FTD-3. However, in some studies, CSF levels of YKL-40 have been shown to be increased in patients with AD and FTD when compared to cognitively healthy controls.^{31,32} Other studies have found that CSF YKL-40 levels in patients with FTD are increased compared to patients with AD.^{33,34} Since FTD is a clinically and pathologically heterogeneous disease, differences in the individual cohorts analyzed could explain why CSF YKL-40 concentrations vary among different studies (for review, see reference 35).

To our knowledge, all studies of YKL-40 as a CSF biomarker in patients with FTD have reported on sporadic cases and therefore might not reflect the pathology of genetic FTD. In spite of our finding that YKL-40 cannot be used as a marker to differentiate mutation carriers from controls, these results do not necessarily imply that neuroinflammation is not an important aspect of FTD-3 pathology. Recently, microglial activation was studied thoroughly in a FTD-3 mouse model. Early microglial proliferation and a clear proinflammatory phenotype was observed in the *CHMP2B* mutant mice compared to wild-types, which is similar to the inflammatory profile found in FTD-3 brains.^{4,36} These data suggest that neuroinflammation could be a pathologic driver of FTD-3 disease and that it might be valuable to measure CSF biomarkers of neuroinflammation in patients with FTD-3. However, in order to differentiate between disease development and normal aging, a more specific CSF marker than YKL-40 should be used.

Taken together, our results suggest that the unchanged levels of tau in CSF corroborate the absence of tau pathology in

these patients. The decreased levels of $A\beta_{38}$ and $A\beta_{40}$ may be due to a general downregulation of APP processing; however, this needs to be further investigated. Our observations of increase in NfL levels in presymptomatic individuals and even further increase of NfL levels in affected patients with FTD-3 indicate a continuous process of neurodegeneration during the presymptomatic to symptomatic state. Furthermore, our study implies that NfL is a sensitive marker of the widespread neurodegeneration occurring in patients with FTD-3, even decades before symptom onset.

Author contributions

Nina Rostgaard: study concept and design, analysis and interpretation of data, drafting/revising the manuscript for content, and obtaining funding. Peter Roos: study concept and design, statistical analysis, analysis and interpretation of data, drafting/revising the manuscript for content, and obtaining funding. Erik Portelius: study concept and design, acquisition of data, interpretation of data, drafting/revising the manuscript for content, and obtaining funding. Kaj Blennow: study concept and design, acquisition of data, interpretation of data, drafting/revising the manuscript for content, and obtaining funding. Henrik Zetterberg: study concept and design, acquisition of data, interpretation of data, drafting/revising the manuscript for content, and obtaining funding. Anja H. Simonsen: study concept and design, analysis and interpretation of data, and drafting/revising the manuscript for content. Jørgen E. Nielsen: study concept and design, analysis and interpretation of data, drafting/revising the manuscript for content, and obtaining funding.

Acknowledgment

FReJA members: Adrian Isaacs, Anders Gade, Elisabet Englund, Elizabeth Fisher, Jeremy Brown, Jette Stockholm, John Collinge, Peter Johannsen, Susanne Gydesen, Suzanne

Lindquist, Tove Thusgaard, and Troels Tolstrup Nielsen. The authors thank Dr. Jonathan Henry Wardman for linguistic proofreading and FTD-3 family members for continued dedication and support.

Study funding

This study was supported by grants from The Jascha Foundation, Desiree & Niels Ydes Foundation, P.A. Messerschmidts and Wife's Foundation, Director Jacob Madsen and Wife Olga Madsens Foundation, the Novo Nordisk Foundation, the Innovation Fund Denmark ("Brainstem" grant no. 4108-00008B), the Lundbeck Foundation (grant no. R167-2013-15940), the Swedish Research Council, the European Research Council, the Torsten Söderberg Foundation, the Knut and Alice Wallenberg Foundation, and Swedish State Support for Clinical Research (ALFGBG).

Disclosure

N. Rostgaard, P. Roos, and E. Portelius report no disclosures relevant to the manuscript. K. Blennow has served as a consultant or on advisory boards for Alzheon, BioArctic, Biogen, Eli Lilly, Fujirebio Europe, IBL International, Pfizer, and Roche Diagnostics, and is co-founder of Brain Biomarker Solutions in Gothenburg AB, a GU Venture-based platform company at the University of Gothenburg. H. Zetterberg has served on advisory boards of Roche Diagnostics, Eli Lilly, and Pharmasum Therapeutics, and is co-founder of Brain Biomarker Solutions in Gothenburg AB, a GU Venture-based platform company at the University of Gothenburg. A. Simonsen and J. Nielsen report no disclosures relevant to the manuscript. Go to Neurology.org/N for full disclosures.

Received May 21, 2017. Accepted in final form September 28, 2017.

References

1. Skibinski G, Parkinson NJ, Brown JM, et al. Mutations in the endosomal ESCRTIII-complex subunit CHMP2B in frontotemporal dementia. *Nat Genet* 2005;37:806–808.
2. Gydesen S, Brown JM, Brun A, Chakrabarti L, Gade A, Johannsen P, et al. Chromosome 3 linked frontotemporal dementia (FTD-3). *Neurology* 2002;59:1585–1594.
3. Isaacs AM, Johannsen P, Holm I, Nielsen JE. Frontotemporal dementia caused by CHMP2B mutations. *Curr Alzheimer Res* 2011;8:246–251.
4. Clayton EL, Mizielinska S, Edgar JR, et al. Frontotemporal dementia caused by CHMP2B mutation is characterised by neuronal lysosomal storage pathology. *Acta Neuropathol* 2015;130:511–523.
5. Blennow K, Hampel H, Weiner M, Zetterberg H. Cerebrospinal fluid and plasma biomarkers in Alzheimer disease. *Nat Rev Neurol* 2010;6:131–144.
6. Hales CM, Hu WT. From frontotemporal lobar degeneration pathology to frontotemporal lobar degeneration biomarkers. *Int Rev Psychiatry* 2013;25:210–220.
7. Rostgaard N, Waldemar G, Nielsen JE, Simonsen AH. Cerebrospinal fluid biomarkers in familial forms of Alzheimer's disease and frontotemporal dementia. *Dement Geriatr Cogn Disord* 2015;40:54–62.
8. del Campo M, Mollenhauer B, Bertolotto A, et al. Recommendations to standardize preanalytical confounding factors in Alzheimer's and Parkinson's disease cerebrospinal fluid biomarkers: an update. *Biomark Med* 2012;6:419–430.
9. Isaacs AM, Johannsen P, Holm I, et al. Frontotemporal dementia caused by CHMP2B mutations. *Curr Alzheimer Res* 2011;8:246–251.
10. Bateman RJ, Xiong C, Benzinger TLS, et al. Clinical and biomarker changes in dominantly inherited Alzheimer's disease. *N Engl J Med* 2012;367:795–804.
11. Meeter LH, Doppert EG, Jiskoot LC, et al. Neurofilament light chain: a biomarker for genetic frontotemporal dementia. *Ann Clin Transl Neurol* 2016;1–14.
12. Perrot R, Berges R, Bocquet A, Eyer J. Review of the multiple aspects of neurofilament functions, and their possible contribution to neurodegeneration. *Mol Neurobiol* 2008;38:27–65.
13. Petzold A, Keir G, Warren J, Fox N, Rossor MN. A systematic review and meta-analysis of CSF neurofilament protein levels as biomarkers in dementia. *Neurodegener Dis* 2007;4:185–194.
14. Landqvist Waldö M, Frizell Santillo A, Passant U, et al. Cerebrospinal fluid neurofilament light chain protein levels in subtypes of frontotemporal dementia. *BMC Neurol* 2013;13:54.
15. de Jong D, Jansen RWMM, Pijnenburg YAL, et al. CSF neurofilament proteins in the differential diagnosis of dementia. *J Neurol Neurosurg Psychiatry* 2007;78:936–938.
16. Pijnenburg Y, Janssen JC, Schoonenboom NSM, et al. CSF neurofilaments in frontotemporal dementia compared with early onset Alzheimer's disease and controls. *Dement Geriatr Cogn Disord* 2007;23:225–230.
17. Sjogren M, Rosengren L, Minthon L, Davidsson P, Blennow K, Wallin A. Cytoskeleton proteins in CSF distinguish frontotemporal dementia from AD. *Neurology* 2000;54:1960–1964.
18. Scherling CS, Hall T, Berisha F, et al. Cerebrospinal fluid neurofilament concentration reflects disease severity in frontotemporal degeneration. *Ann Neurol* 2014;75:116–126.
19. Skillback T, Farahmand B, Bartlett JW, et al. CSF neurofilament light differs in neurodegenerative diseases and predicts severity and survival. *Neurology* 2014;83:1945–1953.
20. Holm IE, Englund E, Mackenzie IR, Johannsen P, Isaacs AM. A reassessment of the neuropathology of frontotemporal dementia linked to chromosome 3. *J Neuropathol Exp Neurol* 2007;66:884–891.
21. Portelius E, Andreasson U, Ringman JM, et al. Distinct cerebrospinal fluid amyloid beta peptide signatures in sporadic and PSEN1 A431E-associated familial Alzheimer's disease. *Mol Neurodegener* 2010;5:2.
22. Augustis K, Axelsson M, Portelius E, et al. Cerebrospinal fluid biomarkers of β -amyloid metabolism in multiple sclerosis. *Mult Scler J* 2013;19:543–552.
23. Cirrito JR, Yamada KA, Finn MB, et al. Synaptic activity regulates interstitial fluid amyloid- β levels in vivo. *Neuron* 2005;48:913–922.
24. Pomara N, Bruno D, Sarreal AS, et al. Lower CSF amyloid beta peptides and higher F2-Isoprostanes in cognitively intact elderly individuals with major depressive disorder. *Am J Psychiatry* 2012;169:523–530.
25. Eskildsen SF, Østergaard LR, Rodell AB, et al. Cortical volumes and atrophy rates in FTD-3 CHMP2B mutation carriers and related non-carriers. *Neuroimage* 2009;45:713–721.
26. Rohrer JD, Guerreiro R, Vandrovova J, et al. The heritability and genetics of frontotemporal lobar degeneration. *Neurology* 2009;73:1451–1456.
27. Baldacci F, Toschi N, Lista S, et al. Two-level diagnostic classification using cerebrospinal fluid YKL-40 in Alzheimer's disease. *Alzheimers Dement* 2017;13:1–11.
28. Rejman JJ, Hurley WL. Isolation and characterization of a novel 39 kilodalton whey protein from bovine mammary secretions collected during the nonlactating period. *Biochem Biophys Res Commun* 1988;150:329–334.
29. Bonneh-Barkay D, Bissel SJ, Wang G, et al. YKL-40, a marker of simian immunodeficiency virus encephalitis, modulates the biological activity of basic fibroblast growth factor. *Am J Pathol* 2008;173:130–143.
30. Bonneh-Barkay D, Wang G, Starkey A, Hamilton RL, Wiley CA. In vivo CHI3L1 (YKL-40) expression in astrocytes in acute and chronic neurological diseases. *J Neuroinflammation* 2010;7:34.
31. Janelidze S, Hertz J, Zetterberg H, et al. Cerebrospinal fluid neurogranin and YKL-40 as biomarkers of Alzheimer's disease. *Ann Clin Transl Neurol* 2016;3:12–20.
32. Alcolea D, Carmona-Iragui M, Suárez-Calvet M, et al. Relationship between beta-secretase, inflammation and core cerebrospinal fluid biomarkers for Alzheimer's disease. *J Alzheimers Dis* 2014;42:157–167.
33. Craig-Schapiro R, Perrin RJ, Roe CM, et al. YKL-40: a novel prognostic fluid biomarker for preclinical Alzheimer's disease. *Biol Psychiatry* 2010;68:903–912.
34. Teunissen CE, Elias N, Koel-Simmelmink MJA, et al. Novel diagnostic cerebrospinal fluid biomarkers for pathologic subtypes of frontotemporal dementia identified by proteomics. *Alzheimer's Dement Diagnosis, Assess Dis Monit* 2016;2:86–94.
35. Baldacci F, Lista S, Cavado E, Bonuccelli U, Hampel H. Diagnostic function of the neuroinflammatory biomarker YKL-40 in Alzheimer's disease and other neurodegenerative diseases. *Expert Rev Proteomics* 2017;14:285–299.
36. Clayton EL, Mancuso R, Nielsen TT, et al. Early microgliosis precedes neuronal loss and behavioural impairment in mice with a frontotemporal dementia-causing CHMP2B mutation. *Hum Mol Genet* 2017;26:ddx003.

CSF neurofilament light concentration is increased in presymptomatic *CHMP2B* mutation carriers

Nina Rostgaard, MSc, Peter Roos, MD, Erik Portelius, PhD, Kaj Blennow, Henrik Zetterberg, Anja H. Simonsen, PhD, and Jørgen E. Nielsen, MD, PhD, MS

Neurology® 2018;90:65. doi:10.1212/WNL.0000000000004799

Correspondence

Dr. Simonsen
anja.hviid.simonsen@regionh.dk

Study question

Can a biomarker predict the onset of neurodegeneration and clinical symptoms in patients with chromosome-3-linked frontotemporal dementia (FTD-3)?

Summary answer

CSF neurofilament light chain (NfL) was elevated in patients with FTD-3 (*CHMP2B* mutation carriers) before the age at clinical onset and further increased in symptomatic carriers.

What is known and what this article adds

Several studies of familial FTD have described predictive changes in core CSF biomarkers of dementia, including total and phosphorylated tau. This is the first study to examine CSF biomarkers in FTD-3, a rare form of early-onset FTD caused by a truncating mutation in the gene *CHMP2B* on chromosome 3.

Participants and setting

CSF samples were collected from FTD-3 family members including 10 symptomatic carriers, 6 presymptomatic carriers, and 14 noncarriers.

Design, size, and duration

The study used an exploratory cross-sectional design that evaluated CSF biomarkers of FTD pathology among individuals with *CHMP2B* mutation.

Primary outcomes

Primary outcomes were CSF markers including total tau and NfL (indicators of neurodegeneration), phosphorylated tau (indicator of tau pathology), β -amyloid 38, 40, and 42 (indicators of amyloid processing), and YKL-40 (a neuroinflammatory marker).

Main results and the role of chance

After adjustment of the generalized linear model for age-dependency, A β 38 and A β 40 but not A β 42 were significantly decreased in mutation carriers compared to noncarriers. In both the unadjusted and adjusted models, NfL was higher in mutation carriers compared to noncarriers ($F_{3,26} = 29.44$, $p = 0.021$) and higher in symptomatic carriers compared to presymptomatic carriers ($F_{3,12} = 7.02$, $p = 0.0064$). NfL levels were elevated in carrier individuals prior to the age of expected clinical onset, 58 years ($F_{3,26} = 29.44$, $p = 0.021$), supporting the utility of NfL as a biomarker for neurodegeneration.

Bias, confounding, and other reasons for caution


The study was inherently limited by the characteristics and size of the target population and a small sample size.

Generalizability to other populations

The findings are generalizable to individuals with FTD-3.

Study funding/potential competing interests

The study was funded by several government and foundation grants. Go to Neurology.org/N for full disclosures.



A draft of the short-form article was written by A. Symons, a writer with Editage, a division of Cactus Communications. The authors of the full-length article and the journal editors edited and approved the final version.

Neurology®

CSF neurofilament light concentration is increased in presymptomatic *CHMP2B* mutation carriers

Nina Rostgaard, Peter Roos, Erik Portelius, et al.

Neurology 2018;90:e157-e163 Published Online before print December 13, 2017

DOI 10.1212/WNL.0000000000004799

This information is current as of December 13, 2017

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/90/2/e157.full
References	This article cites 35 articles, 5 of which you can access for free at: http://n.neurology.org/content/90/2/e157.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Cognitive Disorders/Dementia http://n.neurology.org/cgi/collection/all_cognitive_disorders_dementia All Genetics http://n.neurology.org/cgi/collection/all_genetics Cerebrospinal Fluid http://n.neurology.org/cgi/collection/cerebrospinal_fluid Frontotemporal dementia http://n.neurology.org/cgi/collection/frontotemporal_dementia
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2017 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

