

TITLE PAGE

Decreased antibodies against Hepatitis A in previously vaccinated treatment naïve
jSLE patients: a prospective case control study

Despoina N. Maritsi^{1,2}, Margarita Onoufriou², George Vartzelis¹

1. Second Department of Pediatrics, "P. & A. Kyriakou" Children's Hospital, Medical
Faculty, University of Athens, Greece

2. "Makarios III" Children's' Hospital, Nicosia, Cyprus

Corresponding author: Despoina N. Maritsi, MD, MSc, PhD, MRCPCH Second
Department of Paediatrics, "P. & A. Kyriakou" Children's Hospital, Medical Faculty,
University of Athens, Greece

Email: dmaritsi@gmail.com, tel: +302130051684, fax: +302118007798

Type of article: Letter

Key words:

Inactivated Hepatitis A vaccine

Juvenile SLE

Immune status

Immunosuppressive treatment

Sir,

Juvenile SLE (jSLE) patients are susceptible to infections due to their defective immune system and the immunosuppressive treatment they receive [1]. However we lack data regarding response to certain vaccine preventable diseases. Hepatitis A is an indolent disease with a varying clinical spectrum and consequences and also a vaccine avoidable disease with intermediate endemicity in Mediterranean countries [2].

In this study we aimed to define the immune status against Hepatitis A virus (HAV) in previously vaccinated jSLE patients, prior to commencement of treatment and at one and three years, and compare this to healthy controls.

This was a prospective matched controlled study including 21 newly diagnosed jSLE patients and 76 healthy controls. The control group consisted of healthy children matched for ethnic origin, age and gender to the control group, attending the Pediatric Outpatients Department for routine checks; the same exclusion criteria applied. All participants had two doses of the inactivated Hepatitis A vaccine in early childhood. Exclusion criteria were underlying immunodeficiency, recent blood transfusion (<6 months) and previous treatment with immunomodulating agents. Demographic, clinical and laboratory data as well as data regarding immunization status, vaccine history and mean time between the doses of the vaccine were collected. Seroprotection rates and anti-HAV-IgG titers were measured at enrollment and at timely intervals afterwards (0,12,36 months). Total IgG levels were measured simultaneously. HAV-IgG antibodies were assessed by ELISA. The cutoff value for seroprotection was deemed at 20IU/ml [3]. The Hospital's Research and Ethics' Committee approved the study; written informed consent was obtained. Statistical significance was set at $p < 0.05$ and analyses were conducted using SPSS (version 19.0).

The two groups had similar demographic characteristics, vaccination history and immunization status. No significant differences were detected in terms of vaccine type, time interval between the two groups as well as mean time from last vaccination to blood sampling. Seroprotection rates were adequate for both groups. Nonetheless, the jSLE group had consistently inferior (but not statistically meaningful) seroprotection rates at all time-points. Mean anti-HAV IgG antibodies were significantly lower in the SLE compared to the control group ($p < 0.01$). Similar results were found at one and three years' follow up (table 1). None of the participants had hypogamaglobulinaemia at the time of blood sampling. During the follow up period the jSLE group had greater decrease in antibody levels as indicated from the significant interaction effect of analysis. There was no association detected between degree of antibody loss and type of treatment or jSLE disease activity.

This was a novel study concerning antibody status against Hepatitis A in children with jSLE. Fully vaccinated participants showed satisfactory seroprotection rates and antibody titers. Nonetheless, we detected increased antibody loss against Hepatitis A in children with jSLE, which further accelerated once treatment is initiated. Low antibody titers could be attributed to the disease[4], its activity[5] or medications[6]. Studies concerning other vaccines, performed in children with IBD, JIA and jSLE showed matching results[7,8,9]. The majority of patients with jSLE receive long-term immunosuppressive treatment. B cell depletion observed in patients under immunosuppressive treatment is a well-documented factor leading to antibody depletion; hence the accelerated rate of antibody loss can be justified. Immunity against certain viruses is severely affected in patients receiving immunosuppressive treatment, rendering these patients susceptible to HAV infection[10]. In fact, corticosteroids and immunosuppressants are associated with decreased vaccine serological response whereas hydroxychloroquine seems to improve vaccine immunogenicity[4]. Additionally, steroid use seems to hamper immunogenicity in a dose dependent manner[6].

SLE is partially a lymphocyte-mediated autoimmune disease with abnormality in the adaptive immune system. Self-antigens set off reactive T cells including Th1 cells with production of pro-inflammatory cytokines. Defective Th1 reaction has been repeatedly reported in vaccine non-responders, rendering this hypothesis challenging for further insight[11]. Our study was underpowered to bring up differences amongst the different treatment groups and correlate disease activity to antibody loss. Finally, although baseline immunology tests were carried out, further studies on B cell function were not performed.

In conclusion, although seroprotection rates were similar between the two groups, mean anti-HAV-IgG titers were significantly lower in the jSLE group at all time-points. Further studies are required to address the question of long-term immunity conveyed by immunizations given at an early stage in children with rheumatic diseases. However, evaluation of immunization status against all vaccine preventable diseases in such patients may be beneficiary.

Key messages:

- Children with autoimmune diseases show evidence of immune dysregulation.
- Accelerated antibody loss against Hepatitis A is noted in children with jSLE

Table 1: Demographic characteristics, seroprotection rates and mean anti-HAV IgG titers for the SLE and the control group

Parameters	SLE group	Control group	P value
------------	-----------	---------------	---------

Study sample, n	21	76	0.9*
Age, years, mean (SD)	11.3 (2.3)	10 (2.7)	0.8*
Gender n (%)			
female	20 (95%)	72 (95%)	0.83+
male	1 (5%)	4 (5%)	
Fully vaccinated	16 (71%)	62 (83%)	0.4+
Partially vaccinated (other than HAV)	5 (29%)	14 (17%)	
Type of vaccine given			
• HAVRIX	34 (81%)	119 (78%)	0.84+
• VAQTA	8 (19%)	33 (22%)	
Mean interval between doses (months)	19	23	0.3*
Mean time from last dose to sampling (years)	6	5.4	0.7*
Steroids (21/21)		NA	-
• mean dose	10mg		
• mean duration of treatment	17 months		
HCQ (21/21)			
• mean dose	200mg		
• mean duration of treatment	36 months		
Azathioprine (9/21)			
• mean dose	75mg		
• mean duration of treatment	18 months		
SLEDAI score			
• Enrolment	7		
• 1 year	1		
• 3 years	0		
Seroprotection rate at diagnosis (%)	95	98	0.4+
Seroprotection rate at 1 years(%)	92	97	0.1+
Seroprotection rate at 3 years (%)	89	96	0.06+
Mean IgG titers at diagnosis, mIU/ml	167	249	< 0.01*
Mean IgG titers at 1 year, mIU/ml	131	218	< 0.01*
Mean IgG titers at 3 years, mIU/ml	134	202	< 0.01*

P-value: * Student t-test, +Pearson's chi-square test

Acknowledgements: The authors wish to thank the families for their participation and support

Conflict of interest: None

Source of funding: None required

REFERENCES:

1. Atzeni F, Bendtzen K, Bobbio-Pallavicini F, Conti F, Cutolo M, Montecucco C, et al. Infections and treatment of patients with rheumatic diseases. *Clin Exp Rheumatol*. 2008;48:S67–73.
2. Fiore AE, Wasley A, Bell BP. Prevention of hepatitis A through active or passive immunization recommendations of the Advisory Committee on Immunization Practice(ACIP). *MMWR Recommend Rep*. 2006;55:1–23
3. Karpinski KF, Hayward S, Tryphonas H. Statistical considerations in the quantitation of serum immunoglobulin levels using the enzyme-linked immunosorbent assay(ELISA). *J Immunol Methods* 1987;103:189-94
4. Pasoto SG, Ribeiro AC, Bonfa E. Update on infections and vaccinations in systemic lupus erythematosus and Sjögren's syndrome. *Curr Opin Rheumatol*. 2014;26:528-37.
5. Campos LM, Silva CA, Aikawa NE, Jesus AA, Moraes JC, Miraglia J, Ishida MA, Bueno C, Pereira RM, Bonfa E. High disease activity: an independent factor for reduced immunogenicity of the pandemic influenza a vaccine in patients with juvenile systemic lupus erythematosus. *Arthritis Care Res*. 2013;65:1121-7
6. Aikawa NE, Campos LM, Silva CA, Carvalho JF, Saad CG, Trudes G, Duarte A, Miraglia JL, Timenetsky Mdo C, Viana VS, França IL, Bonfa E, Pereira RM. Glucocorticoid: major factor for reduced immunogenicity of 2009 influenza A (H1N1) vaccine in patients with juvenile autoimmune rheumatic disease. *J Rheumatol*. 2012;39:167-73.
7. Maritsi D, Vartzelis G, Soldatou A, Garoufi A, Spyridis N. Markedly decreased antibody titers against hepatitis B in previously immunized children presenting with juvenile idiopathic arthritis. *Clin Exp Rheumatol*. 2013; 31: 969-73
8. Radzikowski A, Banaszkiwicz A, Łazowska-Przeorek I, Grzybowska-Chlebowczyk U, Woś H, Pytrus T, Iwańczak B, Kowalska-Dupłaga K, Fyderek K, Gawrońska A, Karolewska-Bochenek K, Kotowska M, Albrecht P. Immunogenicity of hepatitis A vaccine in pediatric patients with inflammatory bowel disease. *Inflamm Bowel Dis*. 2011;17: 1117-24
9. Heijstek MW, Van Gageldonk PG, Ber-Bers GA *et al.*: Differences in persistence of measles, mumps, rubella, diphtheria and tetanus antibodies between children with

rheumatic disease and healthy controls: a retrospective cross-sectional study. *Ann Rheum Dis* 2012; 71: 948-54.

10. Ramos-Casals M, Cuadrado MJ, Alba P, Sanna G, Brito-Zerón P, Bertolaccini L, Babini A, Moreno A, D'Cruz D, Khamashta MA. Acute viral infections in patients with systemic lupus erythematosus: description of 23 cases and review of the literature. *Medicine*. 2008;87:311-8.

11. Jafarzadeh A, Shokri F: The antibody response to HBs antigen is regulated by coordinated Th1 and Th2 cytokine production in healthy neonates *Clin Exp Immunol* 2003; 131: 451-6.