

# ExoMol molecular line lists XXX: a complete high-accuracy line list for water

Oleg L. Polyansky,<sup>1,2</sup> Aleksandra A. Kyuberis,<sup>2</sup> Nikolai F. Zobov,<sup>2</sup>  
Jonathan Tennyson,<sup>1\*</sup> Sergei N. Yurchenko<sup>1</sup> and Lorenzo Lodi<sup>1</sup>

<sup>1</sup>*Department of Physics and Astronomy, University College London, London WC1E 6BT, UK*

<sup>2</sup>*Institute of Applied Physics, Russian Academy of Sciences, Ulyanov Street 46, Nizhny Novgorod 603950, Russia*

Accepted 2018 July 9. Received 2018 July 06; in original form 2018 April 3

## ABSTRACT

A new line list for H<sub>2</sub><sup>16</sup>O is presented. This line list, which is called POKAZATEL, includes transitions between rotational–vibrational energy levels up to 41 000 cm<sup>−1</sup> and is the most complete to date. The potential energy surface (PES) used for producing the line list was obtained by fitting a high-quality *ab initio* PES to experimental energy levels with energies of 41 000 cm<sup>−1</sup> and for rotational excitations up to  $J = 5$ . The final line list comprises all energy levels up to 41 000 cm<sup>−1</sup> and rotational angular momentum  $J$  up to 72. An accurate *ab initio* dipole moment surface was used for the calculation of line intensities and reproduces high-precision experimental intensity data with an accuracy close to 1 per cent. The final line list uses empirical energy levels, whenever they are available, to ensure that line positions are reproduced as accurately as possible. The POKAZATEL line list contains over 5 billion transitions and is available from the ExoMol website ([www.exomol.com](http://www.exomol.com)) and the CDS data base.

**Key words:** molecular data; opacity; planets and satellites: atmospheres; stars: atmospheres; stars: low-mass – stars: brown dwarfs. astronomical data bases: miscellaneous.

## 1 INTRODUCTION

Water is prevalent in the Universe. In particular, the existence of water in a wide range of hot astronomical environments has led to the computation of very extensive line lists of rotational–vibrational transitions both for the main H<sub>2</sub><sup>16</sup>O isotopologue (Allard et al. 1994; Viti, Tennyson & Polyansky 1997; Partridge & Schwenke 1997; Barber et al. 2006) and for its minor isotopologues (Voronin et al. 2010; Partridge & Schwenke 1997; Shirin et al. 2008; Polyansky et al. 2017). The most widely used water line lists are probably those of Partridge & Schwenke (1997), henceforth referred to as the Ames line list, and of Barber et al. (2006), henceforth BT2. The Ames line list contains approximately 300 million lines, while BT2 contains 500 million lines. BT2 provided the main input for water in the 2010 release of the HITEMP data base (Rothman et al. 2010); it has since been subject to a number of validations by comparison with laboratory measurements (Bordbar, Wecel & Hyppanen 2014; Alberti et al. 2015; Melin & Sanders 2016). Such comparisons have shown that the Ames line list is often more accurate than BT2 for transitions with wavelengths longer than 1 μm (wavenumbers < 10 000 cm<sup>−1</sup>) but drops in accuracy at shorter wavelengths. By virtue

of its greater number of lines, BT2 gives an excellent coverage for high temperatures up to 3000 K but it is missing significant flux at higher temperatures and for shorter wavelengths. Thus neither of these line lists can be considered to be fully satisfactory.

The BT2 line list has been used as the basis for a number of atmospheric models, such as the widely used BT-Settl model of Allard (2014). However, there is increasing evidence of the presence of water on objects hotter than 3000 K, in which case the coverage offered by BT2 is inadequate. For example, water has been observed at an effective temperature of over 4000 K in sunspots (Sonnabend et al. 2006) and in a variety of giant stars with temperatures between 3500 and 5000 K (Jennings & Sada 1998; Tsuji 2001; Ryde et al. 2006; Abia et al. 2012; Ryde et al. 2015), in dwarf stars with temperatures up to 4000 K (Rajpurohit et al. 2014) and in variable stars whose atmospheres can also reach these temperatures (Banerjee et al. 2005; Pavlenko et al. 2008).

At slightly lower temperatures, water was the first molecule to be observed in exoplanetary atmospheres (Tinetti et al. 2007) and it is now known to be a common constituent of hot Jupiters (Beaulieu et al. 2010; Iyer et al. 2016) and other exoplanets (Fraine et al. 2014). Some of these exoplanet observations require high-accuracy laboratory data (Birkby et al. 2013; Brogi et al. 2014).

Apart from their use in modelling hot objects, transitions involving highly excited energy levels may be important in environments

\* E-mail: [j.tennyson@ucl.ac.uk](mailto:j.tennyson@ucl.ac.uk)

**Table 1.** Comparison of calculated and experimentally derived (Tennyson et al. 2013) vibrational term values in  $\text{cm}^{-1}$  for  $\text{H}_2^{16}\text{O}$  for four PESs: POKAZATEL (this work), BT2 (Barber et al. 2006), Bubukina (Bubukina et al. 2011) and PS (Partridge & Schwenke 1997); levels marked with a star were not included by PS in their fit. For computed energy levels, we report the difference (observed – calculated).

$v_1$	$v_2$	$v_3$	Obs.	Obs. – Calc.			
				POKAZATEL	BT2	Bubukina	PS
0	1	0	1594.746	0.020	-0.13	-0.016	-0.03
0	2	0	3151.630	0.039	-0.05	-0.004	0.00
1	0	0	3657.053	0.004	-0.10	0.038	0.01
0	0	1	3755.929	-0.006	0.01	-0.003	-0.03
0	3	0	4666.790	0.041	0.07	0.000	0.00
1	1	0	5234.976	0.021	-0.24	0.012	-0.05
0	1	1	5331.267	0.006	-0.01	-0.002	0.05
0	4	0	6134.015	0.025	0.18	-0.007	-0.02
1	2	0	6775.094	-0.013	-0.15	0.011	-0.01
0	2	1	6871.520	0.011	-0.01	0.011	0.02
2	0	0	7201.540	-0.011	0.01	0.003	-0.01
1	0	1	7249.817	-0.066	-0.04	-0.004	-0.04
0	0	2	7445.056	0.003	-0.04	0.023	-0.06
0	5	0	7542.372	-0.009	0.16	0.024	-0.14*
1	3	0	8273.976	-0.018	-0.12	-0.001	-0.07
0	3	1	8373.851	0.003	0.00	0.011	-0.05
2	1	0	8761.582	-0.004	-0.12	-0.004	-0.08
1	1	1	8806.999	-0.040	-0.10	-0.004	-0.04
0	6	0	8869.950	-0.162	-0.32	-0.212	-0.64*
0	1	2	9000.136	0.000	-0.02	0.017	0.03
0	4	1	9833.583	-0.008	0.08	0.004	-0.05
2	2	0	10284.364	-0.025	0.01	0.000	0.01
1	2	1	10328.729	-0.047	0.03	0.001	0.06
0	2	2	10521.758	-0.017	-0.05	0.031	-0.01*
3	0	0	10599.686	-0.025	0.09	-0.003	0.01
2	0	1	10613.356	-0.076	-0.03	-0.003	-0.04
1	0	2	10868.875	-0.022	0.03	0.005	-0.02
0	0	3	11032.404	-0.050	-0.05	0.016	-0.06
0	5	1	11242.776	-0.008	0.16	0.002	0.00*
2	3	0	11767.389	-0.008	-0.09	0.004	-0.13*
1	3	1	11813.207	-0.034	0.00	-0.007	-0.02
0	3	2	12007.774	-0.043	-0.14	0.011	-0.15*
3	1	0	12139.315	-0.031	-0.01	0.003	-0.04
2	1	1	12151.254	-0.057	-0.10	0.010	-0.07
1	1	2	12407.662	-0.016	0.00	0.010	0.01
0	1	3	12565.006	-0.043	-0.02	0.010	0.00
3	2	0	13640.717	-0.023	0.25	0.077	0.14*
2	2	1	13652.655	-0.033	0.21	0.021	0.19
4	0	0	13828.275	-0.001	-0.01	0.013	0.11
3	0	1	13830.938	-0.034	-0.07	0.001	0.10
0	7	1	13835.373	0.035	-0.52	-0.022	-0.48*
1	2	2	13910.894	-0.036	0.08	0.013	0.10
0	2	3	14066.194	-0.056	0.01	0.014	-0.02
2	0	2	14221.159	-0.024	0.10	0.004	0.03
1	0	3	14318.813	-0.053	-0.01	0.010	0.06
1	5	1	14647.971	-0.066	-0.17	0.002	-0.22*
2	3	1	15119.028	-0.008	0.02	0.014	0.00
4	1	0	15344.503	-0.012	-0.09	0.005	0.09
3	1	1	15347.956	-0.035	-0.10	0.001	0.11
0	3	3	15534.709	-0.050	-0.06	0.008	-0.12*
2	1	2	15742.797	-0.062	-0.01	-0.021	0.01
1	1	3	15832.766	-0.053	0.00	0.016	0.09
2	4	1	16546.319	-0.010	-0.10	0.017	-0.12*
3	2	1	16821.631	0.007	0.39	-0.000	0.54*
4	2	0	16823.319	0.031	0.20	0.024	0.35*
4	0	1	16898.842	0.000	0.31	0.001	0.56*
2	2	2	17227.380	0.034	0.14	0.051	0.26*
1	2	3	17312.551	-0.077	0.25	0.005	0.31*
3	0	2	17458.213	0.060	0.08	0.014	0.29*
2	0	3	17495.528	-0.038	-0.07	0.033	0.27*

Table 1 – continued

$v_1$	$v_2$	$v_3$	Obs.	Obs. – Calc.		PS
				POKAZATEL	BT2 Bubukina	
1	0	4	17748.107	−0.032	−0.22 0.010	0.14*
3	3	1	18265.821	−0.031	0.04 −0.008	0.12*
5	1	0	18392.778	0.021	0.08 −0.008	0.84*
4	1	1	18393.315	0.003	0.10 −0.015	0.84*
1	3	3	18758.636	−0.055	0.04 0.018	0.17*
2	1	3	18989.960	−0.038	−0.06 0.019	0.30*
5	0	1	19781.103	0.009	−0.20 0.002	1.06*
6	0	0	19781.323	0.028	−0.69 0.013	0.73*
4	2	1	19865.285	0.011	0.74 0.014	1.79*
2	2	3	20442.777	−0.011	0.29 0.031	
3	0	3	20543.129	−0.024	−0.14 0.034	
5	1	1	21221.827	−0.038	−0.52 0.024	
4	3	1	21314.448	0.040	1.14 0.055	
7	0	0	22529.295	−0.054	0.06 −0.005	2.51*
6	0	1	22529.441	−0.054	0.08 0.004	2.50*
7	0	1	25120.278	−0.067	0.52 0.015	4.25*
5	3	2	27502.660	−0.232	−0.85	
9	0	0	27540.690	0.148	1.11	4.23*
6	1	2	27574.910	−0.069	0.50	
9	1	0	28934.140	0.833	3.20	
10	0	0	29810.850	0.417		7.30*
8	0	2	31071.570	−0.034		
10	1	0	31207.090	−1.724		
11	0	0	31909.679	0.673		9.14*
11	1	0	33144.709	−0.080		
11	0	1	33835.222	0.164		15.33*
12	0	0	33835.249	0.193		15.36*
13	0	0	35585.957	0.407		44.65*
12	0	1	35586.007	−0.565		43.14*
12	2	0	36179.317	−5.567		
13	1	0	36684.047	−2.843		
12	1	1	36684.877	−2.052		
9	1	3	36739.777	−2.312		
10	1	2	36740.597	−1.841		
14	0	0	37122.697	0.517		108.47*
13	0	1	37122.717	0.556		108.50*
11	3	1	37309.847	0.221		
12	3	0	37311.277	−0.593		
13	2	0	37765.647	0.186		
14	1	0	38153.247	0.029		
13	1	1	38153.307	0.045		
15	0	0	38462.517	1.094		283.17*
14	0	1	38462.537	1.112		283.11*
14	2	0	39123.767	0.240		
14	1	1	39390.217	1.220		
15	1	0	39390.257	1.258		
15	0	1	39574.537	0.249		
16	0	0	39574.547	0.218		
12	1	2	40044.567	−3.005		
11	1	3	40044.671	2.611		
14	2	1	40226.261	−4.185		
13	3	1	40262.001	−5.579		
16	1	0	40370.547	−0.921		
15	1	1	40370.781	−0.764		
16	0	1	40437.211	1.941		
12	0	3	40704.156	−2.329		
17	0	1	40945.693	0.270		
18	0	1	41100.053	1.468		
19	0	0	41101.337	−0.554		
15	2	1	41121.606	4.564		

**Table 2.** Comparison of observed – calculated rms deviations ( $\sigma$ ) in  $\text{cm}^{-1}$  as a function of rotational ( $J$ ) states for calculations using our new POKAZATEL PES and using the PES by Bubukina et al. (2011). The reported rms deviations are relative to energy levels up to  $E_{\text{max}}$ .  $N_J$  is the number of levels considered in each case.

PES $E_{\text{max}}$ $J$	POKAZATEL 26 000 $\text{cm}^{-1}$		Bubukina 25 000 $\text{cm}^{-1}$		POKAZATEL 41 000 $\text{cm}^{-1}$	
	$N_J$	$\sigma$	$N_J$	$\sigma$	$N_J$	$\sigma$
0	77	0.034	78	0.023	86	0.113
1	274	0.039	270	0.022	306	0.112
2	482	0.040	475	0.032	548	0.112
3	679	0.037	673	0.029	719	0.081
4	841	0.039	842	0.033	875	0.065
5	994	0.047	997	0.031	1025	0.075
6	1064	0.055	1069	0.032	1074	0.064
7	1110	0.070	1104	0.035	1112	0.072
8	1035	0.087	1037	0.037		
9	953	0.109	950	0.039		
10	857	0.125	851	0.044		
11	750	0.134	752	0.054		
12	676	0.150	672	0.063		
13	615	0.157	605	0.069		
14	575	0.169	560	0.075		
15	527	0.179	519	0.082		
16	494	0.183	503	0.097		
17	474	0.193	480	0.108		
18	450	0.194	460	0.116		
19	442	0.206	450	0.124		
20	432	0.210	420	0.126		
21	399	0.203	396	0.131		
22	366	0.209	358	0.134		
23	344	0.212	329	0.138		
24	316	0.219	306	0.140		
25	283	0.207	268	0.142		
26	251	0.193	234	0.154		
27	238	0.189	228	0.162		
28	218	0.185	193	0.163		
29	181	0.194	172	0.163		
30	143	0.190	124	0.165		
31	110	0.198				
32	81	0.185				
33	46	0.213				
34	24	0.223				
35	19	0.198				
36	17	0.217				
37	11	0.199				
38	10	0.170				
39	4	0.186				

far from thermodynamic equilibrium. For example, water fluorescence on comets can occur from very highly excited levels (Dello Russo et al. 2004, 2005; Barber et al. 2007), and such observations are not always well understood (Barber et al. 2007). Similarly, observations with the Atacama Large Millimeter Array (ALMA) are beginning to probe maser emission from vibrationally excited water (Hirota, Kim & Honma 2012, 2016); modelling water maser emission requires very extensive transition data sets (Gray et al. 2016). Finally, extensive water line lists are important for many terrestrial applications, such as modelling and monitoring emissions from combustion engines (Kranendonk et al. 2007; Rein & Sanders 2010) or studying high-explosive blast waves (Carney et al. 2011).

There have been a number of important developments since the computation of the Ames and the BT2 line lists, which suggests that we are now in a position to compute an  $\text{H}_2^{16}\text{O}$  line list that is

both more comprehensive, indeed effectively complete, and more accurate than either of these earlier lists. In particular, improved theoretical methods have led to the development of both potential energy surfaces (PESs) (Barletta et al. 2006; Lamouroux, Tashkun & Tyuterev 2008; Lodi & Tennyson 2008; Bubukina et al. 2011; Shirin et al. 2006; Császár et al. 2010; Polyansky et al. 2013; Mizus et al. 2018) and dipole moment surfaces (DMSs) (Lamouroux et al. 2008; Lodi et al. 2008; Lodi, Tennyson & Polyansky 2011) with significantly improved accuracy, and nuclear motion calculations that extend all the way to dissociation (Mussa & Tennyson 1998; Li & Guo 2001; Császár et al. 2010). Indeed, our ability to compute high-accuracy transition intensities is leading to such computations (Pavanello et al. 2012; Kyuberis et al. 2017; Birk et al. 2017) replacing measurements in standard compilations such as HITRAN (Gordon et al. 2017). In addition, work by an IUPAC (International

**Table 3.** Extracts from the final states file for the POKAZATEL line list showing portions with full quantum number assignments (upper part) and only rigorous quantum numbers given (lower part).

$i$	$\bar{E}$	$g_{\text{tot}}$	$J$	$K_a$	$K_c$	$\nu_1$	$\nu_2$	$\nu_3$	$S$
2	1594.746306	1	0	0	0	0	1	0	A1
3	3151.629850	1	0	0	0	0	2	0	A1
4	3657.053255	1	0	0	0	1	0	0	A1
5	4666.790461	1	0	0	0	0	3	0	A1
6	5234.975555	1	0	0	0	1	1	0	A1
7	6134.015008	1	0	0	0	0	4	0	A1
8	6775.093508	1	0	0	0	1	2	0	A1
9	7201.539855	1	0	0	0	2	0	0	A1
10	7445.056211	1	0	0	0	0	0	2	A1
11	7542.372492	1	0	0	0	0	5	0	A1
12	8273.975695	1	0	0	0	1	3	0	A1
13	8761.581581	1	0	0	0	2	1	0	A1
14	8869.950054	1	0	0	0	0	6	0	A1
15	9000.136035	1	0	0	0	0	1	2	A1
16	9724.179914	1	0	0	0	1	4	0	A1
17	10085.961796	1	0	0	0	0	7	0	A1
18	10284.364368	1	0	0	0	2	2	0	A1
19	10521.757715	1	0	0	0	0	2	2	A1
20	10599.685969	1	0	0	0	3	0	0	A1
100	21703.511719	1	0	-2	-2	-2	-2	-2	A1
101	21764.097656	1	0	-2	-2	-2	-2	-2	A1
102	21844.693359	1	0	-2	-2	-2	-2	-2	A1
103	21916.152344	1	0	-2	-2	-2	-2	-2	A1
104	21972.789062	1	0	-2	-2	-2	-2	-2	A1
105	22006.955078	1	0	-2	-2	-2	-2	-2	A1
106	22127.925781	1	0	-2	-2	-2	-2	-2	A1
107	22166.060547	1	0	-2	-2	-2	-2	-2	A1
108	22326.316406	1	0	-2	-2	-2	-2	-2	A1
109	22376.539062	1	0	-2	-2	-2	-2	-2	A1
110	22385.830078	1	0	-2	-2	-2	-2	-2	A1

Notes:  $i$ , state counting number;

$\bar{E}$ , state energy in  $\text{cm}^{-1}$ ;

$g_{\text{tot}}$ , total state degeneracy;

$J$ , total angular momentum;

$K_a$ , projection of the angular momentum in the prolate symmetric top limit;

$K_c$ , projection of the angular momentum in the oblate symmetric top limit;

$\nu_1$ , symmetric stretch quantum number;

$\nu_2$ , bending quantum number;

$\nu_3$ , asymmetric stretch quantum number;

$S$ , state symmetry in  $C_{2v}$ .

Union of Pure and Applied Chemistry) task group (Tennyson et al. 2009, 2010, 2013, 2014a, 2014b) has led to the determination of accurate experimental water energy levels; such very accurate, experimentally derived energy levels can be used to replace computed ones, resulting both in the near-perfect reproduction of laboratory transition frequencies and in the prediction of many unobserved line positions with similar accuracy.

In this work we exploit these advances to produce a new line list for the main water isotopologue  $\text{H}_2^{16}\text{O}$ . This work is performed as part of the ExoMol project (Tennyson & Yurchenko 2012), which aims to provide molecular line lists for exoplanet and other hot atmospheres. A unique feature of the resulting line list, which we call POKAZATEL, is that it is not simply complete up to some given temperature as is usual for ExoMol line lists of polyatomic systems (Tennyson & Yurchenko 2018). Instead, the aim is to capture *all* bound-to-bound transitions in the system, which implies considering all energy levels lying below the dissociation limit. POKAZATEL is thus the first complete rotational-vibrational line list for a polyatomic molecule. Besides covering

all temperatures for which the water molecule exists, completeness significantly extends the wavelength range of the line list. In this context, we note the recent detection of near-ultraviolet (near-UV) water absorptions in the Earth's atmosphere by Lampel et al. (2017).

Our ability to calculate an accurate and complete water line list is based on the five following factors: (i) the availability of a spectroscopically accurate *ab initio* potential energy surfaces (PESs) describing energies up to the lowest dissociation pathway (Polyansky et al. 2013; Császár et al. 2010); (ii) the ability to fit this *ab initio* PES to empirical energy levels, which significantly improves the accuracy of computed line positions; (iii) an efficient program suite, DVR3D (Tennyson et al. 2004; Tennyson & Yurchenko 2017), which allows us to compute accurate energies, wavefunctions, dipole moment integrals and intensities of transitions up to dissociation; (iv) the availability of spectroscopic data covering not only the conventional infrared and optical regions below  $26\,000\text{ cm}^{-1}$  (Tolchenov et al. 2005; Polyansky et al. 1998, 1997; Schermaul et al. 2002) but also multiphoton spectra probing the region up to (Grechko

**Table 4.** Extract from the transitions file for the POKAZATEL line list.

$f$	$i$	$A_{fi}$
596233	571007	1.9373e-02
725029	732339	1.0832e-02
329530	297534	8.0899e-04
790239	794617	1.8347e-02
221420	214352	2.3773e-02
421277	402418	6.1962e-02
438351	418788	1.2862e-01
472230	500166	2.6986e-04
442671	459574	1.7567e-03
208210	178893	3.1161e-02
380584	398311	5.4103e-03
437709	442656	4.0364e-03
623411	618141	1.3136e-02
41424	44438	6.1976e-04
638780	616418	2.2414e-04
478821	448210	7.2435e-03
92899	71149	1.3900e-03
190855	172844	3.2447e-02
429814	398308	4.1444e-03
78888	100775	3.7271e-05
735537	742327	2.4587e-04

Notes:  $f$ , upper state counting number;  
 $i$ , lower state counting number;  
 $A_{fi}$ , Einstein-A coefficient in  $s^{-1}$ .

et al. 2009) and even exceeding (Zobov et al. 2011) dissociation; such experimental input allowed us to accurately characterize our new water PES up to dissociation; (v) advances in computer hardware, especially in terms of storage, which allowed us to undertake comprehensive computations that were previously impractical; for example, nuclear-motion wavefunctions relative to a single value of the  $J$  angular momentum often occupy more than 1 Tb of storage.

This paper is organized as follows. In Section 2 we describe the PES used in this work. Section 3 presents details on line position and line intensity calculations. In Section 4 we compare our calculated energy levels with experimental values and with existing line lists. Section 5 presents the POKAZATEL line list and discusses some ways of using it for modelling water spectra. Section 6 concludes the paper.

## 2 CALCULATION OF THE PES

The major, distinguishing feature of POKAZATEL with respect to its predecessors (Partridge & Schwenke 1997; Viti et al. 1997; Barber et al. 2006) is its completeness in both vibrational and rotational states. The  $J=0$  vibrational energy levels included in POKAZATEL reach energies of  $40\,000\text{ cm}^{-1}$ , just below the  $\text{H}_2^{16}\text{O}$  dissociation energy  $D_0=41\,146.1\text{ cm}^{-1}$  (Boyarkin et al. 2013). To ensure completeness in rotational levels, we determined the highest  $J$  for which the lowest rotational energy of the ground vibrational state is below  $41\,000\text{ cm}^{-1}$ . This  $J$  turns out to be  $J=72$ . In other words, for  $J=73$  all the levels lie above  $41\,000\text{ cm}^{-1}$  and were not considered. The inclusion of metastable levels beyond dissociation is, in principle, also possible but would require a significant extension of the methodology employed in this work. Besides, such states are only likely to be important only at temperatures at which water is effectively decomposed and so they are not expected to contribute significantly to the molecular opacity.

In order to calculate energy levels up to  $41\,000\text{ cm}^{-1}$  accurately we require two things. First, a computer program for solving the

rotational–vibrational Schrödinger equation capable of computing all the required states to the necessary accuracy. Second, an accurate PES capable of reproducing to high accuracy (to approximately  $0.05\text{ cm}^{-1}$ ) experimentally known energy levels and covering all energies up to dissociation. The DVR3D program of Tennyson et al. (2004) satisfies our first requirement and has recently benefitted from a number of algorithmic improvements as part of the ExoMol project (Azzam et al. 2016; Underwood et al. 2016; Tennyson & Yurchenko 2017), which proved vital for completing the necessary calculations.

A water PES fulfilling our second requirement was not available, so we produced one as part of this work. Our previous, spectroscopically determined water PESs (Polyansky, Jensen & Tennyson 1996; Shirin et al. 2003; Bubukina et al. 2011) combined with new *ab initio* calculations of the  $\text{H}_2\text{O}$  PES provided us with a very good starting point for constructing our new PES. However, even with these surfaces available, it was not easy to produce a PES that both extends up to dissociation and provides near-spectroscopic accuracy. Our fitted PES is based on two separate data sets of water energy levels. The first set consists of conventional spectroscopic data up to  $25\,000\text{ cm}^{-1}$ , also used in our previous fits (Shirin et al. 2003; Bubukina et al. 2011); the second set comprises energy levels from  $27\,000\text{ cm}^{-1}$  up to  $D_0$  obtained using two-photon and three-photon action spectroscopy (Maksyutenko et al. 2007; Grechko et al. 2008, 2009).

Initially, we tried to produce a single PES reproducing the data belonging to both sets; this proved to be impossible, as any attempt to reproduce the high-lying energy levels to better than  $1\text{ cm}^{-1}$  resulted in an unacceptable deterioration in low-energy levels. Eventually, we decided to follow Varandas (1996) and use two separate PES representations joined by a switching function:

$$V_{\text{glob}} = V_{\text{low}}f(E) + V_{\text{up}}(1 - f(E)), \quad (1)$$

$$f = 0.5[1 + \tanh(\gamma \Delta E)],$$

$$\gamma = \gamma_0 + \gamma_1 \Delta E^2,$$

$$\Delta E = V_u - E_0,$$

$$\gamma_0 = 1/500, \gamma_1 = 1/500^3, E_0 = 35000, \quad (2)$$

where the values for the constants are appropriate for energies in wavenumbers.

The upper and lower surfaces employ the same functional form but have different coefficients:

$$V(S_1, S_2, S_3) = V_0 + \left[ \sum_{i,j,k} f_{ijk} S_1^i S_2^j S_3^k \right] x_3 + V_{\text{HH}} + x_1 + x_2, \quad (3)$$

$$x_1 = D[\exp(-2\alpha \Delta r_1) - 2 \exp(-\alpha \Delta r_1)] + D,$$

$$x_2 = D[\exp(-2\alpha \Delta r_2) - 2 \exp(-\alpha \Delta r_2)] + D,$$

$$x_3 = \exp[-b_1(\Delta r_1^2 + \Delta r_2^2)],$$

$$V_{\text{HH}} = 82000 \exp(-6.2r_{\text{HH}}), \quad (4)$$

$$S_1 = (r_1 + r_2)/2 - r_e, S_2 = (r_1 - r_2)/2, S_3 = \cos \theta - \cos \theta_e, \\ \Delta r_i = r_i - r_e, \quad (5)$$



**Table 5.** Comparison of partition functions: VT (Vidler & Tennyson 2000), definitive (Furtenbacher et al. 2016), BT2 (Barber et al. 2006) and Ames (Partridge & Schwenke 1997).

T(K)	VT	POKAZATEL	Definitive	BT2	Ames
100	35.153	35.15320	35.153 12	35.15451	35.15279
300	178.122	178.1210	178.120 6	178.1279	178.1175
500	386.333	386.3309	386.330 0	386.3446	386.3224
800	823.791	823.7822	823.780 1	823.8028	823.7627
1000	1218.319	1 218.276	1218.273	1 218.299	1 218.247
1200	1717.126	1 717.092	1717.087	1 717.114	1 717.052
1500	2713.816	2 713.061	2713.052	2 713.078	2 713.002
1800	4093.180	4 091.037	4091.024	4 091.046	4 090.949
2000	5279.984	5 276.323	5276.309	5 276.322	5 276.186
2500	9465.976	9 456.146	9456.14	9 455.912	9 455.016
3000	15981.08	15 961.28	15961.3	15 956.96	15 949.21
3200	19433.68	19 408.59	19408.7	19 397.33	19 380.95
3400	23467.56	23 436.15	23436.4	23 409.67	23 377.37
3500	25725.04	25 690.04	25690.4	25 650.77	25 606.48
3600	28155.76	28 116.74	28117.2	28 059.71	27 999.90
3800	33577.48	33 529.18	33530.1	33 415.36	33 310.73
4000	39818.16	39 758.33	39760	39 545.59	39 371.57
4200	46969.6	46 895.09	46899	46 519.39	46 242.48
4400	55128.8	55 036.08	55045	54 404.58	53 980.91
4500	59618.4	59 514.84	59527	58 709.55	58 192.42
4600	64399.2	64 283.29	64300	63 266.73	62 640.65
4800	74888.4	74 743.42	74775	73 168.15	72 271.03
5000	86709.2	86 527.25	86584	84 166.99	82 916.31
5200	99976.4	99 748.69	99847	96 316.52	94 615.14
5400	114808.8	114 523.9	114687	109 664.5	107 400.3
5500	122848.8	122 530.8	122739	116 801.3	114 208.9
5600	131324.8	130 970.1	131234	124 252.9	121 298.7
5800	149644.8	149 204.9	149619	140 117.5	136 330.8
6000	169887.2	169 344.6	169977	157 287.6	152 511.6

where units of Å and  $\text{cm}^{-1}$  are used throughout.  $r_1$ ,  $r_2$  and  $\theta$  are the standard bond lengths and bond angle of water, and  $r_e = 0.9586$  Å and  $\theta_e = 104.48^\circ$  are fixed to reference equilibrium values (Császár et al. 2005). The functions  $x_1$  and  $x_2$  are Morse potentials for each of the OH bonds,  $x_3$  is a damping function, and the term  $V_{\text{HH}}$  is a function of  $r_{\text{HH}}$  representing the distance between the H atoms and is introduced to avoid artificial minima in the region where the H atoms are close to each other (Choi & Light 1992). Other non-linear constants were fixed as follows:  $b_1 = 2.15$  Å $^{-1}$ ,  $D = 43\,900$   $\text{cm}^{-1}$ ,  $\alpha = 2.2668$  Å $^{-1}$ . The coefficients  $f_{ijk}$  were determined as discussed below.

The switch between two potentials ( $E_0 = 35\,000$   $\text{cm}^{-1}$ ) was chosen significantly above  $26\,000$   $\text{cm}^{-1}$  in order to minimize the influence of the upper PES,  $V_{\text{up}}$ , on low-lying levels. The starting point for  $V_{\text{low}}$  was the potential of Bubukina et al. (2011), while the starting point for  $V_{\text{up}}$  was the *ab initio* PES of Császár et al. (2010).  $V_{\text{low}}$  was then determined using a fitting procedure similar to the one employed by Bubukina et al. (2011) for the levels below  $26\,000$   $\text{cm}^{-1}$ ; in this region, a set of 1562 empirical energy levels with  $J = 0, 2, 5$  was used.

For the fit for  $V_{\text{up}}$  we took as the starting point the *ab initio* surface of Császár et al. (2010) and then performed a fit of 734 levels with  $J = 0, 2$  up to  $41\,000$   $\text{cm}^{-1}$ . Of the 41 known  $J = 0$  empirical levels above  $27\,000$   $\text{cm}^{-1}$ , 15 levels had to be excluded from the fit. For  $J = 2$  levels, about 25 per cent of the high-lying levels were excluded. Low-energy levels below  $27\,000$   $\text{cm}^{-1}$  were included in the fit but they were assigned a weight 5 to 10 times lower than high-energy ones.

All fits used the approach of Yurchenko et al. (2003), in which the fitted PES is partially constrained by a set of reference *ab initio* data in order to prevent the emergence of non-physical behaviour such as artificial peaks or troughs. Specifically, the following functional was minimized:


$$F = \sum_i (E_i^{(\text{obs})} - E_i^{(\text{calc})})^2 w_i^{\text{en}} + d \sum_k (V_k^{(\text{ab initio})} - V_k^{(\text{calc})})^2 w_k^{\text{PES}}, \quad (6)$$

where  $E_i$  are the rotational–vibrational energy term values,  $V_k$  represents the value of the *ab initio* PES at the  $k$ th geometry,  $w_i$  are the corresponding weight factors for the individual energies/geometries normalized to one, and  $d$  is a further factor defining the relative importance of the *ab initio* PES relative to the experimental energies. A total of 1460 *ab initio* points were chosen for this constraint, covering the energy region up to about  $50\,000$   $\text{cm}^{-1}$ , which corresponds to O–H bond lengths and H–O–H interbond angles ranging from 0.65 to 2.7 Å and from  $35^\circ$  to  $177^\circ$ , respectively. The final PES was found to deviate from the *ab initio* set of points by less than  $40$   $\text{cm}^{-1}$  for all geometries, which indicates that it preserves a physically correct behaviour also for geometries uncharacterized by the available experimental energy levels.

The weights in equation (6) are normalized as follows:

$$\sum_i w_i^{\text{en}} + \sum_k d w_k^{\text{PES}} = 1. \quad (7)$$

The minimum of the function  $F$  was then found using a simple steepest-descent algorithm by simultaneously fitting the



**Figure 1.** Comparison of BT2 (Barber et al. 2006), POKAZATEL and HITRAN 2016 (Gordon et al. 2017) at blue and near-UV wavelengths. The upper plot is for room temperature ( $T = 296$  K); the lower plot is for  $T = 2000$  K.

potential parameters both to the experimental energies and to *ab initio* points (Yurchenko et al. 2003). The required derivatives of the energies with respect to the potential parameters were computed using the Hellmann–Feynman theorem. In our analytical representation, the potential parameters are included linearly, which simplifies the evaluation of the integrals:

$$\frac{\partial E_n}{\partial f_{ijk}} = \langle \psi_n | \frac{\partial \Delta V}{\partial f_{ijk}} | \psi_n \rangle = \langle \psi_n | S_1^i S_2^j S_3^k x_3 | \psi_n \rangle, \quad (8)$$

where  $E_n$  and  $\psi_n$  are the eigenvalues and eigenfunctions of the rotational–vibrational Hamiltonian  $H$ , respectively,  $n$  is a running index, and the  $f_{ijk}$  are the coefficients of PES expansion.

For the fits we used the fitting wrapper `DVR3D_SFIT` (simultaneous fit) written for `DVR3D` and also used by Bubukina et al. (2011). This is a Fortran 95 program which automatically calls the necessary programs from the `DVR3D` suite, namely `DVR3DRJZ`, `Rotlev3B`, and `Xpect3`. This method was used also to refine an  $\text{H}_2\text{S}$  *ab initio* PES by Azzam et al. (2016). The wrapper can be downloaded from [github.com/exomol](https://github.com/exomol) as part of the `DVR3D` project.

Initially, before performing any fit, our initial, composite PES gave observed minus calculated (obs. – calc.) residues of about

$0.1 \text{ cm}^{-1}$  for levels below  $25\,000 \text{ cm}^{-1}$ . We then re-fitted  $V_{\text{low}}$  using the lower levels while keeping  $V_{\text{up}}$  constant. This led to these levels being reproduced with a standard deviation,  $\sigma$ , of  $0.04 \text{ cm}^{-1}$ . After this, the upper part of the potential  $V_{\text{up}}$  was fitted using the levels between  $25\,000$  and  $41\,000 \text{ cm}^{-1}$ . This final fit gave  $\sigma = 0.13 \text{ cm}^{-1}$  for all levels and  $\sigma = 0.04 \text{ cm}^{-1}$  for the levels below  $26\,000 \text{ cm}^{-1}$ .

In order to improve the accuracy of the calculated energy levels at high angular momentum  $J$ s, we used rotational non-adiabatic correction based on those corrections due to Schwenke (2001). Specifically, we used additional operators  $J_{xx}$ ,  $J_{yy}$  and  $J_{zz}$  to take into account the influence of non-adiabatic effects on highly excited rotational states. The coefficients in front of these operators were treated as additional adjustable parameters and were optimized in calculations of the energy levels with rotational quantum number  $J = 20$ ; this optimization fixed their values at 0.194, 0.194 and 0.14, respectively.

The final potential contains two sets of 246 constants and is given in the Supporting Information as a Fortran program. Section 4 below gives a comparison between energy levels computed using the PES  $V_{\text{glob}}$  of equation (1) both with experimentally derived ones and with previous calculations.

### 3 NUCLEAR MOTION CALCULATIONS


The PES described in the previous section was used to calculate energy levels up to the energy of  $40\,000 \text{ cm}^{-1}$  and for angular momentum up to  $J = 72$ . The programs `DVR3DRJZ` and `ROTLEV3B` from the `DVR3D` program suite (Tennyson et al. 2004) were used to perform the nuclear motion calculations. The energy levels and corresponding wavefunctions were then used to calculate dipole matrix elements using the program `DIPOLE3`. In turn, these matrix elements were then used in the program `SPECTRA` (Tennyson, Miller & Le Sueur 1993) to calculate the line positions and intensities for water transitions in the region from 0 to  $40\,000 \text{ cm}^{-1}$ . Nuclear masses were used in all calculations.

The `DVR3DRJZ` calculations used Radau coordinates with 60 radial grid points and 40 angular grid points. The radial coordinates were represented using Morse-like oscillators with parameters  $r_e = 3.0$ ,  $D_e = 0.25$ ,  $\omega_e = 0.007$  in atomic units; associated Legendre polynomials were used for the angular coordinate. Final vibrational matrices of dimension 5500 were diagonalized to give basis functions for the full rotational–vibrational calculation. For the rotational problem, the dimensions of the final matrices were fixed at  $400(J + 1 - p)$ , where  $J$  is the total angular momentum quantum number and  $p$  is the value of parity. Our final energy levels converge to better than  $0.1 \text{ cm}^{-1}$  at energies around  $40\,000 \text{ cm}^{-1}$ , and are significantly better than this below  $40\,000 \text{ cm}^{-1}$ .

### 4 RESULTS OF THE ENERGY-LEVEL CALCULATIONS

In order to illustrate the accuracy of the energy levels associated with the line list calculations, we compare the computed energy levels with a representative sample of experimentally derived ones for  $\text{H}_2\text{O}$ . Of particular significance is the comparison of  $J = 0$  vibrational term values, as usually discrepancies between observed and calculated (obs. – calc.) energy levels for any  $J$  can be decomposed in a major  $J$ -independent vibrational contribution and in a much smaller  $J$ -dependent rotational one (Polyansky et al. 1997). Table 1 gives a comparison for all experimentally known vibrational term values. These data are representative of the general accuracy of all levels.





**Figure 2.** Comparison of absorption by  $\text{H}_2^{16}\text{O}$  predicted by BT2 and POKAZATEL at  $T = 4000$  K; note the increased absorption with POKAZATEL and flattened structures.

From Table 1 we can draw the following conclusions. First, the POKAZATEL vibrational energy levels up to  $18\,000\text{ cm}^{-1}$  are more accurate than those of previous line lists (Partridge & Schwenke 1997; Shirin et al. 2003). Levels marked with stars were not used in the fit of Partridge & Schwenke (1997) as they were experimentally unknown at that time and show the largest differences with experiment in that line list. In particular, un-starred levels below  $16\,000\text{ cm}^{-1}$  are reproduced in the PS line list with a root-mean-square (rms) deviation of  $0.06\text{ cm}^{-1}$ , while the deviation for starred levels is  $0.41\text{ cm}^{-1}$  for levels up to  $20\,000\text{ cm}^{-1}$  (above this energy the PS line list quickly becomes completely unreliable). Note that the (un-starred) levels included in the fit of Partridge & Schwenke (1997) are reproduced better by both POKAZATEL and Bubukina et al. (2011), with a deviation of, respectively,  $0.03$  and  $0.01\text{ cm}^{-1}$ , while BT2 is comparable with PS with a mean average deviation for those levels of  $0.08\text{ cm}^{-1}$ . Overall levels below  $25\,000\text{ cm}^{-1}$  are best reproduced by the PES of Bubukina et al. (2011), although POKAZATEL follows very closely.

In the energy range between  $18\,000$  and  $30\,000\text{ cm}^{-1}$  one finds discrepancies of up to  $10\text{ cm}^{-1}$  for Partridge & Schwenke (1997) and of up to  $5\text{ cm}^{-1}$  for the BT2 line lists, which used a PES by Shirin et al. (2003). The largest improvement of POKAZATEL with respect to previous line lists is for energies approaching  $40\,000\text{ cm}^{-1}$ , for which Partridge & Schwenke (1997) gives discrepancies of up to  $300\text{ cm}^{-1}$  while our new line lists give about  $0.1\text{ cm}^{-1}$  for levels included in the fit and about  $1\text{ cm}^{-1}$  for levels not included in the fit. Of course, Partridge & Schwenke (1997) did not attempt to fit this region and these results simply illustrate the well-known unreliability of the extrapolation of spectroscopically determined PESs.

Another way to present the comparisons between calculated and experimental values is standard deviations for separate  $J$ s. This gives more detail than the overall standard deviation for all the levels. Thus in Table 2 we report rms deviations for energy levels of given rotational angular momentum  $J$  relative to the PES by Bubukina et al. (2011) and to our new POKAZATEL one. It can

be seen that for POKAZATEL energy levels up to  $25\,000\text{ cm}^{-1}$  the deviations with respect to experiment increase with  $J$ , going from about  $0.04\text{ cm}^{-1}$  for  $J=0-5$  to about  $0.2\text{ cm}^{-1}$  for  $J=35-40$ . For the levels up to  $41\,000\text{ cm}^{-1}$ , which are only known for  $J \leq 7$ , deviations are somewhat larger, namely up to  $0.1\text{ cm}^{-1}$ . The PES by Bubukina et al. (2011), which extends only up to  $26\,000\text{ cm}^{-1}$ , gives deviations that are about half those from POKAZATEL for energy levels below  $25\,000\text{ cm}^{-1}$ .

For intensity calculations, wavefunctions obtained with the POKAZATEL PES were used in all cases. However, for our final set of energy levels given in the states file (see below) the following strategy was used. Where available, empirical energies obtained using the MARVEL procedure by the IUPAC task group (Tennyson et al. 2013) were used. Transitions between these levels should give line positions with experimental accuracy even when these levels are yet to be observed. Unknown energy levels below  $18\,000\text{ cm}^{-1}$  and with  $J \leq 50$  were generated using the PES of Bubukina et al. as these better reproduce observed line positions in this low-energy range. Otherwise, levels from the POKAZATEL PES were used. This strategy gives the best available estimate for each energy level; our data structure allows the states file to be further updated in the eventuality that better (empirical) energies become available in the future (see, for example, Barber et al. 2014).

Our nuclear-motion calculations assign to energy levels only exact quantum numbers, namely  $J$ , parity and the ortho/para symmetry label. Furthermore, energy levels within a given  $J$ -parity-symmetry subset are labelled in increasing order of energy with a counting index  $i$ .

However, it is convenient and standard practice to label energy levels with approximate (normal mode) vibrational  $\nu_1\nu_2\nu_3$  and (rigid rotor) rotational  $JK_aK_c$  quantum numbers. Assigning such labels to every level up to dissociation is difficult (Császár et al. 2010) and probably formally impossible (Child, Weston & Tennyson 1999). Nevertheless, many energy levels can indeed be successfully labelled and such labelling is useful in some applications, for

example when considering pressure broadening. For low  $v$  and  $J$  the labelling procedure is straightforward, while for higher excitations a variety of methods were used, which will be discussed elsewhere. In our line list, vibrational ( $v_1v_2v_3$ ) and rotational ( $J_{K_aK_c}$ ) labels were assigned to more than 72 000 H<sub>2</sub>O energy levels with energies up to 20 000 cm<sup>-1</sup> and  $J \leq 28$ .

## 5 CALCULATION AND REPRESENTATION OF THE LINE LIST

The program suite DVR3D calculates the bound rotational–vibrational energy levels and the corresponding wavefunctions on a three-dimensional grid. Using these wavefunctions and the LTP2011S DMS (Lodi et al. 2011) we computed the Einstein A coefficients,  $A_{if}$ , for transitions up to  $J = 72$  and energies up to 40 000 cm<sup>-1</sup>.

The ExoMol data base uses a condensed format that separates transitions into a states file (which includes quantum labels where available) and a transitions file (Tennyson, Hill & Yurchenko 2013). Extracts from these two files are given in Tables 3 and 4, respectively. These files, which contain 810 269 states and 5 745 071 340 transitions, can be obtained from <ftp://cdsarc.u-strasbg.fr/pub/cats/J/MNRAS/xxx/yy> or <http://cdsarc.u-strasbg.fr/viz-bin/qcat?J/MNRAS/xxx/yy> as well as the ExoMol website, [www.exomol.com](http://www.exomol.com).

In order to further improve the accuracy of the line positions presented in the POKAZATEL line list, we produced two additional sets of energy levels, presented as two different states files in ExoMol format, which are made available in the Supporting Information. In the first additional file we substituted the POKAZATEL energy levels up to 20 000 cm<sup>-1</sup> with those calculated with the PES by Bubukina et al. (2011). A comparison between the POKAZATEL PES and the PES by Bubukina et al. (2011) is reported in Tables 1 and 2 and shows that energy levels produced by Bubukina et al. (2011) are more accurate than those resulting from POKAZATEL for energies up to about 20 000 cm<sup>-1</sup>.

A second set of energy levels was produced exclusively from the experimentally derived ones. This set of levels is significantly more limited but much more accurate, as its accuracy corresponds to that of experimental observations.

Using our calculations we also computed the partition function for H<sub>2</sub><sup>16</sup>O for a wide range of temperatures. The partition function of water is important for a variety of applications, and high-accuracy studies are available concentrating solely on this quantity (Vidler & Tennyson 2000; Furtenbacher et al. 2016). Table 5 compares our partition function with those from various previous studies. All partition functions are computed using the HITRAN convention (Gamache et al. 2017), adopted by the ExoMol project, which explicitly includes the spin degeneracy of all particles. As the value of the partition function at a given temperature always increases when more energy levels are included in its calculation, such a value can be used as a measure of the completeness of a given line list at that temperature (Neale, Miller & Tennyson 1996).

Table 5 shows that the POKAZATEL partition function gives excellent agreement with the recent ‘definitive’ partition function of Furtenbacher et al. (2016). The agreement between these and the older partition function of Vidler & Tennyson (2000) is also excellent; Vidler & Tennyson also considered all states up to dissociation with  $J \leq 72$  but used a rather crude model for the high-lying energies. This illustrates an important point: for accurate partition sums at high temperatures, completeness of the energy level list is more important than accuracy of individual levels. Conversely, both the

BT2 and the Ames partition sums are too low at high temperatures, which reflects the incompleteness of these line lists. Our partition function is given in the supplementary data on a grid of 1 K.

To illustrate our results, Figs 1 and 2 present plots of H<sub>2</sub>O spectra in various spectral regions and for various temperatures. Below about 2500 K, at the low resolution of the plots, POKAZATEL coincides quite closely with BT2 for wavenumbers up to 25 000 cm<sup>-1</sup>.

BT2 uses an energy limit of 30 000 cm<sup>-1</sup> and was designed to be complete for transitions below 20 000 cm<sup>-1</sup>: as a result, its predictions are very different from those by POKAZATEL in the near-UV region above 25 000 cm<sup>-1</sup>, see Fig. 1. In particular, at room temperature the BT2 line list predicts a much larger absorption in the near-UV region than POKAZATEL. The quality of the POKAZATEL line list in this region has been demonstrated in a recent analysis of UV terrestrial atmospheric absorption (Lampel et al. 2017), so we can conclusively say that BT2 overestimates absorption in this region.

At room temperature and for visible wavelengths, POKAZATEL and the recent release of HITRAN (Gordon et al. 2017) give reasonable agreement, but HITRAN contains no data on near-UV transitions. At high temperatures, as expected, HITRAN significantly underestimates the absorption. We note that in this region HITEMP (Rothman et al. 2010) corresponds to BT2.

Fig. 2 shows a comparison between BT2 and POKAZATEL for the high temperature of 4000 K. It can be seen that the absorption spectrum predicted by BT2 is more structured than that of POKAZATEL. The flattening of the spectrum is characteristic of a more complete treatment, including high- $J$  states and vibrational hot bands. High-temperature ( $T > 3000$  K) models relying on BT2 are therefore missing significant opacity.


Fig. 3 presents cross-sections computed using ExoCross (Yurchenko, Al-Refaie & Tennyson 2018) for various temperatures using the POKAZATEL line list. This illustrates the change in the absorption spectra with increasing temperature.

## 6 CONCLUSION

We present in this work a new, very complete water line list, which we call POKAZATEL. The line list includes vibrational and rotational energies up to 40 000 cm<sup>-1</sup> and a maximum rotational angular momentum  $J = 72$ . Our calculations are based on a newly developed PES for water that extends all the way up to the lowest-energy dissociation pathway. The accuracy of the computed energy levels is about 0.1 cm<sup>-1</sup> for all the energies up to dissociation. In the lower-energy region up to 25 000 cm<sup>-1</sup>, accuracy is better. The line list comprises nearly 6 billion lines, an order of magnitude more than any previous line list, and can be used for the modelling of very hot water spectra up to the UV region.

For infrared frequencies and temperatures up to 2000 K, the overall absorption modelled by POKAZATEL is very similar to that by BT2 (Barber et al. 2006), although the accuracy of individual lines is significantly improved, as illustrated in Tables 1 and 2. In particular, we note that recent, independent laboratory studies at room temperature (Campargue et al. 2017; Kassi et al. 2018) and in flames (Rutkowski et al. 2018) have strongly endorsed the accuracy of the POKAZATEL predictions. At short wavelengths and higher temperatures, the completeness of the present line list results in significant opacity differences from BT2.

An important aspect of comprehensive line lists is the treatment of pressure broadening. The ExoMol project recently developed a pressure-broadening diet (Barton et al. 2017) aimed at including the effect of broadening by H<sub>2</sub> and He at high temperatures. Particular



**Figure 3.** Temperature dependence of the absorption cross-sections of the POKAZATEL line list. Cross-sections become increasingly flattened with increasing temperature.

attention has been paid to the broadening of water spectra (Faure et al. 2013; Barton et al. 2017). We note that the implementation used within the ExoMol project allows for the treatment of pressure broadening to be transferred between line lists and hence also to the POKAZATEL line list. All data, including pressure-broadening ones, can be found in the ExoMol data base (Tennyson et al. 2016).

## ACKNOWLEDGEMENTS

This work was supported by the ERC Advanced Investigator Project 267219, the UK Natural Environment Research Council and the Russian Fund for Fundamental Studies (grant 18-02-0057).

## REFERENCES

Abia C., Palmerini S., Busso M., Cristallo S., 2012, *A&A*, 548, A55  
 Alberti M., Weber R., Mancini M., Fateev A., Clausen S., 2015, *J. Quant. Spectrosc. Radiat. Transf.*, 157, 14  
 Allard F., 2014, in Booth M., Matthews B. C., Graham J. R., eds, Proc. IAU Symp. 299, The BT-Settl Model Atmospheres for Stars, Brown Dwarfs and Planets, Kluwer, Dordrecht. p. 271  
 Allard F., Hauschildt P. H., Miller S., Tennyson J., 1994, *ApJ*, 426, L39  
 Azzam A. A. A., Yurchenko S. N., Tennyson J., Naumenko O. V., 2016, *MNRAS*, 460, 4063  
 Banerjee D. P. K., Barber R. J., Ashok N. K., Tennyson J., 2005, *ApJ*, 627, L141  
 Barber R. J., Tennyson J., Harris G. J., Tolchenov R. N., 2006, *MNRAS*, 368, 1087  
 Barber R. J., Miller S., Stallard T., Tennyson J., Hirst P., Carroll T., Adamson A., 2007, *Icarus*, 187, 167  
 Barber R. J., Strange J. K., Hill C., Polyansky O. L., Mellau G. C., Yurchenko S. N., Tennyson J., 2014, *MNRAS*, 437, 1828  
 Barletta P., Shirin S. V., Zobov N. F., Polyansky O. L., Tennyson J., Valeev E. F., Császár A. G., 2006, *J. Chem. Phys.*, 125, 204307  
 Barton E. J., Hill C., Yurchenko S. N., Tennyson J., Dudaryonok A., Lavrentieva N. N., 2017, *J. Quant. Spectrosc. Radiat. Transf.*, 187, 453

Barton E. J., Hill C., Czurylo M., Li H.-Y., Hyslop A., Yurchenko S. N., Tennyson J., 2017, *J. Quant. Spectrosc. Radiat. Transf.*, 203, 490  
 Beaulieu J. P. et al., 2010, *MNRAS*, 409, 963  
 Birk M., Wagner G., Loos J., Lodi L., Polyansky O. L., Kyuberis A. A., Zobov N. F., Tennyson J., 2017, *J. Quant. Spectrosc. Radiat. Transf.*, 203, 88  
 Birkby J. L., de Kok R. J., Brogi M., de Mooij E. J. W., Schwarz H., Albrecht S., Snellen I. A. G., 2013, *MNRAS*, 436, L35  
 Bordbar M. H., Weceł G., Hyppänen T., 2014, *Cumbust. Flame*, 161, 2435  
 Boyarkin O. V. et al., 2013, *Chem. Phys. Lett.*, 568-569, 14  
 Brogi M., de Kok R. J., Birkby J. L., Schwarz H., Snellen I. A. G., 2014, *A&A*, 565, A124  
 Bubukina I. I., Polyansky O. L., Zobov N. F., Yurchenko S. N., 2011, *Optics Spectrosc.*, 110, 160  
 Campargue A. et al., 2017, *J. Quant. Spectrosc. Radiat. Transf.*, 189, 407  
 Carney J. R., Lightstone J. M., Piecuch S., Koch J. D., 2011, *Measure. Sci. Tech.*, 22, 045601  
 Child M. S., Weston T., Tennyson J., 1999, *Mol. Phys.*, 96, 371  
 Choi S. E., Light J. C., 1992, *J. Chem. Phys.*, 97, 7031  
 Császár A. G., Czako G., Furtenbacher T., Tennyson J., Szalay V., Shirin S. V., Zobov N. F., Polyansky O. L., 2005, *J. Chem. Phys.*, 122, 214305  
 Császár A. G., Mátyus E., Szidarovszky T., Lodi L., Zobov N. F., Shirin S. V., Polyansky O. L., Tennyson J., 2010, *J. Quant. Spectrosc. Radiat. Transf.*, 111, 1043  
 Dello Russo N., DiSanti M. A., Magee-Sauer K., Gibb E. L., Mumma M. J., Barber R. J., Tennyson J., 2004, *Icarus*, 168, 186  
 Dello Russo N., Bonev B. P., DiSanti M. A., Gibb E. L., Mumma M. J., Magee-Sauer K., Barber R. J., Tennyson J., 2005, *ApJ*, 621, 537  
 Faure A., Wiesenfeld L., Tennyson J., Drouin B. J., 2013, *J. Quant. Spectrosc. Radiat. Transf.*, 116, 79  
 Fraine J. et al., 2014, *Nature*, 513, 526  
 Furtenbacher T., Szidarovszky T., Hruby J., Kyuberis A. A., Zobov N. F., Polyansky O. L., Tennyson J., Császár A. G., 2016, *J. Phys. Chem. Ref. Data*, 45, 043104  
 Gamache R. R. et al., 2017, *J. Quant. Spectrosc. Radiat. Transf.*, 203, 70  
 Gordon I. E. et al., 2017, *J. Quant. Spectrosc. Radiat. Transf.*, 203, 3  
 Gray M. D., Baudry A., Richards A. M. S., Humphreys E. M. L., Sobolev A. M., Yates J. A., 2016, *MNRAS*, 456, 374

- Grechko M., Maksyutenko P., Zobov N. F., Shirin S. V., Polyansky O. L., Rizzo T. R., Boyarkin O. V., 2008, *J. Phys. Chem. A*, 112, 10539
- Grechko M. et al., 2009, *J. Chem. Phys.*, 131, 221105
- Hirota T., Kim M. K., Honma M., 2012, *ApJ*, 757, L1
- Hirota T., Kim M. K., Honma M., 2016, *ApJ*, 817, 168
- Iyer A. R., Swain M. R., Zellel R. T., Line M. R., Roudier G., Rocha G., Livingston J. H., 2016, *ApJ*, 823, 109
- Jennings D. E., Sada P. V., 1998, *Science*, 279, 844
- Kassi S., Stoltmann T., Casado M., Daeron M., Campargue A., 2018, *J. Chem. Phys.*, 148, 054201
- Kranendonk L. A. et al., 2007, *Optics Express*, 15, 15115
- Kyuberis A. A. et al., 2017, *J. Quant. Spectrosc. Radiat. Transf.*, 203, 175
- Lamouroux J., Tashkun S. A., Tyuterev V. G., 2008, *Chem. Phys. Lett.*, 452, 225
- Lampel J. et al., 2017, *Atmos. Chem. Phys.*, 17, 1271
- Li G., Guo H., 2001, *J. Mol. Spectrosc.*, 210, 90
- Lodi L., Tennyson J., 2008, *J. Quant. Spectrosc. Radiat. Transf.*, 109, 1219
- Lodi L. et al., 2008, *J. Chem. Phys.*, 128, 044304
- Lodi L., Tennyson J., Polyansky O. L., 2011, *J. Chem. Phys.*, 135, 034113
- Maksyutenko P., Muenter J. S., Zobov N. F., Shirin S. V., Polyansky O. L., Rizzo T. R., Boyarkin O. V., 2007, *J. Chem. Phys.*, 126, 241101
- Melin S. T., Sanders S. T., 2016, *J. Quant. Spectrosc. Radiat. Transf.*, 180, 184
- Mizus I. I., Kyuberis A. A., Zobov N. F., Makhnev V. Y., Polyansky O. L., Tennyson J., 2018, *Phil. Trans. Roy. Soc. London A*, 376, 20170149
- Mussa H. Y., Tennyson J., 1998, *J. Chem. Phys.*, 109, 10885
- Neale L., Miller S., Tennyson J., 1996, *ApJ*, 464, 516
- Partridge H., Schwenke D. W., 1997, *J. Chem. Phys.*, 106, 4618
- Pavanello M. et al., 2012, *Phys. Rev. Lett.*, 108, 023002
- Pavlenko Y. V. et al., 2008, *A&A*, 485, 541
- Polyansky O. L., Jensen P., Tennyson J., 1996, *J. Chem. Phys.*, 105, 6490
- Polyansky O. L., Zobov N. F., Tennyson J., Lotoski J. A., Bernath P. F., 1997, *J. Mol. Spectrosc.*, 184, 35
- Polyansky O. L., Zobov N. F., Viti S., Tennyson J., Bernath P. F., Wallace L., 1997, *ApJ*, 489, L205
- Polyansky O. L., Zobov N. F., Viti S., Tennyson J., 1998, *J. Mol. Spectrosc.*, 189, 291
- Polyansky O. L., Ovsyannikov R. I., Kyuberis A. A., Lodi L., Tennyson J., Zobov N. F., 2013, *J. Phys. Chem. A*, 117, 9633
- Polyansky O. L., Kyuberis A. A., Lodi L., Tennyson J., Ovsyannikov R. I., Zobov N., 2017, *MNRAS*, 466, 1363
- Rajpurohit A. S., Reyle C., Allard F., Scholz R. D., Homeier D., Schultheis M., Bayo A., 2014, *A&A*, 564, A90
- Rein K. D., Sanders S. T., 2010, *Appl. Optics*, 49, 4728
- Rothman L. S. et al., 2010, *J. Quant. Spectrosc. Radiat. Transf.*, 111, 2139
- Rutkowski L. et al., 2018, *J. Quant. Spectrosc. Radiat. Transf.*, 205, 213
- Ryde N., Richter M. J., Harper G. M., Eriksson K., Lambert D. L., 2006, *ApJ*, 645, 652
- Ryde N. et al., 2015, *A&A*, 573, A28
- Schermaul R., Learner R. C. M., Canas A. A. D., Brault J. W., Polyansky O. L., Belmiloud D., Zobov N. F., Tennyson J., 2002, *J. Mol. Spectrosc.*, 211, 169
- Schwenke D. W., 2001, *J. Phys. Chem. A*, 105, 2352
- Shirin S. V., Polyansky O. L., Zobov N. F., Barletta P., Tennyson J., 2003, *J. Chem. Phys.*, 118, 2124
- Shirin S. V., Polyansky O. L., Zobov N. F., Ovsyannikov R. I., Császár A. G., Tennyson J., 2006, *J. Mol. Spectrosc.*, 236, 216
- Shirin S. V., Zobov N. F., Ovsyannikov R. I., Polyansky O. L., Tennyson J., 2008, *J. Chem. Phys.*, 128, 224306
- Sonnabend G., Wirtz D., Schieder R., Bernath P., 2006, *Solar Phys.*, 233, 205
- Tennyson J., Yurchenko S. N., 2012, *MNRAS*, 425, 21
- Tennyson J., Yurchenko S. N., 2017, *Intern. J. Quantum Chem.*, 117, 92
- Tennyson J., Yurchenko S. N., 2018, *Atoms*, 6, 26
- Tennyson J., Miller S., Le Sueur C. R., 1993, *Comput. Phys. Commun.*, 75, 339
- Tennyson J., Kostin M. A., Barletta P., Harris G. J., Polyansky O. L., Ramanalal J., Zobov N. F., 2004, *Comput. Phys. Commun.*, 163, 85
- Tennyson J. et al., 2009, *J. Quant. Spectrosc. Radiat. Transf.*, 110, 573
- Tennyson J. et al., 2010, *J. Quant. Spectrosc. Radiat. Transf.*, 111, 2160
- Tennyson J. et al., 2013, *J. Quant. Spectrosc. Radiat. Transf.*, 117, 29
- Tennyson J., Hill C., Yurchenko S. N., 2013, in Gillaspay J. D., Weise W. L., Podpaly Y. A., AIP Vol. 1545, Data Structures for ExoMol: Molecular Line Lists for Exoplanet and Other Atmospheres. AIP, New York, p. 186
- Tennyson J. et al., 2014a, *Pure Appl. Chem.*, 86, 71
- Tennyson J. et al., 2014b, *J. Quant. Spectrosc. Radiat. Transf.*, 142, 93
- Tennyson J. et al., 2016, *J. Mol. Spectrosc.*, 327, 73
- Tinetti G. et al., 2007, *Nature*, 448, 169
- Tolchenov R. N. et al., 2005, *J. Mol. Spectrosc.*, 233, 68
- Tsuji T., 2001, *A&A*, 376, L1
- Underwood D. S., Tennyson J., Yurchenko S. N., Huang X., Schwenke D. W., Lee T. J., Clausen S., Fateev A., 2016, *MNRAS*, 459, 3890
- Varandas A. J. C., 1996, *J. Chem. Phys.*, p. 3524
- Vidler M., Tennyson J., 2000, *J. Chem. Phys.*, 113, 9766
- Viti S., Tennyson J., Polyansky O. L., 1997, *MNRAS*, 287, 79
- Voronin B. A., Tennyson J., Tolchenov R. N., Lugovskoy A. A., Yurchenko S. N., 2010, *MNRAS*, 402, 492
- Yurchenko S. N., Carvajal M., Jensen P., Herregodts F., Huet T. R., 2003, *Chem. Phys.*, 290, 59
- Yurchenko S. N., Al-Refaie A. F., Tennyson J., 2018, *A&A*, 614, A131
- Zobov N. F., Shirin S. V., Lodi L., Silva B. C., Tennyson J., Császár A. G., Polyansky O. L., 2011, *Chem. Phys. Lett.*, 507, 48

## SUPPORTING INFORMATION

Supplementary data are available at *MNRAS* online.

Please note: Oxford University Press is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

This paper has been typeset from a  $\text{\TeX}/\text{\LaTeX}$  file prepared by the author.



# List of astronomical key words (Updated on 2017 March)

This list is common to *Monthly Notices of the Royal Astronomical Society*, *Astronomy and Astrophysics*, and *The Astrophysical Journal*. In order to ease the search, the key words are subdivided into broad categories. No more than *six* subcategories altogether should be listed for a paper.

The subcategories in boldface containing the word ‘individual’ are intended for use with specific astronomical objects; these should never be used alone, but always in combination with the most common names for the astronomical objects in question. Note that each object counts as one subcategory within the allowed limit of six.

The parts of the key words in italics are for reference only and should be omitted when the keywords are entered on the manuscript.

## **General**

editorials, notices  
errata, addenda  
extraterrestrial intelligence  
history and philosophy of astronomy  
miscellaneous  
obituaries, biographies  
publications, bibliography  
sociology of astronomy  
standards

## **Physical data and processes**

acceleration of particles  
accretion, accretion discs  
asteroseismology  
astrobiology  
astrochemistry  
astroparticle physics  
atomic data  
atomic processes  
black hole physics  
chaos  
conduction  
convection  
dense matter  
diffusion  
dynamo  
elementary particles  
equation of state  
gravitation  
gravitational lensing: micro  
gravitational lensing: strong  
gravitational lensing: weak  
gravitational waves  
hydrodynamics  
instabilities  
line: formation  
line: identification  
line: profiles  
magnetic fields  
magnetic reconnection  
(*magnetohydrodynamics*) MHD  
masers  
molecular data  
molecular processes  
neutrinos  
nuclear reactions, nucleosynthesis, abundances  
opacity  
plasmas  
polarization

radiation: dynamics  
radiation mechanisms: general  
radiation mechanisms: non-thermal  
radiation mechanisms: thermal  
radiative transfer  
relativistic processes  
scattering  
shock waves  
solid state: refractory  
solid state: volatile  
turbulence  
waves

## **Astronomical instrumentation, methods and techniques**

atmospheric effects  
balloons  
instrumentation: adaptive optics  
instrumentation: detectors  
instrumentation: high angular resolution  
instrumentation: interferometers  
instrumentation: miscellaneous  
instrumentation: photometers  
instrumentation: polarimeters  
instrumentation: spectrographs  
light pollution  
methods: analytical  
methods: data analysis  
methods: laboratory: atomic  
methods: laboratory: molecular  
methods: laboratory: solid state  
methods: miscellaneous  
methods: numerical  
methods: observational  
methods: statistical  
site testing  
space vehicles  
space vehicles: instruments  
techniques: high angular resolution  
techniques: image processing  
techniques: imaging spectroscopy  
techniques: interferometric  
techniques: miscellaneous  
techniques: photometric  
techniques: polarimetric  
techniques: radar astronomy  
techniques: radial velocities  
techniques: spectroscopic  
telescopes



**Astronomical data bases**

astronomical data bases: miscellaneous  
 atlases  
 catalogues  
 surveys  
 virtual observatory tools

**Astrometry and celestial mechanics**

astrometry  
 celestial mechanics  
 eclipses  
 ephemerides  
 occultations  
 parallaxes  
 proper motions  
 reference systems  
 time

**The Sun**

Sun: abundances  
 Sun: activity  
 Sun: atmosphere  
 Sun: chromosphere  
 Sun: corona  
 Sun: coronal mass ejections (CMEs)  
 Sun: evolution  
 Sun: faculae, plages  
 Sun: filaments, prominences  
 Sun: flares  
 Sun: fundamental parameters  
 Sun: general  
 Sun: granulation  
 Sun: helioseismology  
 Sun: heliosphere  
 Sun: infrared  
 Sun: interior  
 Sun: magnetic fields  
 Sun: oscillations  
 Sun: particle emission  
 Sun: photosphere  
 Sun: radio radiation  
 Sun: rotation  
 (*Sun*:) solar–terrestrial relations  
 (*Sun*:) solar wind  
 (*Sun*:) sunspots  
 Sun: transition region  
 Sun: UV radiation  
 Sun: X-rays, gamma-rays

**Planetary systems**

comets: general

**comets: individual: . . .**

Earth  
 interplanetary medium  
 Kuiper belt: general

**Kuiper belt objects: individual: . . .**

meteorites, meteors, meteoroids  
 minor planets, asteroids: general

**minor planets, asteroids: individual: . . .**

## Moon

Oort Cloud  
 planets and satellites: atmospheres  
 planets and satellites: aurorae  
 planets and satellites: composition  
 planets and satellites: detection  
 planets and satellites: dynamical evolution and stability  
 planets and satellites: formation  
 planets and satellites: fundamental parameters  
 planets and satellites: gaseous planets  
 planets and satellites: general

**planets and satellites: individual: . . .**

planets and satellites: interiors  
 planets and satellites: magnetic fields  
 planets and satellites: oceans  
 planets and satellites: physical evolution  
 planets and satellites: rings  
 planets and satellites: surfaces  
 planets and satellites: tectonics  
 planets and satellites: terrestrial planets  
 planet–disc interactions  
 planet–star interactions  
 protoplanetary discs  
 zodiacal dust

**Stars**

stars: abundances  
 stars: activity  
 stars: AGB and post-AGB  
 stars: atmospheres  
 (*stars*:) binaries (*including multiple*): close  
 (*stars*:) binaries: eclipsing  
 (*stars*:) binaries: general  
 (*stars*:) binaries: spectroscopic  
 (*stars*:) binaries: symbiotic  
 (*stars*:) binaries: visual  
 stars: black holes  
 (*stars*:) blue stragglers  
 (*stars*:) brown dwarfs  
 stars: carbon  
 stars: chemically peculiar  
 stars: chromospheres  
 (*stars*:) circumstellar matter  
 stars: coronae  
 stars: distances  
 stars: dwarf novae  
 stars: early-type  
 stars: emission-line, Be  
 stars: evolution  
 stars: flare  
 stars: formation  
 stars: fundamental parameters  
 (*stars*:) gamma-ray burst: general  
 (*stars*:) **gamma-ray burst: individual: . . .**  
 stars: general  
 (*stars*:) Hertzsprung–Russell and colour–magnitude diagrams  
 stars: horizontal branch  
 stars: imaging  
**stars: individual: . . .**  
 stars: interiors

stars: jets  
 stars: kinematics and dynamics  
 stars: late-type  
 stars: low-mass  
 stars: luminosity function, mass function  
 stars: magnetars  
 stars: magnetic field  
 stars: massive  
 stars: mass-loss  
 stars: neutron  
 (*stars:*) novae, cataclysmic variables  
 stars: oscillations (*including pulsations*)  
 stars: peculiar (*except chemically peculiar*)  
 (*stars:*) planetary systems  
 stars: Population II  
 stars: Population III  
 stars: pre-main-sequence  
 stars: protostars  
 (*stars:*) pulsars: general  
 (*stars:*) **pulsars: individual: . . .**  
 stars: rotation  
 stars: solar-type  
 (*stars:*) starspots  
 stars: statistics  
 (*stars:*) subdwarfs  
 (*stars:*) supergiants  
 (*stars:*) supernovae: general  
 (*stars:*) **supernovae: individual: . . .**  
 stars: variables: Cepheids  
 stars: variables: Scuti  
 stars: variables: general  
 stars: variables: RR Lyrae  
 stars: variables: S Doradus  
 stars: variables: T Tauri, Herbig Ae/Be  
 (*stars:*) white dwarfs  
 stars: winds, outflows  
 stars: Wolf–Rayet

### Interstellar medium (ISM), nebulae

ISM: abundances  
 ISM: atoms  
 ISM: bubbles  
 ISM: clouds  
 (*ISM:*) cosmic rays  
 (*ISM:*) dust, extinction  
 ISM: evolution  
 ISM: general  
 (*ISM:*) HII regions  
 (*ISM:*) Herbig–Haro objects

### ISM: individual objects: . . .

(*except planetary nebulae*)  
 ISM: jets and outflows  
 ISM: kinematics and dynamics  
 ISM: lines and bands  
 ISM: magnetic fields  
 ISM: molecules  
 (*ISM:*) photodissociation region (PDR)  
 (*ISM:*) planetary nebulae: general  
 (*ISM:*) **planetary nebulae: individual: . . .**  
 ISM: structure  
 ISM: supernova remnants

### The Galaxy

Galaxy: abundances  
 Galaxy: bulge  
 Galaxy: centre  
 Galaxy: disc  
 Galaxy: evolution  
 Galaxy: formation  
 Galaxy: fundamental parameters  
 Galaxy: general  
 (*Galaxy:*) globular clusters: general  
 (*Galaxy:*) **globular clusters: individual: . . .**  
 Galaxy: halo  
 Galaxy: kinematics and dynamics  
 (*Galaxy:*) local interstellar matter  
 Galaxy: nucleus  
 (*Galaxy:*) open clusters and associations: general  
 (*Galaxy:*) **open clusters and associations: individual: . . .**  
 (*Galaxy:*) solar neighbourhood  
 Galaxy: stellar content  
 Galaxy: structure

### Galaxies

galaxies: abundances  
 galaxies: active  
 (*galaxies:*) BL Lacertae objects: general  
 (*galaxies:*) **BL Lacertae objects: individual: . . .**  
 galaxies: bulges  
 galaxies: clusters: general

### galaxies: clusters: individual: . . .

galaxies: clusters: intracluster medium  
 galaxies: distances and redshifts  
 galaxies: dwarf  
 galaxies: elliptical and lenticular, cD  
 galaxies: evolution  
 galaxies: formation  
 galaxies: fundamental parameters  
 galaxies: general  
 galaxies: groups: general

### galaxies: groups: individual: . . .

galaxies: haloes  
 galaxies: high-redshift

### galaxies: individual: . . .

galaxies: interactions  
 (*galaxies:*) intergalactic medium  
 galaxies: irregular  
 galaxies: ISM  
 galaxies: jets  
 galaxies: kinematics and dynamics  
 (*galaxies:*) Local Group  
 galaxies: luminosity function, mass function  
 (*galaxies:*) Magellanic Clouds  
 galaxies: magnetic fields  
 galaxies: nuclei  
 galaxies: peculiar  
 galaxies: photometry  
 (*galaxies:*) quasars: absorption lines  
 (*galaxies:*) quasars: emission lines  
 (*galaxies:*) quasars: general

*(galaxies:)* **quasars: individual: . . .**

*(galaxies:)* quasars: supermassive black holes  
galaxies: Seyfert  
galaxies: spiral  
galaxies: starburst  
galaxies: star clusters: general

**galaxies: star clusters: individual: . . .**

galaxies: star formation  
galaxies: statistics  
galaxies: stellar content  
galaxies: structure

### **Cosmology**

*(cosmology:)* cosmic background radiation  
*(cosmology:)* cosmological parameters  
*(cosmology:)* dark ages, reionization, first stars  
*(cosmology:)* dark energy  
*(cosmology:)* dark matter  
*(cosmology:)* diffuse radiation  
*(cosmology:)* distance scale  
*(cosmology:)* early Universe  
*(cosmology:)* inflation  
*(cosmology:)* large-scale structure of Universe  
cosmology: miscellaneous  
cosmology: observations  
*(cosmology:)* primordial nucleosynthesis  
cosmology: theory

### **Resolved and unresolved sources as a function of wavelength**

gamma-rays: diffuse background  
gamma-rays: galaxies  
gamma-rays: galaxies: clusters  
gamma-rays: general  
gamma-rays: ISM  
gamma-rays: stars  
infrared: diffuse background  
infrared: galaxies  
infrared: general  
infrared: ISM  
infrared: planetary systems  
infrared: stars  
radio continuum: galaxies  
radio continuum: general  
radio continuum: ISM  
radio continuum: planetary systems  
radio continuum: stars  
radio continuum: transients  
radio lines: galaxies  
radio lines: general  
radio lines: ISM  
radio lines: planetary systems  
radio lines: stars  
submillimetre: diffuse background  
submillimetre: galaxies  
submillimetre: general  
submillimetre: ISM  
submillimetre: planetary systems  
submillimetre: stars  
ultraviolet: galaxies

ultraviolet: general  
ultraviolet: ISM  
ultraviolet: planetary systems  
ultraviolet: stars  
X-rays: binaries  
X-rays: bursts  
X-rays: diffuse background  
X-rays: galaxies  
X-rays: galaxies: clusters  
X-rays: general  
**X-rays: individual: . . .**  
X-rays: ISM  
X-rays: stars