

Autism
The International Journal of Research and Practice

Current Profiles and Early Predictors of Reading Skills in School-Age Children with Autism Spectrum Disorders: A longitudinal, retrospective population study

Journal:	<i>Autism</i>
Manuscript ID	AUT-18-0103.R2
Manuscript Type:	Original Article
Keywords:	Autism spectrum disorders, Communication and language, School-age children, Literacy, Reading, Longitudinal
Abstract:	<p>This study explores current reading profiles and concurrent and early predictors of reading in children with ASD. Before the age of 3 years, the study cohort underwent a neurodevelopmental assessment following identification in a population-based autism screening. At age 8 years, reading, language and cognition were assessed. Approximately half of the sample (n=25) were "poor readers" at age 8 years, meaning that they scored below the normal range on tests of single word reading and reading comprehension. Eighteen were "skilled readers" performing above cut-offs. The final subgroup (n=10) presented with a "hyperlexic/poor comprehenders" profile of normal word reading, but poor reading comprehension. The "poor readers" scored low on all assessments, as well as showing more severe autistic behaviors than "skilled readers". Group differences between "skilled readers" and "hyperlexics/poor comprehenders" were more subtle: These subgroups did not differ on autistic severity, phonological processing or nonverbal IQ, but the "hyperlexics/poor comprehenders" scored significantly lower on tests of oral language. When data from age 3 were considered, no differences were seen between the subgroups in social skills, autistic severity or IQ. Importantly, however, it was possible to identify oral language weaknesses in those that five years later presented as "poor readers" or "hyperlexics".</p>

SCHOLARONE™
Manuscripts

Current Profiles and Early Predictors of Reading Skills in School-Age Children with Autism Spectrum Disorders: A longitudinal, retrospective population study

This study explores current reading profiles and concurrent and early predictors of reading in children with ASD. Before the age of 3 years, the study cohort underwent a neurodevelopmental assessment following identification in a population-based autism screening. At age 8 years, reading, language and cognition were assessed. Approximately half of the sample (n=25) were “poor readers” at age 8 years, meaning that they scored below the normal range on tests of single word reading and reading comprehension. Eighteen were “skilled readers” performing above cut-offs. The final subgroup (n=10) presented with a “hyperlexic/poor comprehenders” profile of normal word reading, but poor reading comprehension. The “poor readers” scored low on all assessments, as well as showing more severe autistic behaviors than “skilled readers”. Group differences between “skilled readers” and “hyperlexics/poor comprehenders” were more subtle: These subgroups did not differ on autistic severity, phonological processing or nonverbal IQ, but the “hyperlexics/poor comprehenders” scored significantly lower on tests of oral language. When data from age 3 were considered, no differences were seen between the subgroups in social skills, autistic severity or IQ. Importantly, however, it was possible to identify oral language weaknesses in those that five years later presented as “poor readers” or “hyperlexics”.

Introduction

Literacy skills are important for lifelong learning, employment and independence in our society and studies exploring the reading profiles and reading difficulties of children with

1
2
3 autism spectrum disorders (ASD) has grown in recent years. Much of this research has
4
5 focused on the mismatch between stronger single word reading/decoding and weaker reading
6
7 comprehension observed in some children with ASD (e.g., Huemer & Mann, 2010), with the
8
9 term “hyperlexia” sometimes being used to describe an extreme version of this profile (c.f.,
10
11 Aaron, 2012; Grigorenko, Klin, & Volkmar, 2003; Nation, 1999; Ostrolenk, d’Arc, Jelenic,
12
13 Samson, & Mottron, 2017). Other studies have, however, noted a very considerable
14
15 heterogeneity in reading capacity among children with ASD (Åsberg, Kopp, Berg-Kelly, &
16
17 Gillberg, 2010; Brown, Oram-Cardy, & Johnson, 2013; Nation, Clarke, Wright & Williams,
18
19 2006; Norbury & Nation, 2011; White et al, 2006).

20
21
22 Nation et al. (2006) utilized a subgrouping procedure to characterize this heterogeneity. In a
23
24 sample of 41 children (aged 6-15 years, M=10.33), 21 children scored either below the
25
26 normal range (i.e. standard score below 85), at floor levels or were non-readers in terms of
27
28 single word reading/decoding skills. Another subgroup (n = 10) were classified as skilled in
29
30 word reading/decoding and reading comprehension on the basis of their performance on age-
31
32 referenced tests scores. Finally, another ten children displayed the “hyperlexic”-profile
33
34 described above. Of particular interest to Nation et al. (2006) were the differences (and
35
36 similarities) between the two latter subgroups; results showed that participants with the
37
38 “hyperlexic/poor comprehender”-profile had difficulties not only with *reading* comprehension
39
40 but also with *language (listening)* comprehension more generally, relative to “skilled readers”
41
42 with ASD. In contrast, the “skilled readers” and the “hyperlexic” subgroups were not
43
44 differentiated by nonverbal cognitive ability. These cross-sectional results fit well with the
45
46 generally accepted conclusion in non-ASD reading research: reading comprehension builds
47
48 on a foundation of oral language (i.e. listening) comprehension (Hoover & Gough, 1990;
49
50 Hulme & Snowling, 2013). In particular, the so-called “Simple view of Reading” proposes
51
52 that reading comprehension is the product of decoding skills and oral language/listening
53
54
55
56
57
58
59
60

1
2
3 comprehension; deficits in oral language therefore place a primary constraint on reading
4
5 comprehension in fluent word readers (Hoover & Gough, 1990).
6
7
8
9

10 Previous research has also explored the extent to which word reading development in ASD is
11 coupled with phonological processing capacity, a lower level linguistic skill often impaired in
12 poor word readers without ASD (i.e. dyslexics) (Hulme & Snowling, 2013; White et al.,
13 2006). Conversely, phonological processing capacity is well developed in skilled word
14 readers, with a presumed causal influence on word reading development (Hulme, Bowyer-
15 Crane, Carroll, Duff, & Snowling, 2012), although the influence between phonology and
16 word reading is likely bidirectional (Nation & Hulme, 2011; Peterson et al., 2018). Nation et
17 al. (2006) did not explicitly assess phonological processing in their participants with ASD, but
18 suggested that the “hyperlexic” group may not present with phonological difficulties despite
19 being weak in non-phonological language domains (e.g., semantic processing). A dissociation
20 between phonological and non-phonological language skills may therefore explain why these
21 children were able to develop skilled word reading while failing to develop age appropriate
22 reading comprehension. This hypothesis is also in line with the two-dimensional model of
23 language and reading difficulties proposed by Bishop and Snowling (2004). This model
24 extends the Simple view of reading in the sense that both phonological and broader language
25 (listening) skills are put forward as differentially important factors in children’s literacy
26 development. Children with impaired phonological and listening comprehension capacity will
27 accordingly be challenged in both single word reading/decoding and reading comprehension,
28 whereas some children with language comprehension difficulties without phonological
29 difficulties will have relatively more selective deficits in reading comprehension. Studies have
30 empirically confirmed that phonology underpins word reading/decoding capacity in ASD as it
31 does in non-ASD individuals (Åsberg & Dahlgren Sandberg, 2012; Newman et al., 2007;
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 White et al., 2006) but it is less clear if this is the case when comprehension difficulties are
4
5 also evident (Bishop & Snowling, 2004).
6
7
8
9

10 A third research strand has addressed the association(s) between autistic symptom severity
11 and different reading skills. A few studies have reported an association between increased
12 autistic symptoms and reduced reading comprehension (Åsberg et al., 2010; McIntyre et al.,
13 2017; Ricketts, Jones, Happé, & Charman, 2012; Westerveld, Paynter, Trembath, Webster,
14 Hodge, & Roberts, 2017). Researchers have suggested that poor reading comprehension in
15 ASD might reflect underlying social-communicative difficulties that by definition are integral
16 to ASD. For instance, impaired coherence making and inferencing skills (e.g., Ricketts et al.,
17 2013) and/or difficulties in identifying author intentions and tracking the mental states of
18 characters in the texts (cf., Hulme & Snowling, 2013) could affect reading comprehension
19 beyond the role of oral language comprehension. However, another study showed that this
20 influence of autistic severity on reading comprehension attenuated when considered in
21 addition to the contributions of word reading/decoding and language capacity in a
22 multifactorial analysis (Lucas & Norbury, 2014). Yet another study only found an association
23 between increased social impairment and alphabetic knowledge (but not with reading
24 comprehension) (Davidson & Ellis Weismar, 2014). Hence, existing results are mixed to date,
25 and may reflect differences in samples in terms of age, sample size, and population
26 representativeness. Elucidating the extent to which reading comprehension difficulties in ASD
27 can be explained by word reading/decoding and/or language (listening) comprehension
28 difficulties, as stipulated by the Simple View (Hoover & Gough, 1990) is of both practical
29 and theoretical importance.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Drawing any causal conclusions based on the cross-sectional results should be approached
4 with caution. Longitudinal studies provide evidence of developmental primacy that can
5 inform causal theories. Davidson and Ellis Weismer (2014) assessed various cognitive,
6 linguistic and autism severity measures at mean age 2 ½ years in 101 children with ASD, all
7 of which were longitudinally related to early literacy skills when the children entered
8 kindergarten. In keeping with previous research, the sample tended to perform better on tasks
9 tapping alphabetic and early decoding skills than on comprehension and meaning. Moreover,
10 the results showed that the best longitudinal prediction model of reading comprehension (the
11 “Meaning subtest”) at age 5.5 years was multifactorial. Of these 2.5 year variables, nonverbal
12 cognition and expressive language stood out as the most significant contributors to later
13 reading comprehension. Similarly, Miller et al. (2017) demonstrated that language ability at
14 age 2 years predicted school-age reading comprehension in 26 children with ASD. Both of
15 these studies concluded that oral language skills provide a foundation for later reading
16 comprehension, although there appeared to be additional influences of other early predictors,
17 including nonverbal cognitive ability and autistic severity. Neither of these two longitudinal
18 studies applied a subgrouping procedure; instead, they predicted the full variation of reading
19 skills in collapsed convenience samples.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40 Although dimensional approaches are typically preferred because they increase statistical
41 power, it is more difficult to identify cases in which reading comprehension and word
42 reading/decoding capacity are decoupled. Consequently, the current paper considers early
43 precursors to qualitatively different reading profiles, including the hyperlexic/poor
44 comprehender profile (e.g., Nation et al., 2006).
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Furthermore, with the exception of the population-based sample from the SNAP study (Jones
4 et al., 2009; Ricketts et al., 2013), previous research on reading and ASD has utilized
5
6 convenience and/or clinical samples, making it hard interpret how common different reading
7
8 profiles are within the ASD population. Therefore, the current study employed a population-
9
10 based sample and a longitudinal, retrospective study design to examine concurrent reading
11
12 profiles and their early predictors based on assessment at the age of 3 years. Children had
13
14 been identified via population-based screening (Kantzer, Fernell, Gillberg, & Miniscalco,
15
16 2013; Kantzer, Fernell, Westerlund, Hagberg, Gillberg, & Miniscalco, 2018). At age 8 years,
17
18 we used a subgrouping procedure based on individual children's scores on standardized tests
19
20 of word reading and reading comprehension to determine the prevalence of different reading
21
22 profiles. The identified subgroups were then compared both concurrently and retrospectively
23
24 on a range of linguistic, cognitive and social skills/autistic severity data.
25
26
27
28

29 Three research questions were posed:
30
31

- 32 1. Which reading profiles can be identified within the sample?
33
- 34 2. To what extent are the reading profiles associated with concurrent measures of
35 language, phonological processing, nonverbal cognitive ability and autistic severity?
36
37
- 38 3. To what extent are the reading profiles associated with language, cognition,
39 communication and social functioning, and autistic severity measures taken at age 3
40
41 years?
42
43
44
45
46
47
48
49

50 **Methods**

51 *Participants*

52
53
54
55
56
57
58
59
60

1
2
3 More than 100 children in Gothenburg, Sweden, were identified as having a suspected autism
4 spectrum disorder (ASD) by a population screening at 2.5 years of age (Nygren, Sandberg, et
5 al., 2012), at child health care centers. According to Magnusson (1997), 97.5% of all Swedish
6 children participate in screening at these centers. The project was called *AUTism Detection and*
7 *Intervention in Early life* (AUDIE) and ran from 2009-2011. Of the children who screened
8 positive for ASD, parents of 107 children gave their consent to participate in the current
9 research project. These children were assessed in depth by a multi-disciplinary team at the
10 Child Neuropsychiatry Clinic (CNC) around 3 years of ages (Kantzer et al., 2013; Kantzer et
11 al., 2018) focusing on autism diagnostics/assessment, language ability, cognitive level, and
12 adaptive functioning. All involved professionals then met and made a consensus diagnosis
13 based on all available information i.e. test results, observation data, parental questionnaires
14 and interviews.

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29 The current follow-up assessment took place when the children were between 5.9 and 9.8
30 years old ($n=85$) (15 girls; 70 boys) and included measures of oral language, non-verbal
31 functioning, and literacy. Assessments were made by two speech and language pathologists
32 (SLP) during one or two sessions at the CNC. Inclusion criteria stipulated that children were
33 in the first or second grade of school (7-8 years of age); 27 children of the 85 were excluded
34 since they had either not started first grade ($n=24$) or entered third grade at the time for
35 assessment (see supplementary online material for further information). Another five
36 participants were excluded because reading assessments were not completed due to time
37 constraints and/or errors during administration. Thus, the total number of participating
38 children was 53 with a mean age of 8.0 (6.6 – 9.8) years old (8 girls; 45 boys).

39
40
41
42
43
44
45
46
47
48
49
50
51
52
53 Of these 53 children, there were five who did not meet full DSM-IV criteria for an ASD
54 diagnosis at their latest full autism assessment, but all were identified with autistic traits. We

1
2
3 chose to include all participants in the current study, due to 1) the unique recruitment
4
5 procedure and 2) that the eventual influence of “autistic severity” (measured both
6
7 dimensionally and categorically) on reading was one of the main issues we wished to explore.
8
9
10
11
12

13 *Tests and material*

14 **Data from the assessment at school age follow-up**

15 Oral language skills (comprehension and production)

16
17
18 Language comprehension was assessed using the Test for Reception of Grammar-2 (TROG-2)
19
20 (Bishop, 2003; Swedish version, 2009) and receptive vocabulary using the Peabody Picture
21
22 Vocabulary Test-, PPVT-III (Dunn & Dunn, 1997). TROG-2 involves matching orally
23
24 presented sentences to the correct picture of a choice of four. The results are presented as both
25
26 raw scores (number of correctly solved blocks out of a maximum of 20), and standard scores
27
28 (M = 100, SD = 15) based on Swedish norms. The Cronbach’s alpha is reported to be .89. On
29
30 the PPVT, the child listens to a word uttered by the assessor and then selects one of four
31
32 pictures that best describes the word's meaning. The test is not standardized for Swedish
33
34 children, and therefore the original American norms were used. The Cronbach’s alpha is
35
36 reported to be .95 in the manual.
37
38
39
40
41
42
43
44

45 We indexed language production (Klem, et al., 2015) using the “Recalling Sentences” subtest
46
47 from the Clinical Evaluation of Language Fundamentals – 4 (CELF 4; Semel, Wiig, &
48
49 Secord, 2003; Swedish version, 2013). “Recalling sentences” consists of 24 sentences which
50
51 are repeated verbatim by the child and scored on a 4-point scale depending on the number of
52
53 errors present in the child’s repetition. We present the results as both raw scores and scaled
54
55
56
57
58
59
60

1
2
3 scores (around a normative $M = 10$, $SD = 3$ based on Swedish norms). The Cronbach's alpha
4
5 is reported to be .89.
6
7
8
9

10 Phonological processing/non-word repetition

11
12 Non-word repetition was assessed with 30, one to five syllable non-words that conform to
13
14 Swedish phonotactics (Radeborg, Barthelom, Sjöberg, & Sahlén, 2006). The children
15
16 repeated the non-words after the SLP's oral presentation. Norms are available for children
17
18 aged 4-6 years. The SLP marked the responses online using broad phonetic transcription
19
20 according to the International Phonetic Alphabet (IPA, 2005). Each repeated non-word was
21
22 immediately scored as correct or incorrect. Radeborg et al. report a Cronbach's alpha of 0.74.
23
24
25
26
27
28

29 Letter knowledge

30
31 The child was asked to name the letters of the Swedish alphabet (which includes three letters
32
33 besides those represented in English), written on two sheets of paper in both uppercase and
34
35 lowercase form. The max score for each form is 24.
36
37

38 Single word reading/decoding

39
40
41 The LÄST test (Elwér, Fridolfsson, Samuelsson & Wiklund, 2009) was used to examine
42
43 participants' single word reading/decoding ability. The task is to read as many words as
44
45 possible in 45 seconds, from two lists of words. A total score is created by summarizing the
46
47 number of correctly read words. Such efficiency measures – rather than separate accuracy and
48
49 fluency measures – are typically used in Sweden and other semi-consistent orthographies.
50
51 Swedish norms are available in the manual based on the stanine scale (i.e., around a mean of
52
53
54
55
56
57
58
59
60

1
2
3 M = 5, SD = 2) for each grade year. The test-retest reliability is reported to be $r = .93$ for both
4
5 lists in the Swedish manual.
6
7
8
9

10 Reading comprehension

11
12 The DLS Bas test (Järpsten, 2004) was used in order to assess the child's reading
13
14 comprehension. The test comprises 20 sentences intertwined into a small story. For each
15
16 sentence, there are five pictures and the child should mark the picture that best can be linked
17
18 with the written content. The child is asked to read as many items as possible in 7 minutes (for
19
20 7 year-old children, i.e. school year one in Sweden) or in 5 minutes (for 8 year-old children,
21
22 i.e. school year two in Sweden), with a maximum possible score of 20. Swedish norms are
23
24 available in the manual based on stanine scores for each grade year. The test-retest reliability
25
26 is reported to be $r = .78$.
27
28
29
30

31 Non-verbal cognitive ability

32
33 The matrix reasoning subtest of Wechsler abbreviated scales of intelligence (WASI) was used
34
35 as a measure of nonverbal cognitive ability (WASI: Wechsler, 1999). Results are expressed in
36
37 raw and T-scores (M = 50, SD = 10) based on American norms. No Swedish norms are
38
39 available.
40
41
42
43
44
45

46 Autism symptomatology

47
48 The Autism Spectrum Screening Questionnaire (ASSQ, Ehlers, Gillberg, & Wing, 1999) uses
49
50 parent ratings on 27 items with a three-point Likert scale to measure autistic symptomatology.
51
52 Test-retest reliability is reported to be high ($r = .96$), and validity was established by Ehlers et
53
54 al. (1999) and by Posserud, Lundervold, and Gillberg (2009), showing a clear correspondence
55
56
57

1
2
3 between total score on the ASSQ and a clinical consensus diagnosis of ASD. A screening cut-
4
5 off for ASD of > 18 has been suggested (Ehlers et al., 1999).
6
7

8 9 **Data from the first assessment at age 3 years**

10
11 Not every child could participate in all tasks described below, therefore the n on each task is
12
13 presented in table 2.
14
15

16 17 Assessment of cognitive/developmental level

18
19 A psychologist assessed children's general cognitive and developmental level using the
20
21 Griffiths' developmental scales (GDS) (Alin-Åkerman & Norberg, 1991). The test includes
22
23 six subscales; the total score ($M = 100$, $SD = 15$) from the subscales provides a developmental
24
25 quotient (DQ) which is used here. McLean, McCormick and Baird (1991) reports adequate
26
27 psychometric properties, both in terms of internal consistency reliability (Cronbach's $\alpha >$
28
29 $.96$) and construct validity according to correlation patterns with other tests of cognitive
30
31 functioning.
32
33
34
35
36

37 38 Autism symptomatology

39
40 The Autism Diagnostic Observation Schedule – Generic (ADOS) (Lord, Risi, Lambrecht,
41
42 Cook, Leventhal, & DiLavore, 2000) is a standardized, semi structured play-based assessment
43
44 of communication, reciprocal social interaction, play, and behaviour. Either module 1 or 2
45
46 was administered, based on the expressive language level of the child. From these data
47
48 calibrated severity scores were calculated (scores from 1-10) (Hus, Gotham & Lord, 2014).
49
50 Higher scores indicate increased autistic symptom severity.
51
52
53

54 55 Adaptive communicative and social functioning

1
2
3 The Communication and Socialization domains of the Vineland Adaptive Behaviour Scales
4 (VABS) (Sparrow, Cicchetti, & Balla, 2005) was administered by a child neuropsychologist
5
6 in a face-to-face interview with one or both parents. Results are expressed in standard scores
7
8 around a normative $M = 100$, $SD = 15$.
9
10

11 12 13 14 Oral comprehension and production language skills 15

16
17 Language comprehension was assessed with the Reynell Developmental Language Scales III
18 (RDSL) (Edwards, Fletcher, Garman, Hughes, & Letts, 1997) which has Swedish norms
19 (Eriksson & Grundström 2000; Lindström & Åström, 2000). The Kuder-Richardsson
20
21 reliability coefficient is reported to be .97 in the manual from Great Britain. In addition, the
22
23 expressive language level of each child was rated by the SLP on a scale from 1-to-5 using the
24
25 PARIS scale (Philippe, Martinez, Guilloud-Bataille, Gillberg, Råstam et al., 1999): 1 = no
26
27 words at all; 2 = a few single words; 3 = a few communicative sentences; 4 = talks a great
28
29 deal, mostly echolalia, or 5 = talks a great deal, mostly in a communicative fashion.
30
31
32
33
34
35

36 ***Statistical analyses*** 37

38 The variables were subject to normality checks. First, the reading scores (single word
39
40 reading/decoding, and reading comprehension) analysed dimensionally in the full group were
41
42 found to diverge from the normal distribution (with many scores at floor); for this reason, we
43
44 cannot complement the following results based on subgrouping with dimensional analyses
45
46 across the full range of abilities. The other variables that were compared across subgroups
47
48 displayed kurtosis and skewness statistics indicative of approximately normally distribution
49
50 (values < 1.2), with the possible exception of the PARIS scale. A histogram inspection
51
52 revealed that this variable seemed to be bimodal, with few scores of 3. Therefore, we chose to
53
54
55
56
57
58
59
60

1
2
3 transform the PARIS scale into a dichotomous variable, with the scores 1 and 2 representing
4
5 less/minimally verbal, and 3, 4 and 5 representing relatively more verbal.
6
7
8

9 10 **Results**

11 12 *Attrition from intake at age 3 years to the current school-age follow-up*

13
14
15 In order to evaluate the population representativeness of the sample, we compared those who
16
17 participated in the reading assessments at school-age follow-up with the rest of the screen
18
19 positive cohort on the assessments of interest from intake. Those who did not participate were
20
21 on average three months younger when they were assessed for the first time ($p = .023$). On all
22
23 other assessments – for the Griffiths’ developmental scales (DQ total score), the RDLS test of
24
25 language comprehension, the ADOS (severity total score), the Vineland socialization, and the
26
27 Vineland communication scores – no significant differences were observed (all $p > .20$) (see
28
29 supplementary online material for details).
30
31
32
33
34
35

36 *RQ1: Which reading profiles can be identified within the sample?*

37
38 Subgrouping were performed based on a cut off of a stanine score of ≤ 2 on standardized
39
40 assessments on single word reading/decoding and reading comprehension (which corresponds
41
42 to the $\sim 10^{\text{th}}$ percentile); this is a common cut off in both research and in Swedish schools to
43
44 identify children with reading difficulties. Results showed that almost half of the sample
45
46 (25/53) were classified into a subgroup that will henceforward be called “poor readers”,
47
48 meaning that they scored below cut off on both single word reading/decoding ($M_{\text{stanine}} = 1.0$;
49
50 $SD = 0.2$) and reading comprehension ($M_{\text{stanine}} = 1.0$, $SD = 0.2$). Another 10 participants were
51
52 assigned in the “hyperlexics/poor comprehenders” group, meaning that they scored above cut
53
54
55
56
57
58
59
60

1
2
3 off in word reading/decoding ($M_{\text{stanine}} = 5.9, SD = 1.0$) but below cut off in reading
4
5 comprehension ($M_{\text{stanine}} = 1.7, SD = 0.5$). For all “hyperlexics/poor comprehenders” there was a
6
7 substantial discrepancy in the word reading versus reading comprehension performance of at
8
9 least 3 stanine scores. Finally, 18/53 was assigned as belonging to a “skilled readers”
10
11 subgroup since they performed above cut off on both word reading/decoding ($M_{\text{stanine}} = 7.1,$
12
13 $SD = 1.3$) and reading comprehension ($M_{\text{stanine}} = 5.8, SD = 1.6$). Note that none showed the
14
15 profile of poor word decoding and relatively better reading comprehension (that sometimes is
16
17 seen in higher-functioning samples of dyslexic readers); hence, only three subgroups of
18
19 interest were identified here.
20
21
22
23
24
25

26 The letter knowledge task was not used for subgroup assignment, but demonstrated that the
27
28 majority of children in the “poor readers” groups were preliterate. As a group, they were only
29
30 able to identify half of the letters in the alphabet, with 11 children not being able to recognize
31
32 any letters at all (see table 1 descriptive data).
33
34
35
36
37

38 ***RQ2: Are the reading profiles associated with concurrent measures of language,***
39 ***phonological processing, nonverbal cognitive ability and autistic severity?***
40
41
42

43 Table 1 present the descriptive data for each of the three reading groups on all measures of
44
45 interest from the school-age follow-up assessment. Since the groups differed on age ($F [2, 50]$
46
47 $= 8.36, p = .001, \eta_p^2 = .251$) with post hoc showing that the “skilled readers” subgroup was
48
49 somewhat older ($p < .01$) than the other two subgroups (who in turn did not differ from one
50
51 another, $p = .616$), we use ANCOVA with control for chronological age in all subsequent
52
53 group comparisons. Raw-scores rather than age-standardized scores were used in these
54
55
56
57
58
59
60

1
2
3 analyses in order not to control for age at multiple times; however, for descriptive purposes
4
5 age-standardized scores values are also reported in Table 1.
6
7

8 >>Table 1. Insert about here<<
9

10 In terms of oral language, there were significant group differences on TROG-2 ($F [2, 49] =$
11 $17.95, p < .001, \eta_p^2 = .423$), PPVT III ($F [2, 49] = 9.24, p < .001, \eta_p^2 = .274$.) and Recalling
12 Sentences (repetition) results ($F [2, 49] = 13.87, p < .001, \eta_p^2 = .361$). Post hoc comparisons
13 on TROG-2 showed that the “skilled readers” performed significantly better than both the
14 “hyperlexics/poor comprehenders” ($p < .01$) and the “poor readers” ($p < .001$). In turn,
15 “hyperlexics/poor comprehenders” performed better than the “poor readers” ($p = .032$). A
16 slightly different pattern was obtained on PPVT III and Recalling Sentences tasks, in which
17 “skilled readers” performed significantly better than both the “hyperlexics/poor
18 comprehenders” (both $p \leq .01$) and the “poor readers” (both $p < .001$), who in turn did not
19 differ from one another on either measure ($p = .201$ and $p = .786$, respectively). Mean scores
20 in the different groups revealed that the “skilled readers” scored within the normal range on
21 the standardized measures, whereas the “poor readers” and the “hyperlexics/poor
22 comprehenders” scored substantially below the normal range.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

39 Turning to phonological processing, a significant group difference was found on the non-word
40 repetition task ($F [2, 48] = 18.85, p < .001, \eta_p^2 = .440$). Post hoc tests showed that the “skilled
41 readers” and the “hyperlexics/poor comprehenders” did not differ from one another ($p = .786$)
42 while both these groups outperformed the “poor readers” (both $p < .001$). There are no norms
43 available for the current age group, but a comparison with norms for four-to-six year-olds
44 shows that the “poor readers” perform below the 25th percentile for the four-year olds. By
45 contrast, the “hyperlexic/poor comprehenders” and the “skilled readers” scored equivalent to
46 the 95th percentile for six-year olds.
47
48
49
50
51
52
53
54
55
56

1
2
3
4
5 Comparing the groups on the Matrix reasoning subtest of nonverbal cognitive ability revealed
6 a significant group difference ($F [2, 48] = 12.00, p < .001, \eta_p^2 = .333$), with post hoc analyses
7 showing no significant difference between the “skilled readers” and the “hyperlexics/poor
8 comprehenders” ($p = .604$), while the “poor readers” scored lower than both the “skilled
9 readers” ($p < .001$) and the “hyperlexics/poor comprehenders” ($p < .01$). Mean scores in the
10 different groups revealed that “skilled readers” and “hyperlexics/poor comprehenders” scored
11 within the normal range, whereas the “poor readers” approx. scored 1.5 standard deviations
12 below the normative mean.
13
14
15
16
17
18
19
20
21
22
23
24
25

26 With regard to autistic severity, the groups differed on ASSQ total scores ($F [2, 47] = 7.49, p$
27 $< .01, \eta_p^2 = .242$), with post hoc comparisons revealing that “poor readers” had higher
28 symptom severity scores than either of the other groups (both $p < .01$), who did not differ
29 from one another ($p = .807$). Although the mean scores on the ASSQ fell below the suggested
30 screening cut-off for ASD in the “hyperlexics/poor comprehenders” ($\bar{x} = 14.6$) and the
31 “skilled readers” ($\bar{x} = 16.9$), it should be stressed that these scores were well over 2 standard
32 deviations above the population mean for 7-9 year olds as described by Posserud, Lundervold
33 and Gillberg (2006). We also analyzed whether the prevalence of a clinical autism spectrum
34 diagnosis differed between the subgroups, with the results revealing a possible trend ($\chi^2 =$
35 $5.01, p = .082$). The five participants who did not receive an ASD diagnosis were all classified
36 as “skilled readers” ($n = 3$) or as “hyperlexics/poor comprehenders” ($n = 2$).
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RQ3: Are the reading profiles associated retrospectively with language, cognition, communication and social functioning, and autistic severity at age 3 years?

Descriptive data for reading subgroups based on means and standard deviations taken when the children were aged 3 are reported in table 2. Chronological age at the 3-years assessment did not differ across the subgroups ($F [2, 50] = 2.03, p = .143, \eta_p^2 = .075$).

>>Table 2. Insert about here<<

Groups did not differ on cognitive ability assessed with Griffiths' developmental scales ($F [2, 43] = .097, p = .908, \eta_p^2 = .005$), degree of autistic severity (as assessed with the ADOS-2 severity scores ($F [2, 49] = 2.22, p = .119, \eta_p^2 = .083$) nor on the Vineland socialization scores ($F [2, 47] = 1.15, p = .324, \eta_p^2 = .047$)). Mean scores indicated equal levels of impairment in all groups.

As predicted, however, when comparing language ability at intake, significant differences were evident on RDLS language comprehension ($F [2, 49] = 14.47, p < .001, \eta_p^2 = .371$), the Vineland communication scale ($F [2, 47] = 3.36, p < .05, \eta_p^2 = .125$), and the SLP rating of expressive language using the PARIS scale ($\chi^2 = 13.52, p = .001$). Post hoc (Tukey) tests showed that the "poor readers" and the "hyperlexics/poor comprehenders" did not differ on the RDLS or Vineland communication ($p = .520$ and $p = 1.00$, respectively). By contrast, the "skilled readers" outperformed the "poor readers" on both measures (all $p < .05$). In terms of PARIS ratings, inspection of adjusted standardized residuals showed that "skilled readers" were more likely to be verbal as toddlers relative to children with poor reading skills who were less/minimally verbal. The skilled readers also outperformed the "hyperlexics/poor comprehenders" on the RDLS comprehension test ($p < .01$), whereas scores from the Vineland communication subscale was not significantly different ($p = .139$).

Discussion

This study aimed to address the heterogeneity present among children with ASD in their early reading skills. Specifically, we investigated individual differences in language comprehension and production, phonological deficits, cognitive level and autistic symptom severity in subgroups identified by different profiles of word reading/decoding skills and reading comprehension. The study allowed us to identify factors related to reading ability profiles both concurrently (mean age 8 years) and retrospectively (around the age of 3). In addition, our participants were identified by screening and are thus population representative to a greater extent than is the case in previous research, reducing ascertainment bias in establishing reading profiles.

The most common reading profile in our sample was that of generally poor reading skills. By contrast, the “hyperlexic/poor comprehenders” profile was relatively less common, whereas a sizable minority (ca 20%) performed age-adequate on both single word reading/decoding and reading comprehension. Hence, our results replicate in a Swedish population sample the great heterogeneity of reading skills and profiles within ASDs (e.g., Åsberg et al., 2010; Jones et al., 2009; Nation et al., 2006).

Our study also confirms the close relationship between reading comprehension and oral language comprehension skills. The “poor readers” subgroup was found to score low on most assessment, including nonverbal cognitive ability, as well as showing a more severe autistic presentation. Hence, the poor reading and language abilities of this group occur in the context

1
2
3 of more general delay. This means that their reading comprehension and oral language
4
5 weaknesses were accompanied by greater impairments across several developmental domains,
6
7 making it difficult to propose a distinct relation between reading comprehension and language
8
9 comprehension deficits in ASD if only this subgroup is considered. By contrast, in the
10
11 “hyperlexics/poor comprehenders” the difficulty in oral language and reading comprehension
12
13 was much more selective when compared to the “skilled readers”, pointing more clearly to a
14
15 distinct association (cf., Nation et al., 2006). This finding is in keeping with the “Simple View
16
17 of Reading” (Hoover & Gough, 1990).
18

19
20
21
22 Another important feature of this study was the longitudinal (retrospective) study design.
23
24 Only two previous studies (Davidson and Ellis Weismer, 2014; Miller et al., 2017) have
25
26 included analyses of developmental precursors to reading ability in ASD. However, neither of
27
28 these distinguished between different reading profiles. Hence, it has not been fully clear to
29
30 what extent identified predictors are important for reading comprehension per se, as opposed
31
32 to aspects of reading comprehension difficulties accompanied by single word
33
34 reading/decoding difficulties. Nonetheless, our results are broadly in line with these previous
35
36 studies, demonstrating that before the age of 3 it may be possible to identify oral language
37
38 weaknesses in both of the subgroups that presented five years later with poor reading
39
40 comprehension. This result suggests that supporting oral language skills in ASD could be one
41
42 way to support reading comprehension in children with ASD, similar to intervention
43
44 approaches targeting poor reading comprehenders without ASD (e.g., Clarke, Snowling
45
46 Truelove, & Hulme, 2010). Interestingly, preliminary intervention studies including children
47
48 with ASD lends initial support for this idea (Bailey, Arciuli, & Stancliffe, 2017).
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Consistent with Nation et al. (2006), our results demonstrate that children with the
4
5 “hyperlexic/poor comprehender” profile had age appropriate phonological processing (as
6
7 measured by the non-word repetition task), even though they performed poorly on tests of oral
8
9 language comprehension. Our results are thus in line with the suggestion by Bishop and
10
11 Snowling in their two-dimensional model of language and reading difficulties (2004), stating
12
13 that the dissociation between (intact) phonological and (poor) non-phonological (semantic and
14
15 syntactic) language skills might explain why some children are able to develop skilled word
16
17 reading/decoding but poor reading comprehension. However, we need to highlight that in this
18
19 study we were only able to measure this association cross-sectional. Indeed, whereas
20
21 phonology traditionally has been interpreted as a “predictor” of word reading development,
22
23 recent work (e.g., Nation & Hulme, 2011; Peterson et al., in press) has shown that phonology,
24
25 including non-word repetition and related skills, may be causally dependent on word reading
26
27 level, not only vice versa. It is fully possible that this is the case also in readers with ASD, and
28
29 hence the causal mechanism in the phonology-word reading link observed here needs to be
30
31 specified in future research. Nevertheless, disentangling the complex relation between letter
32
33 knowledge, phonological skills and single word reading might not be critical for being able to
34
35 provide effective interventions for poor word readers with ASD, at least if intervention
36
37 approaches used in dyslexia can be transferred to poor word readers with ASD. Indeed,
38
39 phonics-based training – where the reading intervention includes both letter-sound and
40
41 phonological knowledge training – has been shown to be effective, whereas a pure
42
43 phonological approach has gained less clear effects (SBU, 2014). Hence, phonology and word
44
45 reading seems developmentally intertwined, and this fact also affects efficient
46
47 teaching/intervention practices.
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 The association between ASD severity and reading profile varied depending on the time point
4 considered. In early school-age, we found evidence of more severe autistic symptoms in the
5 “poor readers” subgroup, which we interpret as an additional sign that this group has more
6 pervasive developmental challenges. However, at age 3 years these differences in autistic
7 severity were not obvious in the ADOS, nor in parent report of social skills. These results
8 suggest that it is probably hard or impossible to predict early reading skills from autistic
9 behaviours/social difficulties observed in toddlers alone.
10
11

12
13
14
15
16
17
18 A related issue is to what extent autistic severity impacts reduced reading comprehension.
19
20 Several authors (e.g., Åsberg Johnels, Gillberg, & Kopp, 2017; Jones et al., 2009; Ricketts et
21 al., 2013; Westerveld, et al., 2017) have suggested that impairments in social-communicative
22 functioning and flexibility, on the one hand, and reading comprehension, on the other, may be
23 coupled, possibly independently of basic language and cognitive skills. For instance, failing to
24 understand social and communicative norms may hamper a reader’s ability to make
25 inferences and, consequently, constrain the processing of the text content (e.g., Ricketts et al,
26 2013). The results presented here do not easily support this idea, since we could not find any
27 difference in autistic severity between the skilled and the “hyperlexic/poor comprehenders”
28 subgroups. This might mean that we need to reconsider the hypothesis that ASD
29 characteristics “per se” influence text comprehension negatively. In terms of theoretical
30 implications, the results thus suggest that the Simple View of Reading can be applied to
31 explain the reading comprehension difficulties in this cohort of readers with ASD. Still, we
32 believe that the age of the study sample needs to be considered more in future research. Our
33 participants attended first or second grade; during these first years in school, the texts and
34 tasks used in reading comprehension assessment are typically quite basic, and this was
35 arguably the case in the current study. One possibility is that with increasing age, the child
36 meets more complex texts and assessment procedures, which place greater demands on the
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 flexible usage of both language and social knowledge and strategies, such as making
4 coherence inferences and interpretations of author intentions (cf., Norbury & Nation, 2011).
5 Hence, we propose that future research should consider whether the association between
6
7 reduced reading comprehension and autistic severity is modulated by age, as well as the text
8
9 type/context that is used in assessment.
10
11
12

13 14 ***Limitations and conclusions of the current study*** 15

16
17 There are a number of weaknesses of the current study. First, although the sample size is
18 fairly large for a population-based study of autism, it is nonetheless clearly the case that the
19 subgroups are small. Hence, the results should be further corroborated in larger samples.
20
21 Another weakness is that no assessment of phonology was done at age 3 years. That would
22 have allowed us to assess the predictive value of early phonological skills on single word
23 reading/decoding. Finally, the current study only focused on reading whereas writing skills
24 were not assessed. It has been argued that spelling performance and analyses of error patterns
25 provide important insight into the mechanisms of literacy acquisition (Perfetti & Hart, 2002).
26
27 Despite these caveats, the results of this unique population-based study of children with ASD
28 confirm a high degree of heterogeneity in reading skills in ASD. Given the importance of
29 literacy for lifelong learning, employment and independence, it is important that practitioners
30 in the field of ASD are aware of these various profiles of strengths and difficulties. Results
31 also show that the profiles appeared to be predictable from and align well with established
32 findings in general reading research: that word reading is strongly associated with
33 phonological skills, whereas reading comprehension builds on a foundation of oral language
34 skills. By age 3 years it appears to be possible to identify oral language weaknesses in
35 children with ASD that five years later present as “poor readers” or “hyperlexics/poor
36 comprehenders”.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56

Acknowledgement

The study was performed with support from the Swedish Research Council for Health, Working Life and Welfare (project no 2013-00092) and Queen Silvia's Jubilee Foundation. We thank SLP Anna-Clara Reinholdson for assistance during data collection.

References

- Aaron, P. G. (2012). *Dyslexia and hyperlexia: Diagnosis and management of developmental reading disabilities* (Vol. 1). Springer Science & Business Media.
- Alin-Åkerman, B., & Norberg, L. (1991). *Griffiths' Development Scales I and II*. Stockholm: Psykologiförlaget AB. (Swedish Version).
- Åsberg, J., Kopp, S., Berg-Kelly, K., & Gillberg, C. (2010). Reading comprehension, word decoding and spelling in girls with autism spectrum disorders (ASD) or attention-deficit/hyperactivity disorder (AD/HD): Performance and predictors. *International Journal of Language & Communication Disorders, 45*(1), 61-71.
- Åsberg, J., & Dahlgren Sandberg, A. (2012). Dyslexic, delayed, precocious or just normal? Word reading skills of children with autism spectrum disorders. *Journal of Research in Reading, 35*(1), 20-31.
- Åsberg Johnels, J., Gillberg, C., & Kopp, S. (2017). A Hyperlexic-Like Reading Style Is Associated With Increased Autistic Features in Girls With ADHD. *Journal of Attention Disorders, 1087054716685838*.
- Bailey, B., Arciuli, J., & Stancliffe, R. J. (2017). Effects of ABRACADABRA literacy instruction on children with autism spectrum disorder. *Journal of Educational Psychology, 109*(2), 257.
- Baird, G., Simonoff, E., Pickles, A., Chandler, S., Loucas, T., Meldrum, D., & Charman, T. (2006). Prevalence of disorders of the autism spectrum in a population cohort of children in South Thames: the Special Needs and Autism Project (SNAP). *The lancet, 368*(9531), 210-215.
- Bishop, D.V.M. (2003). *Test of Reception of Grammar-2* (T.R.O.G, Swedish). Pearson: Stockholm.
- Bishop, D. V., & Snowling, M. J. (2004). Developmental dyslexia and specific language impairment: Same or different?. *Psychological bulletin, 130*(6), 858.
- Brown, H. M., Oram-Cardy, J., & Johnson, A. (2013). A meta-analysis of the reading comprehension skills of individuals on the autism spectrum. *Journal of autism and developmental disorders, 43*(4), 932-955.
- Clarke, P. J., Snowling, M. J., Truelove, E., & Hulme, C. (2010). Ameliorating children's reading-comprehension difficulties: A randomized controlled trial. *Psychological Science, 21*(8), 1106-1116.

- 1
2
3 Davidson, M. M., & Weismer, S. E. (2014). Characterization and prediction of early reading
4 abilities in children on the autism spectrum. *Journal of autism and developmental*
5 *disorders*, 44(4), 828-845.
- 6
7 Dunn, L. M., & Dunn, L. M. (1997). *PPVT-III: Peabody picture vocabulary test*. American
8 Guidance Service.
- 9
10 Edwards, S., Fletcher, P., Garman, M., Hughes, A., Letts, C., & Sinka, I. (1997). *Reynell*
11 *developmental language scales: Manual*. Berkshire, UK: The NferNelson Publishing
12 Company Limited.
- 13
14 Ehlers, S., Gillberg, C., & Wing, L. (1999). A screening questionnaire for Asperger syndrome
15 and other high-functioning autism spectrum disorders in school age children. *Journal of*
16 *autism and developmental disorders*, 29(2), 129-141.
- 17
18 Elwér, Å., Fridolfsson, I., Samuelsson, S., & Wiklund, C. (2009). *LäSt*. Stockholm: Hogrefe
19 Psykologiförlaget.
- 20
21 Eriksson, L. & Grundström, P. (2000). *Reynell Developmental Language Scales III*
22 *språkförståelsedelen. Översättning och normering, samt studie över samband mellan*
23 *testresultat och föräldrars utbildningsnivå*. Examensarbete i Logopedi, Umeå Universitet.
- 24
25 Grigorenko, E. L., Klin, A., & Volkmar, F. (2003). Annotation: Hyperlexia: disability or
26 superability?. *Journal of Child Psychology and Psychiatry*, 44(8), 1079-1091.
- 27
28 Hoover, W. A., & Gough, P. B. (1990). The simple view of reading. *Reading and Writing: An*
29 *Interdisciplinary Journal*, 2, 127-160.
- 30
31 Huemer, S. V., & Mann, V. (2010). A comprehensive profile of decoding and comprehension
32 in autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 40(4), 485-
33 493.
- 34
35 Hulme, C., Bowyer-Crane, C., Carroll, J. M., Duff, F. J., & Snowling, M. J. (2012). The
36 causal role of phoneme awareness and letter-sound knowledge in learning to read:
37 Combining intervention studies with mediation analyses. *Psychological Science*, 23(6),
38 572-577.
- 39
40 Hulme, C., & Snowling, M. J. (2013). *Developmental disorders of language learning and*
41 *cognition*. John Wiley & Sons.
- 42
43 Hus, V., Gotham, K., & Lord, C. (2014). Standardizing ADOS domain scores: Separating
44 severity of social affect and restricted and repetitive behaviours. *Journal of Autism and*
45 *Developmental Disorders*, 44(10), 2400-2412.
- 46
47 Jones, C. R., Happé, F., Golden, H., Marsden, A. J., Tregay, J., Simonoff, E., ... & Charman,
48 T. (2009). Reading and arithmetic in adolescents with autism spectrum disorders: peaks
49 and dips in attainment. *Neuropsychology*, 23(6), 718.
- 50
51 Järpsten, B. (2004). *DLS bas för skolår 1 och 2 [DLS Bas for Grades 1 and 2]*. Stockholm:
52 Hogrefe Psykologiförlaget AB.
- 53
54 Kantzer, A. K., Fernell, E., Gillberg, C., & Miniscalco, C. (2013). Autism in community pre-
55 schoolers: developmental profiles. *Research in developmental disabilities*, 34(9), 2900-
56 2908.

- 1
2
3 Kantzer, A. K., Fernell, E., Westerlund, J., Hagberg, B., Gillberg, C., & Miniscalco, C.
4 (2018). Young children who screen positive for autism: Stability, change and
5 “comorbidity” over two years. *Research in Developmental Disabilities*.
- 6
7 Klem, M., Melby-Lervåg, M., Hagtvet, B., Lyster, S. A. H., Gustafsson, J. E., & Hulme, C.
8 (2015). Sentence repetition is a measure of children's language skills rather than working
9 memory limitations. *Developmental Science*, *18*(1), 146-154.
- 10
11 Lindström, M. & Åström, A-L. (2000). *Reynell Developmental Language Scales III.*
12 *Normering och studie över samband mellan barns testresultat och kön, respektive typ av*
13 *barnomsorg*. Examensarbete i Logopedi, Umeå Universitet.
- 14
15 Lord, C., Risi, S., Lambrecht, L., Cook, E. H., Leventhal, B. L., DiLavore, P. C., ... & Rutter,
16 M. (2000). The Autism Diagnostic Observation Schedule—Generic: A standard measure
17 of social and communication deficits associated with the spectrum of autism. *Journal of*
18 *autism and developmental disorders*, *30*(3), 205-223.
- 19
20 Lucas, R., & Norbury, C. F. (2014). Levels of text comprehension in children with autism
21 spectrum disorders (ASD): The influence of language phenotype. *Journal of autism and*
22 *developmental disorders*, *44*(11), 2756-2768.
- 23
24 Magnusson, M. (1997). Rationality of routine health examinations by physicians of 18-
25 month-old children: Experiences based on data from a Swedish county. *Acta Paediatrica*,
26 *86*, 881–887.
- 27
28 McIntyre, N. S., Solari, E. J., Gonzales, J. E., Solomon, M., Lerro, L. E., Novotny, S., ... &
29 Mundy, P. C. (2017). The Scope and Nature of Reading Comprehension Impairments in
30 School-Aged Children with Higher-Functioning Autism Spectrum Disorder. *Journal of*
31 *Autism and Developmental Disorders*, 1-23.
- 32
33 McLean, M. E., McCormick, K., & Baird, S. M. (1991). Concurrent validity of the Griffiths'
34 Mental Development Scales with a population of children under 24 months. *Journal of*
35 *Early Intervention*, *15*(4), 338-344.
- 36
37 Miller, L. E., Burke, J. D., Troyb, E., Knoch, K., Herlihy, L. E., & Fein, D. A. (2017).
38 Preschool predictors of school-age academic achievement in autism spectrum disorder. *The*
39 *Clinical Neuropsychologist*, *31*(2), 382-403.
- 40
41 Newman, T. M., Macomber, D., Naples, A. J., Babitz, T., Volkmar, F., & Grigorenko, E. L.
42 (2007). Hyperlexia in children with autism spectrum disorders. *Journal of autism and*
43 *developmental disorders*, *37*(4), 760-774.
- 44
45 Nation, K., & Hulme, C. (2011). Learning to read changes children’s phonological skills:
46 Evidence from a latent variable longitudinal study of reading and nonword repetition.
47 *Developmental science*, *14*(4), 649-659.
- 48
49 Nation, K., Clarke, P., Wright, B., & Williams, C. (2006). Patterns of reading ability in
50 children with autism spectrum disorder. *Journal of autism and developmental disorders*,
51 *36*(7), 911.
- 52
53 Norbury, C., & Nation, K. (2011). Understanding variability in reading comprehension in
54 adolescents with autism spectrum disorders: Interactions with language status and
55 decoding skill. *Scientific Studies of Reading*, *15*(3), 191-210.
- 56
57 Nygren, G., Sandberg, E., Gillstedt, F., Ekeröth, G., Arvidsson, T., & Gillberg, C. (2012). A
58 new screening programme for autism in a general population of Swedish
59 toddlers. *Research in developmental disabilities*, *33*(4), 1200-1210.
- 60

- 1
2
3 Ostrolenk, A., d'Arc, B. F., Jelenic, P., Samson, F., & Mottron, L. (2017). Hyperlexia:
4 Systematic review, neurocognitive modelling, and outcome. *Neuroscience &*
5 *Biobehavioral Reviews*.
- 6
7 Perfetti, C. A., & Hart, L. (2002). The lexical quality hypothesis. In L. Vehoeven, C. Elbro, &
8 P. Reitsma (Eds.), *Precursors of functional literacy* (pp. 189–213). Amsterdam,
9 Netherlands: John Benjamins.
- 10
11 Peterson, R. L., Arnett, A. B., Pennington, B. F., Byrne, B., Samuelsson, S., & Olson, R. K.
12 (2018). Literacy acquisition influences children's rapid automatized naming.
13 *Developmental Science*, *21*(3), e12589.
- 14
15 Philippe, A., Martinez, M., Guilloud-Bataille, M., Gillberg, C., Råstam, M., Sponheim, E., ...
16 & Penet, C. (1999). Genome-wide scan for autism susceptibility genes. *Human molecular*
17 *genetics*, *8*(5), 805-812.
- 18
19 Posserud, M. B., Lundervold, A. J., & Gillberg, C. (2006). Autistic features in a total
20 population of 7–9-year-old children assessed by the ASSQ (Autism Spectrum Screening
21 Questionnaire). *Journal of Child Psychology and Psychiatry*, *47*(2), 167-175.
- 22
23 Radeborg, K., Barthelom, E., Sjöberg, M., & Sahlén, B. (2006). A Swedish non-word
24 repetition test for preschool children. *Scandinavian Journal of Psychology*, *47*(3), 187-192.
- 25
26 Ricketts, J., Jones, C. R., Happé, F., & Charman, T. (2013). Reading comprehension in autism
27 spectrum disorders: The role of oral language and social functioning. *Journal of Autism*
28 *and Developmental Disorders*, *43*(4), 807-816.
- 29
30 Semel, E., Wiig, E., & Secord, W. (2003). *Clinical evaluation of language fundamentals-IV*.
31 Marickville: Harcourt Assessment. Swedish version, 2013.
- 32
33 SBU (Statens beredning för medicinsk och social utvärdering). (2014). *Dyslexi hos barn och*
34 *ungdomar – tester och insatser. En systematisk litteraturöversikt*. Stockholm: SBU-rapport
35 nr 225.
- 36
37 Sparrow, S. S., Balla, D. A., & Cicchetti, D. V. (2005). *Vineland II: Vineland adaptive*
38 *behavior scales*. American Guidance Service.
- 39
40 Wechsler, D. (1999). Manual for the Wechsler abbreviated intelligence scale (WASI). *San*
41 *Antonio, TX: The Psychological Corporation*.
- 42
43 Westerveld, M. F., Paynter, J., Trembath, D., Webster, A. A., Hodge, A. M., & Roberts, J.
44 (2017). The Emergent Literacy Skills of Preschool Children with Autism Spectrum
45 Disorder. *Journal of autism and developmental disorders*, *47*(2), 424-438.
- 46
47 White, S., Frith, U., Milne, E., Rosen, S., Swettenham, J., & Ramus, F. (2006). A double
48 dissociation between sensorimotor impairments and reading disability: A comparison of
49 autistic and dyslexic children. *Cognitive Neuropsychology*, *23*(5), 748-761.
- 50
51
52
53
54
55
56
57
58
59
60

Table 1. Outcome measures for the three groups: “poor readers”, “hyperlexic/poor comprehenders”, and “skilled readers” at the school-year assessment

	Mean (SD) [min-max]			
	Poor Readers <i>n</i> = 25	Hyperlexic/ Poor Comprehenders <i>n</i> = 10	Skilled readers <i>n</i> = 18	Group differences using ANCOVA with correction for age
Age (years)	7.8 (0.7)	7.6 (0.4)	8.4 (0.5)	Poor Readers = Hyperlexic < Skilled Readers
Language ability				
PPVT III (raw score)	51.9 (38.9) [0-126]	70.1 (12.7) [41-91]	103.7 (39.5) [0-172]	Poor Readers = Hyperlexic < Skilled Readers
PPVT III (ss) ¹	63.5 (26.8) [40-125]	74.1 (9.0) [59-91]	94.4 (26.8) [40-150]	
TROG-2 (block score)	3.9 (4.7) [0-15]	7.4 (3.8) [3-13]	13.5 (4.4) [6-19]	Poor Readers < Hyperlexic < Skilled Readers
TROG-2 (ss) ¹	61.6 (14.1) [55-110]	68.3 (15.1) [55-93]	89.7 (21.4) [58-116]	
Recalling Sentences CELF-4 (raw score)	10.1 (13.3) [0-39]	19.6 (11.0) [6-21]	32.8(14.8) [1-60]	Poor Readers = Hyperlexic < Skilled Readers
Recalling Sentences CELF-4 (scs) ²	3.5 (4.3) [1-16]	6.1 (4.3) [1-14]	10.1 (5.0) [1-19]	
Non word repetition (max 30) (raw score)	7.9 (8.8) [0-23]	21.0 (3.6) [15-26]	21.1 (8.2) [0-30]	Poor Readers < Hyperlexic = Skilled Readers
Autism symptomatology				
ASSQ ^a (raw score)	25.1 (9.4) [11-50]	14.6 (9.1) [2-35]	16.9 (7.2) [5-28]	Poor Readers < Hyperlexic = Skilled Readers
Non- verbal ability				
Matrix reasoning (raw score)	5.0 (5.6) [0-20]	12.4(6.3) [7-25]	14.9 (6.6) [0-26]	Poor Readers < Hyperlexic = Skilled Readers
Matrix reasoning (t- score) ³	35.0 (10.1) [23-66]	49.0 (9.2) [40-67]	48.7(10.5) [28-68]	

1
2
3 *Note* ¹⁾ standard scores (M=100, SD=15), ²⁾ scaled scores (M=10, SD=3), ³⁾ T -scores (M=50,
4
5 SD=10), ^{a)} *n* =17 of the “skilled readers” parents completed the ASSQ.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Table 2. Results from the age 3 year assessment in the three subgroups: “poor readers”, “hyperlexic/poor comprehenders” and “skilled readers”

Results from the 3 year assessment	Mean (SD)			Group differences using ANOVA
	Poor Readers <i>n</i> = 25	Hyperlexic/Poor Comprehenders <i>n</i> = 10	Skilled readers <i>n</i> = 18	
Age (months)	35.4 (5.9)	36.1 (6.2)	38.8 (4.8)	Poor Readers = Hyperlexic = Skilled Readers
Language ability				
RDLS (raw score)	8.0 (11.3)	14.0 (11.2)	32.6 ² (20.0)	Poor Readers = Hyperlexic < Skilled Readers
PARIS level (no verbal 3-5 [%])	10 (40%)	5 (50%)	17 (94%)	Poor Readers < Skilled Readers
Autism symptomatology				
ADOS severity total (raw score)	6.1 (2.9)	4.7 (2.4)	4.4 ² (2.7)	Poor Readers = Hyperlexic = Skilled Readers
Developmental Quotient				
Griffiths' developmental scales ¹	79.7 ³ (18.5)	80.1 ³ (11.2)	82.5 ³ (21.8)	Poor Readers = Hyperlexic = Skilled Readers
Adaptive functioning				
VABS Socialization ¹	73.2 ⁴ (7.9)	74.9 (14.3)	78.5 ⁴ (12.4)	Poor Readers = Hyperlexic = Skilled Readers
VABS Communication ¹	70.9 ⁴ (13.4)	71.0 (12.7)	81.8 ⁴ (15.2)	Poor Readers < Skilled Readers

Note ¹ standard scores (M=100; SD=15). Not all children had been assessed with the instrument

²) *n* = 1 of “skilled readers” did not provide a score, ³) *n* = 3 of “poor readers”, *n* = 2 of

1
2
3 hyperlexic/poor comprehenders, and $n = 4$ of “skilled readers” did not provide a score, ⁴⁾ $n = 1$ of
4
5 “poor readers” and $n = 2$ of “skilled readers” did not provide a score.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Supplementary material

Group comparisons on baseline measures for assessments at age 3 using *t*-tests and ANCOVA between the current study sample and the remainder of the AUDIE cohort

	Mean (SD)		Test statistic	Group differences
	Current sample <i>n</i> =53	Remainder of the AUDIE cohort <i>n</i> =54	<i>t</i> -value / F-value	<i>p</i> -value
Age (months)	36.70 (5.73)	33.91 (6.71)	-2.31	*
Developmental Quotient (GDS)	80.67 (18.23)	82.78 (18.37)	.56 ¹	.577
Language comprehension (RDLS)	17.17 (18.19)	13.35 (14.71)	-1.19 ¹	.236
ADOS (severity total)	5.29 (2.82)	5.39 (2.88)	-.18 ¹	.861
Vineland socialization	75.22 (10.94)	77.33 (9.50)	.30 ¹	.302
Vineland communication	74.42 (14.48)	76.08 (12.78)	.61 ¹	.543

¹ Corrected for age (ANCOVA result). * $p < .05$

A relatively large proportion of the original cohort could not be included in the analyses since they had not yet started 1st grade. However, in fact $n = 24$ of these children were administered the tests of literacy, even though they had only received very little or no formal reading instruction. A look at their data revealed that there several cases that displayed some reading ability. When their performance was related to norms for grade 1, six of them could classified as “hyperlexics/poor comprehenders” and one as a “skilled reader. In the hyperlexic group there was one case with particularly precocious reading, obtaining a stanine score of 7 using grade 1 norms.