

Journal of
**Neurology, Neurosurgery
& Psychiatry**

**Intensive Upper Limb Neurorehabilitation in Chronic Stroke
- Outcomes from the Queen Square Programme**

Journal:	<i>Journal of Neurology, Neurosurgery, and Psychiatry</i>
Manuscript ID	jnnp-2018-319954.R2
Article Type:	Research paper
Date Submitted by the Author:	n/a
Complete List of Authors:	Ward, Nick; UCL Institute of Neurology, Department of Clinical and Motor Neurosciences; The National Hospital for Neurology and Neurosurgery Brander, Fran; The National Hospital for Neurology and Neurosurgery Kelly, Kate; The National Hospital for Neurology and Neurosurgery
Keywords:	
Specialty:	

SCHOLARONE™
Manuscripts

Intensive Upper Limb Neurorehabilitation in Chronic Stroke – Outcomes from the Queen Square Programme

Nick S Ward^{1,2,3}, Fran Brander^{2,3}, Kate Kelly^{2,3}

¹Department of Clinical and Motor Neuroscience, UCL Institute of Neurology, Queen Square, London WC1N 3BG.

²The National Hospital for Neurology and Neurosurgery, Queen Square, London WC1N 3BG.

³UCLP Centre for Neurorehabilitation, Queen Square, London WC1N 3BG.

Number of characters in the title (with spaces): 67

Number of words in the abstract: 249

Word count (excluding title page, abstract, references, figures, tables): 3845

Number of figures: 6

Number of tables: 4

Correspondence should be addressed to Professor N S Ward

Department of Clinical and Motor Neuroscience

UCL Institute of Neurology,

33 Queen Square, London WC1N 3BG,

Tel: +44 (0) 20 3448 8762

e-mail n.ward@ucl.ac.uk

Abstract

Objective: Persistent difficulty in using the upper limb remains a major contributor to physical disability post-stroke. There is a nihilistic view about what clinically relevant changes are possible after the early post-stroke phase. The Queen Square Upper Limb Neurorehabilitation programme delivers high quality, high dose, high intensity upper limb neurorehabilitation during a 3-week (90 hour) programme. Here, we report clinical changes made by the chronic stroke patients treated on the programme, factors that might predict responsiveness to therapy, and the relationship between changes in impairment and activity.

Methods: Upper limb impairment and activity were assessed on admission, discharge, 6 weeks and 6 months after treatment, with modified upper limb Fugl-Meyer, (FM-UL, max-54), Action Research Arm Test (ARAT, max-57) and Chedoke Arm and Hand Activity Inventory (CAHAI, max-91). Patient reported outcome measures were recorded with the Arm Activity Measure (ArmA) parts A (0-32) and B (0-52), where lower scores are better.

Results: 224 patients (median time post-stroke 18 months) completed the 6 month programme. Median scores on admission were: FM-UL=26 (IQR 16-37), ARAT=18 (IQR 7-33), CAHAI=40 (28-55), ArmA-A=8 (IQR 4.5-12), ArmA-B=38 (IQR 24-46). The median scores 6 months after the programme were: FM-UL=37 (IQR 24-48), ARAT=27 (IQR 12-45), CAHAI=52 (IQR 35-77), ArmA-A=3 (IQR 1-6.5), ArmA-B=19 (IQR 8.5-32). We found no predictors of treatment response beyond admission scores.

Conclusion: With intensive upper limb rehabilitation, chronic stroke patients can change by clinically important differences in measures of impairment and activity. Crucially, clinical gains continued during the 6 month follow up period.

Keywords: Stroke, rehabilitation, upper limb

INTRODUCTION

Stroke remains common¹ and persistent difficulty in using the upper limb is a major contributor to ongoing physical disability². The general consensus remains that most spontaneous recovery of the upper limb occurs over the first three months after stroke and current levels of rehabilitation result in little improvement after that, particularly at the level of impairment³. Improving outcomes through higher dose (time in rehabilitation or number of repetitions) and intensity (dose per session) of rehabilitation is an attractive option⁴. However, clinical trials of higher dose upper limb rehabilitation have generally not produced the magnitude of improvement that will change clinical practice⁵, whether delivered in the early⁶ or chronic stages post-stroke⁷⁻⁹. A common factor in these trials is that the dose (in hours) of additional therapy remained relatively low (18-36 hours). Despite scepticism that stroke patients could tolerate much higher doses⁸, one study managed to deliver 300 hours of upper limb therapy to chronic stroke patients over 12 weeks and reported changes in measures of both impairment and activity that were far greater than those in lower dose studies¹⁰. Three hundred hours represents an order of magnitude higher than any dose of rehabilitation offered in previous upper limb rehabilitation trials and deserves further consideration. However, this idea is challenging because of the logistics of setting up such a trial in health care settings where the ethic of high dose, high intensity rehabilitation is not supported. In this context, it is important to report the findings of clinical services that are able to deliver higher doses than conventionally seen. The Queen Square Upper Limb (QSUL) Neurorehabilitation programme is a single centre clinical service that provides 90 hours of timetabled treatment focusing on the post-stroke upper limb in chronic (> 6 months post-stroke) stroke patients. Here we report (i) outcomes for patients admitted to this programme at the National Hospital for Neurology and Neurosurgery, University College London Hospitals NHS Trust (UCLH), including 6 month follow up data to look at whether any clinical benefits were maintained, (ii) the characteristics of the patients admitted and any predictors of response and (iii) the relationship between changes in impairment and activity.

METHODS

Participants

Patients were referred by primary care physicians for advice about ongoing management of the paretic upper limb following stroke. There were no criteria around time since stroke. Clinical criteria warranting admission to the programme were broad but were focused on whether we felt able to help patients achieve their goals for the affected upper limb. Barriers to admission included factors precluding the ability to work in I upper limb tasks; (i) absent movements, throughout the limb; (ii) a painful shoulder limiting an active forward reach (mostly due to adhesive capsulitis); (iii) severe spasticity or non-neural loss of range; (iv) unstable medical conditions. These patients were not offered admission to the treatment programme, but

1
2
3 were provided with appropriate advice or referred for other treatment approaches. Re-referral to the
4 programme was then considered (for example after successful management of adhesive capsulitis of the
5 shoulder). Features that were more likely to result in admission to the programme included at least some
6 ability to flex the shoulder and at least visible flickers of movement in finger and/or wrist extensors,
7 although this was not always the case as reflected in baseline upper limb scores. Patients were admitted
8 to the programme as day attenders, either from home or from the UCLH dedicated patient accommodation
9 and as such were either self-caring, or self-caring with the support of one other person.
10
11
12
13
14
15

16 **Interventions**

17 Initial assessment consisted of analysis of both movement and performance in activities of daily living.
18 Subsequent treatment was aimed at reducing impairment and promoting re-education of motor control
19 within activities of daily living. Individualised meaningful tasks were practiced repeatedly in order to
20 facilitate task mastery with a focus on quality of movement. This was achieved through (i) adaptation of
21 the task, e.g. decomposing tasks into individual components to be practiced; (ii) adaptation of the
22 environment, e.g. fabrication of functional splints and adaptation of tools such as cutlery or screwdrivers,
23 to enable integration of the affected hand in meaningful activities; (iii) assistance, e.g. de-weighting the
24 arm to allow strengthening and training of movement quality and control through increased range; (iv)
25 independent task practice. Coaching was considered a key component of the programme, used throughout
26 to embed new skills and knowledge into individual daily routines. Consequently, individuals increase
27 participation and confidence in their desired goals, enhancing self-efficacy and motivation to sustain
28 behavioural change beyond the end of the active treatment period.
29
30
31
32
33
34
35
36
37

38 This overall approach was achieved through two daily sessions each of physiotherapy and occupational
39 therapy, supplemented with tailored, individualized interventions, including repetitive practice with a
40 rehabilitation assistant or robotic device, sensory retraining, use of dynamic and functional orthoses,
41 neuromuscular electrical stimulation, and group work. Furthermore, patients were encouraged to work on
42 cardiovascular fitness during the programme. A 6 hour timetable was implemented 5 days a week for 3
43 weeks (total therapy time, 90 hours). Motor tasks could be described as passive or active, assisted or
44 unassisted, functional or non-functional. Over the three weeks, the aim was to increase the time spent on
45 active, unsupported functional tasks, depending on a patient's level of impairment and progress. The
46 programme was staffed with a 1:1 staff to patient ratio (2 physiotherapists, 2 occupational therapists, 2
47 rehabilitation assistants for 6 patients).
48
49
50
51
52
53
54
55
56
57
58
59
60

Quantitative assessment

All patients had the following baseline scores measured on admission (table 1): modified Rankin scale (mRS), Barthel Index (BI), Neurological Fatigue Index (NFI), Hospital Anxiety and Depression Scale (HADS), and sensation (as indexed by light touch on palm as described in Fugl-Meyer assessment of the upper extremity).

The affected upper limb was assessed on admission (T1), discharge (T2), 6 weeks (T3) and 6 months (T4) post-discharge, using the following measures: Fugl-Meyer (upper limb) (FM-UL), Action Research Arm Test (ARAT), Chedoke Arm and Hand Activity Inventory (CAHAI-13), Arm Activity Measure

The FM-UL is a stroke-specific, performance-based impairment index (where impairment refers to loss of body structure and function) with good validity and intra- and inter-rater reliability^{11,12}. Here we have used a modified version of the FM-UL, excluding measures of co-ordination and reflexes based on hierarchical properties of the scale¹³, and as these do not relate to the upper limb synergies of interest¹⁴. The minimum clinically important difference (MCID) has been reported as 5.25 points¹⁵.

The ARAT assesses a patient's ability to handle objects differing in size, weight and shape and therefore can be considered to be an arm-specific measure of activity limitation¹⁶. The MCID has been suggested as 5.7 points¹⁷.

The CAHAI-13 is a validated, upper-limb measure that uses a 13-point quantitative scale in order to assess recovery of the arm and hand in performing activities of daily living after a stroke¹⁸. It is a performance test using day to day items. It is not designed to measure the patient's ability to complete the task using only their unaffected hand, but rather to encourage bilateral upper limb use. No MCID has been reported for CAHAI-13 (although the minimum detectable change is reported as 6.2¹⁸).

The Arm Activity Measure (ArmA) is a patient reported outcome score¹⁹ with two components. The ArmA-A asks patients about their ability to care for their affected arm either themselves with their unaffected arm or by a carer or a combination of both of these. ArmA-B asks patients about how easy or hard it is to use the affected arm in activities of daily living. For ArmA-A and ArmA-B, note that lower scores are better. No MCID has been reported for ArmA-A or ArmA-B.

Analysis plan

Our primary aim was to measure changes in upper limb deficit over time were examined using the Friedman test for all outcome scores. Post-hoc analysis to test differences between individual time points was carried out with the Wilcoxon signed-ranks test. As a secondary aim, we were then interested in whether baseline characteristics correlated with final (T4) outcomes. We initially examined for correlations between individual baseline characteristics and outcomes using Spearman's rank correlation. We also tested whether median T4 outcomes were different depending on gender or side of lesion using Wilcoxon rank

1
2
3 sum test. Time since stroke was not considered as a covariate as once in the chronic phase, there is no
4 indication that time since stroke has a linear effect on outcome. Lastly we performed multiple linear
5 regression to look for predictors of either absolute scores at T4 or the change in scores T1 to T4. Lastly,
6
7 we also examined the relationship between the changes in outcome scores from admission to 6 month
8
9 follow up using Spearman's rank correlation.
10
11

12 13 **RESULTS**

14 **Baseline characteristics**

15
16 Between January 2015 and December 2017, 268 patients were admitted to the QSUL programme
17 (representing 46% of patients assessed in out-patient clinic). 40.4% of patients were admitted from home,
18 whilst 59.6% stayed in the UCLH dedicated patient accommodation. Note, only 16 patients were admitted
19 to the programme prior to 6 months post-stroke, and only 1 prior to 3 months, reflecting overall referral
20 patterns to the programme.
21
22

23
24 Of those, 30 were non-stroke (12 traumatic brain injury, 8 brain tumour, 3 peripheral neurological
25 conditions, 3 spinal cord injury, 4 inflammatory central nervous system conditions) and are excluded from
26 this analysis. A total of 238 stroke patients were admitted, of whom 224 completed follow up assessments
27 at 6 weeks and 6 months after discharge from the programme. Of the 14 incomplete follow ups, 5 patients
28 considered it too far to travel for follow up, 4 patients suffered intercurrent illness (recurrent stroke,
29 fractured hip, complications of surgery, seizures), and 5 could not be contacted. Differences in median
30 values between complete and missing data groups was tested with Wilcoxon Signed Rank test. Where
31 appropriate, differences in proportions between complete and missing data groups were tested with Chi-
32 squared test. There were no differences in baseline characteristics between patients who completed follow
33 up and those who were lost to follow up (table 1). Because (i) there was no systematic difference between
34 subjects with complete data and those with missing data, and (ii) only outcome data (not explanatory
35 variables, which were collected at T1) were missing, then missing data were dealt with by performing a
36 complete case analysis (i.e. only using subjects where complete data were available).
37
38
39
40
41
42
43
44
45
46
47
48

49 **Changes in upper limb deficit**

50
51 The median scores for the affected upper limb at admission (T1), discharge (T2), 6 weeks (T3) and 6 months
52 post-discharge (T4) are shown in table 2. The Friedman test was used to demonstrate a significant effect of
53 time since admission for all outcome scores; FM-UL (χ^2 (3) = 431.8, $p < 0.0001$); ARAT (χ^2 (3) = 383.2,
54 $p < 0.0001$); CAHAI (χ^2 (3) = 371.6, $p < 0.0001$); ArmA-A (χ^2 (3) = 238.4, $p < 0.0001$); ArmA-B (χ^2 (3) = 305.6,
55 $p < 0.0001$). Post hoc analysis with Wilcoxon signed-ranks test was conducted to test the hypothesis that the
56 difference between the paired values (T1-T2, T1-T3, T1-T4, T2-T3, T2-T4, T3-T4) comes from a distribution
57
58
59
60

1
2
3 whose median is zero (i.e. testing whether the change in score is significantly greater than zero). A
4 Bonferroni correction was applied to the significance level for the 6 comparisons made for each of the 5
5 different outcome scores ($p < 0.05/30 = 0.0017$). For each outcome score, there was a significant difference
6 between each pair of time points compared, except T2-T3 for the ArmA-A (table S1). Individual scores at
7 each time point are shown in figures 1 and 2. Individual changes in scores in comparison to admission are
8 shown in table 3 and figures 1 and 2. These results demonstrate improvement during the 3 week
9 programme, but also continued improvement after discharge.

10
11 By 6 months after the programme, 68.3% of patients had achieved greater than the MCID of 5.25 points on
12 the FM-UL. For ARAT, this figure was 61.6%. No published MCID is published for the other scores. MCID is
13 often quoted as 10% of the maximum score. If this were the case, then by 6 months, MCID would have
14 been reached by 59.4% of patients for CAHAI, 53.8% for ArmA-A and 72.3% for ArmA-B.

23 Predictors of outcome

24
25 Firstly, we were interested to see whether any baseline characteristics (T1) correlated with final outcome
26 at T4. Spearman's rank correlation was used to show that admission values at T1 for FM-UL, ARAT, CAHAI,
27 ArmA-A, ArmA-B all correlated with T4 values for FM-UL, ARAT, CAHAI, ArmA-A, ArmA-B. In addition,
28 admission BI correlated with T4 CAHAI; admission mRS correlated with T4 FM-UL, ARAT and CAHAI;
29 admission sensation correlated with T4 CAHAI. Age, NFI and HADS at T1 did not correlate with any scores
30 at T4. All p-values were corrected for multiple comparisons by Bonferroni correction (Table 4). Figure 3
31 helps to visualise the influence of baseline admission scores on final score at 6 month follow up.
32 Furthermore, a Wilcoxon ranked sum test showed that T4 outcomes did not differ by gender or affected
33 hemisphere. There was a trend for patients with non-dominant affected limbs to achieve better (lower)
34 ArmA-B scores in comparison to those with dominant affected limbs (median 17 for non-dominant vs -27
35 for dominant, $p = 0.027$, uncorrected). There were no differences in T4 scores between patients with
36 dominant or non-dominant affected limbs for any of the other outcome scores.

37
38 Secondly, we were interested to see whether any baseline characteristics (T1) correlated with magnitude
39 of change from admission to 6 months follow up (T1 to T4). Spearman's rank correlation was used to show
40 that FM-UL, ARAT, CAHAI and ArmA-B at T1 correlated with change in ARAT and CAHAI (but not FM-UL)
41 from T1 to T4. FM-UL at T1 also correlated with change in ArmA-B from T1 to T4. ArmA-A at T1 correlated
42 with change in both CAHAI and ArmA-A from T1 to T4. ArmA-B at T1 correlated with change in both ArmA-A
43 and ArmA-B from T1 to T4. Age, NFI, HADS, BI and mRS at T1 did not correlate with any scores at T4. All p-
44 values were corrected for multiple comparisons by Bonferroni correction (table 5). We also performed a
45 Wilcoxon ranked sum test to show that change in scores from T1 to T4 were not different depending on
46 gender or affected hemisphere. There was a trend for patients with non-dominant affected limbs to
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 improve on the ArmA-B between T1 and T4 to a greater degree than those with dominant affected limbs
4 (median changes of -17.5 for non-dominant vs -12 for dominant, $p = 0.017$, uncorrected). There were no
5 differences in change scores between patients with dominant or non-dominant affected limbs for any of
6 the other outcome scores.
7
8

9
10 We then performed multiple linear regression to look for predictors of either absolute scores at 6 months
11 after treatment (T4) or the change in scores from admission to 6 months after treatment (T1 to T4) using
12 age, BI, mRS, NFI, HADS and sensation on admission together with the matching initial score on admission
13 (e.g. FM1 if attempting to predict FM4 or change from FM1 to FM4). Models predicting T4 scores for FM-
14 UL, ARAT, CAHAI, ArmA-A and ArmA-B were all highly significant ($p < 0.0001$). Models predicting change
15 from T1 to T4 for ARAT, CAHAI, ArmA-A and ArmA-B (but not FM-UL) were also highly significant ($p < 0.0001$).
16 However, the only factor that significantly contributed to these models was the matched initial score on
17 admission (e.g. ARAT at T1 predicting ARAT at T4), but not age, BI, mRS, NFI, HADS or sensation on
18 admission. As such, this approach did not contribute further to the correlation results shown in table 4 and
19 5, with regression models not able to account for a higher proportion of variance in scores at T4 or change
20 in scores from T1 to T4 than the admission score (see also Figure 3).
21
22
23
24
25
26
27
28

29 30 **Relationship between outcome measures**

31 Each outcome score assesses a different aspect of recovery. Using Spearman's rank correlation, we
32 examined the relationship between the changes in outcome scores from admission to 6 month follow up.
33 A Bonferroni correction was applied to the significance level for the 10 comparisons made between the 5
34 different outcome scores ($p < 0.05/10 = 0.005$) (Table 6 and Figure 4). Changes in FM-UL, ARAT and CAHAI
35 correlated weakly with one another. The ArmA-B change correlated with changes in both ArmA-A and
36 ARAT.
37
38
39
40
41
42
43

44 **DISCUSSION**

45 We report the results from a single centre clinical service delivering high dose and intensity of upper limb
46 focused neurorehabilitation for people with stroke. Almost all of these patients were in the chronic (>6
47 months after stroke) phase. The key messages from our results are (i) patients were able to complete the
48 90 hours of the programme, despite exhibiting a range of impairments and fatigue levels, (ii) large clinical
49 improvements in upper limb impairment (FM-UL) and activity (ARAT, CAHAI, ArmA) were observed, and
50 (iii) these changes were maintained, or even improved upon, 6 months after treatment.
51
52
53
54
55

56 The QSUL programme is based on the hypothesis that a high dose and intensity of upper limb
57 neurorehabilitation can lead to large clinically meaningful improvements in chronic stroke patients. It is
58 worth comparing the magnitude of the changes with previous reported clinical trials of upper limb
59
60

1
2
3 rehabilitation. Most obviously, the study by McCabe et al¹⁰ investigated the effects of 300 hours of upper
4 limb rehabilitation in 48 chronic stroke patients with similar characteristics to those reported here (baseline
5 FM-UL 24, compared to 26 in our patients) and reported an increase of 8-11 points on the FM-UL at the
6 end of 300 hours of upper limb treatment (although later follow up results have not been reported). In the
7 QSUL programme, the median FM-UL (modified) improvement was 6 points after the 90-hour programme,
8 increasing to 9 points by 6 months follow up, achieving a similar overall magnitude of change in comparison
9 to McCabe et al. In addition to reductions in impairment (FM-UL), our patients also improved at the level
10 of activity (ARAT and CAHA) and participation (using the patient reported outcome measure, ArMA).
11 Importantly, we report that these group level effects continued to improve at 6 weeks and then 6 months
12 after treatment had stopped, (figure 1 and 2).

13
14
15
16
17
18
19
20 Other than the study by McCabe et al, the general view has been that large changes, especially in
21 impairment, have not been seen for behavioural interventions of the post-stroke upper limb³. Direct
22 comparison with other studies is difficult because some may have been conducted in less severely affected
23 patients, the dose (measured in time) of the intervention may have been lower, and the numbers of
24 patients enrolled has been much lower than we report here. Whilst this report of our single-centre clinical
25 outcomes is not a replication of the McCabe study, it suggests that the next wave of upper limb
26 rehabilitation trials must investigate much higher doses of the treatment than is currently attempted.
27 Whilst we will hear the refrain that it is not possible to deliver such high doses in current health care settings
28 (the pragmatic view), it is in fact precisely the role of clinical research to challenge what we currently do in
29 order to reshape and improve our clinical services to make them better (the aspirational view). Those
30 involved in stroke medicine will recognise that this is exactly the effect that thrombolysis had on acute
31 management of stroke, leading to the high quality hyperacute stroke services we see today. Post-stroke
32 neurorehabilitation must adopt the same aspirational approach²⁰.

33
34
35
36
37
38
39
40
41
42
43 Whilst the appropriate dose of upper limb rehabilitation is debated, the content of the intervention
44 receives less attention. Our approach was based on analysis of movement and performance in activities of
45 daily living, reduction of impairment and re-education of quality and control of movement within activities
46 of daily living, all performed in an iterative and non-linear fashion. Education, self-efficacy and goal setting
47 were integral components of the programme, and we can speculate that the education and focus on self-
48 efficacy contributed to the continued increase in scores after treatment stopped. The therapeutic approach
49 taken towards specific impairments was the same across patients, but approaches at the activity and
50 participation level will necessarily vary as they are tailored to an individual's specific meaningful goals. We
51 acknowledge that an immediate challenge for us and for the field in general is to be able to define what
52 the 'active ingredients' of neurorehabilitation are, so that these elements can be further tested. Our
53 approach appears to have similarities with that taken by McCabe et al, with an emphasis on the quality of
54
55
56
57
58
59
60

1
2
3 the movement as well as the final goal of the movement. The latter is often the sole focus of task-specific
4 repetition, which has been used in many upper limb rehabilitation studies to date. The evidence for
5 repetitive task-specific training is of low to moderate quality²¹ and the effects have been disappointingly
6 small even when the number of repetitions has been greatly increased⁸. It is difficult to tell whether this
7 lack of effect is specific to task-specific training, because in general the dose (in terms of hours of active
8 training) has been quite low, whether delivered by a therapist²¹ or robotic device²². There have been
9 suggestions that chronic stroke patients might not tolerate more training than has been offered in these
10 trials⁸, but the study of McCabe et al¹⁰ and now our own clinical experience undermine this view.

11
12 Although this result is exciting, we must acknowledge that it is a single centre clinical service with no
13 randomisation and no control group. Although there are some reports of late improvement up to 12
14 months after lacunar stroke²³, we expect that chronic stroke patients who are often several years after
15 their stroke, will not change across this range of outcome scores without treatment. We suggest that it is
16 unlikely that this magnitude of change would be seen without therapeutic intervention, but this assertion
17 requires empirical confirmation. Our work however, provides strong justification for undertaking such high
18 dose clinical trials. Clinic data on which to base these future studies is sorely lacking and this has led to the
19 proliferation of low dose pragmatic rehabilitation studies. However, there is still much to be learned from
20 published data arising from clinical services in neurorehabilitation²⁴, making up for lack of clinical trial rigour
21 with large sample sizes that are rarely achieved in clinical trials. Data from clinical services therefore have
22 the potential to refine what experimental questions are addressed in future randomised clinical trials.
23 Another limitation is the unblinded nature of the assessments, introducing possible bias into the results.
24 All assessors are also therapists on the programme, although we ensure that assessments are carried out
25 by therapists from another team who are not treating the patient. Fidelity of outcome scores is always an
26 important issue in clinical trials. In our programme, we have instituted regular in-service training for FM-
27 UL, ARAT and CAHAI to ensure high levels of reproducibility between assessors.

28
29 A common complaint regarding single centre clinical services is that they select only patients most likely to
30 succeed. It is important to acknowledge that the QSUL programme does not have strict inclusion/exclusion
31 criteria. Although our patients are generally younger than the average stroke patient and have high levels
32 of independence they have a wide range of upper limb impairments. It can be seen from figures 1 and 2
33 that the range of severity of the upper limb on the QSUL programme was very broad, quite unlike a clinical
34 trial, although the median scores suggest that patients were weighted towards the moderate-severe end
35 of upper limb impairment and activity limitation. We would argue that the QSUL programme is aspirational
36 in its goals (high quality, high dose, high intensity treatment) and pragmatic in its application (wide range
37 of patients admitted onto the programme). It's interesting to consider the potential impact of initial upper
38 limb severity on our outcomes, but in fact, although initial score correlates well with the final score at 6
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 months (table 4), the change in score correlates only weakly with initial score (table 5, Figure 3). There are
4 two ways this 'mixed-bag' approach may have potentially reduced the overall magnitude of the clinical
5 effects we observed. Firstly, there may be a ceiling effect from patients with relatively high scores on
6 admission. Secondly, many of our patients were quite severely impaired. Indeed, a quarter of our patients
7 had an admission ARAT below 7, which suggests they have great difficulty in using their hand in any kind of
8 activity of daily living. The treatment goals in these type of patients might initially be very different to the
9 moderate to mild type patients, with benefits less likely to be reflected in changes in FM-UL and ARAT
10 compared to the CAHAI or ArMA for example. However, the wide range of patients treated in the
11 programme means that no clear patterns, in relation to differential effects on various scores, could be seen
12 (Table 6, Figure 4).

13
14
15 Some might argue that the wide range of responses, including those patients who changed very little, points
16 to the need for stratification and selection of those patients most likely to respond, and this is certainly a
17 reasonable view. However, the first point to make is that our results show that with large enough numbers
18 and a high enough dose of the intervention, it is possible to demonstrate large group-level effects in
19 relatively unselected populations of chronic stroke patients. Indeed, if the former (stratification) approach
20 is to be adopted, then these kind of large data sets will be required to determine what the characteristics
21 of the 'responders' are likely to be. The second point to make is that we found no strong clinical indicators,
22 amongst our measured explanatory variables, of which patients were most likely to change, including
23 sensory loss in the hand, fatigue²⁵ and depression (tables 4 and 5). It might be that neurophysiological
24 (presence or absence of motor evoked potentials)²⁶, neuroimaging (assessment of corticospinal tract,
25 whole brain damage or brain function)²⁷⁻³⁰ or cognitive (sustained attention, memory, motivation)^{31,32}
26 measures are required for accurate stratification based on likely response, and this can certainly be tested
27 in future.

28
29
30 In summary, here we present the results from a single centre clinical service dedicated to post-stroke upper
31 limb neurorehabilitation in chronic stroke patients. Despite treating patients with a wide range of
32 impairment, we were able to show a large group-level magnitude of change at both the impairment and
33 activity level. Our experience suggests that much higher doses and intensity of upper limb
34 neurorehabilitation can be delivered and our results should inform future clinical trial design.

53 **ACKNOWLEDGEMENTS**

54
55 Thanks to all the physiotherapists and occupational therapists at The National Hospital for Neurology and
56 Neurosurgery, Queen Square, who have treated patients on this programme. Thanks to UCLH Charities,
57 Friends of UCLH and The National Brain Appeal for funding to purchase equipment used in this programme.
58
59
60

REFERENCES

1. Licher S, Darweesh SKL, Wolters FJ, et al. Lifetime risk of common neurological diseases in the elderly population. *J Neurol Neurosurg Psychiatry*. 2018 Oct 2; doi: 10.1136/jnnp-2018-318650. [Epub ahead of print].
2. Broeks JG, Lankhorst GJ, Rumping K, et al. The long-term outcome of arm function after stroke: results of a follow-up study. *Disabil Rehabil*. 1999;21(8):357–64.
3. Krakauer JW, Carmichael ST, Corbett D, et al. Getting neurorehabilitation right: what can be learned from animal models? *Neurorehabil Neural Repair*. 2012;26(8):923–31.
4. Lohse KR, Lang CE, Boyd LA. Is more better? Using metadata to explore dose-response relationships in stroke rehabilitation. *Stroke*. 2014;45(7):2053–8.
5. Langhorne P, Coupar F, Pollock A. Motor recovery after stroke: a systematic review. *Lancet Neurol*. 2009;8(8):741–54.
6. Winstein CJ, Wolf SL, Dromerick AW, et al. Effect of a Task-Oriented Rehabilitation Program on Upper Extremity Recovery Following Motor Stroke: The ICARE Randomized Clinical Trial. *JAMA*. 2016;315(6):571–81.
7. Lo AC, Guarino PD, Richards LG, et al. Robot-assisted therapy for long-term upper-limb impairment after stroke. *N Engl J Med*. 2010;362(19):1772–83.
8. Lang CE, Strube MJ, Bland MD, et al. Dose-response of task-specific upper limb training in people at least 6 months post stroke: A Phase II, single-blind, randomized, controlled trial. *Ann Neurol*. 2016;80(3):342–54.
9. Klamroth-Marganska V, Blanco J, Campen K, et al. Three-dimensional, task-specific robot therapy of the arm after stroke: a multicentre, parallel-group randomised trial. *Lancet Neurol*. 2014;13(2):159–66.
10. McCabe J, Monkiewicz M, Holcomb J, et al. Comparison of robotics, functional electrical stimulation, and motor learning methods for treatment of persistent upper extremity dysfunction after stroke: a randomized controlled trial. *Arch Phys Med Rehabil*. 2015;96(6):981–90.
11. Duncan PW, Propst M, Nelson SG. Reliability of the Fugl-Meyer assessment of sensorimotor recovery following cerebrovascular accident. *Phys Ther*. 1983;63(10):1606–10.
12. Fugl-Meyer AR, Jääskö L, Leyman I, et al. The post-stroke hemiplegic patient. 1. a method for evaluation of physical performance. *Scand J Rehabil Med*. 1975;7(1):13–31.
13. Crow JL, Harmeling-van der Wel BC. Hierarchical properties of the motor function sections of the Fugl-Meyer assessment scale for people after stroke: a retrospective study. *Phys Ther*. 2008;88(12):1554–67.

14. Hsieh Y-W, Hsueh I-P, Chou Y-T, et al. Development and validation of a short form of the Fugl-Meyer motor scale in patients with stroke. *Stroke*. 2007;38(11):3052–4.
15. Page, S. J., Fulk, G. D. & Boyne, P. Clinically important differences for the upper-extremity Fugl-Meyer Scale in people with minimal to moderate impairment due to chronic stroke. *Phys Ther* **92**, 791–798 (2012).
16. Platz T, Pinkowski C, van Wijck F, et al. Reliability and validity of arm function assessment with standardized guidelines for the Fugl-Meyer Test, Action Research Arm Test and Box and Block Test: a multicentre study. *Clin Rehabil*. 2005;19(4):404–11.
17. Van der Lee, J. H. *et al.* The intra- and interrater reliability of the action research arm test: a practical test of upper extremity function in patients with stroke. *Arch Phys Med Rehabil* **82**, 14–19 (2001).
18. Barreca SR, Stratford PW, Lambert CL, et al. Test-retest reliability, validity, and sensitivity of the Chedoke arm and hand activity inventory: a new measure of upper-limb function for survivors of stroke. *Arch Phys Med Rehabil*. 2005;86(8):1616–22.
19. Ashford S, Turner-Stokes L, Siegert R, et al. Initial psychometric evaluation of the Arm Activity Measure (ArmA): a measure of activity in the hemiparetic arm. *Clin Rehabil*. 2013;27(8):728–40.
20. Bernhardt J, Borschmann K, Boyd L, et al. Moving rehabilitation research forward: Developing consensus statements for rehabilitation and recovery research. *Int J Stroke*. 2016;11(4):454–8.
21. French B, Thomas LH, Coupe J, et al. Repetitive task training for improving functional ability after stroke. *Cochrane Database Syst Rev*. 2016 14;11:CD006073.
22. Veerbeek JM, Langbroek-Amersfoort AC, van Wegen EEH, et al. Effects of Robot-Assisted Therapy for the Upper Limb After Stroke. *Neurorehabil Neural Repair*. 2017;31(2):107–21.
23. Ganesh A, Gutnikov SA, Rothwell PM, Oxford Vascular Study. Late functional improvement after lacunar stroke: a population-based study. *J Neurol Neurosurg Psychiatry*. 2018 Jul 21; doi: 10.1136/jnnp-2018-318434. [Epub ahead of print]
24. Freeman JA, Hobart JC, Playford ED, et al. Evaluating neurorehabilitation: lessons from routine data collection. *J Neurol Neurosurg Psychiatry*. 2005;76(5):723–8.
25. De Doncker W, Dantzer R, Ormstad H, et al. Mechanisms of poststroke fatigue. *J Neurol Neurosurg Psychiatry*. 2018;89(3):287-293.
26. Agarwal S, Koch G, Hillis AE, et al. Interrogating cortical function with transcranial magnetic stimulation: insights from neurodegenerative disease and stroke. *J Neurol Neurosurg Psychiatry*. 2018 Jun 4; doi: 10.1136/jnnp-2017-317371. [Epub ahead of print].
27. Aguilar OM, Kerry SJ, Ong Y-H, et al. Lesion-site-dependent responses to therapy after aphasic stroke. *J Neurol Neurosurg Psychiatry*. 2018 Apr 17; doi: 10.1136/jnnp-2017-317446. [Epub ahead of print].

- 1
2
3 28. Park C-H, Kou N, Ward NS. The contribution of lesion location to upper limb deficit after stroke. *J Neurol*
4 *Neurosurg Psychiatry*. 2016;87(12):1283-1286.
5
6 29. Rondina JM, Park C-H, Ward NS. Brain regions important for recovery after severe post-stroke upper
7 limb paresis. *J Neurol Neurosurg Psychiatry*. 2017;88(9):737-743.
8
9 30. Burke Quinlan E, Dodakian L, See J, et al. Neural function, injury, and stroke subtype predict treatment
10 gains after stroke. *Ann Neurol*. 2015;77(1):132-45.
11
12 31. Robertson IH, Ridgeway V, Greenfield E, et al. Motor recovery after stroke depends on intact sustained
13 attention: a 2-year follow-up study. *Neuropsychology*. 1997;11(2):290-5.
14
15 32. Quattrocchi G, Greenwood R, Rothwell JC, et al. Reward and punishment enhance motor adaptation
16 in stroke. *J Neurol Neurosurg Psychiatry*. 2017;88(9):730-736.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURE LEGENDS

Figure 1. Outcome scores for all patients. Each data point represents a single patient. Top row shows individual scores at admission, discharge, 6 weeks and 6 months after discharge. Bottom row shows the individual difference scores for admission to discharge, admission to 6 weeks post-discharge, and admission to 6 months post-discharge. Scores are shown for modified Fugl-Meyer (upper limb), Action Research Arm Test and Chedoke Arm and Hand Activity Inventory (CAHAI). Median (solid line) and upper and lower quartiles (dotted lines) are shown.

Figure 2. Self-reported outcome scores for all patients. Lower scores are better. Each data point represents a single patient. Top row shows individual scores at admission, discharge, 6 weeks and 6 months after discharge. Bottom row shows the individual difference scores for admission to discharge, admission to 6 weeks post-discharge, and admission to 6 months post-discharge. Scores are shown for the Arm Activity Measure part A and B (ArmA-A and ArmA-B). Median (solid line) and upper and lower quartiles (dotted lines) are shown.

Figure 3. Admission score plotted against final scores at 6 months post-discharge for Fugl-Meyer (FM-UL), Action Research Arm Test (ARAT), Chedoke Arm and Hand Activity Inventory (CAHAI), Arm Activity Assessment A & B (ArmA-A and ArmA-B). The black line represents the line of 'no change'. Patients above this line for FM-UL, ARAT and CAHAI have improved. Patients below this line for ArmA-A and ArmA-B have improved.

Figure 4. Change in scores from admission to 6 months post-discharge for Fugl-Meyer (FM-UL), Action Research Arm Test (ARAT), Chedoke Arm and Hand Activity Inventory (CAHAI), plotted against each other.

Table 1. Demographic and clinical characteristics on admission

Variable	Patients with complete data (n = 224)	Patients with missing data (n=14)	Difference between median values
Age, Median (IQR), yrs	52 (38-60)	54.5 (43-61)	P = 0.29
Gender, Male, n (%)	138 (61.6%)	7 (50%)	P = 0.30
Female, n (%)	86 (38.4%)	7 (50%)	
Affected Limb*			
Left, n (%)	117 (51.3%)	8(57.1%)	P = 0.39
Right, n (%)	107 (48.7%)	6 (42.9%)	
Non-dominant	110 (49.1%)	7 (50.0%)	P = 0.87
Dominant	114 (50.9%)	7 (50.0%)	
Median time since Stroke Months (IQR)	18 (12-51)	16 (12-22)	P = 0.29
Modified Rankin Scale, n (%)			
Median (IQR)	2 (2 - 3)	2 (2 – 2.75)	P = 0.32
No Significant Disability	20 (9.9%)	2 (14.3%)	
Slight Disability	106 (47.4%)	8 (57.1%)	
Moderate Disability	95 (42.4%)	4 (28.6%)	
Moderately Severe	3 (1.3)	0 (0%)	
The Barthel Index, Median (IQR), max = 20	19 (17-19)	19 (18-20)	P = 0.12
The Neurological Fatigue Index, Median (IQR), max = 62	35 (25 - 42)	33 (23-38)	P = 0.34
Hospital Anxiety and Depression Scale, Median (IQR), max = 34	12 (8-16)	13 (8.5-18)	P = 0.64
Sensory Loss, n, %			
0 = Severe	13 (5.8%)	2 (14.3%)	P = 0.36
1 = Mild	111 (49.6%)	7 (50%)	
2 = Normal	100 (44.6%)	5 (35.7%)	
Modified Fugl-Meyer (Upper Limb), Median (IQR) max = 54	26 (16-37)	27.5 (19-34)	P = 0.92
Action Research Arm Test, Median (IQR), Max = 57	18 (7-33)	15 (9.5-23)	P = 0.92
Chedoke Arm and Hand Activity Inventory, Median (IQR), Max = 91	40 (28-55)	33.5 (27-40)	P = 0.37
Arm Activity Measure A, Median (IQR), Max = 28	8 (5-12) [‡]	8 (6-12) ^{**}	P = 0.55
Arm Activity Measure B, Median (IQR), Max = 52	38 (24-46) [‡]	36 (29-41) ^{**}	P = 0.65

Difference in medians tested with Wilcoxon Signed Rank test. Difference in proportions was tested with Chi-squared test. Performed with ‡ 195 patients out of 224 had ArMA-A and ArMA-B measured on admission. † †11 patients out of 14 had ArMA-A and ArMA-B measured on admission. *202 (90.2%) patients were right handed, 22 (9.8%) patients were left handed.

Table 2. Upper limb scores before and after admission

	Admission	Discharge	6 weeks	6 months
FM-UL	26 (16 - 37)	34 (23 - 44)	35 (23 - 46)	37 (24 - 48)
ARAT	18 (7 - 33)	25 (10 - 42)	26 (11 - 44)	27 (11 - 45)
CAHAI	40 (28 - 55)	49 (36 - 70)	50 (35 - 73)	52 (35 - 77)
ArmA-A	8 (4.5 - 12)	5 (2 - 8)	4 (2 - 7)	3 (1 - 6.5)
ArmA-B	38 (24 - 46)	26 (13 - 37)	21 (11 - 35)	19 (8.5 - 32)

All scores given as median (IQR)

Table 3. Changes in upper limb scores

	Adm - Dis	Adm - 6 weeks	Adm - 6 months
FM-UL	6 (3 to 9)	8 (4 to 11)	9 (4 to 12)
ARAT	6 (2 to 8)	7 (2 to 10)	8 (3 to 11)
CAHAI	9 (5 to 13.5)	10 (5 to 16)	12 (6 to 17)
ArmA-A	-2 (-5 to -1)	-3 (-6 to -1)	-4 (-8 to -1)
ArmA-B	-7 (-14 to -3)	-10 (-17 to -4)	-11 (-19 to -5)

All scores given as median (IQR)

Table 4. Correlation between baseline scores and 6 month outcomes

	FM-UL (T4)	ARAT (T4)	CAHAI (T4)	ArmA-A (T4)	ArmA-B (T4)
Age	n.s.	n.s.	n.s.	n.s.	n.s.
Barthel Index	n.s.	n.s.	0.28	n.s.	n.s.
modified Rankin Scale	-0.29	-0.25	-0.32	n.s.	n.s.
Neurological Fatigue Index	n.s.	n.s.	n.s.	n.s.	n.s.
Hospital Anxiety and Depression Scale	n.s.	n.s.	n.s.	n.s.	n.s.
Sensation	n.s.	n.s.	0.22	n.s.	n.s.
FM-UL (T1)	0.92	0.84	0.83	-0.42	-0.63
ARAT (T1)	0.87	0.95	0.91	-0.48	-0.71
CAHAI (T1)	0.84	0.88	0.92	-0.46	-0.71
ArmA-A (T1)	-0.39	-0.44	-0.44	0.56	0.41
ArmA-B (T1)	-0.53	-0.58	-0.61	0.40	0.71

Correlation coefficient (rho) given only when significant at $p < 0.05$ corrected for 55 multiple comparisons ($p < 0.0009$). n.s. = not significant

Table 5. Correlation between baseline scores and change scores from admission to 6 months


	FM-UL (T1-T4)	ARAT (T1-T4)	CAHAI (T1-T4)	ArmA-A (T1-T4)	ArmA-B (T1-T4)
Age	n.s.	n.s.	n.s.	n.s.	n.s.
Barthel Index	n.s.	n.s.	n.s.	n.s.	n.s.
modified Rankin Scale	n.s.	n.s.	n.s.	n.s.	n.s.
Neurological Fatigue Index	n.s.	n.s.	n.s.	n.s.	n.s.
Hospital Anxiety and Depression Scale	n.s.	n.s.	n.s.	n.s.	n.s.
Sensation	n.s.	n.s.	n.s.	n.s.	n.s.
FM-UL (T1)	n.s.	0.40	0.45	n.s.	-0.24
ARAT (T1)	n.s.	0.36	0.52	n.s.	n.s.
CAHAI (T1)	n.s.	0.42	0.35	n.s.	n.s.
ArmA-A (T1)	n.s.	n.s.	-0.31	-0.72	n.s.
ArmA-B (T1)	n.s.	-0.26	-0.38	-0.28	-0.26

Correlation coefficient (rho) given only when significant at $p < 0.05$ corrected for 55 multiple comparisons ($p < 0.0009$). n.s. = not significant

Table 6: Correlations between changes in different outcome scores


	FM-UL (T1-T4)	ARAT (T1-T4)	CAHAI (T1-T4)	ArmA-A (T1-T4)	ArmA-B (T1-T4)
FM-UL (T1-T4)	-	0.32	0.34	n.s.	n.s.
ARAT (T1-T4)	0.32	-	0.37	n.s.	-0.25
CAHAI (T1-T4)	0.34	0.37	-	n.s.	-0.20
ArmA-A(T1-T4)	n.s.	n.s.	n.s.	-	0.29
ArmA-B (T1-T4)	n.s.	-0.25	-0.20	0.29	-

Correlation coefficient (rho) given only when significant at $p < 0.05$ corrected for 10 multiple comparisons ($p < 0.005$). n.s. = not significant




Outcome scores for all patients. Each data point represents a single patient. Top row shows individual scores at admission, discharge, 6 weeks and 6 months after discharge. Bottom row shows the individual difference scores for admission to discharge, admission to 6 weeks post-discharge, and admission to 6 months post-discharge. Scores are shown for modified Fugl-Meyer (upper limb), Action Research Arm Test and Chedoke Arm and Hand Activity Inventory (CAHAI). Median (solid line) and upper and lower quartiles (dotted lines) are shown.

600x250mm (300 x 300 DPI)




Self-reported outcome scores for all patients. Lower scores are better. Each data point represents a single patient. Top row shows individual scores at admission, discharge, 6 weeks and 6 months after discharge. Bottom row shows the individual difference scores for admission to discharge, admission to 6 weeks post-discharge, and admission to 6 months post-discharge. Scores are shown for the Arm Activity Measure part A and B (ArmA-A and ArmA-B). Median (solid line) and upper and lower quartiles (dotted lines) are shown.

399x250mm (300 x 300 DPI)



Admission score plotted against final scores at 6 months post-discharge for Fugl-Meyer (FM-UL), Action Research Arm Test (ARAT), Chedoke Arm and Hand Activity Inventory (CAHAI), Arm Activity Assessment A & B (ArmA-A and ArmA-B). The black line represents the line of 'no change'. Patients above this line for FM-UL, ARAT and CAHAI have improved. Patients below this line for ArmA-A and ArmA-B have improved.

299x199mm (300 x 300 DPI)



Change in scores from admission to 6 months post-discharge for Fugl-Meyer (FM-UL), Action Research Arm Test (ARAT), Chedoke Arm and Hand Activity Inventory (CAHAI), plotted against each other.

299x119mm (300 x 300 DPI)

Table S1. Significance of changes in outcome scores between each time point

	T1-T2	T1-T3	T1-T4	T2-T3	T3-T4	T2-T4
FM-UL	Z=11.9, p<0.0001	Z=11.8, p<0.0001	Z=12.1, p<0.0001	Z=5.8, p<0.0001	Z=8.1, p<0.0001	Z=4.5, p<0.0001
ARAT	Z=11.4, p<0.0001	Z=11.4, p<0.0001	Z=11.4, p<0.0001	Z =5.4, p<0.0001	Z=7.0, p<0.0001	Z=4.2, p<0.0001
CAHAI	Z=12.1, p<0.0001	Z=11.8, p<0.0001	Z=11.9, p<0.0001	Z=3.6, p=0.0003	Z=4.9, p<0.0001	Z=3.2, p=0.0014
ArmA-A	Z=10.1, p<0.0001	Z=10.0, p<0.0001	Z=10.4, p<0.0001	Z=2.6, p=N.S.	Z=5.6, p<0.0001	Z=3.4, p<0.0001
ArmA-B	Z=10.4, p<0.0001	Z=10.9, p<0.0001	Z=11, p<0.0001	Z=4.1, p<0.0001	Z=6.9, p<0.0001	Z=3.8, p=0.0002

Significance determined with Wilcoxon's signed ranks test