

## Archives of Disease in Childhood

### A survey of the definition and monitoring of neonatal hypoglycaemia in English Neonatal Units

Journal:	<i>Archives of Disease in Childhood</i>
Manuscript ID	fetalneonatal-2016-311473.R2
Article Type:	Letter (original)
Edition:	not in use
Date Submitted by the Author:	27-Aug-2016
Complete List of Authors:	Dixon, Kathryn; University of Cambridge, Medical School Ferris, Rebecca; University of Cambridge, Medical School Marikar, Dilshad; Peterborough and Stamford Hospitals NHS Foundation Trust, Paediatrics Chong, Mickey; University College London Medical School, Medical School Mittal, Anita; Bedford Hospitals NHS Trust, Paediatrics Manikam, Logan; University College London Institute of Child Health Rose, Peter; Peterborough and Stamford Hospitals NHS Foundation Trust, Paediatrics
Keywords:	hypoglycaemia, Neonatology, Endocrinology, General Paediatrics, Intensive Care

SCHOLARONE™  
Manuscripts

View Only

1  
2  
3 **Definition and monitoring of neonatal hypoglycaemia: A nationwide survey of NHS England**

4  
5 **Neonatal Units**

6  
7 Corresponding Author:

8  
9 Dilshad Marikar, Department of Paediatrics, Peterborough City Hospital, Edith Cavell Campus,  
10  
11 Bretton Gate, Peterborough, Cambridgeshire PE3 9GZ. Email: dilshad.marikar@nhs.net Tel:  
12  
13 07758892708

14  
15 Co-authors:

16  
17 Kathryn Charlotte Dixon, School of Clinical Medicine, University of Cambridge, Cambridge, UK

18  
19 Rebecca Louise Ferris, School of Clinical Medicine, University of Cambridge, Cambridge, UK,

20  
21 Dilshad Marikar, Department of Paediatrics, Peterborough City Hospital, Edith Cavell Campus,  
22  
23 Bretton Gate, Peterborough, Cambridgeshire PE3 9GZ. Email: dilshad.marikar@nhs.net Tel:  
24  
25 07758892708

26  
27 Mickey Chong, UCL Medical School, London, UK

28  
29 Anita Mittal, Department of Paediatrics, Bedford Hospital, Bedford, UK

30  
31 Dr Logan Manikam, MPH, Population, Policy and Practice, UCL Great Ormond Street Institute of Child  
32  
33 Health, London, UK

34  
35  
36 Peter John Rose, Department of Paediatrics, Peterborough City Hospital, Edith Cavell Campus,  
37  
38 Bretton Gate, Peterborough, Cambridgeshire PE3 9GZ

39  
40  
41  
42 Keywords:

43  
44 Neonatal, NICU, Intensive Care, Hypoglycaemia, Endocrinology

45  
46 Word Count: 464  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Neonatal hypoglycaemia is common; however there is ongoing controversy regarding its operational definition, with recent guidance from the Paediatric Endocrine Society (PES) recommending using significantly higher blood glucose thresholds and longer periods of monitoring than existing guidance [1].

A 2014 survey of Australasian neonatal units identified that the commonest clinical threshold for treatment of hypoglycaemia was 2.6 mmol/L, and some units utilised potentially unreliable near-side blood glucose monitoring devices [2]. The most recent published survey of British neonatologists showed wide variation in definitions of neonatal hypoglycaemia from <1 to <4mmol/L [3].

We surveyed neonatal units in England to assess current working practice on the definition and monitoring of neonatal hypoglycaemia.

All 161 NHS neonatal units in England were surveyed between April and August 2015 via telephone or online questionnaire.

Respondents were asked (1) if a hypoglycaemia guideline was used; (2) definition of hypoglycaemia; (3) methods used to monitor blood glucose; (4) whether hypoglycaemia was confirmed by laboratory samples (5) the number of satisfactory pre-feed blood sugar readings required to discontinue monitoring.

Responses were received from 84% of units (135/161). 1.5% (2/135) reported not having a formal hypoglycaemia guideline.

88.1% of units defined hypoglycaemia as <2.6 mmol/L (119/135). Values ranging from 2.0 to 3.0 mmol/L were used by 11.9% of respondents (16/135), with some units defining hypoglycaemia according to differing neonatal clinical characteristics or feed type (Table 1).

*Table 1: Definition of hypoglycaemia in surveyed English Neonatal Units*

Definition of hypoglycaemia (mmol/L)	Number of units	Percentage of units (%)
<2.0	5	3.7
<2.2	1	0.7
<2.4	1	0.7
<2.5	1	0.7

<2.6	119	88.1
<2.7	3	2.2
<3.0	2	1.5
<2.0 for breast-fed <2.5 for bottle-fed	1	0.7
<3.0 for diabetes in pregnancy <2.0 for other neonates	1	0.7
<4.0 for diabetes in pregnancy <2.6 for other neonates	1	0.7

Glucometers were used by 87.4% of units (118/135) and blood gas machines by 68.1% (92/135).

Both methods were used in 56.3% of units (76/135).

57.7% (78/135) of units confirmed low blood glucose values with a laboratory sample: of these, 34.6% (27/78) of units used a threshold of < 2.6 mmol/L, while 38.5%(30/78) used other thresholds, and 26.9% (21/78) reported that decisions were based on clinical judgement.

43.7% (59/135) of respondents required two and 45.9% (62/135) required three satisfactory pre-feed blood glucose readings to discontinue monitoring. The remaining 10.4% units (14/135) utilised other criteria such as a fixed monitoring period or clinical judgement.

Although the majority of units defined hypoglycaemia as <2.6 mmol/L , this survey identified variation in its definition and monitoring.

Point of care glucometers are the most common measurement method, and laboratory confirmation of low blood glucose readings using these devices is not universal. These findings are significant as there is evidence to suggest some point of care glucose testing devices may overestimate blood glucose levels [4].

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Current practice does not comply with recent guidance from the PES, which aims to help clinicians recognise persistent hypoglycaemia disorders using higher thresholds than existing guidance from the American Academy of Paediatrics[1]. Implementing such guidance may increase neonatal unit admissions, separation from parents and impact on breast feeding, while there is controversy regarding its effect on neurodevelopmental outcomes[1]. The feasibility of adopting nationwide consensus guidelines should be explored.

### References

1. Adamkin, D., & Polin, R. (2016). Imperfect advice: Neonatal hypoglycemia. *The Journal of pediatrics*. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/27297210>
2. Harris, D.L., Weston, P.J., Battin, M.R. and Harding, J.E. (2009) 'A survey of the management of neonatal hypoglycaemia within the Australian and New Zealand neonatal network', *Journal of Paediatrics and Child Health*, 50(10), pp. E55–E62. doi: 10.1111/j.1440-1754.2009.01599.x.
3. Koh, T. and Vong, S. (1996) 'Definition of neonatal hypoglycaemia: Is there a change?', *Journal of paediatrics and child health.*, 32(4), pp. 302–5.
4. Roth-Kleiner M, Stadelmann Diaw C, Urfer J, Ruffieux C, Werner D. Evaluation of different POCT devices for glucose measurement in a clinical neonatal setting. *European Journal of Paediatrics*. 2010; 169(11): 1387-95.

Acknowledgements: The authors would like to thank the neonatal units that participated in the survey.

Contributors: KCD, RLF and DM are joint first authors. DM and AM designed the study and methodology. KCD, RLF, MC and DM conducted the survey. KCD, RLF and DM analysed the data. All authors wrote, edited or approved the final submitted report. DM is guarantor.

Competing interests: None.