

1 **The role of redox dysregulation in the inflammatory response to acute myocardial**
2 **ischaemia-reperfusion injury - adding fuel to the fire**

3 Sauri Hernández-Reséndiz^{a, b, c}, Kroekkiat Chinda^d, Ong Sang Bing^{a, b}, Hector Cabrera-Fuentes^{a, b,}
4 ^e, Cecilia Zazueta C^c, Derek J. Hausenloy^{a, b, f, g, h, i}

5
6 ^aNational Heart Research Institute Singapore, National Heart Centre Singapore, Singapore

7 ^bCardiovascular and Metabolic Disorder Programme, Duke-NUS Medical School, Singapore

8 ^cDepartment of Cardiovascular Biomedicine, National Institute of Cardiology I. Ch, Mexico, Mexico
9 City

10 ^dDepartment of Physiology, Faculty of Medical Science, Naresuan University, Phitsanulok, Thailand

11 ^eInstitute of Biochemistry, Medical School, Justus-Liebig-University, Giessen, Germany

12 ^fYong Loo Lin School of Medicine, National University Singapore, Singapore

13 ^gThe Hatter Cardiovascular Institute, University College London, UK.

14 ^hThe National Institute of Health Research University College London Hospitals Biomedical
15 Research Centre, London, UKi

16 ⁱBarts Heart Centre, St Bartholomew's Hospital, London, UK.

17
18
19
20
21
22 **Corresponding author:**

23 Prof Derek Hausenloy

24 Cardiovascular and Metabolic Disorders Program,

25 Duke-National University of Singapore, 8 College Road Medical School

26 Singapore 169857.

27 Tel +65 66015121

28 E-mail: derek.hausenloy@duke-nus.edu.sg

29 **Abstract**

30 The inflammatory response to acute myocardial ischaemia/reperfusion injury (IRI) plays a critical
31 role in determining myocardial infarct (MI) size and subsequent post-MI left ventricular (LV)
32 remodeling, making it a potential therapeutic target for treating patients presenting with an acute
33 myocardial infarction (AMI). Recent experimental studies using advanced imaging and molecular
34 techniques have yielded new insights into the mechanisms through which reactive oxygen species
35 (ROS) contribute to the inflammatory response during acute myocardial IRI - "*adding fuel to the*
36 *fire*". The infiltration of inflammatory cells into the MI zone, leads to elevated myocardial
37 concentrations of ROS, cytokine release, and activation of apoptotic and necrotic death pathways.
38 Anti-oxidant and anti-inflammatory therapies have failed to protect the heart against acute
39 myocardial IRI. This may be, in part, to a lack of understanding of the time course, nature and
40 mechanisms of the inflammation and redox dysregulation which occur in the setting of acute
41 myocardial IRI. In this article, we will examine the inflammatory response and redox dysregulation
42 induced by acute myocardial IRI, and highlight potential therapeutic options for targeting redox
43 dysregulation in order to attenuate the detrimental effects of the inflammatory response following an
44 AMI so as to reduce MI size and prevent heart failure.

45

46 **Keywords**

47 Myocardial ischaemia/reperfusion injury, Redox dysregulation, Inflammation, Reactive Oxygen
48 Species, Oxidative stress, Neutrophils.

49

50

51

52

53

54

55

56

| | |
|----|--|
| 57 | Contents |
| 58 | 1. Introduction |
| 59 | 2. Oxygen Paradox and ROS formation |
| 60 | 3. Biological roles of ROS in the heart |
| 61 | 4. Sources of ROS and interplay with inflammation during acute myocardial IRI |
| 62 | a) <i>Cytochrome P-450</i> |
| 63 | b) <i>Xanthine oxidase</i> |
| 64 | c) <i>NADPH oxidases</i> |
| 65 | d) <i>Monoamine oxidases</i> |
| 66 | e) <i>Mitochondrial electron transport chain</i> |
| 67 | f) <i>Folding machinery in the sarco-endoplasmic reticulum (SR/ER)</i> |
| 68 | g) <i>Nitric oxide synthase</i> |
| 69 | 5. Dysregulation of myocardial antioxidant pathways during acute myocardial IRI |
| 70 | 6. The acute-phase response: tissue damage, inflammatory response and more ROS |
| 71 | a) <i>How ROS induce the inflammatory response</i> |
| 72 | 7. Physiological consequences of the inflammatory response |
| 73 | a) <i>No-reflow</i> |
| 74 | b) <i>Post-MI left ventricular remodeling</i> |
| 75 | 8. Therapeutic targeting of ROS and inflammation |
| 76 | 9. Conclusion |
| 77 | |
| 78 | |
| 79 | |
| 80 | |
| 81 | |
| 82 | |
| 83 | |
| 84 | |

85 **1. Introduction**

86 Acute myocardial infarction (AMI) and the heart failure which often ensues are one of the leading
87 causes of death and disability worldwide. For patients presenting with an AMI, the treatment of
88 choice is to restore coronary blood flow in the infarct-related artery, to salvage viable myocardium.
89 However, despite optimal therapy the morbidity and mortality of AMI patients remain significant with
90 7% death and 25% heart failure at one year [1]. The reason for this, is in part, due to the presence
91 of 'myocardial reperfusion injury' which refers to the myocardial injury and cardiomyocyte death,
92 that is paradoxically induced by myocardial reperfusion, and which can contribute up to 50% of the
93 final myocardial infarct (MI) size, and for which, there is currently no effective therapy [2] [3]. As
94 such, novel therapies are required to protect the heart against acute myocardial
95 ischaemia/reperfusion injury (IRI), in order to reduce MI size and prevent heart failure.

96 The inflammatory response to acute myocardial IRI plays a critical role in determining MI
97 size and subsequent post-MI left ventricular (LV) remodelling, making it a potential therapeutic
98 target for preventing heart failure following AMI. Experimental studies using molecular techniques
99 and advanced biomedical imaging have yielded new insights into the mechanisms through which
100 reactive oxygen species (ROS) contribute to the inflammatory response during acute myocardial
101 IRI, "*adding fuel to the fire*". The infiltration of inflammatory cells into the MI zone, leads to elevated
102 concentrations of ROS in the myocardium, cytokine release, and the activation of apoptotic and
103 necrotic death pathways. The complex interplay between ROS and inflammation can amplify the
104 effects of ROS as mediators of myocardial injury and determinants of cell death. As such, ROS
105 represent important therapeutic targets for reducing MI size and preventing adverse LV remodelling
106 in AMI patients. In this review article, we highlight the complex interplay between ROS and
107 inflammation in the setting of acute myocardial IRI, and explore emerging therapeutic targets for
108 attenuating ROS and modulating the inflammatory response in patients presenting with AMI.

109

110 **2. Oxygen Paradox and ROS formation**

111 When the myocardium is re-oxygenated after a prolonged period of energy depletion, it rapidly
112 hypercontracts. The hypercontracture and cytolysis induced by reoxygenation have become known

113 as the “Oxygen Paradox” [4], [5], [2]. Bresnahan et al. [6], demonstrated in a canine model that the
114 potentiation of haemorrhage and extension of myocardial infarction is attributable to the re-
115 administration of molecular oxygen. Using the isolated perfused rat heart, in 1973, Hearse and
116 Chain showed that the reoxygenation kills the heart cells and exacerbates cardiac enzyme creatine
117 phosphokinase (CPK), ATP, AMP phosphotransferase (MK) and glyceraldehyde-3-phosphate
118 dehydrogenase (GAPDH) release. The myocardial injury was not identified. However the possible
119 responsible listed was ROS overproduction [4].

120 Following AMI, ROS are generated in the first minute of reperfusion, and peak 4-7 minutes
121 later, although ROS production continues at lower sustained levels for quite some time after [7].
122 Oxidative stress or redox dysregulation occurs in the myocardium when ROS production is
123 enhanced, and the anti-oxidant reserve is exhausted. This highly reactive and unstable group of
124 compounds are formed as a result of the addition of an unpaired electron in the outer orbit of the
125 molecule. Superoxide (O_2^-), hydrogen peroxide (H_2O_2), and the highly reactive hydroxyl radical
126 ($\bullet OH$) are the most prominent free radicals in the pathogenesis of acute myocardial IRI. In the
127 presence of iron, superoxide and H_2O_2 can lead to the formation of highly reactive $\bullet OH$, which can
128 damage cellular proteins, RNA, DNA and lipids. Interaction of ROS with nitrogen monoxide ($NO\bullet$)
129 [8] or fatty acids can produce peroxynitrite or peroxy radicals, respectively. The first ROS produced
130 in response to acute IRI is O_2^- , resulting from the univalent reduction of molecular oxygen.
131 Dismutation of O_2^- produces H_2O_2 , which, in turn, may be entirely reduced to water or partially
132 reduced to $\bullet OH$, one of the strongest pro-oxidants in nature. Also, O_2^- may react with nitric oxide in
133 a reaction controlled by the rate of diffusion of both radicals [9] [10] [11] (fig. 1). The generation of
134 ROS has been connected to stress responses, apoptosis, aging and death. However, the ROS are
135 now being recognized as molecules involved in the cardiac adaptation to different types of
136 physiological stimuli [12] [13] [14] [15] [16].

137

138 **3. Biological roles of ROS in the heart**

139 The most recognized ROS with physiological effects includes the O_2 , $NO\bullet$ and the non-radical
140 specie H_2O_2 . There have been implicated in the regulation of inflammation [17] [18] [19], calcium
141 signaling [20], hypertrophy [21], autophagy [22] and cardioprotection [23].

142 Cysteine (Cys) and methionine (Met) possess reactive sulfur-containing side chains that
143 present targets for ROS [16] [24]. Oxidation of these specific and reactive residues, in turn can, lead
144 to the reversible modification of enzymatic activity. Four oxidation states of Cys can be generated:
145 disulfide (-S-S), sulfenic acid (-SOH) and sulfonic acid (-SO₃H) [16]. Sulfenic acid is readily reduced
146 to cysteine by the cellular reducing agents, glutathione (GSH) and thioredoxin (Trx) [25] [26].
147 Methionine is oxidized to methionine sulfoxide (MetO) by the addition of an extra oxygen atom [27].
148 The Anderson laboratory found that following initial Ca^{2+} -dependent activation of CaMKII
149 (Ca^{2+} /calmodulin-dependent kinase II), the specific oxidation of conserved Met 281/282 residues in
150 the regulatory domain could increase CaMKII activity independent of Ca^{2+} /calmodulin [28].

151 Substantial evidence has revealed that H_2O_2 production has been shown to be a major
152 component of endothelium-derived hyperpolarizing factor to control blood pressure [29] [30]. H_2O_2
153 also caused interprotein disulfide bond in protein kinase G (PKG) which activated the kinase
154 independently of the $NO\bullet$ -cyclic guanosine monophosphate (cGMP) pathway and coupled to
155 vasodilation [31]. In a redox-dead Cys42Ser PKGI- α knock-in mouse, Prisyazhna et al.,
156 demonstrated that H_2O_2 induce an oxidation and activation of PKG which cause vessel
157 hyperpolarization and relaxation [32]. Also, the treatment of endothelial cells or aortic vessels with
158 vascular endothelial growth factor (VEGF) induced growth signaling and angiogenesis dependent of
159 protein kinase A (PKA) oxidation [33].

160 Modulation of the redox potential of reactive thiols may be a general control mechanism by
161 which sarcoplasmic/endoplasmic (SR/ER) reticulum and ryanodine receptor (RyR) controls
162 cytoplasmic Ca^{2+} concentrations in the skeletal muscle [34] and myocardium [35]. Yi X et al.,
163 demonstrated in coronary artery smooth muscle that a local NADPH oxidase system on SR/ER
164 regulates RyR/ Ca^{2+} channel activity and Ca^{2+} release from SR/ER by producing O_2 [20]. The thiol-
165 disulfide exchange model in cardiac muscle has been proposed to describe the mechanism by
166 which O_2 can directly activate the RYR/ Ca^{2+} . In this model, intermolecular thiol-disulfide

167 interexchange reaction within RyR control open or closed states of its Ca²⁺ release channels. When
168 the thiol groups of RyR is in a reduced status (-SOH form), the channel is closed. In contrast, the
169 channel is open when disulfide is formed by oxidation of thiol groups of RyR (-S-S-form) [36] [37]
170 [38] [39].

171 In vascular smooth muscle (VSM), angiotensin II increases NADPH oxidase-dependent
172 ROS production, which is thought to activate signaling pathways involved in the hypertrophic
173 response [40] [41]. The redox signaling modulation of the small G proteins Ras kinases such as
174 ERK1/2, p38MAPK, protein kinase C (PKC) and Akt contribute to the development of GPCR
175 agonist-induced hypertrophy [31] [42].

176 Hydrogen peroxide can induce kinase activation via tyrosine phosphorylation or via the
177 induction of the released zinc from zinc-finger domains of PKC [43]. It has been proposed that O₂
178 and H₂O₂ may play an important signaling role in cardioprotection. Most of the signaling pathways
179 of cardioprotection converge at the mitochondria and the mitochondrial ROS formation mediates
180 signal transduction through post-translational modifications of redox-sensitive proteins [44] [45] [46].
181 Perrelli et al., demonstrated for the first time that the cardioprotective effect of catestatine as a
182 pharmacological postconditioning (CST-Post) depends on the activation of PI3K/Akt, PKCs and
183 mitoKATP channels, which may include a ROS signaling [23].

184 It may seem paradoxical that ROS are essential for promoting normal cellular processes, as
185 opposed to having a toxic effect on the heart. Even cell death that was previously thought to result
186 from oxidative damage is now considered to be the result of ROS triggering a physiological pathway
187 for cell death. Maintaining a basal level of ROS which is above a cytostatic level, but below
188 cytotoxic, therefore enables proper redox biology reactions and the regulation of numerous
189 processes essential for life.

190

191 **4. Sources of ROS and the interplay with inflammation during acute myocardial IRI**

192 A number of different mechanisms and sources are known to underlie ROS generation in the
193 myocardium in the setting of acute IRI. The enzymes systems most commonly implicated in ROS
194 production are cytochrome P-450 (CYP), xanthine oxidase, NADPH oxidase, monoamine oxidases

195 (MAO), uncoupled nitric oxide synthase (NOS), the unfolded-protein response (UPR)-regulated
196 oxidative protein folding machinery in the SR/ER, and the mitochondrial electron transport chain
197 [\[47\]](#) (fig. 2A).

198

199 **a. Cytochrome P-450**

200 The cytochrome P-450 (CYP) family of proteins are mono-oxygenases, which catalyse the oxidation
201 of hydrophobic organic molecules mainly in the liver, but also in the heart [\[48\]](#). The CYP system is
202 known to be a potential source of ROS following reperfusion of the acutely ischaemic myocardium,
203 mainly from endothelial cells [\[49\]](#), macrophages, and neutrophils [\[50\]](#). It has already been shown
204 that ROS can arise from the decay of oxygenated CYP intermediates produced during the catalytic
205 mechanism of mixed-function oxidation. The contents of ROS derived from cytochrome P-450 have
206 been shown to increase in an oxygen concentration-dependent manner as CYP generates O_2 and
207 H_2O_2 through an uncoupling reaction. It is conceivable that the increase in ROS produced by CYP
208 upon reperfusion are due to an increase in uncoupling, concomitant with the increment of oxygen
209 supply to myocardium [\[51\]](#).

210 Members of the cytochrome P-450 2-epoxygenases family (CYP 2), primarily 2C8, 2C9 and
211 2J2, and the hydroxylase CYP 4F are capable of metabolising endogenous arachidonic acid (AA)
212 into vasoactive products such as epoxyeicosatrienoic acids (EETs) and hydroxyeicosatetraenoic
213 acids (HETEs). Although EETs have been reported to play a cardioprotective role, CYP 2C9 can
214 also generate O_2 , H_2O_2 , and $\bullet OH$ during the CYP reaction cycle [\[52\]](#). Using a rat Langendorff
215 preparation, Granville et al., showed that CYP 2C9 is a potent source of ROS during acute
216 myocardial IRI, and contributes to the extension of MI size [\[53\]](#). The O_2 and H_2O_2 produced by CYP
217 2C9 can trigger $NF\kappa B$ activation resulting in the upregulation and secretion of pro-inflammatory
218 cytokines and adhesion molecule expression [\[52\]](#). The selective CYP 2C9 inhibition with
219 sulfaphenazole has been reported to result in a significant reduction in MI size after 2 hours of
220 reperfusion in a rat acute IRI model (fig. 2B) [\[53\]](#).

221 Over-expression of endothelial CYP 2C8 has been shown to increase ROS generation and
222 leukotoxin diols formation, thereby augmenting coronary vasoconstriction and increasing MI size

223 [49] [53]. During acute myocardial ischaemia, AA accumulates, leading to increased generation of
224 20-HETE (20-hydroxy-5,8,11,14-eicosatetraenoic acid) through CYP 4F [54]. 20-HETE acts directly
225 on cardiomyocytes via the stimulation of NADPH oxidase-derived ROS production, and induces
226 cardiomyocyte apoptosis. The treatment of endothelial cells with endogenous 20-HETE leads to an
227 increase in NF κ B activity and endothelial activation, characterised by the increased expression of
228 intracellular adhesion molecules and interleukin-8 (IL-8) levels [55] [56] [57]. Inhibition of ROS
229 production during acute IRI may be more beneficial than a free radical scavenger because such
230 anti-oxidants must compete with cellular targets to protect tissue from ongoing ROS production. For
231 example, the administration of cimetidine upon reperfusion has been demonstrated to reduce MI
232 size, prevent cardiac dysfunction, and attenuate ROS production in the ischaemic region [58]. The
233 inhibition of 20-HETE with HET0016 (N-hydroxy-N'-(4-butyl-2-methylphenyl)-formamidine) prevents
234 the activation of inflammatory genes and the endothelial dysfunction [56] [55]. The selective
235 hydroxylase inhibition with N-methylsulfonyl-12, 12-dibromo-11-enamide (DDMS) 10 minutes before
236 coronary artery occlusion or 5 minutes before reperfusion was found to reduce MI size [59] [60].
237 These data suggest that the inhibition of CYP hydroxylases may induce cardioprotection. However,
238 further studies are warranted to determine whether pharmacological interventions that disrupt CYP
239 2C and CYP 4F signalling prevent the development of inflammation associated with acute
240 myocardial IRI.

241

242 ***b. Xanthine oxidase***

243 Xanthine oxidoreductase catalyses the oxidation of hypoxanthine to xanthine and the latter to uric
244 acid as the final steps of purine degradation [61]. Xanthine oxidoreductase has the peculiar property
245 of existing in two interconvertible forms, xanthine oxidase (XO) and xanthine dehydrogenase (XDH).
246 XO is formed from XDH under ischaemic conditions and upon myocardial reperfusion [62]. It can
247 react with purine substrates (hypoxanthine or xanthine) and O₂ as the terminal electron acceptor,
248 thereby exhibiting the ability to generate O₂⁻ and H₂O₂ [63]. The ·OH and O₂⁻ radicals produced by
249 the enzyme can, in turn, react with cellular proteins and membranes causing cellular injury. XO is
250 present predominantly in the vascular endothelium in the healthy heart, and has been implicated as

251 a primary source of cytotoxic ROS. This is largely based on the observation that allopurinol, an
252 inhibitor of xanthine oxidoreductase, is as effective as an oxygen radical scavenger in attenuating
253 the tissue injury associated with acute IRI [64]. Allopurinol has been shown to decrease MI size
254 and improved the recovery of LV function following acute IRI [65]. Oxypurinol was found to increase
255 cardiac output and improve regional LV function after sustained coronary artery occlusion in the
256 canine heart [66]. Pre-treatment with the XO inhibitor, allopurinol, is effective in inhibiting generation
257 of ROS during reperfusion and improving recovery of LV function [67].

258 XO has also been implicated in the leukocyte recruitment that occurs during reperfusion.
259 Leukocyte-endothelial cell adhesion in post-ischaemic models, and increased neutrophil adhesion
260 after hypoxia have been reported to be significantly attenuated by XO inhibitors [68]. However, one
261 problem inherent to the use of allopurinol is that its therapeutic effect is not dose dependent: at
262 higher doses, it becomes the substrate for XO, which will in turn produce O_2 and thus exacerbate
263 myocardial damage. ROS generated by XO promote the formation of pro-inflammatory stimuli,
264 modify the expression of adhesion molecules on the surface of leukocytes and endothelial cells,
265 and reduce levels of the potent anti-adhesive agent nitric oxide. This latter effect is exacerbated by
266 the decline in nitric oxide synthase (NOS) activity and oxidation of soluble guanylyl cyclase during
267 reperfusion, which serves to amplify the intense inflammatory response [69]. Based on these
268 observations it has been proposed that XO plays an important role in mediating the reperfusion
269 injury response by promoting the recruitment and activation of leukocytes [67], [70].

270

271 **c. NADPH oxidases**

272 The NADPH oxidases (NOX) family comprises seven members, five NOX and two dual oxidases
273 (Duox-1 and Duox-2) [71], [72]. They contain six or seven transmembrane spanning domains,
274 respectively. NADPH oxidase catalyses electron transport from NADPH to molecular oxygen,
275 thereby producing ROS [73]. Among these isoforms, NOX3 is highly expressed in the cochlea [74];
276 NOX1 is expressed in endothelial cells, VSMC and adventitial fibroblasts [75]. NOX2 and NOX 4
277 are abundantly expressed in cardiomyocytes [72]. NOX5 is located in vascular endothelial cells

278 [76], and vascular smooth muscle cells [72], [77] and Duox-1 and Duox-2 are predominantly
279 expressed in epithelial cells [78].

280 The proposition that NOX enzymes contribute to acute myocardial IRI is based on two
281 experimental strands of evidence: (1) the increased expression and activity of NOX in the post-
282 ischaemic myocardium and (2) the attenuation of ROS following pharmacologic inhibition of NOX.
283 Meischl et al. demonstrated that NOX2 is the predominant isoform that is expressed in
284 cardiomyocytes, and its expression is upregulated in response to acute IRI [79]. The use of
285 apocynin and diphenylene iodonium (DPI) (non-specific NOX inhibitors) has been found to reduce
286 the increase in lipid peroxidation, cell death, and apoptosis after simulated IRI in cardiac cells [80].

287 NOX can also indirectly cause damage by enhancing the inflammatory response.
288 Neutrophils that express NOX2 are the primary source of ROS in acute IRI [81], [82]. Some studies
289 have shown that a phagocyte-like NADPH oxidase is the primary source of O₂ in vascular tissue
290 [83]. The potential involvement of neutrophils is supported by the observation that the time course
291 of the inflammatory cell accumulation corresponds with ROS generation and the MI size following
292 acute IRI. The activation of NOX in neutrophils is triggered via PKC-mediated phosphorylation of
293 cytosolic p47phox (for neutrophil cytosolic factor 1), which then binds to membrane-associated
294 gp91phox [84]. ROS generated by NADPH oxidase promote the formation of proinflammatory
295 stimuli, modify the expression of adhesion molecules on the surface of leukocytes and endothelial
296 cells, and reduce levels of the potent anti-adhesive agent nitric oxide. Coincident with these
297 changes, perivascular cells become activated and release another inflammatory mediators such as
298 tumor necrosis factor alpha (TNF- α) and cytokines [85], [86]. The regulation of different cytokines in
299 various organs suggest a cell-specific or organ-specific effect of NOX2. However, with respect to
300 other NOX isoforms, no solid data on their involvement in inflammation and chemotaxis after
301 reperfusion are available.

302

303 ***d. Monoamine oxidases***

304 Monoamine oxidases (MAOs) are flavoenzymes located within the mitochondrial outer membrane,
305 responsible for the oxidative deamination of neurotransmitters and dietary amines [87], [88].

306 Monoamine oxidase A (MAO-A) and B (MAO-B) share 70% amino acid identity, and both contain a
307 covalently bound FAD cofactor attached to an enzyme cysteine via the $\delta\alpha$ -methylene of the
308 isoalloxazine ring [89]. This flavin moiety is the only redox-dependent factor necessary for their
309 activity. The reaction of oxidative deamination occurs in several steps, ultimately resulting in the
310 formation of the aldehyde from the corresponding amine, ammonia and H_2O_2 . MAOs catalyse
311 oxidative deamination of several monoamines (serotonin [5-hydroxytryptamine (5-HT)],
312 noradrenaline, dopamine), resulting in significant ROS production [88]. Recent studies suggest that
313 MAOs contribute to increasing H_2O_2 production and catecholamine release in the early reperfusion
314 period (5-15 minutes) [90]. MAO-A generated H_2O_2 in acute IRI induces sphingosine kinase
315 inhibition, ceramide accumulation, and sphingosine-1-phosphate degradation in cardiomyocytes
316 thereby leading to mitochondria-mediated apoptosis in H9c2 cells [91]. Currently, efforts are
317 underway to investigate the mechanisms underlying the protective effect of MAO inhibitors
318 (selegiline, D-Deprenyl), and the roles of MAO in the setting of acute IRI [92], [93].

319

320 ***e. Mitochondrial electron transport chain***

321 Mitochondria have been implicated as a major source of ROS in acute myocardial IRI. The rapid
322 movement of electrons through the electron transport chain (ETC) of the inner mitochondrial
323 membrane can result in the leakage of electrons, which form O_2^- via univalent reduction of O_2 . All of
324 the ETC complexes have been implicated as both sources and targets of the ROS generated during
325 myocardial IRI, although most evidence supports a role for complexes I and III.

326 Mitochondrial complex I is viewed as a major contributor of ROS [11]. Oxidative impairment
327 of complex I is detected in rat models of acute myocardial IRI [94]. Mitochondrial complex I has two
328 catalytically and structurally distinct forms; one the fully competent, active A-form and the other, the
329 deactivated, D-form. The reversible D-form of complex I predominates under ischaemic conditions,
330 produces O_2^- and H_2O_2 , and may potentially increase the susceptibility of mitochondria to oxidative
331 damage [95]. Reperfusion also induces disruption of complex II; Chen et al. found that ADP-
332 stimulated state 3 respiration driven by succinate was 50% impaired in mitochondria from
333 reperfused hearts, a finding which was attributed to the impairment of complex II. The

334 deglutathionylation of complex II predisposes the 70-kDa flavin binding subunit to oxidative stress
335 induced by ROS during reperfusion injury [96].

336 Complex III is also considered an important source for mitochondrial ROS production in
337 reperfused hearts. In the ischaemic heart, mitochondrial complex activity is reduced by 22%
338 compared with healthy hearts. Increase unstable semiquinone radical ($\cdot Q^-$), is attributed to be the
339 source of O_2^- . Mammalian complex III contains bound cardiolipin molecules that are essential for
340 the catalytic function. The impairment of complex III activity due to the ROS-induced cardiolipin
341 oxidative damage may increase the electron leak from the electron transport chain, generating
342 more O_2^- and perpetuating a cycle of oxygen radical-induced damage, which ultimately leads to an
343 increase in MI size [97]. The burst of ROS from mitochondrial complexes induce the oxidation of
344 cholesterol and the production of oxysterols. Oxysterols can induce interleukin-1 beta (IL-1 β)
345 secretion in vascular endothelial cells and, consequently, the expression of adhesion molecules
346 necessary for the recruitment of immune cells [98]. Additionally, CYPs have been found in the
347 mitochondria of diverse animal species. Mitochondrial CYPs are proteins bound to the inner
348 membrane, and receive electrons for monooxygenation reaction from NADPH via adrenodoxin and
349 NADPH-adrenodoxin reductase [99]. Mitochondrial CYP catalyses the conversion of cholesterol in
350 pregnenolone and play essential roles in cholesterol homeostasis and steroid hormone biosynthesis
351 [100]. Recently, it has been shown that myocardial reperfusion induces mitochondrial cholesterol
352 accumulation. Peradis et al. showed that acute myocardial IRI produces high cholesterol and
353 oxysterol concentrations in the matrix and a simultaneous decrease in mitochondrial membrane
354 fluidity related to oxidative stress [101]. In this setting, the oxysterols 5, 6-epoxycholesterol, 7 β -
355 hydroxycholesterol, 7-ketocholesterol and 25-hydroxycholesterol exert a potent cytotoxic effect by
356 their ability to induce inflammatory effects [102].

357 Liu et al. have demonstrated that the oxysterol, 25-hydroxycholesterol, enhances IL-8
358 production [103]. It is noteworthy to mention that IL-8 is a cytokine which might play an important
359 role in the recruitment of T lymphocytes and monocytes into the arterial subendothelial space [104].
360 As summarised in figure 2C, by inhibiting cholesterol uptake into mitochondria at reperfusion with
361 4'-chlorodiazepam, the accumulation of oxysterols can decrease the inflammatory response and

362 induce cardioprotection. Further investigation is required to explore in more detail the relationship
363 between oxysterols and inflammation in the setting of acute IRI (fig. 2C).

364

365 ***f. UPR-regulated oxidative protein folding machinery in the SR/ER***

366 The cardiomyocyte sarco/endoplasmic reticulum (SR/ER) is an intracellular organelle specialising in
367 the regulation of Ca²⁺ fluxes and different oxidative functions. There is increasing evidence that
368 SR/ER stress plays a crucial role in IRI-induced cell dysfunction. Oxygen starvation during
369 ischaemia and ROS and Ca²⁺ overload during reperfusion results in SR/ER stress and the
370 activation of the pro-inflammatory pathway.

371 Mitochondria and SR/ER are in close apposition and the interface, commonly known as the
372 mitochondrial-associated SR/ER membrane, is believed to act as the focal point for signaling [105].
373 During acute myocardial IRI, both the release and uptake of calcium from the SR/ER are
374 dysregulated, resulting in enhanced Ca²⁺ release [46]. Much of the calcium is taken up by the
375 mitochondria and Ca²⁺ within the mitochondria, and induces the superoxide formation. Several *in*
376 *vitro* studies have demonstrated that the calcium pump on the SR/ER membrane is quite sensitive
377 to oxidative stress and the fact that the SR/ER contains a large amount of lipids and that it produces
378 ROS could also make this organelle very easily damaged by ROS. This vicious cycle of Ca²⁺
379 leakage, calcium overload and ROS generation inhibits cardiac contractility.

380 SR/ER stress initiates the activation of the unfolded protein response (UPR). The UPR
381 increases the capacity of the protein folding machinery resulting in the production of more oxidative
382 equivalents, and further deteriorating the redox state. UPR can induce TNF- α production in
383 response to ER stress through IRE1 α (inositol-requiring transmembrane kinase and endonuclease
384 1 α) and the ER-localised protein kinase PERK pathway [106] [107]. PERK-induced translational
385 arrest leads to the loss of I κ B, thereby activating NF κ B [108]. In addition, the phosphorylation of
386 IRE1 α in response to stress induces a conformational change in its cytosolic domain, which can
387 then bind to the adaptor protein, TNF- α -receptor-associated factor 2 (TRAF2), the receptor that can
388 activate canonical NF κ B JNK MAPK signaling pathway [109]. The efflux of Ca²⁺ from the SR/ER
389 generates ROS and NF κ B activation and gene expression that drive inflammation [110] [111]. NF κ B

390 induction by SR/ER stress is prevented by pre-incubation of cells with intracellular Ca²⁺ chelators,
391 suggesting that Ca²⁺ release precedes ROS formation in the NFκB-mediated SR/ER-nuclear signal
392 transduction pathway [112]. Several studies have demonstrated that hypoxia/reoxygenation is
393 sufficient to induce SR/ER stress [113] [114]. NFκB acts as a link between SR/ER stress and
394 inflammation after hypoxia/reoxygenation. NFκB inhibitors can protect cells against IRI by
395 selectively inhibiting the translocation of NFκB to the nucleus. In this regard, Wu et al. have shown
396 that SN50 can effectively can reduce damage to cardiomyocyte after reoxygenation [107].

397

398 ***g. Nitric oxide synthase***

399 Neuronal nitric oxide synthase (nNOS), inducible NOS (iNOS) and endothelial NOS (eNOS)
400 generate NO• via the oxidation of L-arginine. NOS isoforms contain both an oxygenase and
401 reductase domain. The reductase domain contains flavin adenine dinucleotide (FAD) and flavin
402 mononucleotide (FMN), and binds NADPH, while the oxygenase domain contains heme and
403 tetrahydrobiopterin (BH₄), and binds arginine. Uncoupling of NOS results in the loss of NO•
404 production, and O₂ production. A recent study by Lin et al. demonstrated that phosphorylation of
405 eNOS at threonine 497 mediated the switch between NO• production to superoxide generation
406 [115].

407 However, the most prominent cause of NOS uncoupling is the loss of the critical NOS co-factor,
408 BH₄, either by oxidation or decreased expression of the recycling enzyme dihydrofolate reductase
409 (DHFR) [116]. BH₄ depletion is involved in both endothelial and cardiomyocyte dysfunction in
410 hearts following acute IRI. Myocardial levels of BH₄ levels have been shown to be markedly
411 decreased after 60 minutes of reperfusion, and NOS uncoupling occurs with the increase in
412 myocardial O₂ formation. When electron flow is uncoupled from arginine oxidation, the reduced O₂
413 is released from the heme as O₂.

414 Endothelial NOS is mostly expressed in endothelial cells, cardiomyocytes and platelets. eNOS
415 synthesises NO• in a pulsatile manner with eNOS activity markedly increasing when intracellular
416 Ca²⁺ is increased [117]. Recent evidence has indicated that reversing NOS uncoupling in acute
417 myocardial IRI may be a therapeutic strategy [118]. Also, several studies have demonstrated that

418 low levels of heart BH4 result in myocardial inflammation. Zsuzsna et al. demonstrated that
419 plasma BH4, TNF α - and IL-6 levels showed an inverse correlation with the absolute values of LV
420 function, suggesting that oxidative stress and inflammation may be responsible for LV systolic
421 dysfunction in IRI [119]. However, further studies are warranted to determine whether NOS
422 uncoupling induces inflammation associated with acute myocardial IRI.

423

424 **5. Dysregulation of myocardial anti-oxidant pathways during acute myocardial IRI**

425 Myocardial reperfusion increases the production of ROS and undermines the anti-oxidant defence
426 in heart tissue, cause a redox imbalance. Myocardial anti-oxidants can be divided into the
427 endogenous anti-oxidant system and exogenous anti-oxidants. The first line of endogenous anti-
428 oxidants include anti-oxidant enzymes such as superoxide dismutase (SOD), catalase and
429 glutathione peroxidase, and non-enzymatic anti-oxidants including α -tocopherol (vitamin E),
430 ubiquinol or coenzyme Q₁₀ (Q₁₀), ascorbic acid (vitamin C) and glutathione (GSH) amongst others
431 [120], [121]. In the setting of acute IRI, levels of myocardial non-enzymatic anti-oxidants are
432 suppressed. Total myocardium ascorbate, Q₁₀, and glutathione levels decline as a function of the
433 length of reperfusion period, and the administration of exogenous anti-oxidants can mediate
434 cardioprotection [122].

435 The presence of a higher glutathione peroxidase (GPx) activity is vital for the heart to
436 survive the attack of ROS produced in the reperfused myocardium. GPx catalyses the peroxidation
437 of H₂O₂ in the presence of reduced glutathione (GSH) to form H₂O and oxidised glutathione
438 (GSSG). Cardiomyocytes contain a GSH redox cycle, in which GPx reaction accepts peroxides and
439 peroxide-derived alkoxyl and peroxy radicals as substrates. Glutathione is a tripeptide, γ -L-
440 glutamyl-L-cysteinylglycine, present in the heart at 1-10mM concentrations [123]. GSH reductase
441 replenishes the loss of GSH using NADPH as a donor for reducing equivalents, however, the
442 oxidative stress during reperfusion results in the depletion of myocardial GSH and NADPH and
443 efflux of GSSG [124], [125]. Yoshida et, al, have demonstrated that the GPx knockout (KO) mouse
444 hearts are more susceptible to acute IRI [126].

445 The transcription factor Nrf2 is a master regulator of a spectrum of genes related to GSH
446 metabolism via the anti-oxidant responsive element (ARE) on target genes, and also plays a role in
447 xenobiotic detoxification and proteome maintenance [127], [128]. In response to oxidative stress,
448 Nrf2 dissociates from the inhibitory regulator Keap1, and translocates to the nucleus to induce the
449 transcription of anti-oxidant genes GSH synthetase, glutathione-S-transferase, GSH peroxidase,
450 GSH reductase and NADPH quinone oxidoreductase [127], [129]. The stimulation of Nrf2 is
451 connected with activation of the PI3K/Akt kinase pathway which was shown to play a role in the
452 mechanism of increased myocardial tolerance to acute IRI and reduction of oxidative stress [129] ,
453 [130].

454 Oxidative stress depolarises mitochondria by causing lipid peroxidation, which further leads
455 to mitochondrial dysfunction. Q₁₀ is a well-characterized electron carrier of the respiratory chain,
456 which is mainly localised in the inner mitochondrial membrane where it serves as a highly mobile
457 carrier of electrons and protons between the flavoproteins and the cytochrome system [131].
458 Because of its ability to transfer electrons, it acts as an anti-oxidant. Q₁₀ must be reduced to
459 ubiquinol denoted quinol (QH₂) to yield its maximum anti-oxidative function. In its reduced form
460 (ubiquinol), the Q₁₀ molecule holds electrons loosely and will quite easily give up one or two
461 electrons to neutralise free radicals [131], [132]. The anti-oxidant properties of Q₁₀ and its location
462 within the mitochondria make it an potential therapeutic target for the treatment of acute IRI [133]. In
463 conditions of high oxidative stress, the rate of inactivation of NO• to peroxynitrite by superoxide
464 anions may be reduced by Q₁₀, and reduce the products of lipid peroxidation levels [134].
465 Coenzyme Q₁₀ also decreases blood viscosity, improves coronary vasodilation by protecting the
466 endothelial function in patients with ischaemic heart disease [135], [136]. It decreases inflammatory
467 cytokines and prevents the hyperglycemia-induced endothelial cell damage, monocyte adhesion
468 and evolution of atherosclerotic lesions in diabetic patients [137].

469 SOD is an enzyme that converts superoxide anion to hydrogen peroxide, which is
470 subsequently converted to water by catalase [138]. It protects against oxidative stress and has
471 three isoforms: Cu-Zn SOD (SOD1), located in the cytosol; Mn-SOD (SOD2), located in the
472 mitochondrial matrix; and extracellular SOD (SOD3) [139]. Anti-oxidant enzymes have specific

473 targets, and they function as a sensor of specific types of ROS. For instance catalase and
474 peroxiredoxins target H₂O₂ whereas SODs only target superoxide. Inadequate delivery of anti-
475 oxidants to their target sites within the cell where ROS are produced may be the cause of the
476 controversial results obtained so far because these enzymes can detoxify ROS only at the sites to
477 which they are delivered.

478 Mammals have seven sirtuins (SIRT1-7) that possess NAD⁺-dependent deacetylase,
479 deacetylase, and ADP-ribosyltransferase activities. Sirtuins are found in different subcellular
480 locations, including the nucleus (SIRT1, SIRT6, and SIRT7), cytosol (SIRT2), and mitochondria
481 (SIRT3, SIRT4, SIRT5) [140]. SIRT3 deacetylates several lysine residues of MnSOD thereby
482 increasing MnSOD activity and detoxification of superoxide radicals. The dependence of SIRT3 on
483 the NAD⁺/NAD ratio may determine the function of SIRT3 [141]. During myocardial reperfusion,
484 mitochondrial NAD⁺ levels decrease [142], suggesting that SIRT3 activity may be compromised
485 during reperfusion and may contribute to the extent of acute IRI. In the Langendorff model, seven
486 months old *SIRT3*^{-/-} mice showed impaired recovery of cardiac function and larger MI size following
487 25 minutes of ischaemia [143]. SIRT6 reduced oxidative stress injury via an AMPK-dependent
488 pathway. Under normal nutrients conditions, SIRT6 binds to the promoters of glycolytic genes,
489 keeps histone H3K9 acetylation levels low, and directs glucose into the mitochondria for efficient
490 ATP production and away for glycolysis. SIRT6 deficiency also significantly reduced both the
491 expression and activity of SOD and catalase in ischaemic hearts. *SIRT6*^{-/-} mice showed more
492 severe acute myocardial IRI resulted from the collapse of the endogenous ROS-scavenging
493 enzyme system, which induces ROS accumulation and stronger oxidative stress [144]. Oxidative
494 stress activates FOXOs in cardiomyocytes mediated by AMPK and sirtuins (SIRT1 and SIRT2).
495 SIRT1 protects the heart from acute IRI through upregulation of anti-oxidants and downregulation of
496 proapoptotic molecules. FOXO promotes cardiomyocytes survival upon induction by oxidative
497 stress. SIRT1 enhances transcription factor of some FOXO target genes. Cells lacking Sirt3
498 exhibited altered metabolism, including a significant increase in mitochondrial superoxide levels
499 when exposed to cellular stress [140].

500

501 **6. The acute-phase response: tissue damage, inflammatory response, and more ROS**

502 Myocardial reperfusion is associated with an inflammatory response, which ultimately leads to
503 healing and scar formation. Acute myocardial IRI involves degradation of extracellular matrix
504 components by metalloproteinases (MMPs), oxidative stress, apoptosis and activation of
505 complement system. The inflammatory response in the reperfused myocardium is related to the
506 coordinated activation of a series of cytokine and adhesion molecule genes resulting in loss of
507 barrier integrity and release of ROS into the extracellular matrix. It increases expression of adhesion
508 molecules; acts as a chemoattractant for neutrophils, initiating their recruitment; activates the
509 complement cascade and promotes apoptotic cell death.

510 The production of ROS peaks during the first 2–10 minutes of reperfusion after coronary
511 artery occlusion and has been considered to be the first stimuli from neutrophils that invade the
512 ischaemic region [50]. Necrotic cell death triggers release of cell contents with some of the
513 endogenous compounds being able to activate immune cells. NF- κ B is activated by various local
514 substances including ROS [145], [146]. Upon activation, NF- κ B stimulates inflammatory and
515 immune responses. This factor also triggers gene expression of pro-inflammatory cytokines, such
516 as TNF- α and interleukins, initiating an inflammatory response [147]. Upregulation of chemokines
517 and cytokines results in extravasation of activated blood-derived cells into the infarcted area.
518 Platelets are the first cells recruited to the site of infarct area, as a result of the coagulation process.
519 Subsequently, various subsets of leukocytes infiltrate the myocardium and remove the dead cells
520 and matrix debris [148]. During the adhesion process, the activated platelets release adhesion
521 proteins (fibrinogen, fibronectin, P-selectin, glycoprotein IIb/IIIa), growth factors (PDGF), endothelial
522 growth factor, fibroblast growth factor, chemokines, epithelial neutrophil-activating, cytokine-like
523 factors (interleukins) and coagulation factors into the local environment, thereby altering
524 chemotactic, adhesive and proteolytic properties of endothelial cells and supporting chemotaxis
525 adhesion and transmigration of monocytes to the site of inflammation [149], [150]. Concomitantly,
526 intercellular tight junctions are compromised, which leads to endothelial barrier dysfunction and
527 increased vascular permeability. Also, platelets are capable of initiating complement activation and
528 may play a role in localising the inflammatory response to the area of injury [151].

529 Neutrophils arrive on the scene very early after the tissue damage (4 hours after
530 reperfusion) [152]. Their principal role appears to be mediated by adhesive interactions with
531 activated endothelial cells of the vessels. Neutrophil infiltration into the infarcted area implies the
532 generation of ROS and proteolytic enzymes contributing to the clearance of dead cells and debris
533 from the infarcted area [153]. Also, they may express mediators capable of amplifying cell
534 recruitment. Experiments have suggested that the mechanism of neutrophil-cardiomyocyte
535 adhesion is dependent on CD18 integrin activation on neutrophils and expression of ICAM-1, one of
536 the primary ligands for the CD18 integrins. Neutrophils block capillaries preventing reperfusion of
537 the tissue, which leads to tissue necrosis and an exacerbated immune response. In vitro the
538 mechanism of neutrophil–cardiomyocyte injury was shown to be strictly dependent on CD18 integrin
539 activation and ICAM-1 expression by damaged cardiac cells. A neutrophil NADPH oxidase inhibitor
540 and a monoclonal antibody against the neutrophil CD18 adhesion molecule markedly reduced
541 oxygen radical levels, in addition to reducing MI size and no-reflow [81].

542

543 **7. Physiological consequences of the inflammatory response**

544 **7.1. No-reflow**

545 No-reflow (NR) or microvascular obstruction is the term used to describe the inadequate perfusion
546 of a given coronary segment without angiographic evidence of epicardial vessel obstruction.
547 Between 2 minutes and 8 hours of reperfusion, the area of NR increases 3-fold with most of the
548 expansion occurring within the first 1–2 h of reperfusion [154]. The factors associated with the
549 establishment of NR include endothelial dysfunction, compression of capillaries by swollen
550 myocytes, alteration of the vasoregulation pathways, epicardial spasm, mechanical obstruction from
551 embolization, extrinsic coagulation pathways, leukocyte adherence, microvascular ischaemia,
552 oedema and vasoconstriction mediators [155], [156]. Endothelial cell injury occurs in approximately
553 20% of vessels after 60 minutes of reperfusion, and in 40% of vessels at 20–80 minutes of
554 reperfusion. Indeed, initial reports showed tightly packed erythrocytes and endothelial gaps plugged
555 by platelet and fibrin thrombi with many extravascular red blood cells in capillaries from hearts
556 reperfused only during 20 minutes [157]. Platelet aggregates or fibrin clots could be implicated as

557 factors responsible for obstructing capillaries. Besides mechanical obstruction, leukocytes release a
558 variety of pro-inflammatory cytokines that may contribute to NR by recruiting additional inflammatory
559 cells enhancing leukocyte adhesion to the endothelium, altering coagulation or increasing
560 vasoconstriction (Figure 3).

561

562 **7.2. Post-MI left ventricular remodeling**

563 The inflammatory reaction following AMI contributes to cardiac structural remodelling, followed by
564 scar formation at the site of infarction as well as changes in the non-infarcted myocardium, including
565 interstitial fibrosis and vascular remodelling. The term remodelling was proposed to characterize the
566 response of remote myocardium to regional infarction and the progression from acute myocardial
567 infarction to chronic heart failure.

568 There is growing recognition and experimental evidence that oxidative stress-mediated and
569 inflammation regulate the pathogenesis of myocardial remodelling following AMI. Circulating
570 neutrophils and macrophages arrive at the infarct site after reperfusion. They contribute to the
571 proteolytic digestion and phagocytosis of the infarcted tissue, respectively. The inflammatory
572 response peaks at weeks 1 and 2 post-MI. Collagen synthesis, preferentially mediated by
573 myofibroblasts, is induced in response to different stimuli; these include mechanical stress,
574 vasoactive factors such as angiotensin II and growth factors such as transforming growth factor- β
575 (TGF- β), which can act directly or through the up-regulation of connective tissue growth factor
576 (CTGF). The fibrogenic component, which substitutes for lost parenchymal cells, follows the initial
577 phase of collagen degradation. Collagen degradation is mediated by a family of zinc-containing
578 endoproteinases- matrix metalloproteinases. These enzymes are found in the heart at low levels in
579 normal conditions but can be up-regulated after MI in response to inflammatory cytokines and TGF-
580 β [158], [159].

581

582 **7. Therapeutic targeting of ROS and inflammation**

583 It is not surprising that research over recent years has focused on anti-oxidants as a potential
584 therapy in the setting of acute myocardial IRI [160]. Although many initial studies in cells or animal

585 models have been successful, clinical trials have produced disappointing results, owing to
586 differences between animal models and human disease, the inability of the agents to reach the
587 important cellular locations, and the stage and cell-specific regulation of oxidant and anti-oxidant
588 pathways.

589 According to previously discussed data, it may be preferable to adopt a pharmacological
590 strategy that decreases mitochondrial oxidative damage to reduce acute IRI. The relatively poor
591 efficacy of conventional anti-oxidants may be the consequence of their low penetrance to the
592 mitochondrial matrix, which not only is the main site of ROS production but also suffers from
593 oxidative stress. Pretreatment with or infusion of Q₁₀ soon after coronary artery ligation has been
594 shown to reduce MI size and preserve systolic function in rat models of AMI [161]. Q₁₀ prevents the
595 peroxidation of the cell membrane and subcellular lipids, which occurs during acute IRI [162]. Q₁₀
596 also regulates the release of nitric oxide, and it creates endothelial regeneration and
597 immunostimulation [163], [164] and recovery of LV function after AMI [165]. However, lipophilic
598 cations have a disadvantage. Since the charge accumulation into the matrix leads to mitochondrial
599 membrane depolarisation, at the concentration greater than 10mM, toxicity has been observed, and
600 the low solubility of ubiquinone in water makes it difficult to use *in vitro*, and animals must be fed
601 Q₁₀-enriched diets for several weeks to increase levels in subsequently isolated mitochondria. More
602 hydrosoluble molecules than ubiquinone have been developed and tested in different diseases.
603 Therefore, to manipulate mitochondrial Q₁₀ or ubiquinone Kelso et al. synthesised a ubiquinone
604 analogue selectively targeted to mitochondria by the addition of a lipophilic triphenylphosphonium
605 cation, mitoquinone (MitoQ) [166]. The lead compound, MitoQ, consists of a targeting lipophilic
606 triphenyl phosphonium (TPP) cation linked to a ubiquinone moiety. Using this model, it was
607 demonstrated that the administration of MitoQ before the onset of ischaemia reduced oxidative
608 damage and severity of acute IRI, thereby providing functional protection to the heart [167].
609 Myocardial treatment with MitoQ prevented the initial damage and the activation of the inflammatory
610 response. MitoQ also can reduce Ca²⁺ overload and mPTP opening [168]. These findings suggest
611 that mitochondria-targeted therapies designed to minimise mitochondrial oxidative damage may
612 decrease post-reperfusion dysfunction.

613 A growing number of studies suggest that the compound, Szeto-Schiller-31 (SS-31)
614 peptide, also known as Bendavia may be cardioprotective. SS-peptides were developed by Szeto
615 and Schiller and constitute a series of 4 small, cell-permeable anti-oxidant compounds with three
616 positive charges in homeostatic pH conditions [169]. The SS-31 peptide can scavenge H₂O₂ and
617 ONOO• and inhibit lipid peroxidation, anti-oxidant actions attributed to a tyrosine or dimethyl
618 tyrosine residue in their structure, the latter of the two being more efficient concerning ROS
619 scavenging. In a recent study, Liu et al. showed that SS-31 prevented swelling of mitochondria and
620 protected mitochondrial cristae in both endothelial and epithelial cells. It was associated with a
621 significantly reduced loss of peritubular capillaries and cortical arterioles, interstitial inflammation,
622 and fibrosis four weeks after ischaemia [170]. However, the EMBRACE-MI clinica study failed to
623 demonstrate a reduction in MI size in AMI patients administered Bendavia prior to reperfusion [171].

624 Another strategy to preserve redox balance and maintain mitochondrial function is the
625 induction of endogenous anti-oxidants Trolox (6-hydroxy-2,5,7,8- tetramethylchroman-2-carboxylic
626 acid) is a water-soluble analogue of the free radical scavenger α-tocopherol. Due to its enhanced
627 water solubility, Trolox may function more rapidly during acute oxidative stress, while α-tocopherol
628 requires several days of pretreatment to exhibit anti-oxidant benefits. Du *et al.* demonstrated that
629 chitosan nanoparticles, when used as drug carriers for the delivery of Trolox, exerted a protective
630 effect against hypoxia-mediated oxidative stress and can block the mitochondria-dependent
631 apoptotic pathway through upregulation of Bcl-2 expression and inhibition of Bax activation and
632 Caspase-3 expression [172].

633 Recently, the use of natural molecules with specific physicochemical properties has
634 emerged. Vitamin E, C, A and other agents in complementary and alternative medicine have been
635 studied and whereas some had protective effects in animal models, although none of them has
636 demonstrated clear benefit for patients. Curcumin is the major active component of turmeric, a
637 yellow compound isolated from the plant *Curcuma longa*, used for centuries in traditional medicine
638 [173]. This molecule has shown therapeutic potential against a wide range of diseases, mainly due
639 to its anti-inflammatory [174]. Curcumin exerts both direct and indirect anti-oxidant effects by
640 scavenging ROS. Also, it has been shown that early treatment with curcumin attenuates cardiac

641 hypertrophy and remodelling. Curcumin might have therapeutic potential in the treatment of heart
642 disease by attenuating oxidative stress-related events as cardiac remodelling, mitochondrial
643 dysfunction and cell death [175].

644 The phosphorylated form of GSK-3 correlates with the activity of cardioprotection. Via
645 inhibition of GSK-3, protective signalling pathways act on the end effector mitochondrial
646 permeability transition pore; that is, they prevent the induction of the mitochondrial permeability
647 transition, restore mitochondrial membrane potential, and decrease ROS production. The synthetic
648 17 β -aminoestrogen Prolame [17 β -(3-hydroxy-1-propylamino)- 1,3,5(10)-estratrien-3-ol] is an
649 estradiol analogue in which the C17 position of the steroid nucleus is substituted by an amino-
650 alcohol side chain-NH-(CH₂)₃-OH with three methylenes groups. Prolame might diminish the no-
651 reflow phenomenon and provide cardioprotection in rats with AMI followed by reperfusion [156].

652 Initial success has been established in preclinical models of acute myocardial IRI for a
653 handful of therapeutics that target neutrophils [176] [177]. A monoclonal antibody targeted against
654 the CD11/CD18 integrin showed promising results in animal models, but clinical trials failed to show
655 a significant reduction in MI size [178].

656

657 **CONCLUSION**

658 Following AMI, the production of ROS and the ensuing inflammatory response are critical
659 determinants of myocardial injury, cardiomyocyte death and subsequent LV remodelling in the
660 setting of acute IRI, providing therapeutic targets for cardioprotection. However, a number of anti-
661 oxidants have been tested in the setting of AMI, and despite being positive in the experimental
662 setting, most have failed in the clinical setting. The reasons for this are unclear, but may relate to
663 the inability to achieve sufficient concentrations of anti-oxidant at the site of ROS production.
664 Proteomic and metabolomic approaches may help in discover novel pathways underlying the redox-
665 mediated inflammation progression. Overcoming these issues would greatly enhance the
666 development of successful therapies to combat oxidative stress, inflammation and cell damage,
667 providing new treatments for AMI patients.

668

669 **CONFLICT OF INTEREST**

670 The authors declare that they have no conflict of interest

671

672 **ACKNOWLEDGEMENTS**

673 This work was supported by the British Heart Foundation (grant number FS/10/039/28270), the
674 Rosetrees Trust, the National Institute for Health Research University College London Hospitals
675 Biomedical Research Centre, and Duke-National University Singapore Medical School.

676

677 **REFERENCES**

- 678 [1] Cung, T.T.; Morel, O.; Cayla, G.; Rioufol, G.; Garcia-Dorado, D.; Angoulvant, D.;
679 Bonnefoy-Cudraz, E.; Guerin, P.; Elbaz, M.; Delarche, N.; Coste, P.; Vanzetto, G.; Metge,
680 M.; Aupetit, J.F.; Jouve, B.; Motreff, P.; Tron, C.; Labeque, J.N.; Steg, P.G.; Cottin, Y.;
681 Range, G.; Clerc, J.; Claeys, M.J.; Coussement, P.; Prunier, F.; Moulin, F.; Roth, O.; Belle,
682 L.; Dubois, P.; Barragan, P.; Gilard, M.; Piot, C.; Colin, P.; De Poli, F.; Morice, M.C.; Ider,
683 O.; Dubois-Rande, J.L.; Untersee, T.; Le Breton, H.; Beard, T.; Blanchard, D.; Grollier,
684 G.; Malquarti, V.; Staat, P.; Sudre, A.; Elmer, E.; Hansson, M.J.; Bergerot, C.; Boussaha, I.;
685 Jossan, C.; Derumeaux, G.; Newton, N.; Ovize, M. Cyclosporine before PCI in Patients
686 with Acute Myocardial Infarction. *N. Engl. J. Med.*, **2015**, *373*(11), 1021-1031.
- 687 [2] Yellon, D.M.; Hausenloy, D.J. Myocardial reperfusion injury. *N. Engl. J. Med.*,
688 **2007**, *357*(11), 1121-1135.
- 689 [3] Hausenloy, D.J.; Yellon, D.M. Ischaemic conditioning and reperfusion injury.
690 *Nat. Rev. Cardiol.*, **2016**, *13*(4), 193-209.
- 691 [4] Hearse, D.J.; Humphrey, S.M.; Chain, E.B. Abrupt reoxygenation of the anoxic
692 potassium-arrested perfused rat heart: a study of myocardial enzyme release. *J. Mol.*
693 *Cell. Cardiol.*, **1973**, *5*(4), 395-407.
- 694 [5] Siegmund, B.; Schluter, K.D.; Piper, H.M. Calcium and the oxygen paradox.
695 *Cardiovasc. Res.*, **1993**, *27*(10), 1778-1783.
- 696 [6] Bresnahan, G.F.; Roberts, R.; Shell, W.E.; Ross, J., Jr.; Sobel, B.E. Deleterious
697 effects due to hemorrhage after myocardial reperfusion. *Am. J. Cardiol.*, **1974**, *33*(1),
698 82-86.
- 699 [7] Bolli, R.; Zughuib, M.; Li, X.Y.; Tang, X.L.; Sun, J.Z.; Triana, J.F.; McCay, P.B.
700 Recurrent ischemia in the canine heart causes recurrent bursts of free radical
701 production that have a cumulative effect on contractile function. A pathophysiological
702 basis for chronic myocardial "stunning". *J. Clin. Invest.*, **1995**, *96*(2), 1066-1084.
- 703 [8] Koppenol, W.H.; Traynham, J.G. Say NO to nitric oxide: nomenclature for
704 nitrogen- and oxygen-containing compounds. *Methods Enzymol.*, **1996**, *268*, 3-7.
- 705 [9] Christianson, D.W. Structural chemistry and biology of manganese
706 metalloenzymes. *Prog. Biophys. Mol. Biol.*, **1997**, *67*(2-3), 217-252.
- 707 [10] Forman, H.J.; Maiorino, M.; Ursini, F. Signaling functions of reactive oxygen
708 species. *Biochemistry*, **2010**, *49*(5), 835-842.

- 709 [11] Hernandez-Resendiz, S.; Buelna-Chontal, M.; Correa, F.; Zazueta, C. Targeting
710 mitochondria for cardiac protection. *Curr. Drug Targets*, **2013**, *14*(5), 586-600.
- 711 [12] Rhee, S.G. Cell signaling. H₂O₂, a necessary evil for cell signaling. *Science*, **2006**,
712 *312*(5782), 1882-1883.
- 713 [13] Griendling, K.K.; Touyz, R.M.; Zweier, J.L.; Dikalov, S.; Chilian, W.; Chen, Y.R.;
714 Harrison, D.G.; Bhatnagar, A.; American Heart Association Council on Basic
715 Cardiovascular, S. Measurement of Reactive Oxygen Species, Reactive Nitrogen
716 Species, and Redox-Dependent Signaling in the Cardiovascular System: A Scientific
717 Statement From the American Heart Association. *Circ. Res.*, **2016**, *119*(5), e39-75.
- 718 [14] Kornfeld, O.S.; Hwang, S.; Disatnik, M.H.; Chen, C.H.; Qvit, N.; Mochly-Rosen, D.
719 Mitochondrial reactive oxygen species at the heart of the matter: new therapeutic
720 approaches for cardiovascular diseases. *Circ. Res.*, **2015**, *116*(11), 1783-1799.
- 721 [15] Sun, J.; Xin, C.; Eu, J.P.; Stamler, J.S.; Meissner, G. Cysteine-3635 is responsible
722 for skeletal muscle ryanodine receptor modulation by NO. *Proc. Natl. Acad. Sci. U. S. A.*,
723 **2001**, *98*(20), 11158-11162.
- 724 [16] Finkel, T. Signal transduction by reactive oxygen species. *J. Cell. Biol.*, **2011**,
725 *194*(1), 7-15.
- 726 [17] Kaminski, M.M.; Roth, D.; Krammer, P.H.; Gulow, K. Mitochondria as oxidative
727 signaling organelles in T-cell activation: physiological role and pathological
728 implications. *Arch. Immunol. Ther. Exp. (Warsz)*, **2013**, *61*(5), 367-384.
- 729 [18] Devadas, S.; Zaritskaya, L.; Rhee, S.G.; Oberley, L.; Williams, M.S. Discrete
730 generation of superoxide and hydrogen peroxide by T cell receptor stimulation:
731 selective regulation of mitogen-activated protein kinase activation and fas ligand
732 expression. *J. Exp. Med.*, **2002**, *195*(1), 59-70.
- 733 [19] Chaudhri, G.; Hunt, N.H.; Clark, I.A.; Ceredig, R. Antioxidants inhibit
734 proliferation and cell surface expression of receptors for interleukin-2 and transferrin
735 in T lymphocytes stimulated with phorbol myristate acetate and ionomycin. *Cell.*
736 *Immunol.*, **1988**, *115*(1), 204-213.
- 737 [20] Yi, X.Y.; Li, V.X.; Zhang, F.; Yi, F.; Matson, D.R.; Jiang, M.T.; Li, P.L. Characteristics
738 and actions of NAD(P)H oxidase on the sarcoplasmic reticulum of coronary artery
739 smooth muscle. *Am. J. Physiol. Heart Circ. Physiol.*, **2006**, *290*(3), H1136-1144.
- 740 [21] Xiao, L.; Pimentel, D.R.; Wang, J.; Singh, K.; Colucci, W.S.; Sawyer, D.B. Role of
741 reactive oxygen species and NAD(P)H oxidase in alpha(1)-adrenoceptor signaling in
742 adult rat cardiac myocytes. *Am. J. Physiol. Cell. Physiol.*, **2002**, *282*(4), C926-934.
- 743 [22] Scherz-Shouval, R.; Shvets, E.; Fass, E.; Shorer, H.; Gil, L.; Elazar, Z. Reactive
744 oxygen species are essential for autophagy and specifically regulate the activity of
745 Atg4. *EMBO J.*, **2007**, *26*(7), 1749-1760.
- 746 [23] Perrelli, M.G.; Tullio, F.; Angotti, C.; Cerra, M.C.; Angelone, T.; Tota, B.; Alloatti,
747 G.; Penna, C.; Pagliaro, P. Catestatin reduces myocardial ischaemia/reperfusion injury:
748 involvement of PI3K/Akt, PKCs, mitochondrial KATP channels and ROS signalling.
749 *Pflugers Arch.*, **2013**, *465*(7), 1031-1040.
- 750 [24] Hoshi, T.; Heinemann, S. Regulation of cell function by methionine oxidation
751 and reduction. *J. Physiol.*, **2001**, *531*(Pt 1), 1-11.
- 752 [25] Funato, Y.; Michiue, T.; Asashima, M.; Miki, H. The thioredoxin-related redox-
753 regulating protein nucleoredoxin inhibits Wnt-beta-catenin signalling through
754 dishevelled. *Nat. Cell. Biol.*, **2006**, *8*(5), 501-508.

755 [26] Nordberg, J.; Arner, E.S. Reactive oxygen species, antioxidants, and the
756 mammalian thioredoxin system. *Free Radic. Biol. Med.*, **2001**, *31*(11), 1287-1312.

757 [27] Gao, J.; Yin, D.H.; Yao, Y.; Sun, H.; Qin, Z.; Schoneich, C.; Williams, T.D.; Squier,
758 T.C. Loss of conformational stability in calmodulin upon methionine oxidation.
759 *Biophys. J.*, **1998**, *74*(3), 1115-1134.

760 [28] Erickson, J.R.; Joiner, M.L.; Guan, X.; Kutschke, W.; Yang, J.; Oddis, C.V.; Bartlett,
761 R.K.; Lowe, J.S.; O'Donnell, S.E.; Aykin-Burns, N.; Zimmerman, M.C.; Zimmerman, K.;
762 Ham, A.J.; Weiss, R.M.; Spitz, D.R.; Shea, M.A.; Colbran, R.J.; Mohler, P.J.; Anderson, M.E.
763 A dynamic pathway for calcium-independent activation of CaMKII by methionine
764 oxidation. *Cell*, **2008**, *133*(3), 462-474.

765 [29] Shimokawa, H.; Morikawa, K. Hydrogen peroxide is an endothelium-derived
766 hyperpolarizing factor in animals and humans. *J. Mol. Cell. Cardiol.*, **2005**, *39*(5), 725-
767 732.

768 [30] Miura, H.; Bosnjak, J.J.; Ning, G.; Saito, T.; Miura, M.; Gutterman, D.D. Role for
769 hydrogen peroxide in flow-induced dilation of human coronary arterioles. *Circ. Res.*,
770 **2003**, *92*(2), e31-40.

771 [31] Burgoyne, J.R.; Madhani, M.; Cuello, F.; Charles, R.L.; Brennan, J.P.; Schroder, E.;
772 Browning, D.D.; Eaton, P. Cysteine redox sensor in PKGI α enables oxidant-induced
773 activation. *Science*, **2007**, *317*(5843), 1393-1397.

774 [32] Prysyzhna, O.; Rudyk, O.; Eaton, P. Single atom substitution in mouse protein
775 kinase G eliminates oxidant sensing to cause hypertension. *Nat. Med.*, **2012**, *18*(2),
776 286-290.

777 [33] Burgoyne, J.R.; Rudyk, O.; Cho, H.J.; Prysyzhna, O.; Hathaway, N.; Weeks, A.;
778 Evans, R.; Ng, T.; Schroder, K.; Brandes, R.P.; Shah, A.M.; Eaton, P. Deficient
779 angiogenesis in redox-dead Cys17Ser PKAR1 α knock-in mice. *Nat. Commun.*, **2015**,
780 6, 7920.

781 [34] Xia, R.; Stangler, T.; Abramson, J.J. Skeletal muscle ryanodine receptor is a redox
782 sensor with a well defined redox potential that is sensitive to channel modulators. *J.*
783 *Biol. Chem.*, **2000**, *275*(47), 36556-36561.

784 [35] Kumasaka, S.; Shoji, H.; Okabe, E. Novel mechanisms involved in superoxide
785 anion radical-triggered Ca²⁺ release from cardiac sarcoplasmic reticulum linked to
786 cyclic ADP-ribose stimulation. *Antioxid. Redox Signal.*, **1999**, *1*(1), 55-69.

787 [36] Pessah, I.N.; Kim, K.H.; Feng, W. Redox sensing properties of the ryanodine
788 receptor complex. *Front. Biosci.*, **2002**, *7*, a72-79.

789 [37] Dulhunty, A.; Haarmann, C.; Green, D.; Hart, J. How many cysteine residues
790 regulate ryanodine receptor channel activity? *Antioxid. Redox Signal.*, **2000**, *2*(1), 27-
791 34.

792 [38] Feng, W.; Liu, G.; Xia, R.; Abramson, J.J.; Pessah, I.N. Site-selective modification
793 of hyperreactive cysteines of ryanodine receptor complex by quinones. *Mol.*
794 *Pharmacol.*, **1999**, *55*(5), 821-831.

795 [39] Feng, W.; Pessah, I.N. Detection of redox sensor of ryanodine receptor
796 complexes. *Methods Enzymol.*, **2002**, *353*, 240-253.

797 [40] Ushio-Fukai, M.; Zafari, A.M.; Fukui, T.; Ishizaka, N.; Griending, K.K. p22phox is
798 a critical component of the superoxide-generating NADH/NADPH oxidase system and
799 regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells. *J. Biol.*
800 *Chem.*, **1996**, *271*(38), 23317-23321.

801 [41] Bendall, J.K.; Cave, A.C.; Heymes, C.; Gall, N.; Shah, A.M. Pivotal role of a
802 gp91(phox)-containing NADPH oxidase in angiotensin II-induced cardiac hypertrophy
803 in mice. *Circulation*, **2002**, *105*(3), 293-296.

804 [42] Sugden, P.H.; Clerk, A. Cellular mechanisms of cardiac hypertrophy. *J. Mol. Med.*
805 *(Berl)*, **1998**, *76*(11), 725-746.

806 [43] Tullio, F.; Angotti, C.; Perrelli, M.G.; Penna, C.; Pagliaro, P. Redox balance and
807 cardioprotection. *Basic Res. Cardiol.*, **2013**, *108*(6), 392.

808 [44] Heinzl, F.R.; Luo, Y.; Li, X.; Boengler, K.; Buechert, A.; Garcia-Dorado, D.; Di Lisa,
809 F.; Schulz, R.; Heusch, G. Impairment of diazoxide-induced formation of reactive
810 oxygen species and loss of cardioprotection in connexin 43 deficient mice. *Circ. Res.*,
811 **2005**, *97*(6), 583-586.

812 [45] Boengler, K.; Dodoni, G.; Rodriguez-Sinovas, A.; Cabestrero, A.; Ruiz-Meana, M.;
813 Gres, P.; Konietzka, I.; Lopez-Iglesias, C.; Garcia-Dorado, D.; Di Lisa, F.; Heusch, G.;
814 Schulz, R. Connexin 43 in cardiomyocyte mitochondria and its increase by ischemic
815 preconditioning. *Cardiovasc. Res.*, **2005**, *67*(2), 234-244.

816 [46] Ruiz-Meana, M.; Abellan, A.; Miro-Casas, E.; Agullo, E.; Garcia-Dorado, D. Role of
817 sarcoplasmic reticulum in mitochondrial permeability transition and cardiomyocyte
818 death during reperfusion. *Am. J. Physiol. Heart Circ. Physiol.*, **2009**, *297*(4), H1281-
819 1289.

820 [47] Zweier, J.L.; Talukder, M.A. The role of oxidants and free radicals in reperfusion
821 injury. *Cardiovasc. Res.*, **2006**, *70*(2), 181-190.

822 [48] Thum, T.; Borlak, J. Gene expression in distinct regions of the heart. *Lancet*,
823 **2000**, *355*(9208), 979-983.

824 [49] Edin, M.L.; Wang, Z.; Bradbury, J.A.; Graves, J.P.; Lih, F.B.; DeGraff, L.M.; Foley,
825 J.F.; Torphy, R.; Ronnekleiv, O.K.; Tomer, K.B.; Lee, C.R.; Zeldin, D.C. Endothelial
826 expression of human cytochrome P450 epoxygenase CYP2C8 increases susceptibility
827 to ischemia-reperfusion injury in isolated mouse heart. *FASEB J.*, **2011**, *25*(10), 3436-
828 3447.

829 [50] Vinten-Johansen, J. Involvement of neutrophils in the pathogenesis of lethal
830 myocardial reperfusion injury. *Cardiovasc. Res.*, **2004**, *61*(3), 481-497.

831 [51] Loida, P.J.; Sligar, S.G. Molecular recognition in cytochrome P-450: mechanism
832 for the control of uncoupling reactions. *Biochemistry*, **1993**, *32*(43), 11530-11538.

833 [52] Fleming, I.; Michaelis, U.R.; Bredenkotter, D.; Fisslthaler, B.; Deghani, F.;
834 Brandes, R.P.; Busse, R. Endothelium-derived hyperpolarizing factor synthase
835 (Cytochrome P450 2C9) is a functionally significant source of reactive oxygen species
836 in coronary arteries. *Circ. Res.*, **2001**, *88*(1), 44-51.

837 [53] Granville, D.J.; Tashakkor, B.; Takeuchi, C.; Gustafsson, A.B.; Huang, C.; Sayen,
838 M.R.; Wentworth, P., Jr.; Yeager, M.; Gottlieb, R.A. Reduction of ischemia and
839 reperfusion-induced myocardial damage by cytochrome P450 inhibitors. *Proc. Natl.*
840 *Acad. Sci. U. S. A.*, **2004**, *101*(5), 1321-1326.

841 [54] Ivanics, T.; Miklos, Z.; Dezsi, L.; Ikrenyi, K.; Toth, A.; Roemen, T.H.; Van der
842 Vusse, G.J.; Ligeti, L. Concomitant accumulation of intracellular free calcium and
843 arachidonic acid in the ischemic-reperfused rat heart. *Mol. Cell. Biochem.*, **2001**, *226*(1-
844 2), 119-128.

845 [55] Ishizuka, T.; Cheng, J.; Singh, H.; Vitto, M.D.; Manthati, V.L.; Falck, J.R.; Laniado-
846 Schwartzman, M. 20-Hydroxyeicosatetraenoic acid stimulates nuclear factor-kappaB

847 activation and the production of inflammatory cytokines in human endothelial cells. *J.*
848 *Pharmacol. Exp. Ther.*, **2008**, *324*(1), 103-110.

849 [56] Toth, P.; Csiszar, A.; Sosnowska, D.; Tucsek, Z.; Cseplo, P.; Springo, Z.; Tarantini,
850 S.; Sonntag, W.E.; Ungvari, Z.; Koller, A. Treatment with the cytochrome P450 omega-
851 hydroxylase inhibitor HET0016 attenuates cerebrovascular inflammation, oxidative
852 stress and improves vasomotor function in spontaneously hypertensive rats. *Br. J.*
853 *Pharmacol.*, **2013**, *168*(8), 1878-1888.

854 [57] Cheng, J.; Ou, J.S.; Singh, H.; Falck, J.R.; Narsimhaswamy, D.; Pritchard, K.A., Jr.;
855 Schwartzman, M.L. 20-hydroxyeicosatetraenoic acid causes endothelial dysfunction
856 via eNOS uncoupling. *Am. J. Physiol. Heart Circ. Physiol.*, **2008**, *294*(2), H1018-1026.

857 [58] Ishihara, Y.; Sekine, M.; Nakazawa, M.; Shimamoto, N. Suppression of
858 myocardial ischemia-reperfusion injury by inhibitors of cytochrome P450 in rats. *Eur.*
859 *J. Pharmacol.*, **2009**, *611*(1-3), 64-71.

860 [59] Gross, E.R.; Nithipatikom, K.; Hsu, A.K.; Peart, J.N.; Falck, J.R.; Campbell, W.B.;
861 Gross, G.J. Cytochrome P450 omega-hydroxylase inhibition reduces infarct size during
862 reperfusion via the sarcolemmal KATP channel. *J. Mol. Cell. Cardiol.*, **2004**, *37*(6),
863 1245-1249.

864 [60] Fan, F.; Ge, Y.; Lv, W.; Elliott, M.R.; Muroya, Y.; Hirata, T.; Booz, G.W.; Roman, R.J.
865 Molecular mechanisms and cell signaling of 20-hydroxyeicosatetraenoic acid in
866 vascular pathophysiology. *Front. Biosci. (Landmark Ed)*, **2016**, *21*, 1427-1463.

867 [61] Battelli, M.G.; Bolognesi, A.; Polito, L. Pathophysiology of circulating xanthine
868 oxidoreductase: new emerging roles for a multi-tasking enzyme. *Biochim. Biophys.*
869 *Acta*, **2014**, *1842*(9), 1502-1517.

870 [62] Thompson-Gorman, S.L.; Zweier, J.L. Evaluation of the role of xanthine oxidase
871 in myocardial reperfusion injury. *J. Biol. Chem.*, **1990**, *265*(12), 6656-6663.

872 [63] Chambers, D.E.; Parks, D.A.; Patterson, G.; Roy, R.; McCord, J.M.; Yoshida, S.;
873 Parmley, L.F.; Downey, J.M. Xanthine oxidase as a source of free radical damage in
874 myocardial ischemia. *J. Mol. Cell. Cardiol.*, **1985**, *17*(2), 145-152.

875 [64] Pisarenko, O.I.; Lakomkin, V.L.; Studneva, I.M.; Timoshin, A.A.; Kuzmin, A.I.;
876 Ruuge, E.K.; Kapelko, V.I. Allopurinol-enhanced postischemic recovery in the isolated
877 rat heart involves repletion of high-energy phosphates. *Biochem. Med. Metab. Biol.*,
878 **1994**, *51*(1), 16-26.

879 [65] Miwa-Nishimura, N.; Kanaide, H.; Abe, S.; Nakamura, M. Does allopurinol
880 prevent myocardial injury as a result of hypoxia-re-oxygenation in rats? *Int. J. Exp.*
881 *Pathol.*, **1990**, *71*(5), 727-739.

882 [66] Holzgrefe, H.H.; Gibson, J.K. Beneficial effects of oxypurinol pretreatment in
883 stunned, reperfused canine myocardium. *Cardiovasc. Res.*, **1989**, *23*(4), 340-350.

884 [67] Guan, W.; Osanai, T.; Kamada, T.; Hanada, H.; Ishizaka, H.; Onodera, H.; Iwasa,
885 A.; Fujita, N.; Kudo, S.; Ohkubo, T.; Okumura, K. Effect of allopurinol pretreatment on
886 free radical generation after primary coronary angioplasty for acute myocardial
887 infarction. *J. Cardiovasc. Pharmacol.*, **2003**, *41*(5), 699-705.

888 [68] Matsumura, F.; Yamaguchi, Y.; Goto, M.; Ichiguchi, O.; Akizuki, E.; Matsuda, T.;
889 Okabe, K.; Liang, J.; Ohshiro, H.; Iwamoto, T.; Yamada, S.; Mori, K.; Ogawa, M. Xanthine
890 oxidase inhibition attenuates kupffer cell production of neutrophil chemoattractant
891 following ischemia-reperfusion in rat liver. *Hepatology*, **1998**, *28*(6), 1578-1587.

892 [69] Jones, A.W.; Durante, W.; Korthuis, R.J. Heme oxygenase-1 deficiency leads to
893 alteration of soluble guanylate cyclase redox regulation. *J. Pharmacol. Exp. Ther.*, **2010**,
894 335(1), 85-91.

895 [70] Ichikawa, H.; Flores, S.; Kvietys, P.R.; Wolf, R.E.; Yoshikawa, T.; Granger, D.N.;
896 Aw, T.Y. Molecular mechanisms of anoxia/reoxygenation-induced neutrophil
897 adherence to cultured endothelial cells. *Circ. Res.*, **1997**, 81(6), 922-931.

898 [71] Borejdo, J.; Ushakov, D.S.; Akopova, I. Regulatory and essential light chains of
899 myosin rotate equally during contraction of skeletal muscle. *Biophys. J.*, **2002**, 82(6),
900 3150-3159.

901 [72] Braunersreuther, V.; Montecucco, F.; Asrih, M.; Pelli, G.; Galan, K.; Frias, M.;
902 Burger, F.; Quindere, A.L.; Montessuit, C.; Krause, K.H.; Mach, F.; Jaquet, V. Role of
903 NADPH oxidase isoforms NOX1, NOX2 and NOX4 in myocardial ischemia/reperfusion
904 injury. *J. Mol. Cell. Cardiol.*, **2013**, 64, 99-107.

905 [73] Cave, A.C.; Brewer, A.C.; Narayanapanicker, A.; Ray, R.; Grieve, D.J.; Walker, S.;
906 Shah, A.M. NADPH oxidases in cardiovascular health and disease. *Antioxid. Redox*
907 *Signal.*, **2006**, 8(5-6), 691-728.

908 [74] Banfi, B.; Malgrange, B.; Knisz, J.; Steger, K.; Dubois-Dauphin, M.; Krause, K.H.
909 NOX3, a superoxide-generating NADPH oxidase of the inner ear. *J. Biol. Chem.*, **2004**,
910 279(44), 46065-46072.

911 [75] Wingler, K.; Wunsch, S.; Kreutz, R.; Rothermund, L.; Paul, M.; Schmidt, H.H.
912 Upregulation of the vascular NAD(P)H-oxidase isoforms Nox1 and Nox4 by the renin-
913 angiotensin system in vitro and in vivo. *Free Radic. Biol. Med.*, **2001**, 31(11), 1456-
914 1464.

915 [76] BelAiba, R.S.; Djordjevic, T.; Petry, A.; Diemer, K.; Bonello, S.; Banfi, B.; Hess, J.;
916 Pogrebniak, A.; Bickel, C.; Gorlach, A. NOX5 variants are functionally active in
917 endothelial cells. *Free Radic. Biol. Med.*, **2007**, 42(4), 446-459.

918 [77] Jay, D.B.; Papaharalambus, C.A.; Seidel-Rogol, B.; Dikalova, A.E.; Lassegue, B.;
919 Griendling, K.K. Nox5 mediates PDGF-induced proliferation in human aortic smooth
920 muscle cells. *Free Radic. Biol. Med.*, **2008**, 45(3), 329-335.

921 [78] van der Vliet, A. NADPH oxidases in lung biology and pathology: host defense
922 enzymes, and more. *Free Radic. Biol. Med.*, **2008**, 44(6), 938-955.

923 [79] Meischl, C.; Krijnen, P.A.; Sipkens, J.A.; Cillessen, S.A.; Munoz, I.G.; Okroj, M.;
924 Ramska, M.; Muller, A.; Visser, C.A.; Musters, R.J.; Simonides, W.S.; Hack, C.E.; Roos, D.;
925 Niessen, H.W. Ischemia induces nuclear NOX2 expression in cardiomyocytes and
926 subsequently activates apoptosis. *Apoptosis*, **2006**, 11(6), 913-921.

927 [80] Borch, E.; Parri, M.; Papucci, L.; Becatti, M.; Nassi, N.; Nassi, P.; Nediani, C. Role
928 of NADPH oxidase in H9c2 cardiac muscle cells exposed to simulated ischaemia-
929 reperfusion. *J. Cell. Mol. Med.*, **2009**, 13(8B), 2724-2735.

930 [81] Duilio, C.; Ambrosio, G.; Kuppusamy, P.; DiPaula, A.; Becker, L.C.; Zweier, J.L.
931 Neutrophils are primary source of O₂ radicals during reperfusion after prolonged
932 myocardial ischemia. *Am. J. Physiol. Heart Circ. Physiol.*, **2001**, 280(6), H2649-2657.

933 [82] Romson, J.L.; Hook, B.G.; Kunkel, S.L.; Abrams, G.D.; Schork, M.A.; Lucchesi, B.R.
934 Reduction of the extent of ischemic myocardial injury by neutrophil depletion in the
935 dog. *Circulation*, **1983**, 67(5), 1016-1023.

936 [83] Bayraktutan, U.; Draper, N.; Lang, D.; Shah, A.M. Expression of functional
937 neutrophil-type NADPH oxidase in cultured rat coronary microvascular endothelial
938 cells. *Cardiovasc. Res.*, **1998**, *38*(1), 256-262.

939 [84] Patterson, C.; Ruef, J.; Madamanchi, N.R.; Barry-Lane, P.; Hu, Z.; Horaist, C.;
940 Ballinger, C.A.; Brasier, A.R.; Bode, C.; Runge, M.S. Stimulation of a vascular smooth
941 muscle cell NAD(P)H oxidase by thrombin. Evidence that p47(phox) may participate
942 in forming this oxidase in vitro and in vivo. *J. Biol. Chem.*, **1999**, *274*(28), 19814-
943 19822.

944 [85] Rey, F.E.; Cifuentes, M.E.; Kiarash, A.; Quinn, M.T.; Pagano, P.J. Novel
945 competitive inhibitor of NAD(P)H oxidase assembly attenuates vascular O₂(⁻) and
946 systolic blood pressure in mice. *Circ. Res.*, **2001**, *89*(5), 408-414.

947 [86] Kleikers, P.W.; Wingler, K.; Hermans, J.J.; Diebold, I.; Altenhofer, S.;
948 Radermacher, K.A.; Janssen, B.; Grolach, A.; Schmidt, H.H. NADPH oxidases as a source
949 of oxidative stress and molecular target in ischemia/reperfusion injury. *J. Mol. Med.*
950 *(Berl)*, **2012**, *90*(12), 1391-1406.

951 [87] Edmondson, D.E.; Mattevi, A.; Binda, C.; Li, M.; Hubalek, F. Structure and
952 mechanism of monoamine oxidase. *Curr. Med. Chem.*, **2004**, *11*(15), 1983-1993.

953 [88] Kaludercic, N.; Carpi, A.; Menabo, R.; Di Lisa, F.; Paolocci, N. Monoamine
954 oxidases (MAO) in the pathogenesis of heart failure and ischemia/reperfusion injury.
955 *Biochim. Biophys. Acta*, **2011**, *1813*(7), 1323-1332.

956 [89] Bach, A.W.; Lan, N.C.; Johnson, D.L.; Abell, C.W.; Bembenek, M.E.; Kwan, S.W.;
957 Seeburg, P.H.; Shih, J.C. cDNA cloning of human liver monoamine oxidase A and B:
958 molecular basis of differences in enzymatic properties. *Proc. Natl. Acad. Sci. U. S. A.*,
959 **1988**, *85*(13), 4934-4938.

960 [90] Kunduzova, O.R.; Bianchi, P.; Parini, A.; Cambon, C. Hydrogen peroxide
961 production by monoamine oxidase during ischemia/reperfusion. *Eur. J. Pharmacol.*,
962 **2002**, *448*(2-3), 225-230.

963 [91] Pchejetski, D.; Kunduzova, O.; Dayon, A.; Calise, D.; Seguelas, M.H.; Leducq, N.;
964 Seif, I.; Parini, A.; Cuvillier, O. Oxidative stress-dependent sphingosine kinase-1
965 inhibition mediates monoamine oxidase A-associated cardiac cell apoptosis. *Circ. Res.*,
966 **2007**, *100*(1), 41-49.

967 [92] Qin, F.; Shite, J.; Mao, W.; Liang, C.S. Selegiline attenuates cardiac oxidative
968 stress and apoptosis in heart failure: association with improvement of cardiac
969 function. *Eur. J. Pharmacol.*, **2003**, *461*(2-3), 149-158.

970 [93] Simon, L.; Szilagy, G.; Bori, Z.; Orbay, P.; Nagy, Z. (-)-D-Deprenyl attenuates
971 apoptosis in experimental brain ischaemia. *Eur. J. Pharmacol.*, **2001**, *430*(2-3), 235-
972 241.

973 [94] Lee, H.L.; Chen, C.L.; Yeh, S.T.; Zweier, J.L.; Chen, Y.R. Biphasic modulation of the
974 mitochondrial electron transport chain in myocardial ischemia and reperfusion. *Am. J.*
975 *Physiol. Heart Circ. Physiol.*, **2012**, *302*(7), H1410-1422.

976 [95] Gorenkova, N.; Robinson, E.; Grieve, D.J.; Galkin, A. Conformational change of
977 mitochondrial complex I increases ROS sensitivity during ischemia. *Antioxid. Redox*
978 *Signal.*, **2013**, *19*(13), 1459-1468.

979 [96] Chen, Y.R.; Chen, C.L.; Pfeiffer, D.R.; Zweier, J.L. Mitochondrial complex II in the
980 post-ischemic heart: oxidative injury and the role of protein S-glutathionylation. *J.*
981 *Biol. Chem.*, **2007**, *282*(45), 32640-32654.

982 [97] Petrosillo, G.; Ruggiero, F.M.; Di Venosa, N.; Paradies, G. Decreased complex III
983 activity in mitochondria isolated from rat heart subjected to ischemia and
984 reperfusion: role of reactive oxygen species and cardiolipin. *FASEB J.*, **2003**, *17*(6),
985 714-716.

986 [98] Lemaire, S.; Lizard, G.; Monier, S.; Miguet, C.; Gueldry, S.; Volot, F.; Gambert, P.;
987 Neel, D. Different patterns of IL-1beta secretion, adhesion molecule expression and
988 apoptosis induction in human endothelial cells treated with 7alpha-, 7beta-
989 hydroxycholesterol, or 7-ketocholesterol. *FEBS Lett.*, **1998**, *440*(3), 434-439.

990 [99] Ou, W.J.; Ito, A.; Morohashi, K.; Fujii-Kuriyama, Y.; Omura, T. Processing-
991 independent in vitro translocation of cytochrome P-450(SCC) precursor across
992 mitochondrial membranes. *J. Biochem.*, **1986**, *100*(5), 1287-1296.

993 [100] Pikuleva, I.A. Cholesterol-metabolizing cytochromes P450. *Drug. Metab. Dispos.*,
994 **2006**, *34*(4), 513-520.

995 [101] Paradis, S.; Leoni, V.; Caccia, C.; Berdeaux, A.; Morin, D. Cardioprotection by the
996 TSPO ligand 4'-chlorodiazepam is associated with inhibition of mitochondrial
997 accumulation of cholesterol at reperfusion. *Cardiovasc. Res.*, **2013**, *98*(3), 420-427.

998 [102] Lemaire-Ewing, S.; Prunet, C.; Montange, T.; Vejux, A.; Berthier, A.; Bessedé, G.;
999 Corcos, L.; Gambert, P.; Neel, D.; Lizard, G. Comparison of the cytotoxic, pro-oxidant
1000 and pro-inflammatory characteristics of different oxysterols. *Cell. Biol. Toxicol.*, **2005**,
1001 *21*(2), 97-114.

1002 [103] Liu, Y.; Hultén, L.M.; Wiklund, O. Macrophages isolated from human
1003 atherosclerotic plaques produce IL-8, and oxysterols may have a regulatory function
1004 for IL-8 production. *Arterioscler. Thromb. Vasc. Biol.*, **1997**, *17*(2), 317-323.

1005 [104] Terkeltaub, R.; Banka, C.L.; Solan, J.; Santoro, D.; Brand, K.; Curtiss, L.K. Oxidized
1006 LDL induces monocytic cell expression of interleukin-8, a chemokine with T-
1007 lymphocyte chemotactic activity. *Arterioscler. Thromb.*, **1994**, *14*(1), 47-53.

1008 [105] de Brito, O.M.; Scorrano, L. Mitofusin 2 tethers endoplasmic reticulum to
1009 mitochondria. *Nature*, **2008**, *456*(7222), 605-610.

1010 [106] Hu, P.; Han, Z.; Couvillon, A.D.; Kaufman, R.J.; Exton, J.H. Autocrine tumor
1011 necrosis factor alpha links endoplasmic reticulum stress to the membrane death
1012 receptor pathway through IRE1alpha-mediated NF-kappaB activation and down-
1013 regulation of TRAF2 expression. *Mol. Cell. Biol.*, **2006**, *26*(8), 3071-3084.

1014 [107] Wu, Q.; Wang, Q.; Guo, Z.; Shang, Y.; Zhang, L.; Gong, S. Nuclear factor-kappaB as
1015 a link between endoplasmic reticulum stress and inflammation during cardiomyocyte
1016 hypoxia/reoxygenation. *Cell Biol. Int.*, **2014**, *38*(7), 881-887.

1017 [108] Jiang, H.Y.; Wek, S.A.; McGrath, B.C.; Scheuner, D.; Kaufman, R.J.; Cavener, D.R.;
1018 Wek, R.C. Phosphorylation of the alpha subunit of eukaryotic initiation factor 2 is
1019 required for activation of NF-kappaB in response to diverse cellular stresses. *Mol. Cell.*
1020 *Biol.*, **2003**, *23*(16), 5651-5663.

1021 [109] Urano, F.; Wang, X.; Bertolotti, A.; Zhang, Y.; Chung, P.; Harding, H.P.; Ron, D.
1022 Coupling of stress in the ER to activation of JNK protein kinases by transmembrane
1023 protein kinase IRE1. *Science*, **2000**, *287*(5453), 664-666.

1024 [110] Zhong, Y.; Li, J.; Chen, Y.; Wang, J.J.; Ratan, R.; Zhang, S.X. Activation of
1025 endoplasmic reticulum stress by hyperglycemia is essential for Muller cell-derived
1026 inflammatory cytokine production in diabetes. *Diabetes*, **2012**, *61*(2), 492-504.

1027 [111] Li, J.; Wang, J.J.; Yu, Q.; Wang, M.; Zhang, S.X. Endoplasmic reticulum stress is
1028 implicated in retinal inflammation and diabetic retinopathy. *FEBS Lett.*, **2009**, *583*(9),
1029 1521-1527.

1030 [112] Pahl, H.L.; Baeuerle, P.A. Activation of NF-kappa B by ER stress requires both
1031 Ca²⁺ and reactive oxygen intermediates as messengers. *FEBS Lett.*, **1996**, *392*(2), 129-
1032 136.

1033 [113] Fan, T.; Huang, Z.; Chen, L.; Wang, W.; Zhang, B.; Xu, Y.; Pan, S.; Mao, Z.; Hu, H.;
1034 Geng, Q. Associations between autophagy, the ubiquitin-proteasome system and
1035 endoplasmic reticulum stress in hypoxia-deoxygenation or ischemia-reperfusion. *Eur.*
1036 *J. Pharmacol.*, **2016**, *791*, 157-167.

1037 [114] Wang, Z.; Huang, Y.; Cheng, Y.; Tan, Y.; Wu, F.; Wu, J.; Shi, H.; Zhang, H.; Yu, X.;
1038 Gao, H.; Lin, L.; Cai, J.; Zhang, J.; Li, X.; Cai, L.; Xiao, J. Endoplasmic reticulum stress-
1039 induced neuronal inflammatory response and apoptosis likely plays a key role in the
1040 development of diabetic encephalopathy. *Oncotarget*, **2016**, *7*(48), 78455-78472.

1041 [115] Lin, M.I.; Fulton, D.; Babbitt, R.; Fleming, I.; Busse, R.; Pritchard, K.A., Jr.; Sessa,
1042 W.C. Phosphorylation of threonine 497 in endothelial nitric-oxide synthase
1043 coordinates the coupling of L-arginine metabolism to efficient nitric oxide production.
1044 *J. Biol. Chem.*, **2003**, *278*(45), 44719-44726.

1045 [116] Forstermann, U.; Schmidt, H.H.; Pollock, J.S.; Sheng, H.; Mitchell, J.A.; Warner,
1046 T.D.; Nakane, M.; Murad, F. Isoforms of nitric oxide synthase. Characterization and
1047 purification from different cell types. *Biochem. Pharmacol.*, **1991**, *42*(10), 1849-1857.

1048 [117] Forstermann, U.; Sessa, W.C. Nitric oxide synthases: regulation and function.
1049 *Eur. Heart J.*, **2012**, *33*(7), 829-837, 837a-837d.

1050 [118] Moens, A.L.; Champion, H.C.; Claeys, M.J.; Tavazzi, B.; Kaminski, P.M.; Wolin,
1051 M.S.; Borgonjon, D.J.; Van Nassauw, L.; Haile, A.; Zviman, M.; Bedja, D.; Wuyts, F.L.;
1052 Elsaesser, R.S.; Cos, P.; Gabrielson, K.L.; Lazzarino, G.; Paolocci, N.; Timmermans, J.P.;
1053 Vrints, C.J.; Kass, D.A. High-dose folic acid pretreatment blunts cardiac dysfunction
1054 during ischemia coupled to maintenance of high-energy phosphates and reduces
1055 postreperfusion injury. *Circulation*, **2008**, *117*(14), 1810-1819.

1056 [119] Szelenyi, Z.; Fazakas, A.; Szenasi, G.; Kiss, M.; Tegze, N.; Fekete, B.C.; Nagy, E.;
1057 Bodo, I.; Nagy, B.; Molvarec, A.; Patocs, A.; Pepo, L.; Prohaszka, Z.; Vereckei, A.
1058 Inflammation and oxidative stress caused by nitric oxide synthase uncoupling might
1059 lead to left ventricular diastolic and systolic dysfunction in patients with
1060 hypertension. *J. Geriatr. Cardiol.*, **2015**, *12*(1), 1-10.

1061 [120] Das, D.K.; Maulik, N.; Moraru, I.I. Gene expression in acute myocardial stress.
1062 Induction by hypoxia, ischemia, reperfusion, hyperthermia and oxidative stress. *J. Mol.*
1063 *Cell. Cardiol.*, **1995**, *27*(1), 181-193.

1064 [121] Dhalla, N.S.; Elmoselhi, A.B.; Hata, T.; Makino, N. Status of myocardial
1065 antioxidants in ischemia-reperfusion injury. *Cardiovasc. Res.*, **2000**, *47*(3), 446-456.

1066 [122] Haramaki, N.; Stewart, D.B.; Aggarwal, S.; Ikeda, H.; Reznick, A.Z.; Packer, L.
1067 Networking antioxidants in the isolated rat heart are selectively depleted by ischemia-
1068 reperfusion. *Free Radic. Biol. Med.*, **1998**, *25*(3), 329-339.

1069 [123] Lu, S.C. Glutathione synthesis. *Biochim. Biophys. Acta*, **2013**, *1830*(5), 3143-
1070 3153.

1071 [124] Arreguin, F.; Garcia, N.; Hernandez-Resendiz, S.; Buelna-Chontal, M.; Correa, F.;
1072 Olin-Sandoval, V.; Medina-Campos, O.N.; Pedraza-Chaverri, J.; Zazueta, C. Attenuation

1073 of oxidant damage in the postconditioned heart involves non-enzymatic response and
1074 partial catalytic protection. *Exp. Physiol.*, **2012**, *97*(10), 1119-1130.

1075 [125] Das, D.K.; Engelman, R.M.; Rousou, J.A.; Breyer, R.H.; Otani, H.; Lemeshow, S.
1076 Pathophysiology of superoxide radical as potential mediator of reperfusion injury in
1077 pig heart. *Basic Res. Cardiol.*, **1986**, *81*(2), 155-166.

1078 [126] Yoshida, T.; Maulik, N.; Engelman, R.M.; Ho, Y.S.; Magnenat, J.L.; Rousou, J.A.;
1079 Flack, J.E., 3rd; Deaton, D.; Das, D.K. Glutathione peroxidase knockout mice are
1080 susceptible to myocardial ischemia reperfusion injury. *Circulation*, **1997**, *96*(9 Suppl),
1081 II-216-220.

1082 [127] Buelna-Chontal, M.; Guevara-Chavez, J.G.; Silva-Palacios, A.; Medina-Campos,
1083 O.N.; Pedraza-Chaverri, J.; Zazueta, C. Nrf2-regulated antioxidant response is activated
1084 by protein kinase C in postconditioned rat hearts. *Free Radic. Biol. Med.*, **2014**, *74*, 145-
1085 156.

1086 [128] Chan, K.; Han, X.D.; Kan, Y.W. An important function of Nrf2 in combating
1087 oxidative stress: detoxification of acetaminophen. *Proc. Natl. Acad. Sci. U. S. A.*, **2001**,
1088 *98*(8), 4611-4616.

1089 [129] Huang, X.S.; Chen, H.P.; Yu, H.H.; Yan, Y.F.; Liao, Z.P.; Huang, Q.R. Nrf2-
1090 dependent upregulation of antioxidative enzymes: a novel pathway for hypoxic
1091 preconditioning-mediated delayed cardioprotection. *Mol. Cell. Biochem.*, **2014**, *385*(1-
1092 2), 33-41.

1093 [130] Klaassen, C.D.; Reisman, S.A. Nrf2 the rescue: effects of the
1094 antioxidative/electrophilic response on the liver. *Toxicol. Appl. Pharmacol.*, **2010**,
1095 *244*(1), 57-65.

1096 [131] Crane, F.L. Biochemical functions of coenzyme Q10. *J. Am. Coll. Nutr.*, **2001**,
1097 *20*(6), 591-598.

1098 [132] Mellors, A.; Tappel, A.L. Quinones and quinols as inhibitors of lipid
1099 peroxidation. *Lipids*, **1966**, *1*(4), 282-284.

1100 [133] Portakal, O.; Inal-Erden, M. Effects of pentoxifylline and coenzyme Q10 in
1101 hepatic ischemia/reperfusion injury. *Clin. Biochem.*, **1999**, *32*(6), 461-466.

1102 [134] Erol, B.; Bozlu, M.; Hanci, V.; Tokgoz, H.; Bektas, S.; Mungan, G. Coenzyme Q10
1103 treatment reduces lipid peroxidation, inducible and endothelial nitric oxide synthases,
1104 and germ cell-specific apoptosis in a rat model of testicular ischemia/reperfusion
1105 injury. *Fertil. Steril.*, **2010**, *93*(1), 280-282.

1106 [135] Ozalp, B.; Elbey, H.; Aydin, H.; Tekkesin, M.S.; Uzun, H. The effect of coenzyme
1107 Q10 on venous ischemia reperfusion injury. *J. Surg. Res.*, **2016**, *204*(2), 304-310.

1108 [136] Belardinelli, R.; Tiano, L.; Littarru, G.P. Oxidative stress, endothelial function
1109 and coenzyme Q10. *Biofactors*, **2008**, *32*(1-4), 129-133.

1110 [137] Tsuneki, H.; Sekizaki, N.; Suzuki, T.; Kobayashi, S.; Wada, T.; Okamoto, T.;
1111 Kimura, I.; Sasaoka, T. Coenzyme Q10 prevents high glucose-induced oxidative stress
1112 in human umbilical vein endothelial cells. *Eur. J. Pharmacol.*, **2007**, *566*(1-3), 1-10.

1113 [138] Spitz, D.R.; Oberley, L.W. An assay for superoxide dismutase activity in
1114 mammalian tissue homogenates. *Anal. Biochem.*, **1989**, *179*(1), 8-18.

1115 [139] Wang, V.; Chen, S.Y.; Chuang, T.C.; Shan, D.E.; Soong, B.W.; Kao, M.C. Val-9Ala
1116 and Ile+58Thr polymorphism of MnSOD in Parkinson's disease. *Clin. Biochem.*, **2010**,
1117 *43*(12), 979-982.

1118 [140] Matsushima, S.; Sadoshima, J. The role of sirtuins in cardiac disease. *Am. J.*
1119 *Physiol. Heart Circ. Physiol.*, **2015**, *309*(9), H1375-1389.

1120 [141] Koentges, C.; Pfeil, K.; Schnick, T.; Wiese, S.; Dahlbock, R.; Cimolai, M.C.; Meyer-
1121 Steenbuck, M.; Cenkerova, K.; Hoffmann, M.M.; Jaeger, C.; Odening, K.E.; Kammerer, B.;
1122 Hein, L.; Bode, C.; Bugger, H. SIRT3 deficiency impairs mitochondrial and contractile
1123 function in the heart. *Basic Res. Cardiol.*, **2015**, *110*(4), 36.

1124 [142] Di Lisa, F.; Menabo, R.; Canton, M.; Barile, M.; Bernardi, P. Opening of the
1125 mitochondrial permeability transition pore causes depletion of mitochondrial and
1126 cytosolic NAD⁺ and is a causative event in the death of myocytes in postischemic
1127 reperfusion of the heart. *J. Biol. Chem.*, **2001**, *276*(4), 2571-2575.

1128 [143] Porter, G.A.; Urciuoli, W.R.; Brookes, P.S.; Nadochiy, S.M. SIRT3 deficiency
1129 exacerbates ischemia-reperfusion injury: implication for aged hearts. *Am. J. Physiol.*
1130 *Heart Circ. Physiol.*, **2014**, *306*(12), H1602-1609.

1131 [144] Wang, X.X.; Wang, X.L.; Tong, M.M.; Gan, L.; Chen, H.; Wu, S.S.; Chen, J.X.; Li, R.L.;
1132 Wu, Y.; Zhang, H.Y.; Zhu, Y.; Li, Y.X.; He, J.H.; Wang, M.; Jiang, W. SIRT6 protects
1133 cardiomyocytes against ischemia/reperfusion injury by augmenting FoxO3alpha-
1134 dependent antioxidant defense mechanisms. *Basic Res. Cardiol.*, **2016**, *111*(2), 13.

1135 [145] Schoonbroodt, S.; Piette, J. Oxidative stress interference with the nuclear factor-
1136 kappa B activation pathways. *Biochem. Pharmacol.*, **2000**, *60*(8), 1075-1083.

1137 [146] Lu, L.; Chen, S.S.; Zhang, J.Q.; Ramires, F.J.; Sun, Y. Activation of nuclear factor-
1138 kappaB and its proinflammatory mediator cascade in the infarcted rat heart. *Biochem.*
1139 *Biophys. Res. Commun.*, **2004**, *321*(4), 879-885.

1140 [147] Nichols, T.C. NF-kappaB and reperfusion injury. *Drug News Perspect.*, **2004**,
1141 *17*(2), 99-104.

1142 [148] Bonaventura, A.; Montecucco, F.; Dallegri, F. Cellular recruitment in myocardial
1143 ischaemia/reperfusion injury. *Eur. J. Clin. Invest.*, **2016**, *46*(6), 590-601.

1144 [149] Gawaz, M. Role of platelets in coronary thrombosis and reperfusion of ischemic
1145 myocardium. *Cardiovasc. Res.*, **2004**, *61*(3), 498-511.

1146 [150] Mehta, J.L.; Nicolini, F.A.; Donnelly, W.H.; Nichols, W.W. Platelet-leukocyte-
1147 endothelial interactions in coronary artery disease. *Am. J. Cardiol.*, **1992**, *69*(7), 8B-
1148 13B.

1149 [151] Del Conde, I.; Cruz, M.A.; Zhang, H.; Lopez, J.A.; Afshar-Kharghan, V. Platelet
1150 activation leads to activation and propagation of the complement system. *J. Exp. Med.*,
1151 **2005**, *201*(6), 871-879.

1152 [152] Horckmans, M.; Ring, L.; Duchene, J.; Santovito, D.; Schloss, M.J.; Drechsler, M.;
1153 Weber, C.; Soehnlein, O.; Steffens, S. Neutrophils orchestrate post-myocardial
1154 infarction healing by polarizing macrophages towards a reparative phenotype. *Eur.*
1155 *Heart J.*, **2016**, *38*(3), 187-197.

1156 [153] Ge, L.; Zhou, X.; Ji, W.J.; Lu, R.Y.; Zhang, Y.; Zhang, Y.D.; Ma, Y.Q.; Zhao, J.H.; Li,
1157 Y.M. Neutrophil extracellular traps in ischemia-reperfusion injury-induced myocardial
1158 no-reflow: therapeutic potential of DNase-based reperfusion strategy. *Am. J. Physiol.*
1159 *Heart Circ. Physiol.*, **2015**, *308*(5), H500-509.

1160 [154] Reffelmann, T.; Kloner, R.A. The no-reflow phenomenon: A basic mechanism of
1161 myocardial ischemia and reperfusion. *Basic Res. Cardiol.*, **2006**, *101*(5), 359-372.

1162 [155] Heusch, G. The Coronary Circulation as a Target of Cardioprotection. *Circ. Res.*,
1163 **2016**, *118*(10), 1643-1658.

1164 [156] Hernandez-Resendiz, S.; Palma-Flores, C.; De Los Santos, S.; Roman-Anguiano,
1165 N.G.; Flores, M.; de la Pena, A.; Flores, P.L.; Fernandez, G.J.; Coral-Vazquez, R.M.;
1166 Zazueta, C. Reduction of no-reflow and reperfusion injury with the synthetic 17beta-
1167 aminoestrogen compound Prolame is associated with PI3K/Akt/eNOS signaling
1168 cascade. *Basic Res. Cardiol.*, **2015**, *110*(2), 1.

1169 [157] Kloner, R.A.; Ganote, C.E.; Jennings, R.B. The "no-reflow" phenomenon after
1170 temporary coronary occlusion in the dog. *J. Clin. Invest.*, **1974**, *54*(6), 1496-1508.

1171 [158] Roberts, C.S.; Maclean, D.; Maroko, P.; Kloner, R.A. Early and late remodeling of
1172 the left ventricle after acute myocardial infarction. *Am. J. Cardiol.*, **1984**, *54*(3), 407-
1173 410.

1174 [159] Heusch, G.; Libby, P.; Gersh, B.; Yellon, D.; Bohm, M.; Lopaschuk, G.; Opie, L.
1175 Cardiovascular remodelling in coronary artery disease and heart failure. *Lancet*, **2014**,
1176 *383*(9932), 1933-1943.

1177 [160] Apostolova, N.; Victor, V.M. Molecular strategies for targeting antioxidants to
1178 mitochondria: therapeutic implications. *Antioxid. Redox Signal.*, **2015**, *22*(8), 686-729.

1179 [161] Eleawa, S.M.; Alkhateeb, M.; Ghosh, S.; Al-Hashem, F.; Shatoor, A.S.; Alhejaily, A.;
1180 Khalil, M.A. Coenzyme Q10 protects against acute consequences of experimental
1181 myocardial infarction in rats. *Int. J. Physiol. Pathophysiol. Pharmacol.*, **2015**, *7*(1), 1-13.

1182 [162] Ivanov, A.V.; Gorodetskaya, E.A.; Kalenikova, E.I.; Medvedev, O.S. Single
1183 intravenous injection of coenzyme Q10 protects the myocardium after irreversible
1184 ischemia. *Bull. Exp. Biol. Med.*, **2013**, *155*(6), 771-774.

1185 [163] Kuo, Y.R.; Wang, F.S.; Jeng, S.F.; Lutz, B.S.; Huang, H.C.; Yang, K.D.
1186 Nitrosoglutathione improves blood perfusion and flap survival by suppressing iNOS
1187 but protecting eNOS expression in the flap vessels after ischemia/reperfusion injury.
1188 *Surgery*, **2004**, *135*(4), 437-446.

1189 [164] Ernster, L.; Dallner, G. Biochemical, physiological and medical aspects of
1190 ubiquinone function. *Biochim. Biophys. Acta*, **1995**, *1271*(1), 195-204.

1191 [165] Huang, C.H.; Kuo, C.L.; Huang, C.S.; Tseng, W.M.; Lian Ie, B.; Chang, C.C.; Liu, C.S.
1192 High plasma coenzyme Q10 concentration is correlated with good left ventricular
1193 performance after primary angioplasty in patients with acute myocardial infarction.
1194 *Medicine (Baltimore)*, **2016**, *95*(31), e4501.

1195 [166] Kelso, G.F.; Porteous, C.M.; Coulter, C.V.; Hughes, G.; Porteous, W.K.;
1196 Ledgerwood, E.C.; Smith, R.A.; Murphy, M.P. Selective targeting of a redox-active
1197 ubiquinone to mitochondria within cells: antioxidant and antiapoptotic properties. *J.*
1198 *Biol. Chem.*, **2001**, *276*(7), 4588-4596.

1199 [167] Dare, A.J.; Logan, A.; Prime, T.A.; Rogatti, S.; Goddard, M.; Bolton, E.M.; Bradley,
1200 J.A.; Pettigrew, G.J.; Murphy, M.P.; Saeb-Parsy, K. The mitochondria-targeted anti-
1201 oxidant MitoQ decreases ischemia-reperfusion injury in a murine syngeneic heart
1202 transplant model. *J. Heart Lung Transplant.*, **2015**, *34*(11), 1471-1480.

1203 [168] Davidson, S.M.; Yellon, D.M.; Murphy, M.P.; Duchon, M.R. Slow calcium waves
1204 and redox changes precede mitochondrial permeability transition pore opening in the
1205 intact heart during hypoxia and reoxygenation. *Cardiovasc. Res.*, **2012**, *93*(3), 445-453.

1206 [169] Zhao, K.; Zhao, G.M.; Wu, D.; Soong, Y.; Birk, A.V.; Schiller, P.W.; Szeto, H.H. Cell-
1207 permeable peptide antioxidants targeted to inner mitochondrial membrane inhibit
1208 mitochondrial swelling, oxidative cell death, and reperfusion injury. *J. Biol. Chem.*,
1209 **2004**, *279*(33), 34682-34690.

1210 [170] Liu, S.; Soong, Y.; Seshan, S.V.; Szeto, H.H. Novel cardiolipin therapeutic protects
1211 endothelial mitochondria during renal ischemia and mitigates microvascular
1212 rarefaction, inflammation, and fibrosis. *Am. J. Physiol. Renal Physiol.*, **2014**, *306*(9),
1213 F970-980.

1214 [171] Gibson, C.M.; Giugliano, R.P.; Kloner, R.A.; Bode, C.; Tendera, M.; Janosi, A.;
1215 Merkely, B.; Godlewski, J.; Halaby, R.; Korjian, S.; Daaboul, Y.; Chakrabarti, A.K.;
1216 Spielman, K.; Neal, B.J.; Weaver, W.D. EMBRACE STEMI study: a Phase 2a trial to
1217 evaluate the safety, tolerability, and efficacy of intravenous MTP-131 on reperfusion
1218 injury in patients undergoing primary percutaneous coronary intervention. *Eur. Heart*
1219 *J.*, **2016**, *37*(16), 1296-1303.

1220 [172] Du, L.; Miao, X.; Gao, Y.; Jia, H.; Liu, K.; Liu, Y. The protective effects of Trolox-
1221 loaded chitosan nanoparticles against hypoxia-mediated cell apoptosis. *Nanomedicine*,
1222 **2014**, *10*(7), 1411-1420.

1223 [173] Gupta, S.C.; Prasad, S.; Kim, J.H.; Patchva, S.; Webb, L.J.; Priyadarsini, I.K.;
1224 Aggarwal, B.B. Multitargeting by curcumin as revealed by molecular interaction
1225 studies. *Nat. Prod. Rep.*, **2011**, *28*(12), 1937-1955.

1226 [174] Ghosh, S.S.; Massey, H.D.; Krieg, R.; Fazelbhoj, Z.A.; Ghosh, S.; Sica, D.A.; Fakhry,
1227 I.; Gehr, T.W. Curcumin ameliorates renal failure in 5/6 nephrectomized rats: role of
1228 inflammation. *Am. J. Physiol. Renal Physiol.*, **2009**, *296*(5), F1146-1157.

1229 [175] Hernandez-Resendiz, S.; Correa, F.; Garcia-Nino, W.R.; Buelna-Chontal, M.;
1230 Roldan, F.J.; Ramirez-Camacho, I.; Delgado-Toral, C.; Carbo, R.; Pedraza-Chaverri, J.;
1231 Tapia, E.; Zazueta, C. Cardioprotection by curcumin post-treatment in rats with
1232 established chronic kidney disease. *Cardiovasc. Drugs Ther.*, **2015**, *29*(2), 111-120.

1233 [176] Hausenloy, D.J.; Barrabes, J.A.; Botker, H.E.; Davidson, S.M.; Di Lisa, F.; Downey,
1234 J.; Engstrom, T.; Ferdinandy, P.; Carbrera-Fuentes, H.A.; Heusch, G.; Ibanez, B.;
1235 Iliodromitis, E.K.; Inserte, J.; Jennings, R.; Kalia, N.; Kharbanda, R.; Lecour, S.; Marber,
1236 M.; Miura, T.; Ovize, M.; Perez-Pinzon, M.A.; Piper, H.M.; Przyklenk, K.; Schmidt, M.R.;
1237 Redington, A.; Ruiz-Meana, M.; Vilahur, G.; Vinten-Johansen, J.; Yellon, D.M.; Garcia-
1238 Dorado, D. Ischaemic conditioning and targeting reperfusion injury: a 30 year voyage
1239 of discovery. *Basic Res. Cardiol.*, **2016**, *111*(6), 70.

1240 [177] Cabrera-Fuentes, H.A.; Aragones, J.; Bernhagen, J.; Boening, A.; Boisvert, W.A.;
1241 Botker, H.E.; Bulluck, H.; Cook, S.; Di Lisa, F.; Engel, F.B.; Engelmann, B.; Ferrazzi, F.;
1242 Ferdinandy, P.; Fong, A.; Fleming, I.; Gnaiger, E.; Hernandez-Resendiz, S.; Kalkhoran,
1243 S.B.; Kim, M.H.; Lecour, S.; Liehn, E.A.; Marber, M.S.; Mayr, M.; Miura, T.; Ong, S.B.;
1244 Peter, K.; Sedding, D.; Singh, M.K.; Suleiman, M.S.; Schnittler, H.J.; Schulz, R.; Shim, W.;
1245 Tello, D.; Vogel, C.W.; Walker, M.; Li, Q.O.; Yellon, D.M.; Hausenloy, D.J.; Preissner, K.T.
1246 From basic mechanisms to clinical applications in heart protection, new players in
1247 cardiovascular diseases and cardiac theranostics: meeting report from the third
1248 international symposium on "New frontiers in cardiovascular research". *Basic Res.*
1249 *Cardiol.*, **2016**, *111*(6), 69.

1250 [178] Faxon, D.P.; Gibbons, R.J.; Chronos, N.A.; Gurbel, P.A.; Sheehan, F.; Investigators,
1251 H.-M. The effect of blockade of the CD11/CD18 integrin receptor on infarct size in
1252 patients with acute myocardial infarction treated with direct angioplasty: the results
1253 of the HALT-MI study. *J. Am. Coll. Cardiol.*, **2002**, *40*(7), 1199-1204.

1254

1255 **Figure 1.** Formation of different ROS and reactive nitrogen species from dioxygen.
1256 ischaemia/reperfusion injury. Dioxygen (O_2) is shown to undergo reduction to form superoxide (O_2^-).
1257 Superoxide is shown to dismutate to form hydrogen peroxide (H_2O_2), and hydrogen peroxide is
1258 shown to interact with Fe^{2+} and to form hydroxyl radical ($\bullet OH$) via the Fenton reaction. Superoxide
1259 dismutase (SOD), nitrogen monoxide ($NO\bullet$), peroxynitrite ($ONOO\bullet$).

1260

1261 **Figure 2.** Schematic representation of (A) the sources of reactive oxygen species during acute
1262 myocardial ischaemia/reperfusion injury. (B) During acute myocardial ischaemia, AA accumulates,
1263 leading increased generation of 20-HETE through CYP 4F. 20-HETE acts directly via the
1264 stimulation of NADPH oxidase-derived ROS production and induces $NF\kappa B$ activation. (C)
1265 Myocardial reperfusion generates accumulation of cholesterol into mitochondria. This induces the
1266 formation of oxysterols, which can induce IL-1 β and IL-8 secretion.

1267 Cytochrome P-450 (CYP), xanthine oxidase, NADPH oxidase, monoamine oxidases (MAO),
1268 sarco/endoplasmic reticulumun (SR/ES), nitric oxide synthase (NOS), arachidonic acid (AA), 20-
1269 HETE (20-hydroxy-5,8,11,14-eicosatetraenoic acid), EETs (epoxyeicosatrienoic acids), $NF\kappa B$
1270 (nuclear factor kappa-light-chain-enhancer of activated B cells), N-methylsulfonyl-12, 12-dibromo-
1271 11-enamide (DDMS), HET0016 (N-hydroxy-N'-(4-butyl-2-methylphenyl)-formamidine), cholesterol
1272 (chol), IL-1 β (interleukin-1 beta), IL-8 (interleukin-8), IRI (Ischaemia/Reperfusion Injury), ROS
1273 (reactive oxygen species).

1274

1275 **Figure 3.** No-reflow following acute myocardial ischaemia/reperfusion injury. (A) Example of infact
1276 assessment using triphenyl tetrazolium chloride (TTC) staining (B) Example of no-reflow by
1277 transillumination of Microfil-perfused coronary; and (C) Schematic diagram demonstrating the
1278 multiple etiologies of no-reflow in the reperfused coroanry artery.

1279 ROS (reactive oxygen species).

1280

1281


1282

1283

1284

1285 Figure 1

1286



1287

1288

1289

1290

1291

1292

1293

1294

1295

1296

1297

1298

1299 Figure 2

1300

1301


1302

1303

1304

1305

1306



1307

1308

1309

1310

1311

1312


1313

1314

1315

1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327

Figure 3



1328